



One-pot catalytic C–C double bond cleavage of α,β -enones aided by alkyl group-immobilized silica spheres

Dong Hun Lee, Eun-Ae Jo, Jung-Woo Park, Chul-Ho Jun *

Department of Chemistry and Center for Bioactive Molecular Hybrid, Yonsei University, Seoul 120-749, Republic of Korea

ARTICLE INFO

Article history:

Received 17 September 2009

Revised 22 October 2009

Accepted 22 October 2009

Available online 27 October 2009

This Letter is dedicated to the memory of the late Professor Chi Sun Han (1926–2009) of Yonsei University

ABSTRACT


Catalytic C–C double bond cleavage of α,β -enones with a 1-alkene and H_2O was carried out in the presence of a $(Ph_3P)_3RhCl$ catalyst, 2-amino-3-picoline, cyclohexylamine, benzoic acid, and alkyl group-immobilized silica spheres. Upon completion of the reaction, the corresponding ketones were obtained without needing a further hydrolysis step. In this reaction, alkyl group-immobilized silica spheres act as a water reservoir for hydrolysis of an intermediate ketimine and as a phase divider between the organic solution and H_2O .

© 2009 Elsevier Ltd. All rights reserved.

C–H bond activation by transition-metal catalysts is of current interest in organic synthesis in terms of atom economy.¹ We have successfully developed a Rh(I)-catalyzed C–H bond activation using 2-aminopyridine derivatives or benzylamine as a chelating auxiliary to induce facile cyclometallation.² In the course of our studies on chelation-assisted C–H bond activation, the Rh(I)-catalyzed hydroiminoacylation of an alkyne with an allylamine or an aldehyde, followed by retro-Mannich type fragmentation of the resulting α,β -unsaturated ketimine was found.^{3,4} We have also developed a C–C double bond cleavage of α,β -enone using amine and acid to produce ketones after hydrolysis step.⁵ For example, the reaction of 4-phenyl-but-3-en-2-one (**1a**), 3,3-dimethyl-1-butene (**3a**), and cyclohexylamine (**2**, 200 mol % based upon **1a**) was carried out at 130 °C for 4 h in the presence of the catalyst system of $(Ph_3P)_3RhCl$ (**4**), 2-amino-3-picoline (**5**), and benzoic acid (**6**) to give acetone and ketimine **7**, which was hydrolyzed separately by acidic H_2O giving the corresponding ketone **9a** in a 49% isolated yield (Scheme 1a). This reaction requires greater than 100 mol % of cyclohexylamine based on the starting α,β -enone, and a further hydrolysis step for the resulting ketimine **7** to liberate ketone (**9a**). To decrease the amount of **2** used and to eliminate the separate hydrolysis step, H_2O can be added as a reactant to hydrolyze the in situ generated ketimine **7** resulting in the formation of **9a** directly (Scheme 1b). A proposed mechanism for this one-pot reaction is depicted in Scheme 2. Cyclohexylamine **2** undergoes conjugate addition to the α,β -enone **1a**, and the subsequent retro-Mannich type fragmentation of the resulting β -aminoketone **A** generates a mixture of aldimine **B** and ketone **8**. Since aldimine is not much susceptible for hydrolysis compared with ketimine,


the aldimine survives even in the presence of H_2O during the reaction. Therefore, the aldimine **B** undergoes hydroiminoacylation with 1-alkene (**3a**) mediated by the cocatalyst system of $(Ph_3P)_3RhCl$ (**4**) and 2-amino-3-picoline (**5**) to afford ketimine **7**. This type of transimination/hydroiminoacylation protocol has been utilized in the efficient conversion of aldimine to ketimine.⁶ Now the ketimine **7** is readily hydrolyzed by additional H_2O giving ketone **9a** with regeneration of cyclohexylamine (**2**), which can be involved in the catalytic cycle. Based on this it can be said that without additional H_2O no catalytic reaction takes place since cyclohexylamine cannot be regenerated.

However, addition of H_2O afforded a very low yield of **9a** (33% isolated yield, Scheme 1b) from the reaction of **1a** with **3a** and



Scheme 1. (a) Retro-Mannich fragmentation and subsequent hydroacylation of **1a** with **3a**, and 2 equiv of **2** in the presence of **4–6**. (b) Reactions of **1a** with **3a**, a catalytic amount of **2**, and H_2O in the presence of **4–6**.

* Corresponding author. Tel.: +82 2 2123 2644; fax: +82 2 3147 2644.
E-mail address: junch@yonsei.ac.kr (C.-H. Jun).



Scheme 2. Proposed mechanism for retro-Mannich fragmentation and subsequent hydroacylation of **1a** with **3a** and H₂O in the presence of **2**, **4–6**.

H₂O (100 mol %) in the presence of **2** (50 mol %), **4** (3 mol %), **5** (20 mol %), and **6** (15 mol %).⁷ To overcome this problem, it is necessary to separate the H₂O and organic solution phase (Fig. 1). Herein, we wish to report the efficient one-pot amine-assisted C–C double bond cleavage of α,β -enones aided by alkyl group-immobilized silica spheres. Since the alkyl group-immobilized silica spheres act as both a phase separator and a reservoir for water, the activity of the transition-metal catalyst is maintained even in the presence of water, and a separate hydrolysis step is avoided.

Recently, some organometallic catalytic reactions in aqueous phase have been studied based on the heterogeneous system.⁸ We have developed a new method for the surface-fabrication of the high density long chain alkyl group-grafted silica, which can be used as a divider between the organic reaction media and H₂O absorbed on the silica.⁹ Using this protocol, octadecyl group-functionalized silica spheres **11a** (loading rate of octadecyl group = 0.40 mmol g⁻¹)¹⁰ were prepared by the reaction of silanol group on the surface of silica ball (10 μ m in diameter, 11.67 nm pore size, purchased from Kromasil) with octadecyltrimethylsilane (**10a**) in the presence of a Sc(OTf)₃ catalyst (OTf = trifluoromethylsulfonate) at room temperature (Scheme 3a). To prove the effect of **11a**, the catalytic reaction of **1a**, **3a**, and H₂O (100 mol %) with the catalyst system of (Ph₃P)₃RhCl (**4**), 2-amino-3-picolone (**5**), cyclohexylamine (**2**), and benzoic acid (**6**) was carried out in the presence of **11a** for 4 h to give the corresponding ketone **9a** in a 92% isolated yield without any further hydrolysis step.¹¹

For comparison, other alkyl group-immobilized silica spheres with different alkyl group loading rates were applied to this reaction under the identical reaction conditions: reaction of **1a** with **3a** at 130 °C for 4 h affording **9a** (Table 1). Among these modified silica additives, **11a** (loading rate of 0.45 mmol g⁻¹) exhibited better reactivity than **11b** (0.71 mmol g⁻¹), and **11c** (1.06 mmol g⁻¹), prepared from octadecylmethyl-dimethylsilane (**10b**) and octadecyldimethylmethyl-silane (**10c**), respectively (entries 2 and 3). Interestingly, when the reaction of **1a** with **3a** was conducted






Figure 1. Hydrolysis of ketimine on the alkyl group-immobilized silica spheres.



Scheme 3. (a) Preparation of **11a** through the immobilization reaction of **10a** onto silica spheres under Sc(OTf)₃ catalysis. (b) Representative reaction of **1a** with **3a** and H₂O in the presence of **2**, **4–6**, and **11a**.

Table 1

The effect of **2** on the catalytic C–C double bond cleavage of α,β -enone (**1**) and subsequent hydroacylation with **3**




Entry	Octadecyl group-immobilized silica (11)	Isolated yield of 9a (%)
1	 11a , 0.45 mmol g ⁻¹	92
2	 11b , 0.71 mmol g ⁻¹	49
3	 11c , 1.06 mmol g ⁻¹	28
4	 11a-SiMe₃ ^a	20
5	silica sphere (TMS-coated, 1.3 mmol g ⁻¹) ^a	30
6	Silica sphere (bare)	49

^a **11a-SiMe₃** and TMS-coated silica spheres were prepared by the reaction of **11a** or untreated silica spheres with 1,1,1,3,3,3-hexamethyldisilazane, respectively.

using **11a-SiMe₃**, in which the residual silanol groups on **11a** were fully capped with trimethylsilyl groups by treatment of **11a** with 1,1,1,3,3,3-hexamethyldisilazane,¹² compound **9a** was obtained in a low isolated yield (20%) (entry 4), which suggests that the residual silanol groups on **11a** are essential for the in situ hydrolysis

Table 2
Catalytic C–C double bond cleavage of α,β -enone (**1**) and subsequent hydroacylation with **3**



Entry	α,β -Unsaturated ketone (1) (R_1, R_2)	Olefin (3) (R_3)	Rxn time	Yield of 9 ^a (%)	Yield of 9 without using 11a ^a (%)
1	Ph, Me (1a)	$n\text{-C}_4\text{H}_9$ (3b)	4 h	89(98) (9b)	41(51)
2		$n\text{-C}_6\text{H}_{13}$ (3c)		85(98) (9c)	47(56)
3		$c\text{-C}_6\text{H}_{12}$ (3d)		77(80) (9d)	25(33)
4		C_6H_5 (3e)		83(95) (9e)	54(58)
5		SiMe_3 (3f)		89(99) (9f)	30(45)
6		SiEt_3 (3g)		88(97) (9g)	17(33)
7	Me, Me (1b)	3a	45 min	(82) (9h)	(44)
8	Ph, Ph (1c)	3a	14 h	85(89) (9a)	51(61)

^a Isolated yield. The yields in parentheses are determined by GC.

step. Although the reason for this result is not clear, but we speculate that H_2O bound on the surface of silanol groups readily interacts with ketimine **7** to be hydrolyzed in organic solution as shown in Figure 1. However, if silanol group on the silica surface is completely covered with hydrophobic SiMe_3 or octadecylmethylsilyl groups, there is no room for H_2O on the silica surface, and H_2O may smear into the interior of the silica network. This fact was also confirmed by the use of exclusive TMS-coated silica spheres (loading rate: 1.3 mmol g^{-1}), prepared by treatment of silica spheres with 1,1,1,3,3,3-hexamethyldisilazane. With this TMS-coated silica spheres, only a 30% yield of **9a** was isolated (entry 5). When silica spheres untreated with alkylating reagents were used, only a 49% isolated yield of **9a** was obtained (entry 6). It is known that transition-metal phosphine complexes can react with silanol groups on the silica surface to generate inactive metal catalysts due to the formation of P(V) species of the ligand phosphine.¹³

The reactions of various α,β -enones (**1**) with olefins (**3**) are summarized in Table 2. The reaction of 4-phenyl-but-3-en-2-one (**1a**), 1-hexene (**3b**), and H_2O was carried out at 130 °C for 4 h in the presence of **4** (3 mol %), **5** (30 mol %), **2** (50 mol %), and **6** (15 mol %) using the octadecyl group-immobilized silica spheres (**11a**) to afford 1-phenylheptan-1-one (**9b**) in an 89% isolated yield. Under identical reaction conditions without **11a**, the reaction of **1a** and **3b** produced a 41% isolated yield of **9b** (Table 2, entry 1). Similar results were observed in the reactions of other 1-alkenes (**3c–g**), as fairly good yields of the corresponding isolated ketones (**9c–g**) were obtained with **11a** (Table 2, entries 2–6); only low yields of the products were obtained in the absence of C_{18} -immobilized silica **11a**. When other substituted α,β -enones (**1b** and **1c**) were applied to the reaction with **3a**, different reactivities were observed depending on the substrate (Table 2, entries 7 and 8). The reaction of 3-penten-2-one (**1b**) with **3a** proceeded to give an 82% GC yield of **9h** after 45 min using the identical catalytic system; however, the reaction of *trans*-chalcone (**1c**) with **3a** was sluggish, requiring 14 h to get a similar GC yield (89%) of corresponding ketone **9a**. These results about different reactivity based on the substituent of substrate imply that the rate-determining step is probably the cleavage of double bond which proceeds by the addition of cyclohexylamine followed by retro-Mannich type fragmentation.

In conclusion, we have demonstrated the one-pot catalytic C–C double bond cleavage of α,β -enone using alkyl group-immobilized silica spheres and a catalyst system consisting of Rh(I) complex, 2-amino-3-picoline, cyclohexylamine, and benzoic acid. In this

reaction alkyl group-immobilized silica spheres act as a water reservoir for hydrolysis of the ketimine and a phase divider for the organic solution and H_2O on the silica. This one-pot catalytic reaction system is simple and efficient, and practical. Further applications of this one-pot catalytic system with functional group-immobilized silica are currently being investigated.

Acknowledgments

This work was supported by a Korea Research Foundation Grant funded by the Korean Government (MOEHRD) (KRF-2008-313-C00483), the WCU (World Class University) program through the Korea Science and Engineering Foundation funded by the Ministry of Education, Science and Technology (R32-2008-000-10217-0) and the CBMH. D.H.L. and J.W.P. acknowledge the fellowships from the BK21 program of the Ministry of Education and Human Resources Development.

References and notes

- Alberico, D.; Scott, M. E.; Lautens, M. *Chem. Rev.* **2007**, *107*, 174; Seregin, I. V.; Gevorgyan, V. *Chem. Soc. Rev.* **2007**, *36*, 1173; Godula, K.; Sames, D. *Science* **2006**, *312*, 67; Kakiuchi, F.; Chatani, N. *Adv. Synth. Catal.* **2003**, *345*, 1077; Kakiuchi, F.; Murai, S. *Acc. Chem. Res.* **2002**, *35*, 826; Ritleng, V.; Sirlin, C.; Pfeffer, M. *Chem. Rev.* **2002**, *102*, 1731; Labinger, J. A.; Bercaw, J. E. *Nature* **2002**, *417*, 507.
- Jo, E.-A.; Jun, C.-H. *Tetrahedron Lett.* **2009**, *50*, 3338; Park, Y. J.; Park, J.-W.; Jun, C.-H. *Acc. Chem. Res.* **2008**, *41*, 222; Jun, C.-H.; Jo, E.-A.; Park, J.-W. *Eur. J. Org. Chem.* **2007**, 1869; Park, Y. J.; Jun, C.-H. *Bull. Korean Chem. Soc.* **2005**, *26*, 871; Jun, C.-H.; Lee, H.; Hong, J.-B.; Kwon, B.-I. *Angew. Chem., Int. Ed.* **2002**, *41*, 2146; Jun, C.-H.; Chung, K.-Y.; Hong, J.-B. *Org. Lett.* **2001**, *3*, 785; Jun, C.-H.; Lee, D.-Y.; Lee, H.; Hong, J.-B. *Angew. Chem., Int. Ed.* **2000**, *39*, 3070; Jun, C.-H.; Lee, H.; Hong, J.-B. *J. Org. Chem.* **1997**, *62*, 1200.
- Lee, D.-Y.; Hong, B.-S.; Cho, E.-G.; Lee, H.; Jun, C.-H. *J. Am. Chem. Soc.* **2003**, *125*, 6372; Lee, D.-Y.; Kim, I.-J.; Jun, C.-H. *Angew. Chem., Int. Ed.* **2002**, *41*, 3031; Jun, C.-H.; Lee, H.; Moon, C. W.; Hong, H.-S. *J. Am. Chem. Soc.* **2001**, *123*, 8600; Jun, C.-H.; Lee, H.; Park, J.-B.; Lee, D.-Y. *Org. Lett.* **1999**, *1*, 2161.
- Several examples for retro-Mannich type fragmentation of β -aminoketone: Buchanan, A. C. W. *Chem. Commun.* **1966**, 773; Venit, J. J.; Magnus, P. *Tetrahedron Lett.* **1980**, *21*, 4815; Winkler, J. D.; Muller, C. L.; Scott, R. D. *J. Am. Chem. Soc.* **1988**, *110*, 4831; Winkler, J. D.; Haddad, N.; Ogilvie, R. J. *Tetrahedron Lett.* **1989**, *30*, 5703; Risch, N.; Langhals, M.; Hohberg, T. *Tetrahedron Lett.* **1991**, *32*, 4465; Kitayama, T.; Yokoi, T.; Kawai, Y.; Hill, R. K.; Morita, M.; Okamoto, T.; Yamamoto, Y.; Fokin, V. V.; Sharpless, K. B.; Sawada, S. *Tetrahedron* **2003**, *59*, 4857; Cramer, N.; Juretschke, J.; Laschat, S.; Baro, A.; Frey, W. *Eur. J. Org. Chem.* **2004**, 1397; White, J. D.; Ihle, D. C. *Org. Lett.* **2006**, *8*, 1081; Ragains, J. R.; Winkler, J. D. *Org. Lett.* **2006**, *8*, 4437.
- Lim, S.-G.; Jun, C.-H. *Bull. Korean Chem. Soc.* **2004**, *25*, 1623.
- Jun, C.-H.; Hong, J.-B. *Org. Lett.* **1999**, *1*, 887.

7. Herrmann, W. A. In *Aqueous-Phase Organometallic Catalysis—Concepts and Applications*; Cornils, B., Herrmann, W. A., Eds.; Wiley-VCH: Weinheim, 1998; p 39.
8. Arhancet, J. P.; Davis, M. E.; Merola, J. S.; Hanson, B. E. *Nature* **1989**, 339, 454; Wan, K. T.; Davis, M. E. *J. Catal.* **1994**, 148, 1.
9. Yeon, Y.-R.; Park, Y. J.; Lee, J.-S.; Park, J.-W.; Kang, S.-G.; Jun, C.-H. *Angew. Chem., Int. Ed.* **2008**, 47, 109.
10. Loading rate was determined based on the C value of an elemental analysis. Calculation of loading rate of **11a** (C (%) 10.5841) = $(10.5841 \times 10^{-2} \text{ g of C/1 g of sample } \mathbf{11a}) \times (10^3 \text{ mmol of C/12 g of C}) \times (1 \text{ mmol of C in octadecylmethylsilyl group/22 mmol of C}) = 0.40 \text{ mmol octadecyl group/1 g of } \mathbf{11a}$.
11. *General procedure* (Scheme 3b): A screw-capped pressure vial (1 ml) equipped with a magnetic stirring bar was charged with 4-phenylbut-3-en-2-one (**1a**, 29.2 mg, 0.2 mmol), 3,3-dimethyl-1-butene (**3a**, 84.0 mg, 1.0 mmol), water (3.6 mg, 0.2 mmol), $(\text{Ph}_3\text{P})_3\text{RhCl}$ (**4**, 5.5 mg, 0.006 mmol), 2-amino-3-picoline (**5**, 6.5 mg, 0.06 mmol), cyclohexylamine (**2**, 9.9 mg, 0.1 mmol), and **11a** (100 mg) dissolved in 400 mg of toluene, then the reaction mixture was stirred at 130 °C for 4 h. After the reaction, a crude mixture was purified by column chromatography (SiO_2 , *n*-hexane/ethyl acetate = 20:1) to give 43.7 mg (92% yield based on **1a**) of 4,4-dimethyl-1-phenylpentan-1-one (**9a**).
12. Zapilko, C.; Widenmeyer, M.; Nagl, I.; Estler, F.; Anwender, R.; Raudaschl-Sieber, G.; Groeger, O.; Engelhardt, G. *J. Am. Chem. Soc.* **2006**, 128, 16266.
13. Crudden, C. M.; Allen, D. P.; Motorina, D. P.; Fairgrieve, M. In *Nanostructured Catalysts*; Scott, S. L., Crudden, C. M., Jones, C. W., Eds.; Kluwer Academic/Plenum: New York, 2003; p 138; Beml, L.; Clark, H. C.; Davies, H. A.; Fyfe, C. A.; Wasylshen, R. E. *J. Am. Chem. Soc.* **1982**, 104, 438; Blümel, J. *Inorg. Chem.* **1994**, 33, 5050; Behringer, K. D.; Blümel, J. *Inorg. Chem.* **1996**, 35, 1814.