

OLIGOMERIC FLAVANOID. PART 15^a. BASE-CATALYZED PYRAN REARRANGEMENTS OF
PROCYANIDIN B-2, AND EVIDENCE FOR THE OXIDATIVE TRANSFORMATION OF B- TO
A-TYPE PROCYANIDINS

JOHANN F.W. BURGER^b, HERBERT KOLODZIEJ^b, RICHARD W. HEMINGWAY^c, JAN P.
STEYNBERG^b, DESMOND A. YOUNG^b, AND DANEEL FERREIRA^{b*}.

^b Department of Chemistry, University of the Orange Free State, P.O. Box 339,
Bloemfontein, 9300 South Africa

^c Southern Forest Experiment Station, USDA-Forest Service, 2500
Shreveport Highway, Pineville, Louisiana, 71360 U.S.A.


(Received in UK 30 April 1990)

Abstract — Procyanidin B-2 **3** is subject to facile C-ring isomerizations in 0.1M NaHCO₃ solution to form a novel series of 3,4,9,10-tetrahydro-2H,8H-pyrano[2,3-b]chromenes **7**, **9**, and **10**. The low percentage conversion of B- to A-type procyanidin **2** is rationalized in terms of an initial oxidative removal of hydride ion at C-2 (C-ring).

Proanthocyanidin A-2 **1** [(–)-epicatechin(4β → 8, 2β → O → 7)-(–)epicatechin] was first isolated by Mayer *et al.* from the seed of *Aesculus hippocastanum*¹. The structure was deduced by Haslam and his co-workers *via* spectroscopic and chemical evidence² and has, more recently, been unequivocally established by X-ray crystallography³. A variety of proanthocyanidins possessing the doubly-linked unit of type **1** has since been reported^{2,4-10}. Owing to the close structural relationship between proanthocyanidin A-2 **1** and procyanidin B-2 **3**, Porter¹¹ has proposed a biosynthetic pathway for the conversion of B- to A-type procyanidins which involves an enzyme mediated hydroxylation at C-2 (C-ring) of **3**. Despite the considerable progress in the semi-synthetic approach towards condensed tannins over the last decade similar efforts for the A-type procyanidins are limited to a fortuitous reference⁹ to the oxidative conversion of procyanidin B-1 to proanthocyanidin A-1, but with no experimental details. Our recent investigations of the base-catalyzed pyran rearrangements of proflisetinidins^{12,13} and procyanidin B-3¹⁴ which emphasizes the involvement of 7-OH(D) and C-2(C) (*cf.* structure **3**), in conjunction with the industrial interest^{15,16} in the base-simulated reactions of procyanidins, prompted application of a similar protocol to procyanidin B-2 **3**.

Owing to the lability of the interflavanyl bond in procyanidins at alkaline pH^{14,17}, the conditions previously¹²⁻¹⁴ employed were adjusted for procyanidin B-2. Thus, treatment of (4β,8)-bis-(–)-

^aPart 14. J.C.S. Malan, P.J. Steynberg, J.P. Steynberg, D.A. Young, B.C.B. Bezuidenhout, and D. Ferreira, Tetrahedron, 1990, paper AM 9409(2)



Scheme: Proposed routes to the formation of phlobatannins 7, 9, 10, and the C-2(F) epimer 2 (A-4) of procyanidin A-2

epicatechin **3** with 0.1M NaHCO₃ (pH 8.15) for 6.5 h at 40°C under nitrogen containing traces of oxygen, gave complete conversion into a mixture consisting of oligomeric procyanidins (*ca.* 70%) and five mobile fractions following chromatography on Sephadex LH-20 in ethanol. The latter fractions afforded a mixture (47:53 by ¹H NMR analysis) of (-)-epicatechin **4** and (-)-catechin **5** (*J*_{2,3} *ca.* 1.0 and 8.0 Hz respectively), and four pure compounds with modified C-rings (Scheme). These comprised the functionalized 8,9-*cis*-9,10-*trans*-3,4,9,10-tetrahydro-2*H*,8*H*-pyrano[2,3-*h*]chromene **9** as the major product, its C-2(F) epimer **10**, the 8,9-*trans*-9,10-*cis* analogue **7**, and the C-2(F) epimer **2** of proanthocyanidin A-2 **1**. The relative ease of purification of these phenols contrasts with those derived from the proflisetinidins^{12,13} where the additional chromatographic stages offered by successive methylation and acetylation were a prerequisite for compound purity.

The ¹H NMR spectra of the tetrahydropyrano[2,3-*h*]chromenes **7**, **9**, and **10** are conspicuously free of the effects of dynamic rotational isomerism¹⁸ at ambient temperatures hence indicating their pyran rearranged nature¹². ¹H NMR coupling constants (Table) of the heterocyclic proton resonances (*J*_{2,3} *ca.* 1.0 Hz for **7** and **9**, *J*_{2,3} 8.5 Hz for **10**; *J*_{8,9} 7.5, *J*_{9,10} 5.0 Hz for **7**; *J*_{8,9} *ca.* 1.0 Hz for **9** and **10**, *J*_{9,10} 2.5 and 2.0 Hz for **9** and **10** respectively) are in accordance¹² with 2,3-*cis*-8,9-*trans*-9,10-*cis* relative configuration for **7**, 2,3-*cis*-8,9-*cis*-9,10-*trans* for **9**, and 2,3-*trans*-8,9-*cis*-9,10-*trans* for **10**.

Table. ¹H NMR peaks of the tetrahydropyrano[2,3-*h*]chromenes **7**, **9**, and **10**, and the A-type procyanidin **2** in (CD₃)₂CO (23°C) at 300 MHz. Splitting patterns and J-values (Hz) are given in parentheses.

Ring Proton	7	9	10	2	
A	3/5 or 6/8	5.98(br. s)	5.90(d, 2.0) 5.98(d, 2.0)	5.94(d, 2.0) 6.01(d, 2.0)	6.08(d, 2.5) 6.10(d, 2.5)
B	2 5 6	6.86(d, 2.0) 6.74(d, 8.0) 6.67(dd, 2.0, 8.0)	6.65(d, 2.0) 6.70(d, 8.0) 6.46(dd, 2.0, 8.0)	6.63(d, 2.0) 6.68(d, 8.0) 6.47(dd, 2.0, 8.0)	7.17(d, 2.0) 6.81(d, 8.0) 7.03(dd, 2.0, 8.0)
C	8 9 10	4.99(d, 7.5) 4.21(dd, 5.0, 7.5) 4.73(d, 5.0)	5.65(br. s, <i>ca.</i> 1.0) 4.30(dd, 1.0, 2.5) 4.22(d, 2.5)	5.59(br. s, <i>ca.</i> 1.0) 4.15(dd, 1.0, 2.0) 4.19(d, 2.0)	3-H: 4.16(d, 3.5) 4-H: 4.29(d, 3.5)
D	6	6.06(s)	6.00(s)	5.86(s)	5.97(s)
E	2 5 6	6.57(d, 2.0) 6.58(d, 8.0) 6.17(dd, 2.0, 8.0)	6.66(d, 2.0) 6.77(d, 8.0) 6.12(dd, 2.0, 8.0)	6.85(d, 2.0) 6.76(d, 8.0) 6.69(dd, 2.0, 8.0)	6.83(d, 2.0) 6.73(d, 8.0) 6.67(dd, 2.0, 8.0)
F	2 3 4 4 _{ax} 4 _{eq}	4.76(br. s, <i>ca.</i> 1.0) 4.07(m) 2.70(dd, 3.0, 17.0) 2.81(dd, 4.5, 17.0)	4.69(br. s, <i>ca.</i> 1.0) 4.12(m) 2.69(dd, 2.5, 17.0) 2.85(dd, 5.0, 17.0)	4.53(d, 8.5) 3.97(m) 2.49(dd, 8.0, 16.0) 2.83(dd, 5.5, 16.0)	4.56(d, 7.5) 3.95(m) 2.54(dd, 8.0, 17.0) 2.89(dd, 5.5, 17.0)

In contrast to the analogous pyran rearranged products from the base-catalyzed conversion of procyanidin

B-3¹⁴ which were characterized as octamethyl ether diacetates thus facilitating demonstration of a 'liberated' phloroglucinol A-ring by ¹H nuclear Overhauser effect (n.O.e.) difference spectroscopy, a different approach had to be adopted towards the structural elucidation of free phenols 7, 9, and 10. The ¹H NMR spin systems in all three analogues were differentiated and fully analyzed by extensive spin-decoupling experiments using the benzylic 2-, 8-, and 10-H resonances as reference signals. A weak but structurally significant n.O.e. association (0.9, 1.9% resp.)^b of the broadened two-proton singlet (δ 5.98) of the A-ring with 2- and 6-H(E) (δ 6.57, 6.17 resp.) not only confirmed the tetrahydropyrano[2,3-*h*]chromene arrangement for analogue 7 but also established the location of the phloroglucinol moiety at C-10. It furthermore indicated a *cis*-relationship¹³ of the aryl substituents at C-2 and -10. When taken in conjunction with the known α -orientation of the E-ring of the (-)-epicatechin DEF unit, this also indicated an α -orientation for the phloroglucinol A-ring. A strong negative Cotton effect (CE) at 237 nm in the CD spectrum confirmed the 10 α -aryl substituent by application of the aromatic quadrant rule¹⁹. These CD and ¹H NMR (Table) features collectively facilitated definition of the absolute configuration of 7 as 2*R*,3*R*:8*S*,9*R*,10*R*.

The spin-decoupling experiments on analogues 9 and 10 also established a benzylic connection of 2-H(F) (δ 4.69, 4.53 for 9 and 10 resp.) and 10-H(C) (δ 4.22, 4.19 for 9 and 10 resp.) with the 2- and 6-protons of the respective pyrocatechol E- and B-rings. Besides the n.O.e. effect of 8-H(C) (δ 6.65, 5.59 for 9 and 10 resp.) with 2- and 6-H (3.5, 3.8 and 3.9, 4.3% for 9 and 10 resp.) of a pyrocatechol moiety in both 9 and 10, reminiscent of these structural types with a 3,4-disubstituted aryl group at C-10¹², the former proton did not exhibit additional long-range coupling thus reflecting the presence of an *ortho*-disubstituted phenyl residue at C-10. These observations hence implied an 'interchange' of the phloroglucinol A- and pyrocatechol B-rings in analogues 9 and 10 relative to the positions of these rings in the tetrahydropyrano[2,3-*h*]chromene 7^c. The chemical shifts of 8- and 10-H(C) and thus unambiguous proof for such an A-/B-ring interchange in 9 were confirmed by 2D-heteronuclear correlation of these protons with, respectively, C-8 and 10- (δ 69.04, 45.33 resp.). This experiment also facilitated assignment of the chemical shifts of all H-bearing carbons (*cf.* Experimental). Confirmation of both the *cis*-relationship between the C-2 and -10 pyrocatechol E- and B-rings, both α -orientated (*vide supra*), and the tetrahydropyrano[2,3-*h*]chromene arrangement of 9 was again furnished by the n.O.e. associations of both 5- and 6-H(B) (δ 6.70, 6.46 resp.) with 6-H(E) (δ 6.12) (0.7, 1.1% resp.). A high-amplitude negative CE at 238 nm in the CD spectrum confirmed the 10 α -aryl substituent hence facilitating, in conjunction with ¹H NMR coupling constants of C-ring protons, definition of 2*R*,3*R*:8*S*,9*S*,10*R* absolute configuration for 9. Although insufficient sample quantities precluded similar confirmation of the chemical shifts of 8- and 10-H(C) in 10, the very similar shifts of the C-ring protons in 9 and 10 (Table) were taken as sufficient proof of a [2,3-*h*]arrangement for 10. These small chemical shift differences may be attributed to the β -orientated E-ring in the latter compound. The conspicuous absence of n.O.e. associations between the protons of the pyro-

^bApproximated value owing to signal overlap

^cCompare refs. 12-14 for similar phenomena occurring during base-catalyzed conversions of profisetinidins and procyanidin B-3

catechol B- and E-rings in 10 presumably indicated a *trans*-relationship of these rings and hence a 10α B-ring. The CD spectrum, however, exhibited an intense positive CE at 239 nm which apparently reflected a 10β B-ring. Such a contradiction was commonly encountered for proflisetinidin related analogues^{12,13} with 2,10-*trans* aryl groups and had been explained by significant contributions of A-conformers^{20,21} (F-ring) reversing the sign of the low-wavelength CE. We thus favour the 2*S*,3*R*:8*S*,9*S*,10*R* absolute configuration depicted for analogue 10. The inversion of the absolute configuration at C-9 in 9 and 10 relative to that at C-3 in the biflavanoid precursor 3 is explained later.

The ¹H NMR spectrum (Table) of the remaining analogue 2 displayed in the heterocyclic proton region an AB-system [δ 4.16 (3-H), 4.29 (4-H), $J_{3,4}$ 3.5 Hz] characteristic² of the C-ring protons of A-type procyanidins. Confirmation of the chemical shift of 3-H was obtained by its pronounced n.o.e. association with both 2- and 6-H (B; 3.9, 4.3% resp.). Epimerization²³ at 2-C(F) under the mild basic conditions and hence conversion of the (-)-epicatechin DEF unit in precursor 3 to a (-)-catechin moiety in 2 was evident from the coupling constant ($J_{2,3}$ 7.5 Hz) of 2- and 3-H(F). A high-amplitude positive CE at 234 nm in the CD spectrum confirmed the β -orientation at C-4 and, combined with ¹H NMR data and known absolute configuration of procyanidin B-2 3, subsequently also the configuration as is depicted in 2. Analogue 2 therefore represents the C-2(F) epimer of procyanidin A-2 which was previously isolated by Nishioka *et al.*⁹ and designated proanthocyanidin A-4.

Under the mild basic conditions procyanidin B-2 3 is presumably transformed to an intermediate B-ring quinone-methide²² 6^d which then serves as common precursor to the novel tetrahydropyrano[2,3-*h*]-chromenes 7, 9, and 10. Analogue 7 originates *via* stereospecific pyran recyclization¹²⁻¹⁴ involving 7-OH(D) and the *si*-face at C-2 in quinone-methide 6. Migration of the (-)-epicatechin moiety, assisted by the strongly electron-releasing phloroglucinol unit at C-4, to the *si*-face at C-2 in 6 and subsequent recyclization *via* 7-OH(D) and the *si*-face at C-4 in quinone-methide 8^d may feasibly rationalize the genesis of the tetrahydropyrano[2,3-*h*]chromene with its 'interchanged' phloroglucinol A- and pyrocatechol B-rings. Inversion of configuration²¹ at the equivalent of C-3(C) in procyanidin B-2 3 associated with the observed ring interchange is substantiated by CD data (*vide supra*). The susceptibility of the E-ring in *eg.* 3, 6, and 8 to quinone-methide formation at alkaline pH^{12-14,22} presumably also initiates epimerization at C-2(F) and hence formation of the (-)-catechin DEF unit in the ring-interchanged tetrahydropyrano[2,3-*h*]chromene 10. Generation of the (-)-epicatechin/(-)-catechin mixture 4 and 5 is attributable to a similar phenomenon following cleavage of the base-labile¹⁷ interflavanyl bond in procyanidin B-2. A putative A-ring quinone-methide^{14,17} resulting from such a bond rupture may also induce the formation of the condensed analogues of unknown constitution *via* uncontrolled condensation with procyanidin B-2.

Formation of the A-/B-ring interchanged analogues 9 and 10, arising *via* the exclusive 1,3-migration of the (-)-epicatechin moiety in quinone-methide 6, contrasts with results for procyanidin B-3¹⁴ where pre-

^dQuinone-methides 6 and 8 are postulated and have not been isolated.

ferential migration of the phloroglucinol A-ring at 4-C in an analogous quinone-methide was observed. Such an exclusivity in the shift of the (-)-epicatechin unit with its reduced migratory aptitude compared to that of the phloroglucinol moiety at C-4 for quinone-methide **6**, is presumably attributable to steric factors. The *cis*-coplanarity of the β -orientated sp^3 -orbital adjoining C-4 and C-8(D) and the electron-deficient *p*-orbital at the *si*-face of C-2 should provide the additional driving force for the 1,3-migration of the (-)-epicatechin unit in **6**.

The transformation of procyanidin B-2 **3** into the C-2(F) epimer **2** of procyanidin A-2 **1** presumably involves the oxidative removal of hydride ion at C-2(C) as the initial step. The nature of the oxidizing species is, however, not clear. Although the trace amounts of oxygen may effect the transformation **3** \rightarrow **11**, it seems more reasonable to suggest that the prevailing conditions induce oxidation of the *o*-dihydroxy functionality of the pyrocatechol B- or E-rings to an *o*-quinone²³ which subsequently serves as oxidant for the conversion **3** \rightarrow **11**. Experiments aimed at verifying the latter proposal *via* selective protection of the 3'-OH groups of the pyrocatechol rings of **3** and related procyanidins are presently being investigated. These results will be discussed in an impending publication. The results presented here nevertheless provide unequivocal chemical evidence in favour of the β -orientated doubly-linked unit of procyanidin A-2 **1** and of its C-2(F) epimer **2**. The apparently exclusive formation of procyanidin A-4 **2** is explicable in terms of formation of an E-ring quinone-methide of either **3** or **11** and subsequent recyclization to the thermodynamically more stable 2,3-*trans* relative configuration.

Under these extremely mild conditions we could not find evidence of rearrangements of procyanidin B-2 to catechinic acid-type products^{24,25} reputed for either decreasing its reactivity towards aldehydes or enhancing acidity^{24,26,27}. Our results, when considered in conjunction with similar observations for procyanidin B-3¹⁴, hence indicate that with the proper selection of conditions, extraction of conifer barks^{15,16} at alkaline pH may be performed without the adverse effects which have hitherto hampered the successful application of such an approach towards the economically important procyanidins. The 'liberated' phloroglucinol-type A-rings in tetrahydropyrano[2,3-*h*]chromenes **7**, **9**, and **10** should, indeed, lead to increased reactivity towards aldehydes compared to that of procyanidin B-2 **3**.

EXPERIMENTAL

¹H NMR spectra were recorded on a Bruker AM-300 spectrometer, for (CD₃)₂CO solutions (D₂O exchange) with Me₄Si as internal standard. Mass spectral data were obtained with a Kratos MS80 instrument, and CD data in MeOH on a Jasco J-20 spectropolarimeter. TLC was performed on precoated Merck plastic sheets (DC-Plastikfolien Kieselgel 60 PF₂₅₄, 0.25 mm) and compounds were located by H₂SO₄-HCHO (40:1 v/v) spray reagent. CC was done on Sephadex LH-20 in EtOH at a flow rate of 0.5 cm³ min⁻¹. Evaporations were done under reduced pressure at ca. 60°C in a rotary evaporator.

Base-catalyzed Conversion of (4 β ,8)-Bis-(-)-epicatechin **3.** — Procyanidin B-2²⁸ **3** (500 mg) was dissolved in 0.1M NaHCO₃ (200 cm³) and the mixture was stirred for 6.5 h at 40°C under nitrogen containing traces of oxygen. The mixture was chilled with crushed ice, acidified (0.1M HCl) to pH 6, and extracted with EtOAc (8x200 cm³). Drying (Na₂SO₄) of the extract followed by evaporation of the solvent afforded a brown, amorphous solid (330 mg). This was subjected to CC (2.5x90 cm column; 15 cm³/tube; first 400 cm³ of eluant discarded) to give the following fractions: 1(tubes 56-66, 32 mg), 2(114-132, 17 mg), 3(146-150, 6 mg), 4(160-180, 29 mg), 5(211-240, 15 mg), and 6(245-280), 63 mg).

Fraction 1 consisted of a (-)epicatechin/(-)catechin mixture (47:53) and fraction 6 consisted of high-molecular-mass analogues of procyanidin B-2. Owing to its complexity this mixture was not further investigated.

Fraction 2 afforded the C-2(F) epimer 2 (A-4) of procyanidin A-2 as a white solid (Found: M⁺, 576.1260. C₃₀H₂₄O₁₂ requires M, 576.1268); ¹H NMR data (Table); CD [ε]₂₉₇ 0, [ε]₂₇₈ 1.0x10⁴, [ε]₂₅₅ 0, [ε]₂₃₄ 6.9x10⁴, and [ε]₂₀₀ 1.9x10⁴.

Fraction 3 gave (2R,3R:8S,9R,10R)-3,5,9-trihydroxy-2,8-bis-(3,4-dihydroxyphenyl)-10-(2,4,6-trihydroxyphenyl)-2,3-cis-8,9-trans-9,10-cis-3,4,9,10-tetrahydro-2H,8H-pyrano[2,3-h]chromene 7 as a white solid (Found: M⁺, 578.1418. C₃₀H₂₆O₁₂ requires M, 578.1424); ¹H NMR data (Table); CD [ε]₃₂₀ 0, [ε]₂₇₆ -5.3x10⁴, [ε]₂₅₂ -1.4x10⁴, [ε]₂₃₇ -8.9x10⁴, and [ε]₂₂₆ 0.

Fraction 4 afforded (2R,3R:8S,9S,10R)-3,5,9-trihydroxy-2,10-bis-(3,4-dihydroxyphenyl)-8-(2,4,6-trihydroxyphenyl)-2,3-cis-8,9-cis-9,10-trans-3,4,9,10-tetrahydro-2H,8H-pyrano[2,3-h]chromene 9 as a white solid (Found: M⁺ 578.1431. C₃₀H₂₆O₁₂ requires M, 578.1424); ¹H NMR data (Table); δ_c [(CD₃)₂CO; 23°C, 75.4 MHz) 96.07, 94.92 (C-3 + -5, A-ring), 116.27 (C-2, B)^e, 115.95 (C-5, B), 120.24 (C-6, B), 69.04 (C-8, C), 73.99 (C-9, C), 45.33 (C-10, C), 97.55 (C-6, D), 113.14 (C-2, E)^e, 116.05 (C-5, E), 118.28 (C-6, E), 79.31 (C-2, F), 66.79 (C-3, F), and 29.03 (C-4, F); CD [ε]₃₁₉ 0, [ε]₂₇₇ 2.6x10⁴, [ε]₂₄₆ 0, [ε]₂₃₈ -7.6x10⁴, and [ε]₂₂₄ 0.

Fraction 5 consisted of (2S,3R:8S,9S,10R)-3,5,9-trihydroxy-2,10-bis-(3,4-dihydroxyphenyl)-8-(2,4,6-trihydroxyphenyl)-2,3-trans-8,9-cis-9,10-trans-3,4,9,10-tetrahydro-2H,8H-pyrano[2,3-h]chromene 10 as a white solid (Found: M⁺, 578.1429. C₃₀H₂₆O₁₂ requires M, 578.1424); ¹H NMR data (Table); CD [ε]₃₀₂ 0, [ε]₂₇₀ -1.2x10⁴, [ε]₂₅₇ 0, [ε]₂₃₉ 5.6x10⁴, and [ε]₂₀₂ 0.

Acknowledgements

Support by the Foundation for Research Development, C.S.I.R., Pretoria, the Sentrale Navorsingsfonds of this University, and the Marketing Committee, Wattle Bark Industry of South Africa, Pietermaritzburg is acknowledged.

REFERENCES

- Mayer, W.; Goll, L.; Arndt, E.V.; Mannschreck, A. *Tetrahedron Lett.*, 1966, 429.
- Jacques, D.; Haslam, E.; Bedford, G.R.; Greatbanks, D. *J. Chem. Soc., Perkin Trans. 1*, 1974, 2263.
- Otsuka, H.; Fujioka, S.; Komiya, T.; Mizuta, E.; Takemoto, M. *Yakugaku Zasshi*, 1982, 102, 162.
- Weinges, K.; Kaltenhauser, W.; Marx, H.-D.; Nader, E.; Perner, J.; Seiler, D. *Annalen*, 1968, 711, 184.
- Hikino, H.; Shimoyama, N.; Kasahara, Y.; Takahashi, M.; Konno, C. *Heterocycles*, 1982, 19, 1381.
- Hikino, H.; Takahashi, M.; Konno, C. *Tetrahedron Lett.*, 1982, 23, 673.
- Ohigashi, H.; Minami, S.; Fukui, H.; Koshimizu, K.; Mizutani, F.; Sugijara, A.; Tomana, T. *Agric. Biol. Chem.*, 1982, 46, 2555.
- Nonaka, G.; Morimoto, S.; Nishioka, I. *J. Chem. Soc., Perkin Trans. 1*, 1983, 2139.
- Nonaka, G.; Morimoto, S.; Kinjo, J.; Nohara, T.; Nishioka, I. *Chem. Pharm. Bull.*, 1987, 35, 149.
- Morimoto, S.; Nonaka, G.; Nishioka, I. *Chem. Pharm. Bull.*, 1988, 36, 33.
- Porter, L.J. in *The Flavanoids - Advances in Research since 1980*, ed. Harborne, J.B., Chapman and Hall, London, 1988, 59.
- Steynberg, J.P.; Burger, J.F.W.; Young, D.A.; Brandt, E.V.; Steenkamp, J.A.; Ferreira, D. *J. Chem. Soc., Perkin Trans. 1*, 1988, 3323, 3331.
- Burger, J.F.W.; Steynberg, J.P.; Young, D.A.; Brandt, E.V.; Ferreira, D. *J. Chem. Soc., Perkin Trans. 1*, 1989, 671.
- Steynberg, J.P.; Bezuidenhout, B.C.B.; Burger, J.F.W.; Young, D.A.; Ferreira, D. *J. Chem. Soc., Perkin Trans. 1*, 1990, 203.
- Laks, P.; Hemingway, R.W. *J. Chem. Soc., Perkin Trans. 1*, 1987, 465, and refs. cited therein.

^eAssignments may be interchanged

16. Laks, P.E.; Hemingway, R.W.; Conner, A.H. *J. Chem. Soc., Perkin Trans. 1*, 1987, 1875, and refs. cited therein.
17. Hemingway, R.W.; Foo, L.Y. *J. Chem. Soc., Chem. Commun.*, 1983, 1035.
18. Du Preez, I.C.; Rowan, A.C.; Roux, D.G. *Chem. Commun.*, 1971, 315, and refs. cited therein.
19. Van der Westhuizen, J.H.; Ferreira, D.; Roux, D.G. *J. Chem. Soc., Perkin Trans. 1*, 1981, 1220.
20. Porter, L.J.; Wong, R.Y.; Benson, M.; Chan, B.G. *J. Chem. Res.*, 1986, (S) 86; (M) 830.
21. Steynberg, J.P.; Burger, J.F.W.; Young, D.A.; Brandt, E.V.; Ferreira, D. *Heterocycles*, 1989, 28, 923.
22. Whalley, W.B.; Mehta, P.P. *J. Chem. Soc.*, 1963, 5327.
23. Musso, H. in Oxidative Coupling of Phenols, eds. Taylor, W.I. and Battersby, A.R., Marcel Dekker, Inc., New York, 1967, 75.
24. Sears, K.D.; Casebier, R.L.; Hergert, H.L.; Stout, G.H.; McCandlish, L.E. *J. Org. Chem.*, 1974, 39, 3244.
25. Courbat, P.; Weith, A.; Albert, A.; Pelter, A. *Helv. Chim. Acta*, 1977, 60, 1665.
26. Herrick, F.H. *J. Agric. Food Chem.*, 1980, 28, 228.
27. Kiatgrajai, P.; Wellons, J.D.; Gollub, L.; White, J.D. *J. Org. Chem.*, 1982, 47, 2910.
28. Kolodziej, H.; Ferreira, D.; Roux, D.G. *J. Chem. Soc., Perkin Trans. 1*, 1984, 343.