


## Lability of Glycosidic Linkages of 2'-Thio-ribonucleosides

Richard Johnson, Colin B. Reese\* and Zhang Pei-Zhuo


Department of Chemistry, King's College London, Strand, London, WC2R 2LS, England.

**Abstract:** 2'-Thiouridine **2**, 2'-thioadenosine **5** and 2'-thiocytidine **8** undergo glycosidic cleavage in aqueous solution at 25°C at and above pH 6.5. The glycosidic linkages of [1-β-D-(2-thioarabinofuranosyl)]-adenine **7** and the dimeric disulfides **12a**, **12b** and **12c**, derived from the 2'-thio-ribonucleosides **2**, **5** and **8**, respectively, are stable under the same neutral and mildly basic conditions. 2'-Thioadenosine **5** and particularly [1-β-D-(2-thioarabinofuranosyl)]-adenine **7** are more stable than 2'-deoxyadenosine to acid-catalyzed hydrolysis.


2'-Thiouridine **2**, which was the first 2'-thio-ribonucleoside described in the literature, was obtained<sup>1</sup> by Imazawa *et al.* in virtually quantitative yield by treating its 2'-*S*-acetyl-3',5'-di-*O*-acetyl derivative **1** with a large excess of 1.0 mol dm<sup>-3</sup> potassium hydroxide in ethanol-water (1:1 v/v) at 10°C for 30 min. The latter workers also found<sup>1</sup> that uracil **3** was obtained as a major product when the triacetate **1** was treated "with ammonia or metallic alkali at high or moderate temperature". However these authors also reported<sup>1</sup> that isolated 2'-thiouridine **2** was "fairly stable in neutral or slightly acidic aqueous solution at room temperature". When the tetrabenzoyl derivative **4** of 2'-thioadenosine **5** was heated with sodium methoxide in methanol solution (see Scheme 1 and discussion below), no 2'-thioadenosine **5** was detected in the products<sup>2</sup> but adenine was obtained. In contrast with this result, the *arabino*-diastereoisomer **7** of 2'-thioadenosine **5** was obtained<sup>3</sup> in satisfactory yield when its *S*-acetyl derivative was treated with ammonia in methanol solution.



We have recently developed<sup>4,5</sup> what we believe to be a general synthesis of 2'-thio-ribonucleosides in which the 2'-thiol function is protected with an acid-labile protecting group that is removed in the final step. This approach has led not only to an alternative synthesis of 2'-thiouridine<sup>4</sup> **2** but also to the successful synthesis both of 2'-thiocytidine<sup>4</sup> **8** and 2'-thioadenosine<sup>5</sup> **5**. While previous attempts<sup>2,6</sup> to prepare 2'-thioadenosine **5** had all failed, a prior claim had been made for the successful synthesis<sup>7</sup> of 2'-thiocytidine **8**. However, the reported <sup>1</sup>H and <sup>13</sup>C NMR spectra<sup>7</sup> of the latter putative 2'-thiocytidine differed from the NMR spectroscopic data<sup>4</sup> that we obtained. In any case, the extreme ease with which 2'-thiocytidine **8** undergoes aerial oxidative dimerization (see below) must cast further doubt on validity of the earlier synthesis<sup>7</sup>. We originally adopted a strategy involving acidic unblocking conditions in order to avoid the possibility of the 2'-thio-ribonucleosides undergoing oxidative dimerization<sup>8</sup> to give the corresponding disulfides. However, it is clear that our approach to the synthesis of 2'-thio-ribonucleosides has the additional advantage that the possibility of base-promoted cleavage of the glycosidic linkages is also avoided.



The base-lability of the glycosidic linkages of 2'-thio-ribonucleosides recently became particularly apparent to us following two independent studies. First, when 2',3'-dithiouridine<sup>9</sup> **9** was treated with *ca.* 2 mol. equiv. of triethylamine in methanol solution at room temperature for 1 h, uracil was obtained and isolated in 88% yield (see Experimental). The second study was perhaps of greater significance. We<sup>10</sup> and others<sup>11-14</sup> believe that the modification of the 2'-hydroxy functions of oligoribonucleotides might well prove to be a valuable tool in the elucidation of the mechanism of ribozyme action. As the replacement of a hydroxy by a thiol function is an obvious such modification, we undertook<sup>10</sup> the synthesis of 2'-thiouridylyl-(3'→5')-uridine **10**. We found<sup>10</sup> that the 2'-thiol function in the latter dinucleoside phosphate **10** was unable to participate in the cleavage of the internucleotide linkage either under acidic or basic conditions, or in the presence of ribonuclease A. However, the dinucleoside phosphate **10** did show<sup>10</sup> a tendency to undergo aerial oxidative dimerization (to give the corresponding symmetrical 2'→2'-disulfide) under acidic, neutral and basic conditions. Under basic conditions (e.g. at pH 9.0, room temperature) the dinucleoside phosphate **10** also underwent fragmentation to give uracil **3** and uridine 5'-phosphate **11** as identifiable products<sup>10</sup>. The latter compound **11** presumably resulted from the further decomposition of the apyrimidinic acid formed following the cleavage of the bond joining uracil to the anomeric centre adjacent to the 2'-thiol function. We subsequently found that when a *ca.* 0.004 mol dm<sup>-3</sup> solution of 2'-thiouridine<sup>1,4</sup> **2** was allowed to stand at pH 9.0 in 0.1 mol dm<sup>-3</sup> tris hydrochloride buffer at room temperature for 18 h, uracil **3** was obtained and was isolated from the products in 53% yield (see Experimental).

**Table 1.** Reactions<sup>a</sup> of 2'-thio-ribonucleosides in 0.1 mol dm<sup>-3</sup> tris hydrochloride buffer at 25°C


Entry No.	Substrate	pH	% Substrate <sup>b</sup>	% Dimer <sup>b</sup>	% Aglycone <sup>b</sup>
1	2	6.5	6.4 (60.1)	68.8 (8.6)	11.6 (21.0)
2	2	7.0	10.1 (38.5)	44.3 (5.2)	27.4 (43.2)
3	2	8.0	13.7 (17.5)	39.0 (1.1)	34.3 (67.0)
4	2	9.0	8.9 (13.5)	31.6 (0.5)	42.2 (65.9)
5	2	10.0	15.4 (13.9)	14.9 (0.5)	51.1 (62.1)
6	8	6.5	2.4 (59.9)	92.0 (6.9)	4.9 (20.9)
7	8	7.0	2.1 (51.9)	94.7 (5.0)	2.3 (16.2)
8	8	8.0	5.0 (76.3)	85.7 (4.7)	4.9 (9.3)
9	8	9.0	7.6 (69.4)	87.0 (3.7)	3.9 (1.6)
10	8	10.0	6.6 (60.0)	76.5 (2.8)	2.4 (1.3)
11	5	6.5	29.7 (52.9)	49.4 (2.2)	15.7 (24.0)
12	5	7.0	18.5 (60.8)	64.6 (2.1)	10.1 (19.6)
13	5	8.0	28.5 (74.4)	65.2 (2.4)	4.4 (7.1)
14	5	9.0	35.1 (78.9)	58.9 (2.9)	3.6 (4.3)
15	5	10.0	25.0 (77.1)	65.3 (2.8)	6.4 (3.2)

<sup>a</sup>Initial substrate concentrations were *ca.* 0.004 mol dm<sup>-3</sup> and all reactions were allowed to proceed for 20 h. <sup>b</sup>Percentage estimates are based on reverse phase HPLC analyses. Experiments were carried out both in the absence and presence of 0.04 mol dm<sup>-3</sup> 2-mercaptoethanol. The italicized numbers in parentheses represent the percentages obtained in the presence of 2-mercaptoethanol. In the latter experiments other unidentified products, possibly including unsymmetrical disulfides, were detected by HPLC.

These observations have prompted us to investigate the dependence of the base-promoted cleavage of the glycosidic linkages of 2'-thio-ribonucleosides on pH. Solutions (*ca.* 0.004 mol dm<sup>-3</sup>) of 2'-thiouridine 2 in 0.1 mol dm<sup>-3</sup> tris hydrochloride buffer (pH 6.5, 7.0, 8.0, 9.0 and 10.0) were allowed to stand at 25°C for 20 h and were then analyzed by reverse phase HPLC. It can be seen from Table 1 (entries nos. 1-5) that, in all five experiments, little substrate remained. However, while dimer<sup>1,4</sup> 12a predominated at pH 6.5 - 8.0, uracil 3 was the main product at pH 9.0 and 10.0. As the experiments were carried out with only *ca.* 0.001 g (0.004

mmol) of substrate **2**, it did not prove possible to prevent the occurrence of aerial oxidative dimerization. The susceptibility of 2'-thiouridine **2** to undergo glycosidic cleavage was more clearly revealed when a *ca.* tenfold excess of 2-mercaptoethanol over substrate was added to the reaction solution to impede oxidative dimerization (Table 1, entries nos. 1-5, italicized numbers in parentheses).

In the absence of 2-mercaptoethanol, 2'-thiocytidine<sup>4</sup> **8** (Table 1, entries nos. 6-10) showed more tendency to undergo oxidative dimerization (see above) than 2'-thiouridine **2**, and very little cytosine **13** was obtained over the whole pH range examined. As can be seen from the experiments carried out in the presence of 2-mercaptoethanol (entries nos. 6-10, italicized numbers in parentheses), even when oxidative dimerization was largely suppressed, 2'-thiocytidine **8** showed little tendency to undergo glycosidic cleavage except perhaps at pH 6.5 and 7.0. In the absence of 2-mercaptoethanol, 2'-thioadenosine<sup>5</sup> **5** (entries nos. 11-15) also showed more tendency to undergo oxidative dimerization than glycosidic cleavage, and again the corresponding experiments carried out in the presence of 2-mercaptoethanol (entries nos. 11-15, italicized numbers in parentheses) confirmed that glycosidic cleavage did not occur readily, especially at pH 8.0 - 10.0. It is noteworthy that all three dimers<sup>1,4,5</sup> **12a-c** were stable in 0.1 mol dm<sup>-3</sup> tris hydrochloride buffer at 25°C for 20 h over the whole pH range (6.5 - 10.0) examined, and in no case was the corresponding aglycone detected. Finally, while no adenine **6** was obtained when the *arabino*-compound<sup>3,5</sup> **7** was allowed to stand at 25°C for 20 h in 0.1 mol dm<sup>-3</sup> tris hydrochloride buffer solution over the whole pH range (6.5 - 10.0), extensive oxidative dimerization was observed (see Experimental).



**Scheme 1** Reagents and conditions: i, NaOMe, MeOH, reflux, 2 hr; ii, Ac<sub>2</sub>O, C<sub>6</sub>H<sub>5</sub>N, RT

Ryan *et al.* reported<sup>2</sup> that when the tetrabenzoyl derivative **4** was heated, under reflux, with sodium methoxide in methanol solution (Scheme 1) and the products were acetylated after the removal of adenine **6**, the dimeric glycoside **16** was obtained in 57% isolated yield. Although they were unable firmly to establish the intermediacy of 2'-thioadenosine **5** or its 6-*N*-benzoyl derivative, the latter workers suggested<sup>2</sup> that adenine resulted from the internal nucleophilic attack of the released 2'-thiolate ion on the anomeric centre leading, *via* an intermediate episulfide **14** and monomeric glycoside **15**, to the isolated product **16**. In the present study, only the 2'-thio-ribonucleosides **2**, **5** and **8** and neither the dimers **12a-c** nor the *arabino*-2'-thio-nucleoside **7** decomposed to give the corresponding aglycones in the pH range 6.5 - 10.0. It is therefore reasonable to suggest that, under these conditions, the deglycosylation of the above 2'-thionucleosides **2**, **5** and **8** is accompanied by the formation of the intermediate episulfide<sup>2</sup> **14**. It is relevant to add that Imazawa *et al.* estimated<sup>1</sup> that the pK<sub>a</sub> of the thiol function in 2'-thiouridine **2** was 7.3. What is still perhaps difficult to rationalize in connection with the data presented in Table 1 is the observed dependence of the percentage yield of aglycone on pH and on the nature of the aglycone itself.

As would be expected<sup>15</sup>, 2'-thioadenosine **5** [ $t_{1/2} = ca. 62$  min in 0.01 mol dm<sup>-3</sup> hydrochloric acid (pH 2.0) at 60°C] underwent acid-catalyzed cleavage of its glycosidic linkage less readily than 2'-deoxyadenosine [ $t_{1/2} = ca. 22$  min under the same conditions] but much more readily than adenosine. It is noteworthy that 9-[1- $\beta$ -D-(2-thioarabinofuranosyl)]-adenine **7** [ $t_{1/2} = ca. 42$  min in 0.01 mol dm<sup>-3</sup> hydrochloric acid at 100°C] was appreciably more stable to acidic hydrolysis than 2'-thioadenosine **5**. As expected, 2'-thiouridine **2** was very resistant indeed to acidic hydrolysis under the latter conditions [pH 2.0, 100°C].

## EXPERIMENTAL

NMR spectra were measured at 360 MHz with a Bruker AM 360 spectrometer; tetramethylsilane was used as an internal standard. Merck silica gel 60 F<sub>254</sub> TLC plates were developed in solvent system A [chloroform-methanol (9:1 v/v)]. Liquid chromatography (HPLC) was carried out on a Jones APEX Octadecyl 5 $\mu$  column which was eluted with 0.1 mol dm<sup>-3</sup> triethylammonium acetate - acetonitrile mixtures. Tris hydrochloride buffer solutions were prepared by following a literature procedure<sup>16</sup>. The 2'-thionucleoside and disulfide substrates were all prepared by literature procedures<sup>4,5</sup>.

*Reaction Between 2',3'-Dithiouridine 9 and Triethylamine in Methanol Solution.* - 2',3'-Dithiouridine<sup>9</sup> (0.10g, 0.036 mmol) was dissolved in a 0.36 mol dm<sup>-3</sup> solution of triethylamine in methanol (2.0 cm<sup>3</sup>) at room temperature. After 1 h, when TLC (system A) revealed that no starting material remained, the products were filtered and the residue was washed with ether. The dried residue (0.036 g, 88%) was identified as uracil;  $R_f$  0.21 (system A);  $\delta_C$  [(CD<sub>3</sub>)<sub>2</sub>SO] 10.3, 142.3, 151.6, 164.4.

*Decomposition of 2'-Thiouridine 2 in 0.10 mol dm<sup>-3</sup> Tris Hydrochloride Buffer (pH 9.0).* - 2'-Thiouridine (0.10 g, 0.38 mmol) was dissolved in 0.10 mol dm<sup>-3</sup> tris hydrochloride buffer (pH 9.0, 100 cm<sup>3</sup>) at room temperature. After 18 h, the products were concentrated under reduced pressure and the residue was chromatographed on a column of DEAE Sephadex A-25. The column was eluted with a linear gradient (from 0.01 - 0.80 mol dm<sup>-3</sup> over 1.0 dm<sup>3</sup>) of aqueous triethylammonium bicarbonate buffer (pH 7.5). The appropriate fractions were combined and concentrated under reduced pressure. Crystallization of the residue from water gave uracil (0.023 g, 53%), identical [TLC (system A), <sup>13</sup>C NMR] to authentic material.

*Behaviour of 2'-Thionucleosides 2, 5, 8 and 7 and Corresponding Disulfides 12a, 12b and 12c in 0.1 mol dm<sup>-3</sup> Tris Hydrochloride Buffer Solutions (pH 6.5 - 10.0) at 25°C.* - 0.1 mol dm<sup>-3</sup> Tris hydrochloride buffer solutions were prepared. Each of the 7 substrates (**2**, **5**, **8**, **7**, **12a**, **12b** and **12c**; 0.004 mmol) was dissolved in 1.0 cm<sup>3</sup> of each of the 5 buffer solutions (pH 6.5, 7.0, 8.0, 9.0 and 10.0), and the solutions were placed in sealed vessels. All of the solutions were kept at 25°C for 20 h, and were then analyzed by HPLC. The HPLC column was eluted with mixtures of 0.1 mol dm<sup>-3</sup> triethylammonium acetate - acetonitrile, using either isocratic or gradient elution programmes as appropriate. The results obtained with substrates **2**, **5** and **8** are given in Table 1 (entries nos. 1-5, 11-15 and 6-10, respectively). [1- $\beta$ -D-(2-Thioarabinofuranosyl)]-adenine **7** was found to undergo only oxidative dimerization (52, 83, 98, 95 and 97% at pH 6.5, 7.0, 8.0, 9.0 and 10.0, respectively), and the symmetrical disulfides **12a**, **12b** and **12c** remained unchanged.

Reactions involving the three 2'-thio-ribonucleosides **2**, **5** and **8** were carried out in the same way in buffer solutions that were also 0.04 molar with respect to 2-mercaptoethanol. The results of these experiments are also given in Table 1 (entries nos. 1-5 for 2'-thiouridine **2**; entries nos 6-10 for 2'-thiocytidine **8**; entries nos. 11-15 for the 2'-thioadenosine **5**).

*Acid-catalysed Hydrolysis of 2'-Deoxyadenosine, 2'-Thioadenosine 5 and [1-β-D-(2-Thioarabinofuranosyl)]-adenine 7.* - A solution of substrate (0.01 g) in 0.01 mol dm<sup>-3</sup> hydrochloric acid (pH 2.0, 10 cm<sup>3</sup>), contained in a sealed vessel, was heated either at 60°C (for 2'-deoxyadenosine and 2'-thioadenosine 5) or at 100°C (for [1-β-D-(2-thioarabinofuranosyl)]-adenine 7). Aliquots of the hydrolysate were removed after appropriate intervals of time and analyzed by HPLC. Satisfactory straight lines were obtained by plotting log<sub>10</sub> (remaining substrate) against time. The half-times of hydrolysis of 2'-deoxyadenosine and 2'-thioadenosine 5 were found to be *ca.* 22 and 62 min, respectively, at 60°C; the half-time of hydrolysis of [1-β-D-(2-thioarabinofuranosyl)]-adenine 7 was found to be *ca.* 42 min at 100°C. Some concomitant oxidative dimerization both of 2'-thioadenosine 5 and [1-β-D-(2-thioarabinofuranosyl)]-adenine 7 occurred in the course of hydrolysis. The rate of hydrolysis of 2'-thiouridine 2 was too slow to measure with any degree of accuracy under the latter conditions.

One of us (R.J.) thanks the Medical Research Council AIDS Directed Programme for the award of a research studentship.

#### REFERENCES

1. Imazawa, M.; Ueda, T.; Ukita, T. *Chem. Pharm. Bull. Japan*, **1975**, *23*, 604-610.
2. Ryan, K. J.; Acton, E. M.; Goodman, L. *J. Org. Chem.* **1971**, *36*, 2646-2657.
3. Ranganathan, R.; Larwood, D. *Tetrahedron Lett.*, **1978**, 4341-4344.
4. Divakar, K. J.; Mottoh, A.; Reese, C. B.; Sanghvi, Y. S. *J. Chem. Soc. Perkin Trans. 1*, **1990**, 969-974.
5. Marriott, J. H.; Mottahedeh, M.; Reese, C. B. *Carbohydr. Res.*, **1991**, *216*, 257-269.
6. Mengel, R.; Griesser, H. *Tetrahedron Lett.*, **1977**, 1177-1180.
7. Patel, A. D.; Schrier, W. H.; Nagyvary, J. *J. Org. Chem.*, **1980**, *45*, 4830-4834.
8. Reist, E. J.; Benitez, A.; Goodman, L. *J. Org. Chem.*, **1964**, *29*, 554-558.
9. Johnson, R.; Joshi, B. V.; Reese, C. B. *J. Chem. Soc., Chem. Commun.*, **1994**, 133-134.
10. Reese, C. B.; Simons, C.; Zhang, P-Z. *J. Chem. Soc., Chem. Commun.*, **1994**, 1809-1810.
11. Jencks, W. P. *Nature*, **1992**, *358*, 543-544.
12. Picken, W. A.; Olsen, D. B.; Bonsaler, F.; Aurup, H.; Eckstein, F. *Science*, **1991**, *253*, 314-317.
13. Williams, D. M.; Picken, W. A.; Eckstein, F. *Proc. Natl. Acad. Sci USA*, **1992**, *89*, 918-921.
14. Aurup, H.; Tuschl, T.; Bonsaler, F.; Ludwig, J.; Eckstein, F. *Nucleic Acids Res.*, **1994**, *22*, 20-24.
15. Zielonacka-Lis, E. *Nucleosides and Nucleotides*, **1989**, *8*, 383-405.
16. *Data for Biomedical Research, 2nd Edn.*, Dawson, R.M.C.; Elliott, D.C.; Elliott, W. H.; Jones, K. M.; Eds.; Clarendon, Oxford, 1969, p. 494.