

Conformationally Restricted Analogues of Methionine: Synthesis of Chiral 3-Amino-5-methylthio-2-piperidones

Ricardo Rodríguez, M. Angels Estiarte, Anna Diez, and Mario Rubiralta*

Laboratori de Química Orgànica. Facultat de Farmàcia. Universitat de Barcelona, 08028-Barcelona, Spain

Anna Colell, Carmen García-Ruiz, and José C. Fernández-Checa

Unitat Hepatologia. Hospital Clínic Provincial. Universitat de Barcelona, 08036-Barcelona, Spain


Abstract — (αR , $3RS$, $5S$)-3-Amino-*N*-(2-hydroxy-1-phenylethyl)-5-methylthio-2-piperidones **1** have been synthesized from enamide **2** by subsequent free radical addition of methanethiol on position 5 and amination of 3-position.
 Copyright © 1996 Elsevier Science Ltd

INTRODUCTION

Methionine¹ and cystathionine² have shown to inhibit the glutathione (γ -glutamylcysteinylglycine, GSH) efflux of isolated hepatocytes. Hepatic GSH is exclusively synthesized in the cytosol and needs specific carriers for its transport into bile and blood circulation, and thus reach distal extrahepatic organs.³ It has also been reported that cystathionine is a more powerful GSH transport inhibitor than methionine.^{1,2}

In the context of our studies on the synthesis of conformationally restricted pseudopeptides presenting a 3-amino-2-piperidone backbone,⁴ we have envisaged the synthesis of 3-amino-5-methylthio-2-piperidones **1** as conformationally restricted analogues of methionine, to be tested as a potential inhibitor of the hepatic transport of glutathione (GSH).^{5,6}


The synthesis of the methionine-phenylglycinol derivatives **1** was planned by subsequent functionalization of positions 5 and 3 of enamide **2**.



Scheme 1

RESULTS AND DISCUSSION

We first studied the introduction of the amino group on 3-position using 2-piperidone **4**⁷⁻⁹ as a model structure. Treatment of compound **4** with *sec*-BuLi (2.5 equivalents) and Br₂ led to the expected C-3 epimeric mixture of 3-bromolactams **5** (**a**:**b** = 1.3:1). However, the minor isomer **5b** epimerizes to give **5a** on SiO₂ flash column chromatography, and also on standing.^{9,10} Subsequent substitution of the bromine atom through treatment of **5a**¹¹ with NaN₃ or potassium phthalimide led to the corresponding 3-azido- and 3-phthalimido derivatives **6** and **7** as single diastereomers. The reduction of azide **6** was performed by reaction with PPh₃-H₂O at room temperature¹² to give 3-amino-2-piperidone **8** in 16% yield. Amine **8** was also obtained from compound **7** by hydrazinolysis in 65% yield (Scheme 2). The analytical data of bromolactams **5** and aminolactam (*αR,3R*)-**8** were identical to the ones found by Prof. Husson's group.¹⁰




Reagents and conditions: i) 1. *sec*-BuLi (2.5 equivalents), THF, -78°C, 20 min. 2. Br₂ (1.1 equivalent). 1 h 45 min (55%); ii) SiO₂ or spontaneously; iii) NaN₃ (2 equivalents), H₂O-DMF/AcOH, room temperature, 7 h, (78%); iv) potassium phthalimide (2 equivalents), DMF, room temperature, 18 h (56%); v) 1. PPh₃ (1.1 equivalents), THF, room temperature, 6 h. 2. H₂O (1.5 equivalents), 15 h (16%); vi) NH₂NH₂·H₂O (10 equivalents) MeOH, reflux, 1h (70%).

Scheme 2

Enamide **2** was prepared in two steps and 69% yield from imide **3** (Scheme 3).⁷ Thus, Super-Hydride[®] reduction¹³ of **3** gave the epimeric mixture of 6-hydroxylactams **9**, which yielded enamide **2** on TFA treatment. Enamide **2** was easily identified by the presence of the olefin signals in its ¹H and ¹³C NMR spectra (see experimental).

Since enamide **2** can be regarded as a deactivated enamine,¹⁴ its double bond was expected to react as an alkene rather than as an enamine. Indeed, when enamide **2** was treated with electrophiles such as Me₂S₂ or methyl methanethiolsulfonate no reaction was observed, but the AIBN activated free-radical addition of MeSH¹⁵ followed by acid hydrolysis of the acetate group¹⁵ gave the expected anti-Markovnikov diastereomeric 5-methylthiolactams **10a,b** in a ratio of 1:1.2. The most characteristic ¹H NMR data of mercaptane **10a** were a singlet at δ 1.96 (SMe), a double doublet at δ 3.20 (6-H_e), and a double doublet at δ 3.11 (6-H_a) whose

coupling constants ($J = 11$ and 7 Hz) indicated that the C-5 SMe group adopted an axial disposition. The ^1H NMR signals for isomer **10b** were a singlet at δ 1.90 (SMe), a double doublet at δ 3.47 (6- H_e) and a double doublet at δ 2.88 (6- H_a , $J = 12$ and 7 Hz), which indicated that the C-5 SMe group was also axially disposed.




Reagents and conditions: i) LiBHET_3 (1.5 equivalents), THF, -78°C , 2 h; ii) TFA, CH_2Cl_2 , room temperature, 30 min (69% two steps); iii) MeSH (10 equivalents), AIBN (catalytic), THF, -78°C , 15 h (62%); iv) 5% aqueous KOH, THF, reflux, 3 h (98%); v) *sec*-BuLi (2.5 equivalents), THF, -78°C , 20 min. 2. Br_2 (1.1 equivalents), 1 min 45 sec (45%).

Scheme 3

The introduction of the amino group on 3-position of compound **10b** was then carried out following the procedure shown in Scheme 2. Thus, the bromination of compound **10b** yielded a 3:1 diastereomeric mixture of bromides **11a,b** (Scheme 3). The major isomer, **11a**, was assigned as 3,5-*trans* isomer on the basis of its spectral data. Thus, the axial proton on C-6 appeared at δ 2.89 as a double doublet of $J = 12$ and 10 Hz, showing that the C-5 SMe group was equatorial. The signal corresponding to 3-H was a narrow triplet ($J = 5$ Hz) at δ 4.64, indicating that the bromine atom is axially disposed. In addition, no 2D nOe correlation was observed between 3-H and 5-H. Since the bromination gave mainly the *3S* configuration, we deduced that compound **11a** was (*3S,5S*). By comparison, the minor isomer **11b** was assigned as the 3,5-*cis* isomer, i.e. (*3R,5S*). Subsequent substitution of the bromine atom through treatment of **11a** with NaN_3 led to the corresponding 3-azido derivative **12** as a single diastereomer in 70% yield. The ^1H NMR data of compound **12** showed that the substituents on C-3 and C-5 were *cis*, and therefore (*3R,5S*), as expected. The diagnostic signals in this case were the triple doublet corresponding to the axial proton on C-4 (δ 1.65, $J = 12$ and 11 Hz) and the double doublet corresponding to the equatorial C-6 proton (δ 3.42, $J = 12$ and 5 Hz), respectively. The $\text{Ph}_3\text{P-H}_2\text{O}$ reduction of compound **12** yielded the target aminolactam **1a**. The structural assignment of compound **1a** was again inferred from the ^1H NMR data to be the (*3R,5S*)-3-amino-5-methylthio derivative, with both substituents equatorially disposed.

Surprisingly, when bromide **11a** was treated with potassium phthalimide we obtained a 5:1 diastereomeric mixture of phthalimidolactams **13**, from which the major isomer **13b** was isolated pure. According to its ^1H NMR spectrum, compound **13b** showed a 3,5-*trans* relationship. Thus, the axial proton on C-6 appeared as a broad doublet ($J = 13$ Hz), which implied that the SMe group was in an axial disposition.

and 3-H proton was a double doublet at δ 5.17 ($J = 11$ and 8 Hz), indicating that the phthalimido group was equatorial. This result can only be explained if the SMe substituent prefers an axial disposition, as it does in compounds **10**, which would oblige an epimerization of C-3 to give an equatorial disposition to the bulkiest phthalimido substituent, and therefore, a (3*S*,5*S*) stereochemistry. Hydrazinolysis of compound **13b** yielded compound **1b**, which was assigned on the basis of its spectral data



Reagents and conditions: i) NaN_3 (2 equivalents), H_2O -DMF/AcOH, room temperature, 7 h, (70%); ii) 1. PPh_3 (1.1 equivalents), THF, room temperature, 12 h. 2. H_2O (1.5 equivalents), 6 h (30%); iii) potassium phthalimide (1.5 equivalents), DMF, room temperature, 24 h, (64%); iv) $\text{NH}_2\text{NH}_2 \cdot \text{H}_2\text{O}$ (10 equivalents), MeOH, reflux, 3.5 h (73%).

Scheme 4

CONCLUSION

The application of compounds **1a** and **1b** as inhibitors of the hepatic transport of GSH in relation with other known thioether aminoacids is currently under way, both on isolated rat hepatocytes and on oocytes from *Xenopus Laevis* expressing the sinusoidal transport system of GSH.

EXPERIMENTAL

General. Melting points were determined in a capillary tube on a Büchi apparatus. ^1H and ^{13}C NMR spectra were recorded on a Varian Gemini-200 instrument (200 MHz) and 2D NMR COSY experiments were performed on a Varian XL-500 instrument (500 MHz). Unless otherwise noted, NMR spectra were registered in CDCl_3 , and chemical shifts are expressed in parts per million (δ) relative to internal Me_4Si . IR spectra were recorded on a Nicolet FT-IR spectrophotometer. Mass spectra were determined on a Hewlett-Packard 5988A mass spectrometer. Flash column chromatography was carried out on SiO_2 (silica gel 60, 40-63 mm, SDS). TLC was performed on SiO_2 (silica gel 60 F254, Macherey-Nagel) and developed with the solvent

described in each case for flash chromatography. The spots were located by UV light and Dragendorff or hexachloroplatinate reagent. Optical rotations have been measured with a Perkin-Elmer 241 polarimeter. Purification of reagents and solvents was effected according to standard methods. Prior to concentration under reduced pressure, all extracts were dried over anhydrous Na₂SO₄. Microanalyses were performed on a Carlo Erba 1106 analyzer by the Departament de Química Orgànica i Biològica, CID, Barcelona.

($\alpha R, 3RS$)-3-Bromo-*N*-(2-hydroxy-1-phenylethyl)piperidin-2-ones (5a,b). To a solution of piperidone **4**⁷ (500 mg, 2.28 mmol) in dry THF (25 ml), cooled at -78°C, *sec*-Buli (5.3 ml, 6.84 mmol) was added. After 20 min, cold Br₂ (25 ml, 0.45 mmol) was added dropwise. The resulting mixture was stirred for 2 min 15 sec at -78°C and the reaction was quenched with aqueous NH₄Cl. The solvent was evaporated and the residue, dissolved in CH₂Cl₂, was washed with brine. The organic extracts were dried and evaporated to yield a mixture of ($\alpha R, 3S$)-**5a** and ($\alpha R, 3R$)-**5b**, which were isolated pure by flash chromatography (SiO₂, AcOEt-hexane, 1:2). **Bromolactam** ($\alpha R, 3S$)-**5a** (300 mg, 44%): [α]_D = -112 (c = 0.8, MeOH); IR (NaCl) 3400 (OH), 1650 (CO) cm⁻¹; ¹H NMR (500 MHz) 1.75 (dm, *J* = 14 Hz, 1H, 5-H_e), 2.10 (m, 1H, 5-H_a), 2.23 (dm, *J* = 14 Hz, 1H, 4-H_e), 2.31 (tt, *J* = 14 and 4 Hz, 1H, 4-H_a), 3.08 (dt, *J* = 12 and 4 Hz, 1H, 6-H_e), 3.29 (td, *J* = 12 and 4 Hz, 1H, 6-H_a), 4.06 (dd, *J* = 11.5 and 10 Hz, 1H, β -H_A), 4.19 (dd, *J* = 11.5 and 5 Hz, 1H, β -H_B), 4.66 (t, *J* = 4 Hz, 1H, 3-H_e), 5.83 (dd, *J* = 10 and 5 Hz, 1H, α -H), 7.20-7.40 (m, 5H, Ph-H); ¹³C NMR 18.7 (C-5), 30.9 (C-4), 42.3 (C-6), 45.9 (C-3), 57.5 (C- α), 60.3 (C- β), 127.3 (Ph-*o*), 127.7 (Ph-*p*), 128.5 (Ph-*m*), 136.5 (C-*ipso*), 168.0 (C-2); EIMS *m/z* (%) 299 (M⁺+1, 1), 297 (M⁺-1, 1), 268 (99), 266 (100), 188 (54), 186 (60), 91 (60). Anal. Calcd for C₁₃H₁₆BrNO₂: C, 52.52; H, 5.43; N, 4.71; Br, 26.57. Found: C, 52.36; H, 5.37; N, 4.69; Br, 26.82.

Bromolactam ($\alpha R, 3R$)-**5b** (106 mg, 15%): ¹H NMR 1.65 (m, 1H, 5-H_e), 2.15 (m, 3H, 4-H and 5-H_a), 2.95 (m, 1H, 6-H_e), 3.30 (m, 1H, 6-H_a), 3.60 (br s, 1H, OH), 4.05 (m, 2H, β -H), 4.60 (m, 1H, 3-H), 5.60 (dd, *J* = 11 and 9 Hz, 1H, α -H), 7.15-7.35 (m, 5H, Ph-H); ¹³C NMR 19.3 (C-5), 30.8 (C-4), 43.9 (C-6), 45.7 (C-3), 59.9 (C- α), 61.3 (C- β), 127.8 (Ph-*o*), 127.9 (Ph-*p*), 128.7 (Ph-*m*), 136.0 (Ph-*ipso*), 167.9 (C-2); EIMS *m/z* (%) 299 (M⁺+2, 21), 298 (M⁺, 21), 268 (100), 266 (99), 200 (87), 186 (64), 159 (38), 91 (67), 77 (39).

Pure isomers **5a** and **5b** epimerize in solution to give a 2:1 mixture of **5a:5b**. Epimerization of **5b** is particularly quick.

($\alpha R, 3R$)-3-Azido-*N*-(2-hydroxy-1-phenylethyl)piperidin-2-one (6). To a solution of pure bromide **5a** (1.05 g, 3.53 mmol) in DMF (40 ml) containing AcOH (1.8 ml), cooled at 0°C, a solution of NaN₃ (459 mg, 7.1 mmol) in H₂O (1.8 ml) was added. The resulting mixture was stirred for 7 h at room temperature. The layers were separated and the aqueous phase was extracted with CH₂Cl₂. The organic extracts were washed with brine, dried and evaporated to yield azidopiperidone **6** (715 mg, 78%), which were isolated by flash chromatography (SiO₂, AcOEt-hexane, 1:2): [α]_D = +33.5 (c = 1, CHCl₃); IR (NaCl) 3500 (OH), 2103 (N₃), 1639 (CO) cm⁻¹; ¹H NMR 1.65 (m, 1H, 5-H), 1.75 (m, 1H, 5-H), 1.90 (m, 1H, 4-H), 2.05 (m, 1H, 4-H), 2.98 (dt, *J* = 12 and 5 Hz, 1H, 6-H_e), 3.14 (br s, 1H, OH), 3.26 (ddd, *J* = 12, 8 and 4 Hz, 1H, 6-H_a), 4.15 (m, 3H, β -H and 3-H), 5.80 (dd, *J* = 11 and 7 Hz, 1H, α -H), 7.20-7.40 (m, 5H, Ph-H); ¹³C NMR 20.2 (C-5), 27.3 (C-4), 43.3 (C-6), 59.1 (C-3), 59.7 (C- α), 61.3 (C- β), 127.7 (Ph-*o*), 128.0 (Ph-*p*), 128.8 (Ph-*m*), 136.3 (C-*ipso*), 169.2 (C-2); EIMS *m/z* (%) 260 (M⁺, 1), 229 (99), 200 (71), 173 (24), 146 (21), 103 (88), 91 (100). Anal. Calcd for C₁₃H₁₆N₄O₂: C, 59.97; H, 6.20; N, 21.53. Found: C, 59.67; H, 6.34; N, 21.48.

(*αR,3R*)-*N*-(2-Hydroxy-1-phenylethyl)-3-phthalimidopiperidin-2-one (**7**). To a solution of bromide **5a** (100 mg, 0.33 mmol) in DMF (2 ml), a solution of potassium phthalimide (124 mg, 0.67 mmol) in DMF (1.5 ml) was added. After stirring for 18 h at room temperature, the mixture was washed with brine. The aqueous layer was extracted with CH₂Cl₂ and the combined organic extracts, dried and evaporated, were flash chromatographed (AcOEt-hexane, 2:1) to yield phthalimidopiperidone **7** (67 mg, 56%): [α]_D = -49.5 (c = 1, CHCl₃); IR (NaCl) 3453 (OH), 1716 (CO), 1648 (CO) cm⁻¹; ¹H NMR 1.70 (m, 1H, 5-H_a), 1.90 (dm, *J* = 14 Hz, 1H, 5-H_e), 2.65 (m, 1H, 4-H_e), 2.21 (qd, *J* = 12 and 3 Hz, 1H, 4-H_a), 2.93 (br s, 1H, OH), 3.00 (br d, *J* = 12 Hz, 1H, 6-H_e), 3.36 (td, *J* = 12 and 4 Hz, 1H, 6-H_a), 4.10 (m, 2H, β-H), 4.80 (dd, *J* = 11 and 7 Hz, 1H, 3-H_a), 5.80 (dd, *J* = 8 and 6 Hz, 1H, α-H), 7.20-7.40 (m, 5H, Ph-H), 7.70-7.90 (m, 4H, Ar-H); ¹³C NMR 21.8 (C-5), 26.7 (C-4), 43.1 (C-6), 49.9 (C-3), 58.9 (C-α), 61.1 (C-β), 123.4 (Phth-α), 127.6 (Ph-o), 127.8 (Ph-p), 128.6 (Ph-m), 131.9 (Phth-quaternary), 134.0 (Phth-β), 136.5 (C-*ipso*), 167.4 (CO), 167.8 (CO); EIMS *m/z* (%) 365 (M⁺+1, 1), 333 (96), 305 (75), 200 (87), 182 (41), 159 (100). Anal. Calcd for C₂₁H₂₀N₂O₄: C, 69.23; H, 5.49; N, 7.69. Found: C, 69.21; H, 5.51; N, 7.61.

(*αR,3R*)-3-Amino-*N*-(2-hydroxy-1-phenylethyl)piperidin-2-one (**8**). **Procedure A:** To a solution of azide **6** (100 mg, 0.38 mmol) in THF (5 ml), PPh₃ (121 mg, 0.46 mmol) was added. The resulting mixture was stirred for 6 h at room temperature, H₂O (1.5 equivalents) was added, and the solution was stirred at room temperature overnight. The solvent was evaporated and the residue, dissolved in CH₂Cl₂, was washed with ice-cold 2N HCl. The acid aqueous solution was washed with CH₂Cl₂, basified with NaHCO₃, and extracted with CH₂Cl₂. The organic extracts, dried and evaporated, yielded pure 3-aminopiperidone **8** (15 mg, 16%): [α]_D = -72.1 (c = 1, CHCl₃); IR (NaCl) 3350 (OH), 3349 and 3302 (NH₂), 1635 (CO) cm⁻¹; ¹H NMR 1.60 (m, 2H, 5-H), 1.86 (m, 1H, 4-H), 2.14 (m, 1H, 4-H), 2.30 (br s, 3H, NH₂ and OH), 2.93 (dt, *J* = 12 and 3.5 Hz, 1H, 6-H_e), 3.19 (ddd, *J* = 12, 7 and 2 Hz, 1H, 6-H_a), 3.40 (br dd, *J* = 12 and 6 Hz, 1H, 3-H_a), 4.04 (dd, *J* = 12 and 9 Hz, 1H, β-H_A), 4.10 (dd, *J* = 12 and 5 Hz, 1H, β-H_B), 5.73 (dd, *J* = 9 and 5 Hz, 1H, α-H), 7.20-7.40 (m, 5H, Ph-H); ¹³C NMR 21.2 (C-5), 29.4 (C-4), 42.9 (C-6), 52.2 (C-3), 58.0 (C-α), 60.8 (C-β), 127.5 (Ph-o), 127.7 (Ph-p), 128.6 (Ph-m), 137.0 (C-*ipso*), 174.1 (CO); EIMS *m/z* (%) 235 (M⁺+1, 3), 203 (42), 175 (48), 106 (32), 77 (21), 70 (100). Anal. Calcd for C₁₃H₁₈N₂O₂: C, 66.66; H, 7.69; N, 11.96. Found: C, 66.54; H, 7.65; N, 11.91.

Procedure B: To a solution of phthalimide **7** (121 mg, 0.35 mmol) in MeOH (6 ml), NH₂NH₂·H₂O (0.22 ml, 3.54 mmol) was added. The resulting mixture was refluxed for 1 h. The solvent was evaporated and the residue, dissolved in CH₂Cl₂, was washed with 2.6N KOH. The layers were separated and the aqueous phase was extracted with CH₂Cl₂. The organic extracts were dried and evaporated to yield pure amine **8** (55 mg, 70%).

(*αR*)-*N*-(2-Acetoxy-1-phenylethyl)-Δ⁵-piperidin-2-one (**2**). To a solution of imide **3**⁹ (1 g, 3.64 mmol) in dry THF (50 ml) cooled at -78°C, LiBHET₃ (5.5 ml, 5.46 mmol) was added. After stirring for 2.5 h, the reaction was quenched with aqueous NaHCO₃. The solvent was evaporated and the residue, dissolved in dry CH₂Cl₂, was treated with TFA (3 ml) for 1.5 h at room temperature and the reaction was quenched with aqueous NaHCO₃. The resulting mixture was diluted with CH₂Cl₂ and washed with brine. The organic extracts were dried and evaporated to give an oil, which after flash chromatography (AcOEt:hexane, 1:1)

yielded *N*-(2-acetoxy-1-phenylethyl)-6-hydroxypiperidin-2-ones⁹ **9** (150 mg, 15%) and enamide **2** (650 mg, 69%): $[\alpha]_D = -91.8$ ($c = 1$, CHCl_3); IR (NaCl) 1745 (CO), 1675 (CO), 1600 (C=C) cm^{-1} ; $^1\text{H NMR}$ 2.05 (s, 3H, CH_3), 2.30 (m, 2H, 4-H), 2.58 (dd, $J = 8$ and 1.5 Hz, 1H, 3-H), 2.61 (dd, $J = 8$ and 2.5 Hz, 1H, 3-H), 4.51 (dd, $J = 12$ and 8 Hz, 1H, $\beta\text{-H}_A$), 4.65 (dd, $J = 12$ and 5 Hz, 1H, $\beta\text{-H}_B$), 5.15 (dt, $J = 8$ and 4 Hz, 1H, 5-H), 5.95 (d, $J = 8$ Hz, 1H, 6-H), 6.12 (dd, $J = 8$ and 5 Hz, 1H, $\alpha\text{-H}$), 7.20-7.40 (m, 5H, Ph-H); $^{13}\text{C NMR}$ 19.7 (C-4), 20.7 (CH_3CO), 31.5 (C-3), 52.7 (C- α), 62.6 (C- β), 106.8 (C-5), 125.7 (C-6), 127.3 (Ph-*o*), 127.8 (Ph-*p*), 128.6 (Ph-*m*), 136.2 (C-*ipso*), 169.5 (CON), 170.5 (COO); EIMS m/z (%) 259 (M^+ , 10), 215 (49), 199 (67), 186 (84), 159 (89), 91 (100). Anal. Calcd for $\text{C}_{15}\text{H}_{17}\text{NO}_3$: C, 69.49; H, 6.56; N, 5.40. Found: C, 69.40; H, 6.55; N, 5.51.

($\alpha R, 5RS$)-*N*-(2-Hydroxy-1-phenylethyl)-5-methylthiopiperidin-2-ones (10a,b). To a solution of enamide **2** (1.7 g, 6.57 mmol) in dry THF (20 ml) cooled at -78°C , CH_3SH (4.3 ml, 73.4 mmol) and catalytic AIBN were added. The solution was stirred for 30 min at -78°C and at room temperature overnight. The solvent was evaporated and the residue was filtered through SiO_2 to yield the epimeric mixture of the acetates of **10** (1.26 g). To a solution of the epimeric mixture in dry THF (45 ml) 5% aqueous KOH (10 ml) was added, and the mixture was refluxed for 1.5 h and neutralized with 3N HCl. The solvent was evaporated and the residue, dissolved in CH_2Cl_2 , was washed with aqueous NaHCO_3 and with brine. The organic extracts were dried and evaporated to yield, after flash chromatography (AcOEt; AcOEt:MeOH 1:1), a 1:1.2 mixture of mercaptanes **10a** and **10b**. (**$\alpha R, 5R$**)-**10a** (477 mg, 27%): m.p. 54.1-54.3 $^\circ\text{C}$ (*i*PrOH), $[\alpha]_D = -25.3$ ($c = 1$, CHCl_3); IR (NaCl) 3375 (OH), 1623 (CO), 1440 (CS) cm^{-1} ; $^1\text{H NMR}$ 1.70 (m, 1H, 4- H_a), 1.96 (s, 3H, SMe), 2.11 (m, 1H, 4- H_e), 2.51 (ddd, $J = 17, 9$ and 7 Hz, 1H, 3- H_a), 2.68 (ddd, $J = 17, 7$ and 5.5 Hz, 1H, 3- H_e), 2.84 (m, 1H, 5-H), 3.11 (dd, $J = 11$ and 7 Hz, 1H, 6- H_a), 3.20 (dd, $J = 11$ and 4 Hz, 1H, 6- H_e), 4.08 (dd, $J = 10$ and 9 Hz, 1H, $\beta\text{-H}_A$), 4.15 (dd, $J = 10$ and 5 Hz, 1H, $\beta\text{-H}_B$), 5.87 (dd, $J = 9$ and 5 Hz, 1H, $\alpha\text{-H}$), 7.20-7.40 (m, 5H, Ph-H); $^{13}\text{C NMR}$ 13.8 (SMe), 20.8 (C-4), 30.9 (C-3), 40.5 (C-5), 47.3 (C-6), 57.8 (C- α), 61.1 (C- β), 127.6 (Ph-*o*), 127.8 (Ph-*p*), 128.6 (Ph-*m*), 136.6 (C-*ipso*), 170.7 (CO); EIMS m/z (%) 265 (M^+ , 1), 247 (20), 234 (100), 201 (15), 186 (18). Anal. Calcd for $\text{C}_{14}\text{H}_{19}\text{NSO}_2$: C, 63.37; H, 7.22; N, 5.28; S, 12.08. Found: C, 63.40; H, 7.24; N, 5.21; S, 11.90.

($\alpha R, 5S$)-10b (583 mg, 33%): m.p. 95.2-95.4 $^\circ\text{C}$ (*i*PrOH:pentane, 9:1), $[\alpha]_D = -43.8$ ($c = 1$, CHCl_3); IR (NaCl) 3420 (OH), 1614 (CO), 1435 (CS) cm^{-1} ; $^1\text{H NMR}$ 1.80 (m, 1H, 4- H_a), 1.90 (s, 3H, SMe), 2.23 (m, 1H, 4- H_e), 2.50 (dt, $J = 17$ and 7 Hz, 1H, 3- H_a), 2.66 (dt, $J = 17$ and 7 Hz, 1H, 3- H_e), 2.88 (dd, $J = 12$ and 7 Hz, 1H, 6- H_a), 3.01 (m, 1H, 5-H), 3.47 (dd, $J = 12$ and 4 Hz, 1H, 6- H_e), 4.06 (dd, $J = 12$ and 9 Hz, 1H, $\beta\text{-H}_A$), 4.13 (dd, $J = 12$ and 5 Hz, 1H, $\beta\text{-H}_B$), 5.90 (dd, $J = 9$ and 5 Hz, 1H, $\alpha\text{-H}$), 7.20-7.40 (m, 5H, Ph-H); $^{13}\text{C NMR}$ 13.6 (SMe), 26.5 (C-4), 30.5 (C-3), 40.1 (C-5), 46.8 (C-6), 57.5 (C- α), 60.8 (C- β), 127.7 (Ph-*o*), 128.5 (Ph-*p*), 128.9 (Ph-*m*), 136.4 (C-*ipso*), 170.7 (CO); EIMS m/z (%) 265 (M^+ , 1), 247 (25), 234 (100), 201 (20), 186 (15). Anal. Calcd for $\text{C}_{14}\text{H}_{19}\text{NO}_2\text{S}$: C, 63.37; H, 7.22; N, 5.28; S, 12.08. Found: C, 63.40; H, 7.24; N, 5.21; S, 11.90.

($\alpha R, 3RS, 5R$)-3-Bromo-*N*-(2-hydroxy-1-phenylethyl)-5-methylthiopiperidin-2-ones (11a,b). Operating as for the preparation of compounds **5**, from thioether **10b** (550 mg, 2.08 mmol), dry THF (15 ml), *sec*-BuLi (4 ml, 5.19 mmol), and Br_2 (0.13 mL, 2.29 mmol) a mixture of **11a** and **11b** was obtained, which was flash chromatographed (AcOEt-hexane, 2:1). (**$\alpha R, 3S, 5S$**)-**11a** (220 mg, 31%): $[\alpha]_D = -21.7$ ($c = 1.25$, CHCl_3); IR

(NaCl) 3405 (OH), 1648 (CO), 1437 (C-S) cm^{-1} ; ^1H NMR (500 MHz) 1.94 (s, 3H, SMe), 2.20 (ddd, $J = 14, 10$ and 5 Hz, 1H, 4- H_a), 2.45 (dt, $J = 14$ and 4 Hz, 1H, 4- H_e), 2.89 (dd, $J = 12$ and 10 Hz, 1H, 6- H_a), 3.32 (m, 1H, 5-H), 3.52 (dd, $J = 12$ and 5 Hz, 1H, 6- H_e), 4.09 (m, 2H, β -H), 4.64 (t, $J = 5$ Hz, 1H, 3-H), 5.57 (dd, $J = 8$ and 5 Hz, 1H, α -H), 7.20-7.40 (m, 5H, Ph-H); ^{13}C NMR 12.6 (SMe), 35.8 (C-5), 36.1 (C-4), 43.4 (C-3), 47.1 (C-6), 58.4 (C- α), 60.1 (C- β), 126.9 (Ph- o), 127.1 (Ph- p), 127.8 (Ph- m), 134.8 (C- $ipso$), 166.0 (CO); EIMS m/z (%) 345 ($\text{M}^+ + 1$, 1), 343 ($\text{M}^+ - 1$, 1), 314 and 312 (100), 268 (10), 246 (70), 234 (60), 91 (90). Anal. Calcd for $\text{C}_{14}\text{H}_{18}\text{NSBrO}_2$: C, 48.98; H, 5.29; N, 4.08; Br, 23.01. Found: C, 48.78; H, 5.27; N, 4.12; Br, 22.97.

($\alpha\text{R},3\text{R},5\text{S}$)-**11b** (100 mg, 14%): IR (NaCl) 3395 (OH), 1649 (CO), 1450 (C-S) cm^{-1} ; ^1H NMR 1.97 (s, 3H, SMe), 2.15 (m, 1H, 4-H), 2.28 (m, 1H, 4-H), 2.95-3.10 (m, 2H, 6- H_a and 5-H), 3.40 (dd, $J = 12$ and 4 Hz, 1H, 6- H_e), 4.00-4.20 (m, 2H, β -H), 4.59 (dd, $J = 8.5$ and 7 Hz, 1H, 3-H), 5.73 (dd, $J = 9$ and 5 Hz, 1H, α -H), 7.20-7.40 (m, 5H, Ph-H); ^{13}C NMR 13.6 (SMe), 38.5 (C-4), 39.1 (C-5), 44.4 (C-3), 47.9 (C-6), 59.3 (C- α), 60.9 (C- β), 126.9 (Ph- o), 128.1 (Ph- p), 128.7 (Ph- m), 135.8 (C- $ipso$), 167.4 (CO); EIMS m/z (%) 345 ($\text{M}^+ + 1$, 1), 343 ($\text{M}^+ - 1$, 1), 314 and 312 (100), 246 (70), 234 (45), 91 (90).

($\alpha\text{R},3\text{R},5\text{S}$)-**3-Azido-*N*-(2-hydroxy-1-phenylethyl)-5-methylthiopiperidin-2-one (12)**. Operating as for the preparation of compound **6**, from bromide **11a** (70 mg, 0.20 mmol) in DMF-AcOH (6:4, 2.1 ml), NaN_3 (26 mg, 0.40 mmol), and H_2O (0.2 ml), azidolactam **12** (43 mg, 70%) was obtained, after flash chromatography (SiO_2 , AcOEt-hexane, 1:1): $[\alpha]_D = -57.4$ ($c = 1$, CHCl_3); IR (NaCl) 3395 (OH), 2114 (N_3), 1649 (CO), 1445 (C-S) cm^{-1} ; ^1H NMR 1.65 (td, $J = 12$ and 11 Hz, 1H, 4- H_a), 1.99 (s, 3H, SMe), 2.42 (m, 1H, 4- H_e), 2.83 (dd, $J = 12$ and 10 Hz, 1H, 6- H_a), 3.00 (ddt, $J = 12, 11$ and 5 Hz, 1H, 5- H_a), 3.42 (dd, $J = 12$ and 5 Hz, 1H, 6- H_e), 4.15 (d, $J = 7$ Hz, 2H, β -H), 4.19 (dd, $J = 11$ and 7 Hz, 1H, 3- H_a), 5.80 (t, $J = 7$ Hz, 1H, α -H), 7.20-7.40 (m, 5H, Ph-H); ^{13}C NMR 13.5 (SMe), 33.6 (C-4), 37.6 (C-5), 47.5 (C-6), 58.6 (C-3), 58.8 (C- α), 60.9 (C- β), 127.9 (Ph- o), 128.2 (Ph- p), 128.8 (Ph- m), 135.4 (C- $ipso$), 168.9 (CO); EIMS m/z (%) 306 (M^+ , 1), 275 (75), 246 (34), 198 (12), 91 (100). Anal. Calcd for $\text{C}_{14}\text{H}_{18}\text{N}_4\text{SO}_2$: C, 54.88; H, 5.93; N, 18.30. Found: C, 54.79; H, 5.88; N, 18.29.

($\alpha\text{R},3\text{S},5\text{S}$)-***N*-(2-Hydroxy-1-phenylethyl)-5-methylthio-3-phthalimidopiperidin-2-one (13)**. Operating as for the preparation of compound **7**, from bromides **11a** (150 mg, 0.44 mmol) and potassium phthalimide (122 mg, 0.66 mmol) in DMF (3 ml), and after 24 h of reaction, a 1:5 diastereomeric mixture of phthalimidolactams **13a,b** was obtained, from which only isomer **13b** was isolated by flash chromatographed (AcOEt-hexane, 2:1). Compound **13b** (115 mg, 64%) m.p. 199-199.7°C (AcOEt); $[\alpha]_D = -11.5$ ($c = 2$, CHCl_3); IR (NaCl) 3430 (OH), 1717 (CO), 1660 (CO), 1390 (C-S) cm^{-1} ; ^1H NMR 1.84 (s, 3H, SMe), 2.29 (m, 1H, 4- H_e), 2.58 (ddd, $J = 13, 10$ and 4 Hz, 1H, 4- H_a), 3.12 (br d, $J = 13$ Hz, 1H, 6- H_a), 3.28 (m, 1H, 5- H_e), 3.75 (dd, $J = 13$ and 3 Hz, 1H, 6- H_e), 4.23 (m, 2H, β -H), 5.17 (dd, $J = 11$ and 8 Hz, 1H, 3- H_a), 5.98 (dd, $J = 8$ and 5 Hz, 1H, α -H), 7.25-7.40 (m, 5H, Ph-H), 7.70-7.75 and 7.85-7.90 (2 m, 2H each, Phth-H); ^{13}C NMR 13.9 (SMe), 31.2 (C-4), 39.3 (C-5), 45.9 (C-6), 47.1 (C-3), 58.1 (C- α), 60.9 (C- β), 123.5 (Phth- α), 127.9 (Ph- o and Ph- p), 128.6 (Ph- m), 131.8 (Phth-quaternary), 134.1 (Phth- β), 136.1 (C- $ipso$), 166.7 (CO), 167.7 (CO); EIMS m/z (%) 411 ($\text{M}^+ + 1$, 1), 379 (100), 333 (39), 246 (61), 179 (35), 156 (56). Anal. Calcd for $\text{C}_{22}\text{H}_{22}\text{N}_2\text{O}_4\text{S}$: C, 64.37; H, 5.40; N, 6.82; S, 7.81. Found: C, 64.25; H, 5.36; N, 6.76; S, 7.64.

($\alpha R, 3R, 5S$)-3-Amino-*N*-(2-hydroxy-1-phenylethyl)-5-methylthiopiperidin-2-one (1a). Operating as for the preparation of compound **8**, from azide **12** (80 mg, 0.26 mmol) and PPh₃ (68 mg, 0.26 mmol) in THF (3 ml), and after a preparative chromatography (SiO₂, CH₂Cl₂, MeOH, 95:5), 3-aminolactam **1** (22 mg, 30%) was obtained: $[\alpha]_D = -41.7$ ($c = 0.5$, CHCl₃); IR (NaCl) 3356 (OH and NH₂), 1629 (CO), 1463 (C-S) cm⁻¹; ¹H NMR 1.55 (q, $J = 11$ Hz, 1H, 4-H_A), 1.95 (s, 3H, SMe), 2.45 (m, 1H, 4-H_E), 2.75 (br s, 3H, NH₂ and OH), 2.90 (dd, $J = 12$ and 11 Hz, 1H, 6-H_A), 3.03 (m, 1H, 5-H_A), 3.45 (dd, $J = 12$ and 6 Hz, 1H, 6-H_E), 3.50 (dd, $J = 10$ and 7 Hz, 1H, 3-H_A), 4.05 (m, 2H, β -H), 5.70 (dd, $J = 9$ and 5 Hz, 1H, α -H), 7.20-7.40 (m, 5H, Ar-H); ¹³C NMR 13.4 (SMe), 35.8 (C-4), 38.2 (C-5), 47.9 (C-6), 51.7 (C-3), 58.9 (C- α), 60.9 (C- β), 127.6 (Ph-*p*), 128.0 (Ph-*o*), 128.7 (Ph-*m*), 136.1 (C-*ipso*), 173.6 (CO); EIMS m/z (%) 280 (M⁺, 4), 249 (22), 233 (27), 173 (44), 159 (100). Anal. Calcd for C₁₄H₂₀N₂SO₂: C, 59.98; H, 7.20; N, 10.00. Found: C, 60.08; H, 7.21; N, 9.85.

($\alpha R, 3S, 5S$)-3-Amino-*N*-(2-hydroxy-1-phenylethyl)-5-methylthiopiperidin-2-one (1b). Operating as for the preparation of compound **8**, from phthalimide **13** (50 mg, 0.12 mmol) and NH₂NH₂·H₂O (0.1 ml, 1.2 mmol) in MeOH (1 ml), amine **1b** (25 mg, 73%) was obtained: $[\alpha]_D = -25.7$ ($c = 1$, CHCl₃); ¹H NMR 1.86 (s, 3H, SMe), 2.01 (ddd, $J = 13, 10$ and 4 Hz, 1H, 4-H_A), 2.21 (dt, $J = 13$ and 7 Hz, 1H, 4-H_E), 2.41 (br s, 3H, NH₂ and OH), 2.96 (dd, $J = 12.5$ and 6 Hz, 1H, 6-H_A), 3.13 (m, 1H, 5-H), 3.50 (dd, $J = 12.5$ and 4 Hz, 1H, 6-H_E), 3.69 (dd, $J = 10$ and 7 Hz, 1H, 3-H), 4.05 (dd, $J = 11$ and 9 Hz, 1H, β -H_A), 4.15 (dd, $J = 11$ and 5 Hz, 1H, β -H_B), 5.82 (dd, $J = 9$ and 5 Hz, 1H, α -H), 7.20-7.40 (m, 5H, Ph-H); ¹³C NMR 13.9 (SMe), 34.8 (C-4), 38.8 (C-5), 46.4 (C-6), 49.5 (C-3), 50.8 (C- α), 61.1 (C- β), 127.7 (Ph-*o*), 127.9 (Ph-*p*), 128.7 (Ph-*m*), 136.5 (C-*ipso*), and 173.8 (CO); EIMS m/z (%) 280 (M⁺, 1), 249(18), 233 (22), 173 (51), 159 (100). Anal. Calcd for C₁₄H₂₀N₂SO₂: C, 59.98; H, 7.20; N, 10.00; S, 22.83. Found: C, 60.08; H, 7.21; N, 9.85; S, 22.98.

ACKNOWLEDGEMENTS

Support for this research has been provided by the CIRIT (Generalitat de Catalunya) through grants QFN92 4303, QFN95-4703, and by the Ministerio de Educación y Ciencia through grant HF96 312B.

REFERENCES AND NOTES

1. a) Fernández-Checa, J.C.; Yi, J.R.; García-Ruiz, C.; Knezic, Z.; Tahara, S.M.; Kaplowitz N. *J. Biol. Chem.*, **1993**, 268, 2324-2328; b) Fernández-Checa, J.C.; Maddatu, T.; Ookhtens, M.; Kaplowitz, N. *Am. J. Physiol.*, **1990**, 258, G967-G973; c) AW, T.Y.; Ookhtens, M.; Kaplowitz, N. *Am. J. Physiol.*, **1986**, 250, G354-G361; d) Aw, T.Y.; Ookhtens, M.; Kaplowitz, N. *J. Biol. Chem.*, **1984**, 259, 9355-9358.
2. a) Fernández-Checa, J.C.; García-Ruiz, C.; Colell, A.; Yi, J.-R.; Kaplowitz, N. *Am. J. Physiol.*, **1996**, in press; b) Yi, J.R., Lu, S.; Fernández-Checa, J.C.; Kaplowitz, N. *Proc. Natl. Acad. Sci. USA*, **1995**, 92, 1425-1429; c) Yi, J.-R.; Lu, S.; Fernández-Checa, J.C.; Kaplowitz, N. *J. Clin. Invest.*, **1994**, 93, 1841-1845; d) Beatty, P.W.; Reed, D.J. *Arch. Biochem. Biophys.*, **1980**, 204, 80-87.
3. Hepatic GSH is transported by putative plasma membrane carriers, localized in the sinusoidal (basolateral) and canalicular (apical) membrane domains, responsible for the release of GSH into blood circulation and bile, respectively. We have recently cloned both carriers: see references 2b-c.

4. a) Rodríguez, R.; Diez, A.; Rubiralta, M.; Giralt, E., "XIIIth International Symposium on Medicinal Chemistry", Paris, 1994; b) Rodríguez, R.; Diez, A.; Rubiralta, M.; Giralt, E. *Heterocycles*, **1996**, *43*, 000.
5. Kaplowitz, N.; Aw, T.Y.; Ookhtens, M. *Ann. Rev. Pharmacol. Toxicol.*, **1985**, *25*, 715-744.
6. a) Fernández-Checa, J.C.; Yi, J-R.; García-Ruiz, C.; Ookhtens, M.; Kaplowitz, N. *Sem. Liv. Dis.*, **1996**, in press; b) Meister, A.; Anderson, M.E. *Ann. Rev. Biochem.*, **1983**, *52*, 711-760.
7. Micouin, L.; Varea, T.; Riche, C.; Chiaroni, A.; Quirion, J.-C.; Husson, H.-P. *Tetrahedron Lett.*, **1994**, *35*, 2529-2532.
8. For the pharmacological interest of 3-aminolactams, see: a) Shuman, R.T.; Rothenberg, R.B.; Campbell, C.S.; Smith, G.F.; Gifford-Moore, D.S.; Gesellchen, P.D. *J. Med. Chem.*, **1993**, *36*, 314-319; b) Balasubramanian, N.; St. Laurent, D.R.; Federici, M.E.; Meanwell, N.A.; Wnight, J.J.; Schumacher, W.A.; Seiler, S.M. *J. Med. Chem.*, **1993**, *36*, 300-303; c) Nagai, U.; Sato, K.; Nakamura, R.; Kato, R. *Tetrahedron*, **1993**, *49*, 3577-3592; d) Sreenivasan, U.; Mishra, R.K.; Johnson, R.L. *J. Med. Chem.*, **1993**, *36*, 256-263.
9. In this case, compound **4** was prepared from methyl (*R*)-(-)-phenylglycinate and glutaric anhydride according to: Royer, J.; Husson, H.-P. *Heterocycles* **1993**, *36*, 1493-1496.
10. Micouin, L.; Bonin, M.; Cherrier, M.-P.; Mazurier, A.; Thomas, A.; Quirion, J.-C.; Husson, H.-P. *Tetrahedron*, previous paper in this issue.
11. A low level of epimerisation of isomer **5a** in a CH₂Cl₂ solution was observed only after 15 h.
12. a) Rubiralta, M.; Diez, A.; Miguel, D.; Remuson, R.; Gelas-Mialhe, Y. *Synth. Commun.*, **1992**, *22*, 359-367; b) Vaultier, M.; Knouzi, N.; Carrié, R. *Tetrahedron Lett.*, **1983**, *24*, 763-764.
13. Takacs, J.M.; Weidner, J.J. *J. Org. Chem.*, **1994**, *59*, 6480-6483.
14. Lenz, G.R. *Synthesis*, **1978**, 489-518.
15. a) Griesbaum, K. *Angew. Chem. Int. Ed.*, **1970**, *9*, 273-287; b) LeBel, N.A.; DeBoer, A.. *J. Am. Chem. Soc.*, **1967**, *89*, 2784-2785; c) Skell, P.S.; Allen, R.G. *J. Am. Chem. Soc.*, **1960**, *82*, 1511-1512; d) Walling, C.; Helmreich, W. *J. Am. Chem. Soc.*, **1959**, *81*, 1144-1148.

(Received in UK 2 February 1996; revised 26 March 1996; accepted 2 April 1996)