



New Cytotoxic Sesterterpenes From The New Caledonian Marine Sponge *Petrosaspongia nigra* (Bergquist).

Luigi Gomez Paloma^{*}, Antonio Randazzo and Luigi Minale

Dipartimento di Chimica delle Sostanze Naturali, Università degli Studi di Napoli Federico II
via D. Montesano 49, 80131 Napoli, Italy

Cécile Debitus

ORSTOM, Centre de Nouméa, B.P. A5 Nouméa Cedex, New Caledonia

Christos Roussakis

Faculté de Pharmacie, 1 Rue Gaston Veil 44035 Nantes Cedex 01, France

Abstract: Along with two known cheilanthane sesterterpene lactones, **1** and **2**, eight new related sesterterpenes (**3-10**) and two new nor-sesterterpenes (**11** and **12**) have been isolated from the New Caledonian marine sponge *Petrosaspongia nigra* Bergquist 1995 (new genus, new species). Their structures were determined from 1D and 2D NMR studies and mass spectral data. They exhibited cytotoxicity against the NSCLC-N6 human bronchopulmonary non-small-cell-lung carcinoma cell lines.
© 1997 Elsevier Science Ltd.

Marine sponges have provided the largest number of sesterterpenes, a relatively rare group of pentaprenyl terpenoids of widespread distribution, i.e. fungi, lichens, higher plants, insects and various marine organisms, whose structures are derivable from geranyl-farnesyl diphosphate¹⁻³. Sesterterpenes from Dictyoceratic sponges include the linear C-25 furans and their C-21 degradation products, the group of γ -hydroxybutenolides, potent anti-inflammatory agents exemplified by the monocarbocyclic manoalide⁴, the bicarbocyclic cacospongiolide⁵ and the tricarbocyclic luffolide⁶, that are inhibitors of the enzyme phospholipase A₂ (PLA₂)⁷, and the most common tetracarbo-cyclic scalarane sesterterpenoids that also exhibit a variety of biological activities, including anti-inflammatory.

Recently, two new cheilanthane sesterterpene lactones **1** and **2**⁹ (the structure of **1** was determined by X-ray analysis¹⁰) were isolated from a sponge incorrectly assigned to the genus *Dactylospongia*⁹ and then reassigned as a new genus and new species: *Petrosaspongia nigra* (Bergquist 1995)¹⁰.


In our continuing search for cytotoxic substances from New Caledonian marine organisms¹¹, we found potent activity in the dichloromethane extract of the sponge *Petrosaspongia nigra* (Bergquist 1995 sp.nov., Dictyoceraida, Spongidae) collected off New Caledonia, from which we isolated **1** and **2**, eight new related sesterterpenes (**3-10**) and two new nor-sesterterpenes (**11** and **12**). Herein we report their isolation, structure elucidation and cytotoxicity data.

Freeze dried specimens (2.7 Kg, 12 Kg fresh), collected off the southern coral reef of New Caledonia were sequentially extracted with *n*-hexane and dichloromethane in a Soxhlet apparatus and then with methanol at room temperature. The dichloromethane soluble material was fractionated by silica gel flash chromatography (MPLC) eluting with CHCl₃/MeOH mixtures with increasing amounts of MeOH, followed by reverse phase HPLC to yield **1-12**, which we named petrosaspongiolides A-L (yields: 10.1mg, 15.0mg, 3.3mg, 2.5mg, 3.4mg, 2.8mg, 2.5mg, 5.4mg, 0.9mg, 3.1mg, 5.0mg, 2.0mg, respectively).

Petrosaspongiolides A (**1**) and B (**2**) (known compounds): structure analysis was carried out mostly with **2**, *m/z* 460 (M⁺) and 191 (C₁₄H₂₃). The ¹H-NMR spectrum (CDCl₃, 500 MHz) exhibited signals for five methyl

singlets (δ 0.75, 0.77, 0.80 and 0.85 ppm) including one at δ 2.18 ppm for an acetate, and three oxygenated methines (δ 5.70 s, 5.48 br s, and 5.15 dd). The ^{13}C -NMR spectrum (CDCl_3 , 125 MHz) revealed 27 carbons including signals for an acetate unit (δ 170.1 and 21.1 ppm), an ester carbonyl at δ 171.1 ppm and three oxygenated methines, including two acetals, δ 114.1 and 102.8 ppm. Inspection of literature indicated that the same structural units were present in the structure of a sesterterpene lactone, isolated from a sponge of the genus *Dactylosporgia*⁹. Identification of the major compound from the *Petrosaspongia nigra* was confirmed by comparison of ^1H - and ^{13}C -NMR and mass spectral data with those reported for **2**⁹.


The identification of the isomeric second major compound, m/z 460 (M^+) and 191 ($\text{C}_{14}\text{H}_{23}$), with **1** followed in a straightforward manner. The most significant differences observed in the ^1H -NMR data of **1** and **2** were for protons of H25, shielded by *ca.* 0.5 ppm in **2** (δ 5.70 ppm in **2** and 6.21 ppm in **1**), and of H24 which appeared as a singlet in **2** and as a doublet ($J=2$ Hz) coupled to H13 in **1**.



R'


R''

1	CH ₃	OAc
3	CH ₂ OAc	OAc
4	CH ₂ OH	OAc
5	CHO	OAc
6	COOH	OAc
7	CH ₂ OH	OH
8	COOH	OH



R

2	CH ₃
9	CH ₂ OAc

**10**

Petrosaspongiolide C (**3**) had a molecular ion (EIMS) at m/z 518, 58 mass units more than the known **1** and **2**. ^1H -NMR (CDCl_3 , 500 MHz) and ^{13}C -NMR (CDCl_3 , 125 MHz) data were very similar to those of **1**, except for the lack of a methyl group replaced by an acetoxymethyl group in **3**, δ_{H} 4.19 -3.87 (AB system, $J=12.0$ Hz, CH₂) and 2.01s (CH₃), δ_{C} 66.9 (CH₂), 21.0 (CH₃), 171.3 ppm (CO) (Table 1). Significant shifts of C3, C4 and C20 (δ_{C} 36.2, 37.0 and 27.3 ppm) in **3** relative to **2** (δ_{C} 42.0, 33.3 and 33.3 ppm) suggested the location of acetoxyl group at C21¹². This assignment was substantiated by ROESY (mixing time 500ms; CDCl_3) data which indicated that H₂21 was axial, because an intense cross-peak was observed between H₂21 and Me22 at δ_{H} 0.77 ppm. Further ROESY correlations between H14 (δ_{H} 1.01 ppm) and H25 (δ_{H} 6.18 ppm) and between H24 (δ_{H} 5.58 ppm), H13 (δ_{H} 1.54 ppm) and H12 β (δ_{H} 1.66 ppm) were consistent with an α orientation for H25 and a β orientation for H24. The signals for H14 which appeared as a triplet with a large coupling constant ($J=$

10.0 Hz) confirmed that the C/D ring junction was *trans*. Thus, petrosaspongiolide C (**3**) is the 21-acetoxy derivative of petrosaspongiolide A (**1**).

Petrosaspongiolide D (**4**), *m/z* 476, 16 mass units more than **1** and **2**, again showed ^1H (CDCl_3 , 500 MHz) and ^{13}C -NMR (CDCl_3 , 125 MHz) spectra very similar to those of **1**, except for the presence of additional primary hydroxyl group in **4**, δ_{H} 3.72 and 3.45 (AB system, $J=12.0$ Hz, CH_2). A C21 hydroxyl group followed unambiguously from NMR data and was further supported by the absence in **4** of the characteristic high-field methyl carbon signal observed at δ_{C} 21.3 ppm in **1** and replaced by an oxygenated methylene carbon resonating at δ_{C} 65.4 ppm (Table 1).

Petrosaspongiolide E (**5**). NMR spectra (CDCl_3 , 500 MHz) showed low field resonances at δ_{H} 9.70 (s) ppm and δ_{C} 205.5 ppm assigned to a formyl group. Again, comparison of spectral data makes clear that compound **5**, *m/z* 474 (14 mass units more than **1**) is related to the known **1** by oxydation of the Me21 to an aldehyde.

Petrosaspongiolide F (**6**), *m/z* 490 (EIMS), is related to **1** by oxydation of C21 methyl to a carboxyl group, followed unambiguously from NMR data (CDCl_3 , 500 MHz). The only significant differences in the data were for signals corresponding to ring A (Table 1), such as the replacement of methyl carbon signal (δ 21.4 ppm in **1**) by a carbonyl signal at δ 172.0 ppm, accompanied by the downfield shift of C4, δ_{C} 44.9 vs. 33.3 ppm in **1**, and by the highfield shift of C3 and C20, δ_{C} 38.8 and 28.9 vs. 41.9 and 33.3 ppm in **1**.

Petrosaspongiolide G (**7**), *m/z* 434 (EIMS). ^{13}C -NMR (125 MHz, CD_3OD) data of **7** were very similar to those of **4** except for the absence of an acetyl group in **7**, thus indicating that **7** was the deacetyl derivative of **4**. This was substantiated by the significant high field shift of H16 (500 MHz, CD_3OD) (δ_{H} 4.25 vs. 5.42 ppm in **1**) and downfield shifts of C15 (δ_{C} 34.1 vs. 30.1 ppm in **1**) and C17 (δ_{C} 90.4 vs. 86.9 ppm in **1**).

Petrosaspongiolide H (**8**), *m/z* 448 (EIMS), is the more polar compound of this series. The ^1H -NMR spectra (500 MHz, CD_3OD) readily implied that compound **8** is the 16-deacetyl derivative of **6**. Again the only significant differences in the NMR data of **8** relative to those of **6** were for signals corresponding to H16 (δ_{H} 4.29 ppm) and carbons C15 and C17 (125 MHz, CD_3OD) (δ_{C} 34.5 and 90.2 ppm, respectively).

Table 1. ^{13}C -NMR chemical shifts (ppm) of ring A of compounds **1,3-8**

	1 (CDCl_3)	3 (CDCl_3)	4 (CDCl_3)	5 (CDCl_3)	6 (CDCl_3)	7 (CD_3OD)	8 (CD_3OD)
C1	39.9	39.8	40.0	39.2	40.2	41.5	41.9
C2	18.5	18.1	18.2	18.1	19.1	19.4	20.8
C3	41.9	36.2	35.6	34.4	38.8	36.7	40.1
C4	33.3	37.0	38.6	48.3	44.9	38.4	38.1
C5	56.2	56.8	56.8	56.6	56.8	58.5	58.8
C10	37.5	37.4	37.4	37.8	38.0	39.8	38.1
C20	33.3	27.3	26.8	24.1	28.9	27.7	29.9
C21	21.4	66.9	65.4	205.6	172.0	65.2	174.7

Petrosaspongiolide I (**9**), *m/z* 518 (EIMS), is an isomer of **3**. The ^1H (CDCl_3 , 500 MHz) and ^{13}C -NMR (CDCl_3 , 125 MHz) spectra was very similar with **2** except for the presence of an additional acetoxyl group, which was unambiguously located at C21, as followed from the comparison of the proton and carbon signals corresponding to ring A with the same signals in **3**.

Petrosaspongiolide J (**10**). The ^1H (500 MHz, CDCl_3) and ^{13}C -NMR (125 MHz, CDCl_3) spectra confirmed the presence of four tertiary methyls (δ_{H} 0.73, 0.77, 0.78 and 0.80 ppm) two acetate function (δ_{H} 2.09 and 2.18; δ_{C} 170.2, 20.9 and 169.9, 21.15 ppm), three oxygenated methins (δ_{H} 6.20brs, 5.38s and 5.29dd) and one more carbonyl at 172.7 ppm. The electron impact low resolution mass spectrum of **10** showed the base peak at *m/z* 191 ($\text{C}_{14}\text{H}_{23}$), already observed in the spectra of **1** and **2** and interpreted, after accurate measurements, as

originating from cleavage of the ring C. These spectral features indicated that **10** showed the common structure of rings A, B and C of **1** and **2**, and oxygenated functions, including two acetals, were present in the remaining part of the molecule. Interpretation of COSY cross-peaks led to the partial structure C9 to C16. HMBC data connected the oxygenated methine at C16 (δ_{H} 5.29 ppm) to the acetate carbonyl at δ 169.9 ppm, as well as the oxygenated methine at δ_{H} 5.38 (H25) was weakly correlated to the second acetate carbonyl at δ 170.2 ppm; the remaining oxygenated methine at δ_{H} 6.20 (H24) gave HMBC cross-peaks with C13 at δ_{C} 44.0 and C17 at δ_{C} 86.6 ppm. Further HMBC correlations H16/C15, C14 and H25/C24 allowed the construction of the bicyclic oxygenated portion. The remaining CH₂-COOH structural unit was linked to C17 as substantiated by ³J HMBC cross peaks H18/C16 and ²J HMBC correlation H18/CO19. Thus the structure of compound **10** could be determined. The stereochemistry was assigned as in **2**, from the appearance of H24 as a singlet and supported from an intense ROESY (mixing time 500ms; CDCl₃) cross-peak observed between H13 and H25.

Petrosaspongiolide K (**11**). The ¹H (500 MHz, CDCl₃) and ¹³C-NMR (125 MHz, CDCl₃) spectra and the interpretation of the HMBC data (Table 2) indicated that **11** showed a common structure of rings A,B and C of petrosaspongiolide A (**1**) and B (**2**), while the remaining portion was significantly different. In addition to fourteen signals assigned to carbons C1-C14 and to methyl carbons C20-C23 in the rings A,B and C, the ¹³C NMR spectrum revealed six more carbons including signals for an α,β -conjugated ketone (δ_{C} 155.0, 131.4 and 201.3 ppm), two methylenes (δ_{C} 37.6 and 36.2 ppm) and one carbonyl at δ_{C} 174.0 ppm, which accounted in total for 24 carbons and for 35 non exchangeable protons. The presence of an α,β -unsaturated ketone was supported by UV (234 nm, $\epsilon=5545$). The EI mass spectrum showed the molecular ion peak at m/z 372 (HREIMS: m/z 372.269733, required 372.266445), accompanied by a prominent fragment at m/z 328 (loss of CO₂), thereby indicating the presence of a carboxyl group. Thus, the molecular formula C₂₄H₃₆O₃ could be determined. The olefinic singlet at δ_{H} 6.64 ppm (C24, δ_{C} 154.9 ppm) and the methylenes at δ_{H} 2.49dd-2.17t ppm (C15) and 3.27d-3.15d ppm (C18) were connected through non protonated carbons on the basis of HMBC data (Table 2), which included cross peaks: H24/C12,C14,C16,C18; H15 α /C14,C16; H15 β /C14; H13/C24; Me23/C14, thereby constructing the α,β -unsaturated cyclohexanone fused at C13 and C14 of ring C, carrying a side chain CH₂COOH at C17. Thus, the structure of compound **11** could be determined. Rings C/D were assigned as trans-fused junction on the basis of two high coupling constants exhibited by the allylic H14 resonating in a clear region of the ¹H-NMR spectrum at δ 2.28 (brt, J=8Hz) and indicative of the axial orientation of H14 and H13. Petrosaspongiolide K (**11**) is the first example of a 25-norsesterterpene of the cheilantane group.

Petrosaspongiolide L (**12**) gave a FABMS (negative-ion) pseudo molecular ion peak at m/z 368 (M-H)⁻. In the EIMS the molecular ion at m/z 369 (HREIMS: m/z 369.309780, required 369.308731) is accompanied by a prominent peak (base peak) at m/z 324 (M-CO₂H)⁻, indicative for the presence of a carboxyl group. The ¹³C-NMR (125 MHz, CD₃OD) spectrum revealed 24 carbons including four quaternary methyls (δ_{C} 17.1, 21.9, 26.0 and 33.8 ppm) and three quaternary carbons at δ_{C} 34.3 ppm, 39.0 ppm and 39.6 ppm consistent with a tricyclic skeleton with a gem-dimethyl group at C4 and two methyl groups at the ring junction C8 and C10, already observed in **1** and **2**. The high field signals at δ 18.6 ppm, 19.7 ppm and 20.0 ppm, attributed to the methylenes at C11, C6 and C2 lying in position γ with respect to the axial methyls confirmed the transA/B-transB/C skeleton. Interpretation of COSY cross-peaks identified the sequence C1-C3, C5-C7 and C9-C12 and showed that the protons of C12 were downfield shifted to δ 2.78 and 2.98 ppm, characteristic for a benzylic methylene. Also shifted downfield was the C8 methyl signals, δ_{H} 1.25 ppm, δ_{C} 26.0 ppm. Three quaternary and two CH aromatic carbons at δ_{C} 162.6s, 158.4s and 130.7s, 149.4d, 120.5d ppm, respectively, indicated the presence of a trisubstituted pyridine. The UV spectrum (266, $\epsilon=2375$; 210, $\epsilon=5545$) gave support to presence of a pyridine. Two methylene signals at δ 33.6 ppm and 36.0 ppm coupled each to the other (COSY) and one carbonyl signal at δ_{C} 177.7 ppm completed the ¹³C-NMR spectrum of **11**. The two aromatic protons were observed resonating as singlets at δ_{H} 8.18 ppm and 7.26 ppm. These structural units were connected through non protonated carbons on the basis of HMBC data (Table 2). Cross peaks H12/C13, C14 and Me23 /C14 implied

that the pyridine was fused to ring C at C13, C14; H15/C13, C8 and C16, H17/C15 and H24/C13, C14 allowed constructing the pyridine portion with the propionic side chain at C16. Petrosaspongiolide L (**11**), which is apparently a 25-norsesiterpene, could be considered a condensation product with ammonia of a 16keto 24al precursor, similar to molliorin A and B¹³⁻¹⁴, condensation products of scalaradial with amines.




Table 2. ¹H- and ¹³C-NMR chemical shifts and HMBC correlations of the nor-sesiterpenes **11** and **12**

position	11 (CDCl ₃)			12 (CD ₃ OD)		
	$\delta_{\text{H}}^{\text{a}}$	δ_{C}	HMBC ^b	$\delta_{\text{H}}^{\text{a}}$	δ_{C}	HMBC ^b
1	1.97–0.78	39.7			40.8	
2	1.55–1.32	18.6			20.0	
3	1.38–1.12	42.0			43.2	
4		33.0			34.3	
5	0.81	56.6		0.95	57.7	
6	1.59–1.38	18.4			19.7	
7	1.69–0.81	39.8		2.48	40.0	
8		36.8			39.0	
9	0.85	59.2		1.28	56.2	
10		37.5			39.6	
11	1.69–1.39	20.6	C9	1.72–1.95	18.6	C9, C12
12	1.18–2.01dd(4.2, 14.4)	32.8	C11, C13	2.78ddd(16, 11.3, 7.1)- 2.98dd(16, 7.3)	28.0	C13 C13, C14
13	2.38brt(8.0)	36.9	C24		130.7	
14	1.5	54.3			162.6	
15	2.49dd(4.2, 16.7)- 2.17t(16.7)	37.6	C16 C14, C16	7.26s	120.5	C16, C8, C13
16		201.3			158.4	
17		131.4		3.05t(8.9) 2.67t(8.9)	33.6	C15
18	3.15d(16.8)- 3.27d(16.8)	36.2			36.0	
19		174.0			177.7	
20	0.82s	21.0		0.92s	21.9	
21	0.86s	33.0		0.92s	33.8	
22	0.85s	16.3		1.02s	17.1	
23	0.99s	15.4	C14, C9, C8	1.25s	26.0	C14
24	6.64brs	155.0	C12, C14, C16, C18	8.18s	149.4	C13, C14

^aCoupling constants are in parentheses and given in Hz. ¹H assignments aided by COSY experiments. ^bHMBC optimized for ²⁵J_{Ct}=10.0 Hz.

Cytotoxic activity of the sesiterpenoids **1–12** is summarized in Table 3. Petrosaspongiolide A(**1**) was submitted to *in vivo* tests on immunodepressed rats carrying a bronchopulmonary tumor (NSCLC-N6). It was observed an inhibition of tumoral proliferation at 20mg/Kg without significant toxicity.

Table 3. Cytotoxicity against NSLC-N6^a tumor cells (IC₅₀ expressed in µg/ml)

Compounds	IC ₅₀	Compounds	IC ₅₀
1	13.0	7	inactive
2	14.8	8	8.1
3	0.5	9	6.8
4	5.2	10	6.3
5	4.5	11	1.3
6	8.7	12	5.7

^aNSCLC-N6: human bronchopulmonary non-small-cell-lung-carcinoma.

Experimental Section

General Information. For general information see: Zampella A. *et al.*¹⁵

Isolation. The sponge *Petrosaspongia nigra*. (Dictyoceratida, Spongidae) was collected in November 1987 and December 1988 in the waters of the southern coast of New Caledonia. Taxonomic identification was performed by Dr. P.R. Bergquist, and reference specimens are on file (reference 321) at the ORSTOM Centre of Nouméa. Preliminary assays for cytotoxic activity showed marked activities for the dichloromethane extracts (P388 cells, 100% inhibition at 10 µg/ml dose, and KB cells, 78% inhibition at 10 µg/ml dose).

The organism was freeze-dried and the lyophilized material (1.0 Kg) was extracted with *n*-hexane and dichloromethane in a Soxhlet apparatus, then with MeOH (3x2 L) at room temperature. The dichloromethane extracts were filtered and concentrated under reduced pressure to give 30.0 g of a brown oil. The crude dichloromethane extract was chromatographed by MPLC on a silica gel column (150x2 g) using a solvent gradient from CHCl₃ to CHCl₃/MeOH 9:1. MPLC fractions were further purified by HPLC on a semipreparative (7.8x300 mm) µ-Bondapak-C18 column (flow rate 5 ml/min) eluting with MeOH:H₂O mixtures to afford pure compounds **1-12**. The purity of each compound was judged to be > 90% by HPLC and ¹H-NMR.

Petrosaspongiolide A (**1**) and B (**2**): (C₂₇H₄₀O₆, M.W.=460) (**1**, [α]_D = -15°, c=0.02, CHCl₃; **2**, [α]_D = -71°, c=0.03, CHCl₃), t_r=24.0 and 19.2 min, respectively, eluting with MeOH/H₂O 85:15. ¹H and ¹³C-NMR were virtually identical to those reported by Cambie *et al.*⁹.

Petrosaspongiolide C (**3**): (C₂₉H₄₂O₈, M.W.=518) [α]_D = -12.5° (c=0.003, CHCl₃); t_r=7.6 min, eluting with MeOH/H₂O 85:15. ¹H and ¹³C-NMR see text and Table 1.

Petrosaspongiolide D (**4**): (C₂₇H₄₀O₇, M.W.=476) [α]_D = -27° (c=0.003, CHCl₃); t_r=4.8 min, eluting with MeOH/H₂O 85:15. ¹H and ¹³C-NMR see text and Table 1.

Petrosaspongiolide E (**5**): (C₂₇H₃₈O₇, M.W.=474) [α]_D = -19.4° (c=0.003, CHCl₃); t_r=5.6 min, eluting with MeOH/H₂O 85:15. ¹H and ¹³C-NMR see text and Table 1.

Petrosaspongiolide F (**6**): (C₂₇H₃₈O₈, M.W.=490) [α]_D = -17.8° (c=0.002, CH₃OH); t_r=4.0 min, eluting with MeOH/H₂O 85:15. ¹H and ¹³C-NMR see text and Table 1.

Petrosaspongiolide G (**7**): (C₂₅H₃₈O₆, M.W.=434) [α]_D = -25.2° (c=0.002, CH₃OH); t_r=8.0 min, eluting with MeOH/H₂O 70:30 (analytical column). ¹H and ¹³C-NMR see text and Table 1.

Petrosaspongiolide H (**8**): (C₂₅H₃₆O₇, M.W.=448) [α]_D = +4.3° (c=0.003, CH₃OH); t_r=3.6 min, eluting with MeOH/H₂O 70:30 (analytical column). ¹H and ¹³C-NMR see text and Table 1.

Petrosaspongiolide I (**9**): (C₂₉H₄₂O₈, M.W.=518) [α]_D = -28° (c=0.001, CHCl₃); t_r=5.2 min, eluting with MeOH/H₂O 85:15. ¹H and ¹³C-NMR see text and Table 1.

Petrosaspongiolide J (**10**): (C₂₉H₄₄O₈) [α]_D = -14.5° (c=0.003, CHCl₃); t_r=17.6 min, eluting with MeOH/H₂O 80:20. White amorphous solid, EIMS m/z 520 (1, M⁺), 460 (31, M⁺-CH₃COOH), 445 (39, M⁺-CH₃-CH₃COOH), 400 (14, M⁺-2CH₃COOH), 191 (100, C₁₄H₂₃) ¹H-NMR (CDCl₃) δ : 6.20 (1H, br s, H24), 5.38 (1H, s, H25), 5.29 (1H, br d, J=10 Hz, H16), 3.06 and 2.91 (2H, AB, J=16.9 Hz, H18, 18'), 2.18 (3H, s, MeCO), 2.07 (3H, s, MeCO), 0.83 (3H, s, Me-20), 0.79 (3H, s, Me-21), 0.78 (3H, s, Me-23), 0.75 (3H, s, Me-22); ¹³C-NMR (CDCl₃) δ: 172.7 (s, CH₂COOH), 170.2 (s, MeCO), 169.9 (s, MeCO), 11.3 (d, C24), 94.0 (d, C25), 86.6 (s, C17), 71.8 (d, C16), 59.3 (d, C9), 56.6 (d, C5), 45.0 (d, C14), 44.0 (d, C13), 42.0 (t, C3), 39.9 (t, C1), 39.9 (t, C7), 37.5 (s, C10), 37.0 (s, C8), 35.4 (t, C18), 33.3 (q, C20), 33.3 (s, C4), 31.0 (t, C12), 26.3 (t, C15), 21.4 (q, C21), 21.2 (q, MeCO), 20.9 (q, MeCO), 20.6 (t, C11), 18.6 (t, C2), 18.5 (t, C6), 16.1 (q, C23), 15.2 (q, C22).

Petrosaspongiolide K (**11**): (C₂₄H₃₆O₃, M.W.=372) [α]_D = -15.4° (c=0.004, CHCl₃); t_r=24.8 min, eluting with MeOH/H₂O 80:20. ¹H and ¹³C-NMR see text and Table 2.

Petrosaspongiolide L (**12**): (C₂₄H₃₅O₂N, M.W.=369) [α]_D = -33.3° (c=0.001, CH₃OH); t_r=3.2 min, eluting with MeOH/H₂O 80:20 (analytical column). ¹H and ¹³C-NMR see text and Table 2.

Cytotoxic assays. Experiments are performed in 96 wells microtiter plates (2x10⁵ cells/ml). Cell growth is estimated by a colorimetric assay based on the conversion of tetrazolium dye (MTT) to a blue formazan product using live mitochondria.¹⁶ Eight determinations are performed for each concentration. Control growth is estimated for 16 determinations. Optical density at 570 nm corresponding to solubilized formazan is read for each well on Titertek Multiskan MKII.

ACKNOWLEDGMENTS

This contribution is part of the ORSTOM-CNRS SMIB program; we thank the ECC project "Marine Sciences and Technology, MAST III" (Contract MAS 3-CT95-0032) for financial support. Some mass spectra were provided by the "Servizio di Spettrometria di Massa del CNR e dell'Università di Napoli". The undergraduate students C. Viscardi, S. Carbonelli, N. Di Prizito, D. Capozzoli, L. Ciasullo are acknowledged for their kind collaboration. The NMR spectra were recorded at CRIAS "Centro Interdipartimentale di Analisi Strumentale", Faculty of Pharmacy, University of Naples. A specimen of the sponge *Petrosaspongia nigra* is kept at the Queensland Museum in Brisbane (QMG 304685).

REFERENCES AND NOTES

1. Crews, P.; Naylor, S. *Progress in the Chem.Org.Nat.Products* **1985**, *48*, 203
2. Hanson, J.R. *Nat.Prod.Rep.* **1986**, *3*, 123-132; **1992**, *9*, 481-489; **1996**, *13*, 529-535.
3. Faulkner, D.J. *Nat.Prod.Rep* **1996**, *13*, 75-125, and earlier reviews cited therein.
4. de Silva, E.D.; Scheuer P.J. *Tetrahedron Lett.* **1980**, *21*, 1611-12.
5. De Rosa, S.; De Stefano, S.; Zavodnik, N. *J.Org.Chem.* **1988**, *53*, 5020.
6. Kernan, M.R.; Faulkner, D.J.; Parkanyi, L.; Clardy, J.; de Carvalho, M.S.; Jacobs, R.S. *Experientia* **1989**, *45*, 388-390.
7. Glaser, K.B.; de Carvalho, M.S.; Jacobs, R.S.; Kernan, M.R.; Faulkner, D. *J Med.Pharm.* **1989**, *30*, 782.
8. de Carvalho, M.S.; Jacobs, R.S. *J. Biochem. Pharm.* **1991**, *42(B)*, 1621-1626.
9. Lal, A.R.; Cambie, R.C.; Rickard, C.E.F.; Bergquist, P.R. *Tetrahedron Lett.* **1994**, *35*, 2603-2606.
10. Cambie, R.C.; Lal, A.R.; Rickard, E.F. *Acta Cryst.* **1996**, *52*, 709-711.

11. Zampella, A.; D'Auria, M.V.; Minale, L.; Debitus, C.; Roussakis, C. *J. Am. Chem. Soc.* **1996**, *118*, 11085-11088.
12. Fang, J.M.; Lang, C.I.; Chen, W.L.; Cheng, Y.S. *Phytochemistry* **1991**, *30*, 2793-2795.
13. Cafieri, F.; De Napoli, L.; Fattorusso E.; Santacroce C.; Jies, F. *Tetrahedron Lett.* **1977**, 477.
14. Cafieri, F.; De Napoli, L.; Fattorusso E.; Santacroce, C. *Experientia* **1977**, *33*, 994.
15. Zampella, A., D'Auria M.V.; Gomez-Paloma, L.; Minale, L.; Debitus, C.; Henin, Y. *J. Am. Chem. Soc.* **1996**, *118*, 6202.
16. Mosmann, T. *J. Immunol. Methods* **1983**, *65*, 55-68.

(Received in UK 11 March 1997; revised 30 May 1997; accepted 5 June 1997)