



Unusual Reaction of 1,4-Diamino-2-nitrobenzene Derivatives toward Nucleophiles: Catalysis by Sodium Sulphite

Luciano Forlani,^{*,§} Carla Boga[§], Milena Mazza[§], Vanni Cavrini[#] and Vincenza Andrisano[#]


[§] Dipartimento di Chimica Organica, "A. Mangini", viale Risorgimento 4, Università di Bologna, I-40136 Bologna, Italia.

[#] Dipartimento di Scienze Farmaceutiche, via Belmeloro 6, Università di Bologna, I-40126 Bologna, Italia.

Received 21 January 1998; accepted 19 February 1998

Abstract. Unusual substitution of amino group occurs by reactions of some 1,4-diamino-2-nitrobenzenes (semipermanent hair dyes) and nucleophiles (NH₃, H₂O). The reaction is catalyzed by sodium sulfite. The obtained products are suspected of being toxic substances which may be present in cosmetic matrices. Apparently, this reaction is a nucleophilic aromatic substitution but it may be explained by a mechanism involving a tautomeric form of substrate. © 1998 Elsevier Science Ltd. All rights reserved.

1,4-Diamino-2-nitrobenzene derivatives are semipermanent hair dyes,¹ which are useful in cosmetics. Some of their, and other here considered derivatives, are reported as follows.




	R	R ¹	R ²	commercial name
1	NHCH ₂ CH ₂ OH	NO ₂	NH ₂	HCRED 3
2	NH ₂	NO ₂	NH ₂	NP
3	NHCH ₂ CH ₂ OH	NO ₂	NHCH ₂ CH ₂ OH	HCVIOLET
4	NH ₂	NO ₂	NHCH ₂ CH ₂ OH	HCRED 7
5	NHCH ₂ CH ₂ OH	NO ₂	N(CH ₂ CH ₂ OH) ₂	BLUE SOLID
6	NH ₂	NO ₂	N(CH ₂ CH ₂ OH) ₂	HCRED 13
7	NHCH ₂ CH ₂ OH	NH ₂	NO ₂	HCY 5
8	NHCH ₂ CH ₂ OH	NO ₂	NO ₂	
9	C ₅ H ₁₀ N (piperidyl)	NO ₂	NO ₂	
10	SO ₃ H	NO ₂	NO ₂	
11	NH(CH ₂) ₃ CH ₃	NO ₂	NO ₂	
12	NHCH ₂ CH ₂ OH	NO ₂	H	
13	OH	NO ₂	NH ₂	
14	OH	NO ₂	NO ₂	
15	NH ₂	NO ₂	NO ₂	
16	NH ₂	NO ₂	H	
17	OH	NO ₂	NHCH ₂ CH ₂ OH	
18	OH	NO ₂	H	

In principle, the stability of hair dyes is important not only for obvious technical requirements, but also because some of their reaction products may be toxic substances. In the XV European Directive (92/86, 21/10/92) a list of forbidden hair dyes is reported because of their potential toxicity.²

Recently,³ in commercial cosmetic samples (which are multicomponent mixtures) containing 1-(N-hydroxyethyl)-2-nitro-1,4-phenylenediamine **1** (HCRED3), the presence of 2-nitro-1,4-phenylenediamine **2** (NP) was detected. In order to obtain some informations on the origin of NP, we investigated this transformation and the possible action of some components of commercial matrix which contains HCRED3 or other related dyes and the mechanism of formation of NP.

Results and discussion

Commercial samples, containing **1** and NH₃, stored at room temperature for several months, show the presence of large amounts of **2**; when they are stored at 70°C for a month, about 50% of HCRED3 is converted to **2**. This transformation is reported in Scheme 1.



Scheme 1

Apparently, the reaction of Scheme 1 seems to be a nucleophilic aromatic substitution (S_NAr), but it is surprising for two main reasons. The first is that 2-nitro-substituted benzenes (for example 2-nitro-halogenobenzenes) are only moderately activated substrates⁴ toward the attack of the nucleophiles and the presence of the amino group in 4 position reduces the activation of the substrate by his strong electron releasing effect. The second reason is that the amino group is a bad leaving group. In particular, primary amino group may be hardly replaced by amines,⁵ while instances of replacement of secondary amino groups are reported in literature.⁶

In cosmetic matrix samples, when NH₃ is replaced by hydroxyethylamine, (or by other amines) HCRED3 is not converted to NP remaining unchanged in commercial samples after 2-3 months at 70°C. This fact is an indication that the formation of NP from HCRED3 in commercial samples cannot be ascribed to some N-dealkylation reactions of the hydroxy-ethyl group.

After some preliminary experiments on commercial samples and on mixtures of their separate components, we verified the importance of the presence of sodium sulfite which is added as antioxidant in cosmetic matrix. Table 1 reports the main results of the reaction of HCRED3 in the presence of NH₃ (or piperidine).

Table 1
Reactions between HCRED3 (1) and ammonia, at 70°C.

Entry	Solvent	Added substances	Time(days)	Yield %	
				2 (NP)	13
1	H ₂ O	--	40	40	0
2	DMSO/H ₂ O ^a	--	40	5	0
3	H ₂ O ^b	--	40	0	0 ^c
4	DMSO/H ₂ O ^a	Na ₂ SO ₃	1	100	0
5	H ₂ O	Na ₂ SO ₃	3	80	15
6	H ₂ O ^b	Na ₂ SO ₃	20	0	80
7	H ₂ O	Na ₂ SO ₃	5	0	80
8	H ₂ O	Na ₂ SO ₃ (NaOH)	0.2	0	95

a) (10:1 v/v) b) With piperidine. c) HCRED3 was recovered unchanged from the reaction mixtures.

In aqueous solvents, variable amounts of the corresponding phenol (2-nitro-4-aminophenol) 13 were recovered from the reaction mixtures.

Table 2
Reactions of selected substrates related with nucleophiles in MeOH/H₂O (1:1 v/v) at 70°C.

Entry	Substrate	nucleophiles	time ^a	A (yield %) B (yield %)	
				(see Scheme 2)	
1	1	NH ₃ /Na ₂ SO ₃ ^b	3	2(80)	13(15)
2	1	Piperidine/Na ₂ SO ₃ ^b	20	2(0)	13(95) ^c
3	8	Piperidine/Na ₂ SO ₃ ^b	5	15(0)	14(90) ^c
4	1	NaOH	0.2	2(0)	13(100)
5	8	NH ₃	3	15(30)	14(0)
6	8	NH ₃ /Na ₂ SO ₃ ^a	2	15(0)	14(100)
7	9	NH ₃	0.5	15(100)	14(0)
8	7	NH ₃ /Na ₂ SO ₃ ^a	20	----- ^d	
9	5 ^e	NH ₃ /Na ₂ SO ₃ ^a	6	6(80)	14(0)
10	10	NH ₃	15m	15(100)	14(0)
11	11	NH ₃ /Na ₂ SO ₃ ^b	12	15(30-50)	14(30-50) ^f
12	12	NH ₃ /Na ₂ SO ₃ ^a	6	16(80)	18(0)
13	12	NH ₃	10	----- ^d	
14	3 ^g	NH ₃ /Na ₂ SO ₃ ^a	1	6(85)	17(10) ^h

a) Days, unless otherwise indicated. b) Substrate/Na₂SO₃ ratio = 1:10. c) No presence of piperidino-substitution products was detected. d) No reactions. e) BLUE SOLID. f) Variable and poorly reproducible relative yields. g) HC VIOLET. h) No presence of products of substitution of the N(CH₂CH₂OH)₂ group was detected.

An interesting fact is offered by entry 4 (Table 1) which clearly indicates that the formation of NP in commercial samples easy occurs when the ammonium hydroxide and sodium sulfite (or related derivatives such as sodium hydrosulfite) are present.


Piperidine is unable to afford substitution products (see entries 2,3 of Table 2): ammonia and water are the only entering groups. The presence of sodium sulfite strongly reduces the times of conversion.

Sulfite ion is known as a powerful nucleophile,⁷ and a reasonable intermediate of the reaction of Scheme 1 may be a sulphonate derivative. However, this product was not recovered from the reaction mixtures. Some other data of Table 1 are worthy of consideration.

In S_NAr reactions, piperidine is a better nucleophile than NH₃, but, in the present reaction, the piperidine is unreactive toward HCRED 3. (See entry 3). The reaction is depressed by the use of DMSO as solvent (entry 2), while DMSO is well known to favour S_NAr.

The feeble activation of the substrate, the presence of a bad leaving group, the absence of piperidino-substitution product and the negative effect of DMSO are arguments which strongly indicate that the reaction of Scheme 1 cannot occur by the usual two steps mechanism of S_NAr reactions.

Table 2 reports some results obtained for substrates related to HCRED 3. Also in this cases the water may compete with ammonia in displacement reactions, as Scheme 2 illustrates.



Scheme 2


The reactivity of HCRED 3 may be explained by some possible interactions between the *ortho*-nitro group and the hydroxy group of the amino moiety, to facilitate the leaving group departure. Unfortunately our data cannot be decisive on this possibility. A feeble indication supporting this idea, is that N-butyl-2,4-dinitroaniline 11 reacts with the mixture NH₃/Na₂SO₃ slower (entry 11 of Table 2) than the N-(2-hydroxyethyl)-2,4-dinitroaniline 8 (entry 6 of Table 2). Owing the strong activation of dinitroderivatives, probably these reactions are simple S_NAr reactions.

To have some informations on the behaviour of the hydroxy group of the 2-hydroxy-ethylamine, we detected the rate of the reaction of 2-nitro-fluorobenzene and 2-hydroxy-ethylamine in DMSO (at 25°C) to obtain the aniline derivative 12. The rate of this reaction (k in $\text{s}^{-1} \text{ mol}^{-1} \text{ dm}^3 = 1.7 \times 10^{-3}$) is somewhat lower than the value of the similar reactions with 2-nitro-fluorobenzene and 2-N,N-dimethyl-ethylenediamine ($k = 3.7 \times 10^{-3}$)⁴ and ethylenediamine (3.3×10^{-3}).⁴ These feeble differences (referred to an S_NAr reaction) cannot be indicative of presence of particular interactions between *ortho*-nitro and hydroxy groups.

The mechanism of reaction of Scheme 1 may be represented by Scheme 3. The first step of Scheme 3 is a tautomeric equilibrium affording the imino compound 18. The nucleophilic attack of sulfite (or bisulfite) ion on 18 produces the bisulfite adduct 19 which reacts to substitute the NHR group with ammonia or water.


The experimental conditions of reaction of Scheme 1 falls in the field of Bucherer reactions⁸ which are well known in naphthalene serie but which are rarely observed in monocyclic derivatives.⁹ Even if a pathway of reaction parallel to that of the Bucherer reaction cannot be ruled out, no evidences of the presence of sulphonic group in position 3 of the substrate (as usually accepted by the bucherer mechanism) were obtained.

Other unusual activation of strong electron donor groups (amino and hydroxy groups) toward nucleophilic attack were observed in naphthalene¹⁰ and in 2-thiazoleamine derivatives¹¹. In all cases these unexpected reactivities were explained by tautomerism of substrates affording the reactive species.



Scheme 3

The Scheme 3 explains also the reactivity of compound 5 (**BLUE SOLID**) even if it is moderately less reactive than **HCRED 3** (see entry 9). In another mechanistic hypothesis (depicted in Scheme 4), **21** is an intermediate similar to **19**; **21** should be reached from the quinone imine **20**.



Scheme 4

21-like intermediate cannot be obtained from **BLUE SOLID** and so **21** should explain the decrease of reactivity by passing from **HCRED 3** to **BLUE SOLID**.

Usually compounds **20**-like are obtained by oxidizing reagents which are absent in our reaction mixtures. Consequently, Scheme 4, which is a possible reaction step in permanent dyes use,¹² is a poorly probable reaction path-way to explain reaction of **HCRED 3**.

Probably the presence of the nitro group in position 2 favours the formation of tautomer **18** by acidification of the hydrogen atom of the NHR group and it enhances the attack of nucleophiles on the imino tautomer **18**. In fact, in **HCVIOLET (3)** the reaction occurs replacing the $\text{NHCH}_2\text{CH}_2\text{OH}$ group in position 1 (entry 14 of Table 2), while the same group in position 4 is not replaced. Accordingly, **HCY5 7** is unreactive toward the $\text{NH}_3/\text{Na}_2\text{SO}_3$ mixture (see table 2, entry 8).

The presence of the amino group in 4 position is not completely necessary in activating the substrate. When the amino group in 4 position is absent the reactions with ammonia occurs (entry 12 of Table 2) at lower rate than that of **HCRED 3** (entry 12 of Table 2). In the case of 2-nitro-aniline derivative, the reaction appears to be an $\text{S}_{\text{N}}\text{Ar}$ reaction. However, the importance of sodium sulfite catalysis (entry 13) indicates that

the main reaction pathway may be the mechanism depicted in Scheme 3, even if the possibility of the presence of an S_NAr mechanism with an unusual catalysis by sodium sulfite cannot be completely ruled out. In fact, not only sodium sulfite is well known as powerful nucleophile but also it is a powerful leaving group (see entry 10 of Table 2). However, the behaviour of sulfite as nucleophile needs further investigation. Clearly, the use of sodium sulfite to stabilize the commercial matrix of hair dyes produces a dangerous destabilization of the dyes.

Present data well agree with the pathway of Scheme 3. Probably the nitro group in *ortho* position with respect to the reaction centre, enhance the nucleophilic attack of both, sulfite ion and entering group, while the 4 position is less activated (it is a meta position with respect to the nitro group) and consequently unreactive.

Experimental

General conditions

NMR data were recorded on a Varian Gemini 300 MHz spectrometer. Chemical shifts are reported in ppm relative to solvent ($CDCl_3$ ($\delta = 7.27$), d^6 -DMSO ($\delta = 2.60$); d^6 -acetone ($\delta = 2.20$); J values are given in Hz. Signals related to N-H and O-H groups disappear after addition of D_2O .

Mass spectra were recorded with a VG-7070E spectrometer.

IR spectra were recorded (in KBr) with a Perkin Elmer mod. 1600 FT-IR spectrophotometer.

UV/VIS spectra were recorded with a Perkin Elmer mod. Lambda 5.

Melting points were determined with a Büchi apparatus and they are uncorrected.

Materials. Amines were commercial samples (Carlo Erba) purified by usual procedures. 1-(N-piperidyl)-2,4-dinitrobenzene **9**,¹³ N-butyl-2,4-dinitroaniline **11**¹⁴ were prepared by described procedures. Samples of cosmetic matrix and of separate components (including all the commonly used dyes **1-7**) were obtained from Lowenstein dyes cosmetics, inc. (USA), from Hoechst (Germany) and from S.N.P.E. (France) and used as received. Their structure and purity (>95%) was checked by spectral data which are reported as follows.

1-(N-2-hydroxyethyl)-2-nitro-1,4-phenylenediamine (HCRED3) 1 mp 128-130 °C; IR ν_{max}/cm^{-1} 3360, 3340, 3280, 3180, 1560, 1520; 1H NMR (300MHz, $CDCl_3$): δ 8.05-7.85 (1H, m, NH), 7.53 (1H, d, J=4.6, 3-H), 6.99 (1H, dd, J=8.7, J=2.8, 5-H), 6.82 (1H, d, J=8.8, 6-H), 4.00-3.98 (3H, m, CH_2OH and OH), 3.60-3.40 (4H, m, CH_2NH and NH_2); δ_H in d^6 -DMSO 8.05 (1H, t, NH), 7.37 (1H, d, J=2.7, 3-H), 7.14 (1H, dd, J=9.1, J=2.7, 5-H), 7.00 (1H, d, J=9.1, 6-H), 5.06 (1H, t, OH), 5.02 (2H, s, NH_2), 3.80-3.65 (2H, m, CH_2OH), 3.50-3.35 (2H, m, CH_2NH); m/z 197 (M^+ , 47), 166 (100), 136 (14), 119 (27).

2-nitro-1,4-phenylenediamine (NP) 2 mp 137-140 °C; IR ν_{max}/cm^{-1} 3320, 1590, 1560, 1510; 1H NMR (300MHz, $CDCl_3$): δ 7.44 (1H, d, J = 8.7, J=2.8, 3-H), 6.88 (1H, dd, J=8.8, J=2.8, 6-H), 6.70 (1H, d, J=8.9, 5-H), 5.75 (2H, br s, NH_2), 3.50 (2H, br s, NH_2); δ_H in d^6 -DMSO 7.24 (1H, d, J=2.6, 3-H), 7.03-6.90 (4H, m, 5-H, 6-H and NH_2), 4.95 (2H, br s, NH_2); m/z 153 (M^+ , 100), 136 (3), 119 (9), 107 (68).

N,N'-bis(2-hydroxyethyl)-2-nitro-1,4-phenylenediamine (HCVIOLET) 3 mp 106-108 °C; IR ν_{max}/cm^{-1} 3330, 1560, 1520; 1H NMR (300MHz, $CDCl_3$): δ 8.10-8.00 (2H, m, NH), 7.87 (2H, dd, J = 8.0, J = 0.6, 7-H), 7.41 (1H, d, J=2.7, 3-H), 7.00 (1H, dd, J=9.3, J=3.0, 5-H), 6.85 (1H, d, J=9.2, 6-H), 4.00-3.85 (4H, m, CH_2OH), 3.55-3.45 (2H, m, CH_2NH), 3.35-3.25 (2H, m, CH_2NH), 1.80-1.60 (2H, br s, OH); δ_H in d^6 -DMSO 8.16 (1H, t, NH), 7.25-7.15 (2H, m, aromatic), 7.05 (1H, d, J=9.1, aromatic), 5.50 (1H, t, NH), 5.05 (1H, t, OH), 4.83 (1H, t, OH), 3.90-3.65 (4H, m, CH_2), 3.50-3.35 (2H, m, CH_2), 3.10-3.00 (2H, m, CH_2); m/z 241 (M^+ , 55), 210 (100), 197 (16), 189 (5).

1-(N-2-hydroxyethyl)-3-nitro-1,4-phenylenediamine (HCRED7) 4 mp 100-102 °C; IR ν_{max}/cm^{-1} 3520, 3460, 3350, 1570, 1510; 1H NMR (300MHz, $CDCl_3$): δ 7.32 (1H, d, J=2.7, 3-H), 6.90 (1H, dd, J=8.9, J=2.8, 5-H), 6.72 (1H, d, J=8.8, 6-H), 5.90-5.70 (2H, br s, NH_2), 4.00-3.85 (3H, m, CH_2OH and NH),

3.40-3.25, (2H, m, CH₂NH), 1.75 (1H, br s, OH); δ_{H} in d⁶-DMSO 7.20-6.90 (5H, m, aromatic and NH₂), 5.40 (1H, t, NH), 4.83 (1H, t, OH), 3.75-3.55 (2H, m, CH₂OH), 3.20-3.05 (2H, m, CH₂NH); m/z 197 (M⁺,37), 166 (100), 153 (4), 120 (38).

1-(N-2-hydroxyethyl)-N'-(bis-2-hydroxyethyl)-2-nitro-1,4-phenylenediamine 5 (BLUE SOLID) mp 82-84 °C; IR $\nu_{\text{max}}/\text{cm}^{-1}$ 3320, 1510; ¹H NMR (300MHz, d₆-DMSO): δ 8.05 (1H, t, NH), 7.50-6.95 (3H, m, aromatic), 5.05 (1H, br s, OH), 4.85 (2H, br s, OH), 3.85-3.75, 3.70-3.60, 3.55-3.40, 3.15-3.05 (12 H, m, CH₂); m/z 285 (M⁺,26), 254 (100), 241 (10), 210 (47).

1-(N-bis-2-hydroxyethyl)-3-nitro-1,4-phenylenediamine (HCRED13) 6 mp 91-92 °C; IR $\nu_{\text{max}}/\text{cm}^{-1}$ 3340, 1645,1600, 1530, 1450; ¹H NMR (300MHz, d₆-DMSO): δ 8.27 (1H, m, 3-H), 7.90-7.70 (1H, m, 5-H), 7.26 (1H, d, J=9.0, 6-H), 4.50-3.80 (4H, br s, OH and NH₂), 3.70-3.40 (8H, m, CH₂); m/z 241 (M⁺,19), 210 (100), 197 (11), 166 (42)

1-(N-2-hydroxyethyl)-4-nitro-1,2-phenylenediamine (HCY5) 7 mp 131-133 °C; ¹H NMR (300MHz, d₆-DMSO): δ 7.63 (1H, dd, J=8.8, J=2.7, 5-H), 7.53 (1H, d, J=2.7, 3-H), 6.59 (1H, d, J=9.1, 6-H), 6.02 (1H, t, NH), 5.25 (2H, br s, ,NH₂), 4.93 (1H, t, , OH), 3.75-3.65 (2H, m, CH₂OH),3.40-3.30 (2H, m, CH₂NH); m/z 197 (M⁺,30), 166 (100), 120(38).

N-(2,4-dinitrophenyl)-2-aminoethanol 8.

0.68 mL (5.38 mmol) of 2,4-dinitrofluorobenzene were dissolved in 4.0 mL of THF/acetone mixture (1/1 v/v). 2.0 mL (10.8 mmol) of ethanolamine were added and the reaction mixture was stirred at room temperature for 60 min. The solvent was removed under vacuum. 1.13 g (yield = 93%) of **8** are obtained after purification by flash chromatography (F.C.) (eluent: ethyl acetate/petroleum light 1/1). mp 89-90 °C (from MeOH); ¹H NMR (300MHz, CDCl₃): δ 9.13 (1H, d, J=2.7, 3-H), 8.90-8.70 (1H, m, NH), 8.27 (1H, dd, J=9.6, J=2.6, 5-H), 7.00 (1H, d, J=9.6, 6-H), 4.10-3.90 (2H, m, CH₂OH), 3.70-3.55 (2H, m, NHCH₂), 1.94 (1H, br s, OH); m/z 227 (M⁺, 17), 196 (100), 150 (12), 104(11).

N-(2-nitrophenyl)-2-aminoethanol 12.

12 was obtained in high yield (98 %) by the same procedures of **8**. DMSO was the solvent of the reaction and the reaction mixture was poured in water, extracted with CH₂Cl₂ and purified by F.C. mp 71-73 °C (from EtOH) ¹H NMR (300MHz, CDCl₃): δ 8.35-8.20 (1H, m, NH), 8.20 (1H, d, J=8.9, 3-H), 7.48 (1H, t, 4-H), 6.92 (1H, d,J=8.7, 6-H), 6.70 (1H, t, 5-H), 4.05-3.95 (2H, m, CH₂OH), 3.60-3.50 (3H, m, NHCH₂ and OH). m/z 182 (M⁺, 25), 151 (100), 135 (3), 121 (6), 104 (14), 93 (23).

Reactions between 1 or related substrates (see tables 1 and 2) and nucleophiles. Typical procedure.

3.0 mL of aqueous 30% ammonia (or piperidine or aqueous 10% NaOH) and Na₂S₃ (as indicated in tables 1,2) were added to a solution of 0.050 g (0.25mmol) of **1** in 3.0 mL of solvent (molar ratio substrate:Na₂S₃ was about 1:10) The reaction mixture was stirred at 70 °C in a sealed vessel. The reaction was monitored by TLC (eluent ethyl acetate/ petroleum light 8/2) The products of the reaction were isolated and purified by F.C. and they were identified by comparison with authentic samples or by spectroscopic analysis.

Physical-chemical data of compounds **9** and **11** are in agreemen with literature.

4-nitro-2-aminophenol 13; mp 123-125 °C (from EtOH) (lit¹⁵ 125-127 °C) ¹H NMR (300MHz, CDCl₃): δ 10.17 (1H, s, OH), 7.36 (1H, m, 3-H), 7.00 (2H, m, 5-H and 6-H), 3.70-3.65 (2H, m, NH₂,); m/z 154 (M⁺, 100), 137 (26), 124 (6), 108 (20).

2,4-dinitrophenol 14; mp 113-115 °C (from acetone) (lit¹⁶ 112-114 °C) ¹H NMR (300MHz, CD₃COCD₃): δ 10.7-10.2 (1H, br s, OH), 9.08 (1H, d, J=2.8, 3-H), 8.67 (1H, dd, J=9.3, J=2.8, 5-H), 7.60 (1H, d, J=9.2, 6-H); m/z 184 (M⁺,100), 154 (25), 107 (30), 91 (39).

2,4-dinitroaniline 15; mp 176-178 °C (from EtOH) (lit¹⁷ 176-178 °C); ¹H NMR (300MHz, CD₃COCD₃): 9.09 (1H, d, J=2.7, 3-H), 8.37 (1H, dd, J=9.3, J=2.7,H-5), 8.20-8.00 (2H, br s, NH₂), 7.39 (1H, d, J=9.5, 6-H).

2-nitroaniline 16; mp 70-72 °C, (from H₂O) (lit¹⁸ 73-76 °C); ¹H NMR (300MHz, CDCl₃): δ 8.13 (1H, dd, J=8.7, J=1.6, 3-H), 7.37 (1H, m, 4-H), 6.80 (1H, dd, J=1.3, J=8.4, 6-H), 6.72 (1H, m, 5-H), 6.20-5.85 (2H, m, NH₂); m/z 138 (M⁺, 100), 121 (6), 92 (54).

4-N-(2-hydroxyethyl)-2nitrophenol 17; mp 112-114°C (from MeOH); ¹H NMR (300MHz, CDCl₃): δ 10.23 (1H, s, OH), 7.28 (1H, m, 3-H), 7.01 (2H, m, 5-H and 6-H), 4.12-4.05 (2H, m, CH₂OH and NH), 4.00-3.95 (2H, m, CH₂OH), 3.20-3.15 (2H, m, NHCH₂); m/z 198 (M⁺, 82), 167 (100), 121 (27).

Kinetic data were obtained by usual procedures by u.v./vis spectrophotometric method⁴. The reaction is first order in each reagent. In preparative runs, the only derivative **12** was recovered from the reaction mixtures. No presence of product arising from the attack of the hydroxy group on the 2-nitro-fluorobenzene was detected.

Acknowledgements

The authors thank the Ministero dell'Università e della Ricerca Scientifica e Tecnologica, the Consiglio Nazionale delle Ricerche (CNR, Roma) and the University of Bologna (Progetto d'Ateneo: Biomodulatori organici: sintesi, proprietà ed applicazioni).

References

- Knowlton, J.; Pearce, S. *The Handbook of Cosmetic Sciences and Technology*; Elsevier Advanced Technology, London, 1993; pp 201-232
Chemistry and Technology of the Cosmetics and Toiletries Industry, eds. Williams, D. F.; Schmitt, W. H. Blackie, 1966.
Brown, K. *J. Soc. Cosmet. Chem.*, **1982**, *33*, 375.
Corbett, J. F. *J. Soc. Cosmet. Chem.*, **1984**, *35*, 297.
- Cosmetic, Toiletry and Fragrance Association, Inc., *J. Am. Coll. Toxicol.* **1992**, *11(4)*, 381, 423, 447, 521.
- Andrisano, V.; Gotti, R.; Roveri, P.; Cavrini, V. *Chromatographia.*, **1997**, *44*, 431,
- Boga, C.; Forlani L.; Guardia, P. *Gazz. Chim. Ital.*, **1997**, *127*, 259.
- Terrier, F. *Nucleophilic Aromatic Displacement*, VCH Publ. New York, 1991.
- de Vargas, E. B.; de Rossi, R. H. *Tetrahedron Lett.*, **1982**, *23*, 4423.,
de Vargas, E. B.; de Rossi, R. H. *J. Phys. Org. Chem.*, **1989**, *2*, 507.
Sekiguchi, S.; Ishikura, H.; Hirose, Y.; Ono, N. *Tetrahedron*, **1990**, *46*, 5567.
Sekiguchi, S.; Hosokawa, M.; Suzuki, T.; Sato, M. *J. Chem. Soc. Perkin Transaction 2*, **1993**, 1111.
- Edwards, J.O.; Pearson, R. G. *J. Amer. Chem. Soc.*, **1962**, *84*, 16.
Bevan, C. W. L.; Foley, A. J.; Hirst, J.; Uwamu, W. O. *J. Chem. Soc. B*, **1970**, 794.
Crampton, M. R.; Willison, M. J. *J. Chem. Soc. Perkin Transaction 2*, **1976**, 160.
Bernasconi, C. F.; Bergstrom, R. G. *J. Amer. Chem. Soc.*, **1973**, *95*, 3603.
- Seeboth, H. *Angew. Chem.Int. Ed.*, **1967**, *6*, 307.
- Bucherer, H. T. *Berichte*, **1920**, *53*, 1457.
- Bosco, M.; Forlani, L.; Todesco, P. E. *J. Chem. Soc. B*, **1970**, 1742.
- Forlani, L.; Medici, A.; Todesco, P. E. *Tetrahedron Lett.*, **1976**, 201,3.
Forlani, L. *Gazz. Chim. Ital.*, **1984**, *114*, 279.
- Brown, K. C.; *J. Soc. Cosmet. Chem.*, **1982**, *33*, 375.
- Bunnett, J. F. Garst, R. H. *J. Amer. Chem. Soc.*, **1965**, *87*, 3879.
- Ross, S. D. *J. Amer. Chem. Soc.*, **1959**, *81*, 2113.
- Beilsteins Handbuch der Organischen Chemie*, Verlag Springer, Berlin **1930**, *13*, 520.
- Beilsteins Handbuch der Organischen Chemie*, Verlag Springer, Berlin **1930**, *6*, 251.
- Beilsteins Handbuch der Organischen Chemie*, Verlag Springer, Berlin **1930**, *12*, 747.
- Beilsteins Handbuch der Organischen Chemie*, Verlag Springer, Berlin **1930**, *12*, 687.