

Corrigendum

Corrigendum to “Examination of Ligand Effects in the Heck Arylation Reaction”


[Tetrahedron 56 (2000) 7975]<sup>☆</sup>

Maryiam Qadir, Tobias Möchel and King Kuok (Mimi) Hii

*Department of Chemistry, King’s College London, Strand, London WC2R 2LS, United Kingdom*

On page 7976 Table 1 should have appeared as:

**Table 1.** Effect of catalyst with/without phosphine ligands (reactions were carried out in pressure tubes, with 1 mg catalyst, 3 equiv. of methylacrylate with respect to iodobenzene, and 3 equiv. of NEt<sub>3</sub>)



| Entry          | Catalyst  | Reaction time/h | % conversion | TON     |
|----------------|---|-----------------|--------------|---------|
| 1              | Pd(PPh <sub>3</sub> ) <sub>4</sub>                    | 5               | 100          | 109,106 |
| 2              | Pd <sub>2</sub> (dba) <sub>3</sub> ·CHCl <sub>3</sub> | 4               | 100          | 100,000 |
| 3              | Pd(OAc) <sub>2</sub>                                  | 4.5             | 95           | 95,000  |
| 4 <sup>a</sup> | Pd(OAc) <sub>2</sub> / <b>4a</b>                      | 19.5            | 97           | 97,000  |
| 5              | Pd(OAc) <sub>2</sub> /PPh <sub>3</sub>                | 19.5            | 100          | 100,000 |
| 6              | [(dppp)Pd(dba)]                                       | 6               | 100          | 100,000 |

<sup>a</sup> **4b–4e** showed the same reactivity.

<sup>☆</sup> PII of original article: S0040-4020(00)00703-1