

Corrigendum

Corrigendum to “A Convenient Method to Synthesize Phosphinic Peptides Containing an Aspartyl or Glutamyl Aminophosphinic Acid. Use of the Phenyl Group as the Carboxyl Synthons”
[Tetrahedron 55 (1999) 14635][☆]

Dimitris Georgiadis,^a Magdalini Matziari,^a Stamatia Vassiliou,^a Vincent Dive^b and Athanasios Yiotakis^a

^aDepartment of Chemistry, Laboratory of Organic Chemistry, University of Athens, Panepistimiopolis, Zografou, 15771, Athens, Greece

^bDepartment d'Ingenierie et d'Etudes des Proteines, CEA-Direction des Sciences du Vivant, Centre des Etudes de Saclay, 91191 Gif/Yvette Cedex, France

The legend for Scheme 1 should read as:

Scheme 1. (a) Cbz-Cl, MgO, H₂O, 4 h; (b) AdBr, Ag₂O, CHCl₃, reflux, 2 h; (c) *N,N*-dimethylformamide di-*tert*-butyl acetal, benzene, reflux, 30 min.; (d) 5% TFA/CH₂Cl₂, 7 h; (e) 5% TFA/CH₂Cl₂, 3 h; (f) HMDS, 110°C, 1 h or TMS-Cl, *i*-Pr₂EtN, 0°C to rt; (g) H₂C=C(CH₃)COOEt, 90°C, 3.5 h or H₂C=C(CH₃)COOEt, 0°C to rt, 24 h, then EtOH, then deprotection 50% TFA.

[☆] PII of original article: S0040-4020(99)00910-2