


Contents

REPORT

The domino way to heterocycles

Albert Padwa* and Scott K. Bur*

pp 5341–5378




ARTICLES

Synthesis of indole analogues of the anti-*Helicobacter pylori* compounds CJ-13,015, CJ-13,102, CJ-13,104 and CJ-13,108

Zoe E. Wilson, Amanda M. Heapy and Margaret A. Brimble*


pp 5379–5385



Asymmetric organocatalysis of epoxidation by iminium salts under non-aqueous conditions

Philip C. Bulman Page,* Benjamin R. Buckley, David Barros, A. John Blacker, Brian A. Marples and Mark R. J. Elsegood


pp 5386–5393



Effect of substituent structure on pyrimidine electrophilic substitution


pp 5394–5405

Christiaan W. van der Westhuyzen,* Amanda L. Rousseau and Christopher J. Parkinson

**A novel three-component reaction for the synthesis of new 4-azafluorenone derivatives**


pp 5406–5414

Shuijiang Tu,* Bo Jiang, Hong Jiang, Yan Zhang, Runhong Jia, Junyong Zhang, Qingqing Shao, Chunmei Li, Dianxiang Zhou and Longji Cao

**Electroreductive crossed pinacol coupling of aromatic ketones with aliphatic ketones and aldehydes**


pp 5415–5426

Naoki Kise,* Yousuke Shiozawa and Nasuo Ueda


**Effect of extended conjugation with a phenylethynyl group on the fluorescence properties of water-soluble arylboronic acids**

pp 5427–5436

Shi-Long Zheng, Na Lin, Suazette Reid and Binghe Wang*




Synthesis and the effect of alkyl chain length on optoelectronic properties of diarylethene derivatives pp 5437–5449
 Chunhong Zheng, Shouzhi Pu,* Jingkun Xu, Mingbiao Luo, Dechao Huang and Liang Shen



Photochromic bis(2-*n*-alkyl-5-formyl-3-thienyl)perfluorocyclopentene derivatives have been synthesized. The effect of alkyl chain length on their optoelectronic properties was investigated in detail.

Equilibria of formation and dehydration of the carbinolamine intermediate in the reaction of benzaldehyde with hydrazine pp 5450–5454

Melek S. Baymak and Petr Zuman*




A highly efficient three-component coupling of aldehyde, terminal alkyne, and amine via C–H activation catalyzed by reusable immobilized copper in organic–inorganic hybrid materials under solvent-free reaction conditions pp 5455–5459

Pinhua Li and Lei Wang*




A branched, hydrogen-bonded heterodimer: a novel system for achieving high stability and specificity pp 5460–5469
 Jason B. Bialecki,* Li-Hua Yuan and Bing Gong

A 1:1 ratio of synthesized oligoamides **1** and **2** tightly bind in chloroform to form a very stable six hydrogen-bonded heterodimer **1•2**.



Synthesis of asymmetric halomesylate mustards with aziridineethanol/alkali metal halides: application to an improved synthesis of the hypoxia prodrug PR-104


Shangjin Yang,* Graham J. Atwell and William A. Denny



High pressure and thermal Diels–Alder reactions of 5-nitro[2.2]paracyclophanepyrans-6-one

pp 5477–5481


Lucio Minuti,* Assunta Marrocchi, Selvaggia Landi, Mirko Seri and Eszter Gacs-Baitz



Intramolecular addition of carbon radicals to aldehydes: synthesis of enantiopure tetrahydrofuran-3-ols


pp 5482–5489

Marcello Tiecco,* Lorenzo Testaferri, Francesca Marini,* Silvia Sternativo, Claudio Santi, Luana Bagnoli and Andrea Temperini



Efficient dual catalytic enantioselective diethylzinc addition to the exocyclic C=N double bond of some 1,2,4-N-triazinylarylimines using polymer-supported chiral β -amino alcohols derived from norephedrine

Ashraf A. El-Shehawy




Regiochemical control of the ring-opening of aziridines by means of chelating processes. Part 4:

pp 5501–5509


Regioselectivity of the opening reactions with MeOH of 1-(benzyloxy)-2,3- and -3,4-N-(methoxycarbonyl)aziridoalkanes under condensed and gas-phase operating conditions

Paolo Crotti,* Gabriele Renzi,* Graziella Roselli, Valeria Di Bussolo, Sandra Giuli,
Sofia Tsitsigia and Maria Rosaria Romano

**Toward breast cancer resistance protein (BCRP) inhibitors: design, synthesis of a series of new simplified fumitremorgin C analogues**


pp 5510–5528

Guofeng Wu, Jiawang Liu, Lanrong Bi, Ming Zhao,* Chao Wang, Michèle Baudy-Floc'h,
Jingfang Ju* and Shiqi Peng*

**Ferrocenylimidazoline palladacycles: efficient phosphine-free catalysts for Suzuki–Miyaura cross-coupling reaction**


pp 5529–5538

Ji Ma, Xiuling Cui,* Bi Zhang, Maoping Song and Yangjie Wu*

**Synthesis and pH-dependent self-assembly of semifluorinated calix[4]arenes**

pp 5539–5547


Oana M. Martin and Sandro Mecozzi*



Assignment of ^{13}C NMR spectrum for blepharismin C based on biosynthetic studies

pp 5548–5553

Yoshiyuki Uruma, Kei Sakamoto, Kohei Takumi, Matsumi Doe, Yoshinosuke Usuki and Hideo Iio*




Analyses of the HMQC, HMBC, and INADEQUATE spectra of ^{13}C -enriched blepharismin confirmed the assignment of the ^{13}C NMR signals.

A regioselective route to 5-substituted pyrazole- and pyrazoline-3-phosphonic acids and esters

pp 5554–5560

P. Conti,* A. Pinto, L. Tamborini, V. Rizzo and C. De Micheli




R = Ph, CO₂Me, C₄H₉, C₉H₁₉

Conformational heterogeneity of tripeptides containing Boc–Leu–Aib as corner residues in the solid state


pp 5561–5566

Debasish Haldar,* Michael G. B. Drew and Arindam Banerjee*

**Synthesis of tetrahydroisoquinoline-based pseudopeptides and their characterization as suitable reverse turn mimetics**


pp 5567–5578

Giordano Lesma,* Elisa Meschini, Teresa Recca, Alessandro Sacchetti and Alessandra Silvani*



Synthesis of novel 3'-spirocyclic-oxindole derivatives and assessment of their cytostatic activities
 Sarah R. Yong, Alison T. Ung,* Stephen G. Pyne,* Brian W. Skelton and Allan H. White



pp 5579–5586



Synthesis of the Arctic sponge alkaloid viscosaline and the marine sponge alkaloid theonelladin C
 Benjamin J. Shorey, Victor Lee* and Jack E. Baldwin*


pp 5587–5592

The synthesis of two sponge alkaloids, viscosaline and theonelladin C is described.




Iron(III) chloride-catalyzed convenient one-pot synthesis of β -acetamido carbonyl compounds
 Abu T. Khan,* Tasneem Parvin and Lokman H. Choudhury

pp 5593–5601




Monovalent cation-promoted ordering of a glycine-rich cyclic peptide
 K. B. Joshi and Sandeep Verma*

pp 5602–5607



Gadolinium triflate catalyzed alkylation of pyrroles: efficient synthesis of 3-oxo-2,3-dihydro-1*H*-pyrrolizine derivatives


Canan Unaleroglu* and Arife Yazici



Copper(I)-catalysed homo-coupling of aryl diazonium salts: synthesis of symmetrical biaryls

Ivica Cepanec,* Mladen Litvić, Josipa Udiković, Ivan Pogorelić and Marija Lovrić

pp 5614–5621




(i) 1.5 equiv CuOTf; MeCN; 0 °C-rt; overnight; R=EWG(s); (ii) 20 mol % Cu(OTf)₂, bpy; 3 equiv Cu; MeCN; rt; overnight; R=H or EDG(s); 15 examples; yields: 7–96%.

Stereoselective syntheses of 20-*epi* cholanic acid derivatives from 16-dehydropregnенолон acetate

Bapurao B. Shingate, Braja G. Hazra,* Vandana S. Pore, * Rajesh G. Gonnade and Mohan M. Bhadbhade

pp 5622–5635




i+

A facile one-pot access to cone and 1,3-alternate conformers of calix[4]arene-bis(amido)crowns

Har Mohindra Chawla,* Suneel Pratap Singh and Shailesh Upreti


pp. 5636–5642



A straightforward preparation of fluorine-containing 1,2-dihydropyrimidines and pyrimidines with 2,2-dihydropolyfluoroalkylaldehydes

Xian-Jin Yang, Li-Si Zhang and Jin-Tao Liu*


pp 5643–5648



1,2,3,4-Tetrahydro-1,5-naphthyridines and related heterocyclic scaffolds: exploration of suitable chemistry for library development

Grace H. C. Woo, Aaron B. Beeler and John K. Snyder*


pp 5649–5655



A convenient approach to λ^5 -phosphinines via interaction of phosphorylated 3-pyrrolidinocrotonitrile with 2-bromoacetophenones

Yuri V. Svyaschenko, Alexandr N. Kostyuk,* Bogdan B. Barnych and Dmitriy M. Volochnyuk


pp 5656–5664



Computational study on the ring-opening reaction of protonated oxirane and methylpropene


James M. Coxon* and Michael A. E. Townsend

pp 5665–5668




Regioselective synthesis of pyridines and dihydropyridines derived from β -amino acids and aminophosphonates by reaction of *N*-vinylic phosphazenes with α,β -unsaturated ketones
 Francisco Palacios,* Esther Herrán, Gloria Rubiales and Concepción Alonso

pp 5669–5676



Efficient functional molecule incorporation method to functionalized peptide nucleic acid (PNA): use in synthesis of labeled PNA oligomers pp 5677–5689

Hisafumi Ikeda,* Fumihiko Kitagawa and Yushin Nakamura



OTHER CONTENTS

Tetrahedron reports on organic chemistry

pp I–XX

Tetrahedron Perspectives

p XXI

Tetrahedron reports on organic chemistry order form

p XXII

Tetrahedron reports on organic chemistry author index

pp XXIII–XXX

*Corresponding author

①[†] Supplementary data available via ScienceDirectFull text of this journal is available, on-line from **ScienceDirect**. Visit www.sciencedirect.com for more information.

Abstracted/indexed in: AGRICOLA, Beilstein, BIOSIS Previews, CAB Abstracts, Chemical Abstracts, Current Contents: Life Sciences, Current Contents: Physical, Chemical and Earth Sciences, Current Contents Search, Derwent Drug File, Ei Compendex, EMBASE/Excerpta Medica, Medline, PASCAL, Research Alert, Science Citation Index, SciSearch. Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®



ISSN 0040-4020