


Contents

REPORT

Formation of five- and six-membered heterocyclic rings under radical cyclisation conditions

pp 793–826

Krishna C. Majumdar,* Pradipta K. Basu and Sudip K. Chattopadhyay




ARTICLES

Probing the ribosomal RNA A-site with functionally diverse analogues of paromomycin—synthesis of ring I mimetics


pp 827–846

Stephen Hanessian,* Susanta Adhikari, Janek Szychowski, Kandasamy Pachamuthu, Xiaojing Wang, Michael T. Migawa, Richard H. Griffey and Eric E. Swayze

**Synthesis of novel 2,3-substituted quinazolin-4-ones by condensation of alkyl or aromatic diamines with 5-(*N*-arylimino)-4-chloro-5*H*-1,2,3-dithiazoles**


pp 847–854

Maria de Fatima Pereira, Valérie Thiéry and Thierry Besson*



A practical diastereoselective synthesis of β -hydroxy- β -trifluoromethyl imines
Zhen-Jiang Liu, Ying-Qiao Mei and Jin-Tao Liu*


pp 855–860



Asymmetric carbon–carbon bond formation reaction of chiral *N*-(*tert*-butanesulfinyl) ketimines and α -trifluoromethylated ketones was achieved to give chiral β -hydroxy- β -trifluoromethyl imines in moderate to high yields with good diastereoselectivities.

Facile synthesis of azocino[4,3-*b*]indoles by ring-closing metathesis
M.-Lluïsa Bennasar,* Ester Zulaica, Daniel Solé and Sandra Alonso


pp 861–866



Synthesis of latonduine derivatives via intramolecular Heck reaction

pp 867–879


Aurélien Putey, Lionel Joucla, Laurent Picot, Thierry Besson and Benoît Joseph*



Synthesis of a regioisomeric analogue of the 3C-protease inhibitor thysanone via a Hauser annulation strategy

pp 880–887


Margaret A. Brimble,* Scott I. Houghton and Paul D. Woodgate



Highly efficient ring opening reactions of epoxides with deactivated aromatic amines catalyzed by heteropoly acids in water

Najmedin Azizi and Mohammad R. Saidi*


pp 888–891



Microwave-assisted solvent-free synthesis of substituted 2-quinolones

Cheng-Sheng Jia, Ya-Wei Dong, Shu-Jiang Tu and Guan-Wu Wang*


pp 892–897



Introducing difluoromethylene sulfonamide group via nucleophilic addition of difluoromethylene anion with aromatic aldehydes

Jun-Li Li and Jin-Tao Liu*


pp 898–903



1,8-Diazabicyclo[5.4.0]undec-7-ene (DBU)-promoted efficient and versatile aza-Michael addition

Chang-Eun Yeom, Mi Jeong Kim and B. Moon Kim*


pp 904–909



Reaction of chromone-3-carbaldehyde with α -amino acids—syntheses of 3- and 4-(2-hydroxybenzoyl)pyrroles

pp 910–917


Andrea G. P. R. Figueiredo, Augusto C. Tomé, Artur M. S. Silva* and José A. S. Cavaleiro



Threonine aldolases—an emerging tool for organic synthesis

pp 918–926


Johannes Steinreiber, Kateryna Fesko, Christoph Reisinger, Martin Schürmann, Friso van Assema, Michael Wolberg, Daniel Mink and Herfried Griengl*



Cyclic polysulfides fused to polyaromatics

pp 927–933

Ashraful Alam, Masaru Kon-no, Satoshi Ogawa and Ryu Sato*




Syntheses and X-ray crystal structures of cyclic polysulfides fused to naphthalene and phenanthrene have been studied.

Unprecedented formation of stable ketene-*N,O*-acetals and their rearrangement under basic conditions


pp 934–940

Matthias Breuning* and Tobias Häuser




Non-symmetrical oligothiophenes with ‘incompatible’ substituents
Delphine Didier, Sergey Sergeyev and Yves Henri Geerts*

pp 941–946




Isoprene-catalysed lithiation: deprotection and functionalisation of imidazole derivatives
Rosario Torregrosa, Isidro M. Pastor and Miguel Yus*

pp 947–952




Novel illustrations of the specific reactivity of 1,1-diamino-2,2-dinitroethene (DADNE) leading to new unexpected compounds
Grégoire Hervé* and Guy Jacob

pp 953–959



One-pot two-step synthesis of 4-vinylphenols from 4-hydroxy substituted benzaldehydes under microwave irradiation: a new perspective on the classical Knoevenagel–Doebner reaction
Arun K. Sinha,* Anuj Sharma and Bhupendra P. Joshi


pp 960–965



Deoxygenation of sulfoxides, selenoxides, telluroxides, sulfones, selenones and tellurones with Mg–MeOH

pp 966–969


Jitender M. Khurana,* Vandana Sharma and Silvi N. Chacko



New synthetic routes towards various α -fluorinated aryl ketones and their enantioselective reductions using baker's yeast

pp 970–976


Balaka Barkakaty, Yutaka Takaguchi and Sadao Tsuboi*



An efficient synthesis of fused tricycles with a benzene core via intramolecular double ring-closing enyne metathesis

pp 977–985


Zhiqiang Liang, Shengming Ma,* Jihong Yu and Ruren Xu



A fast and highly efficient protocol for Michael addition of *N*-heterocycles to α,β -unsaturated compound using basic ionic liquid [bmIm]OH as catalyst and green solvent

pp 986–990


Jian-Ming Xu, Chao Qian, Bo-Kai Liu, Qi Wu and Xian-Fu Lin*



Stereoselective cycloadditions of (*1Z,4R*,5R)-1-arylmethylidene-4-benzoylamino-5-phenylpyrazolidin-3-on-1-azomethine imines to maleimides**


Lidija Pezdirc, Janez Cerkovnik, Samo Pirc, Branko Stanovnik and Jurij Svetec*

pp 991–999



An efficient chemoselective strategy for the preparation of (*E*)-cinnamic esters from cinnamaldehydes pp 1000–1007 using heterogeneous catalyst and DDQ


Arun K. Sinha,* Anuj Sharma, Anand Swaroop and Vinod Kumar



Structures of four new alkaloids from *Stemona sessilifolia*

Yukio Hitotsuyanagi, Maho Hikita, Takahisa Oda, Daichi Kakuta, Haruhiko Fukaya and Koichi Takeya*


pp 1008–1013



A general route to 5,6-seco-hexahydrodibenzopyrans and analogues: first total synthesis of (+)-Machaeridiol B and (+)-Machaeriol B

Qinggang Huang, Qiaoling Wang, Jiyue Zheng, Jiyong Zhang, Xinfu Pan and Xuegong She*

pp 1014–1021



OTHER CONTENT**Corrigendum****p 1022*****Corresponding author****i[†]** Supplementary data available via ScienceDirect**COVER**

Threonine aldolases have had a rather slow start in the synthesis of β -hydroxy- α -amino acids due to low selectivities and yields. Here we report for the first time on a stereoselective synthesis of D-syn-phenylserine and its derivatives at high conversion rates using D-threonine aldolase from *Alcaligenes xylosoxidans*. The L-diastereomers were obtained at similar rates as L-syn/anti-mixtures using L-threonine aldolase from *Pseudomonas putida*. *Tetrahedron* 2007, 63, 918–926.

© 2006. H. Griengl. Published by Elsevier Ltd.



Full text of this journal is available, on-line from **ScienceDirect**. Visit www.sciencedirect.com for more information.

Abstracted/indexed in: AGRICOLA, Beilstein, BIOSIS Previews, CAB Abstracts, Chemical Abstracts, Current Contents: Life Sciences, Current Contents: Physical, Chemical and Earth Sciences, Current Contents Search, Derwent Drug File, Ei Compendex, EMBASE/Excerpta Medica, Medline, PASCAL, Research Alert, Science Citation Index, SciSearch. Also covered in the abstract and citation database SCOPUS®. Full text available on ScienceDirect®



ISSN 0040-4020