



A convenient synthesis of 2-substituted indoles by the reaction of 2-(chloromethyl)phenyl isocyanides with organolithiums

Kazuhiro Kobayashi*, Daisuke Iitsuka, Shuhei Fukamachi, Hisatoshi Konishi

Division of Applied Chemistry, Department of Chemistry and Biotechnology, Graduate School of Engineering, Tottori University, 4-101 Koyama-minami, Tottori 680-8552, Japan

ARTICLE INFO

Article history:

Received 7 April 2009

Received in revised form 30 June 2009

Accepted 30 June 2009

Available online 4 July 2009

Keywords:

Indoles

Isocyanides

Organolithiums

Intramolecular substitution

Imidoyl anion

ABSTRACT

The reaction of 2-(chloromethyl)phenyl isocyanides, readily available by dehydration of the respective *N*-[2-(chloromethyl)phenyl]formamides, with organolithiums produced 2-substituted indoles in satisfactory yields through addition of organolithiums to the isocyano carbon followed by intramolecular substitution reaction of the resulting imidoyl anion intermediates.

© 2009 Elsevier Ltd. All rights reserved.

1. Introduction

Indoles are undoubtedly very important heterocycles in organic synthesis. Therefore, development of new and efficient methods for the general preparation of indoles has been the subject of intense research.¹ In this paper we wish to describe that a new and convenient method for the preparation of 2-substituted indoles, which is based on reactions of 2-(chloromethyl)phenyl isocyanides with organolithiums, has been developed. The utility of *ortho*-functionalized phenyl isocyanides in heterocycle synthesis is well established.^{2–5} In particular, Fukuyama and Tokuyama have elegantly demonstrated the utility of 2-isocyanostyrene derivatives in the synthesis of indole derivatives including natural products.² A synthesis of 2,3-disubstituted quinolines through cyclization of 2-alkynylisocyanobenzenes has been reported by Ito and Sugimoto.³ Ichikawa has reported syntheses of 3-fluoroquinoline derivatives based on reactions utilizing β -difluoro-2-isocyanostyrenes.⁴ However, there have been few reports on utilization of 2-(chloromethyl)phenyl isocyanides in organic synthesis so far, though 2-(chloromethyl)phenyl isocyanide has been used in the synthesis of 1- and 2-platinum(II)-substituted indole derivatives by Michelin and colleagues.⁶ To the best of our knowledge, the present report is the first example of the practical use of these isocyanide in organic synthesis.



2. Results and discussion

Our preparation of 2-substituted indoles **3** was conducted as illustrated in Scheme 1. The starting materials of the present synthesis, *N*-(2-chlorophenyl)formamides **1**, were easily prepared by formylation of 2-aminobenzyl alcohols with ethyl or butyl formate followed by chlorination with thionyl chloride. These formamides **1** were converted into 2-(chloromethyl)phenyl isocyanides **2** on treatment with phosphorous oxychloride in the presence of triethylamine in THF.⁷ Because these isocyanides **2** were unstable under purification conditions (distillation and chromatography on silica gel), they were used in the next step without any purification after the usual workup as soon as possible. Thus, crude isocyanides **2** were allowed to react with three molar amounts of organolithiums in THF at -78 °C. After stirring for 2 h, the usual workup gave, after purification of preparative TLC on silica gel, the desired products **3**. The results are summarized in Scheme 1, which indicates that overall yields of **3** from **1** are generally moderate-to-fair. When 5-chloro-2-(chloromethyl)phenyl isocyanide (**2b**) was used, somewhat higher yields of the corresponding desired indoles **3** were obtained, compared to those using other 2-(chloromethyl)phenyl isocyanides **2a** and **2c**. These results may be attributed to higher reactivity of the isocyano carbon toward nucleophiles due to the chloro substituent on the benzene ring.


The formation of 2-substituted indoles **3** is induced by attack of an organolithium at the isocyano carbon of 2-(chloromethyl)phenyl isocyanides **2** to generate the imidoyl anion intermediate **4**, as depicted in Scheme 2. This intermediate undergoes intramolecular

* Corresponding author. Tel./fax: +81 857 31 5263.

E-mail address: kkoba@chem.tottori-u.ac.jp (K. Kobayashi).



Scheme 1.



Scheme 2.

S_N2 reaction to give the indolenine intermediate **5**, which, probably, tautomerized during workup and/or purification procedures to afford **3**. The possibility of lithiation of **5** to an indolyl lithium intermediate with excess organolithium is excluded, because addition of iodomethane to the reaction mixture did not give the corresponding *N*-methyl derivatives. The use of three molar amounts of organolithiums is essential for the satisfactory production of the desired products. With less than three molar amounts of organolithiums the reaction sequence did not proceed in a satisfactory manner and gave mixtures of only rather decreased yields of the desired products and the starting materials. One of the probable reasons for the modest yields of the products may be ascribed to somewhat lower purity of 2-(chloromethyl)phenyl isocyanides used without purification; no other products, whose structures could be determined, were isolated. Although the reason for the necessity of excess organolithiums in the present reaction is not clear, the uses of excess nucleophiles have often been reported in the heterocycle synthesis initiated by the attack of nucleophiles on the isocyno carbon.^{4b,5f} When the reaction sequence was carried out in diethyl ether or 1,2-dimethoxyethane, an intractable mixture of products including ones probably arising from deprotonation of the benzyl proton of **2** was obtained in each case. No trace of **3** was isolated from this mixture. Grignard reagents, such as phenylmagnesium bromide and ethylmagnesium bromide did not work well in the present indole formation to give a complex reaction mixture in each case.

In conclusion, the results described above have shown that the synthesis of 2-substituted indoles is conveniently accomplished from readily available 2-(chloromethyl)phenyl isocyanides. We believe that this procedure may be valuable in organic synthesis because of its simplicity.

3. Experimental

3.1. General

All melting points were obtained on a Laboratory Devices MEL-TEMP II melting apparatus and are uncorrected. IR spectra were determined with a Shimadzu FTIR-8300 spectrophotometer. The ¹H NMR spectra were determined in CDCl₃ using TMS as an internal reference with a JEOL ECP500 FT NMR spectrometer operating at 500 MHz. The ¹³C NMR spectra were determined in CDCl₃ using TMS as an internal reference with a JEOL ECP500 FT NMR spectrometer operating at 125 MHz. Low- and high-resolution MS spectra (EI, 70 eV) were measured by a JEOL JMS AX505 HA spectrometer. TLC was carried out on a Merck Kieselgel 60 PF₂₅₄. Column chromatography was performed using Merck Kieselgel 60 (0.063–0.200 mm). All of the organic solvents used in this study were dried over appropriate drying agents and distilled prior to use.

3.2. Starting materials

N-[2-(Chloromethyl)phenyl]formamide (**1a**) was prepared by the reported method¹ from *N*-[2-(hydroxymethyl)phenyl]formamide,⁶ which was prepared by treating 2-aminobenzylalcohol with refluxing ethyl formate. 2-Amino-4-chlorobenzyl alcohol⁸ was prepared by the LAH reduction of ethyl 2-amino-4-chlorobenzoate.⁹ All other chemicals used in this study were commercially available.

3.2.1. 1-Chloromethyl-2-isocyanobenzene (**2a**)^{6,10}

This compound was prepared from *N*-[2-(chloromethyl)phenyl]formamide⁶ by a modified procedure^{7a} of Ugi's method.^{7b} Thus, to a stirred solution of *N*-[2-(chloromethyl)phenyl]formamide (0.30 g, 1.8 mmol) in THF (6 mL) containing Et₃N (1.3 g, 12 mmol) at 0 °C was added POCl₃. After 10 min the mixture was treated with saturated aqueous NaHCO₃ (15 mL) and extracted with Et₂O twice (10 mL each). The combined extracts were washed with saturated aqueous NaHCO₃ and then brine, and dried over anhydrous K₂CO₃. After evaporation of the solvent the residue (a brown liquid) was used in the next step without any purification.

3.2.2. *N*-[5-Chloro-2-(hydroxymethyl)phenyl]formamide

This compound was prepared by treating 2-amino-4-chlorobenzyl alcohol with refluxing ethyl formate in 66% yield; a white solid; mp 124–126 °C (hexane–Et₂O); IR (KBr) 3300, 1682 cm⁻¹; ¹H NMR δ 1.68 (br s, 1H), 4.72 and 4.73 (2s, combined 2H), 7.06–8.68 (m, 5H). Anal. Calcd for C₈H₈ClNO₂: C, 51.77; H, 4.34; N, 7.55. Found: C, 51.70; H, 4.39; N, 7.32.

3.2.3. *N*-[5-Chloro-2-(chloromethyl)phenyl]formamide (**1b**)

This compound was prepared by treating *N*-[5-chloro-2-(hydroxymethyl)phenyl]formamide with SOCl₂ according to the reported procedure¹ in 55% yield; mp 95–97 °C (hexane–Et₂O); IR (KBr) 3279, 1665 cm⁻¹; ¹H NMR δ 4.57 and 4.59 (2s, combined 2H), 7.14–8.61 (m, 5H). Anal. Calcd for C₈H₇Cl₂NO: C, 47.09; H, 3.46; N, 6.86. Found: C, 46.93; H, 3.51; N, 6.61.

3.2.4. 1-Chloro-4-chloromethyl-3-isocyanobenzene (**2b**)

This compound was prepared from *N*-[5-chloro-2-(chloromethyl)phenyl]formamide by a similar procedure as described for the preparation of **2a**, and used in the next step without any purification after the usual workup; a brown liquid; IR (neat) 2121 cm⁻¹; ¹H NMR δ 4.67 (s, 2H), 7.42 (dd, *J*=8.2, 2.3 Hz, 1H), 7.44 (d, *J*=2.3 Hz, 1H), 7.47 (d, *J*=8.2 Hz, 1H); MS *m/z* 185 (M⁺, 100). HRMS Calcd for C₈H₅Cl₂N: M, 184.9799. Found: *m/z* 184.9776.

3.2.5. *N*-[2-(Hydroxymethyl)-4-methylphenyl]formamide

This compound was prepared by treating 2-amino-5-methylbenzyl alcohol with refluxing butyl formate in 60% yield; a white solid; mp 96–98 °C (hexane–CHCl₃); IR (KBr) 3231, 1657 cm⁻¹; ¹H NMR δ 2.08 and 2.20 (2br s, combined 1H), 2.31 and 2.33 (2s, combined 3H), 4.69 and 4.70 (2s, combined 2H), 7.02–8.60 (m, 5H). C₉H₁₁NO₂: C, 65.44; H, 6.71; N, 8.48. Found: C, 65.39; H, 6.77; N, 8.44.

3.2.6. *N*-[2-(Chloromethyl)-4-methylphenyl]formamide (**1c**)

This compound was prepared by treating *N*-[2-(hydroxymethyl)-4-methylphenyl]formamide with SOCl₂ according to the reported procedure¹ in 48% yield; a white solid; mp 117–119 °C (hexane–Et₂O); IR (KBr) 3231, 1661 cm⁻¹; ¹H NMR δ 2.33 and 2.35 (2s, combined 3H), 4.57 and 4.59 (2s, combined 2H), 7.11–8.53 (m, 5H). C₉H₁₀ClNO: C, 58.86; H, 5.49; N, 7.63. Found: C, 58.81; H, 5.51; N, 7.40.

3.2.7. 1-Chloromethyl-5-methyl-2-isocyanobenzene (**2c**)

This compound was prepared from *N*-[2-(chloromethyl)-4-methylphenyl]formamide by a similar procedure as described for the preparation of **2a**, and used in the next step without any purification after the usual workup; a brown liquid; IR (neat) 2120 cm⁻¹; ¹H NMR δ 2.39 (s, 3H), 4.67 (s, 2H), 7.16 (dd, *J*=7.8, 1.4 Hz, 1H), 7.30 (d, *J*=7.8 Hz, 1H), 7.31 (d, *J*=1.4 Hz, 1H); MS *m/z* 165 (M⁺, 100). HRMS Calcd for C₉H₈ClN: M, 165.0345. Found: *m/z* 165.0337.

3.3. Typical procedure for the preparation of 2-substituted indoles **3**

3.3.1. 2-Phenylindole (**3a**)¹¹

To a stirred solution of **2a** (0.15 g, 1.0 mmol) in THF (4 mL) at –78 °C was added dropwise PhLi (3.0 mmol; 1.04 M in cyclohexane–Et₂O); the mixture was stirred at the same temperature for 2 h. The resulting mixture was diluted by adding 15 mL of Et₂O, and saturated aqueous NH₄Cl (10 mL) and water (5 mL) was added. The layers were separated, and the aqueous layer was extracted with Et₂O twice (5 mL each). The combined extracts were washed with brine, dried over anhydrous Na₂SO₄, and the solvent was evaporated. The residue was purified by preparative TLC on silica gel (1:7 THF–hexane) to give **3a** (0.13 g, 67%); colorless needles; mp 189–190 °C (hexane–Et₂O). This product was identified by a direct comparison with a commercially available sample, purchased from Wako Pure Chemical Industries, Ltd.

3.3.2. 2-(4-Methylphenyl)indole (**3b**)¹²

Colorless needles; mp 214–216 °C (hexane–Et₂O) (lit.,¹³ 213–214 °C). The spectral data (IR and ¹H NMR) were identical to those reported previously.¹³

3.3.3. 2-Butylindole (**3c**)¹⁴

A white solid; mp 35–36 °C (hexane–Et₂O) (lit.,¹⁴ 36–37 °C). The spectral data (IR and ¹H NMR) were identical to those reported previously.¹⁵

3.3.4. 6-Chloro-2-phenylindole (**3d**)¹⁶

Colorless needles; mp 180–181 °C (hexane) (lit.,¹⁷ 181–182 °C). The spectral data (IR and ¹H NMR) were identical to those reported previously.¹⁷

3.3.5. 6-Chloro-2-(2-methylphenyl)indole (**3e**)

Colorless needles; mp 105–106 °C (hexane); IR (KBr) 3373, 1602 cm⁻¹; ¹H NMR δ 2.49 (s, 3H), 6.59 (dd, *J*=1.4, 0.9 Hz, 1H), 7.11 (dd, *J*=8.2, 1.8 Hz, 1H), 7.27–7.33 (m, 3H), 7.40 (t, *J*=0.9 Hz, 1H), 7.46 (dd, *J*=8.7, 3.2 Hz, 1H), 7.54 (d, *J*=8.2 Hz, 1H), 8.14 (br s, 1H); MS *m/z* 241 (M⁺, 100). Anal. Calcd for C₁₅H₁₂ClN: C, 74.53; H, 5.00; N, 5.79. Found: C, 74.49; H, 5.13; N, 5.77.

3.3.6. 6-Chloro-2-(3-methylphenyl)indole (**3f**)

Colorless needles; mp 135–136 °C (hexane–Et₂O); IR (KBr) 3433, 1609 cm⁻¹; ¹H NMR δ 2.43 (s, 3H), 6.78 (dd, *J*=1.4, 0.9 Hz, 1H), 7.08 (dd, *J*=8.7, 1.8 Hz, 1H), 7.16 (d, *J*=7.8 Hz, 1H), 7.34 (dd, *J*=7.8, 7.3 Hz, 1H), 7.38 (s, 1H), 7.45 (d, *J*=7.3 Hz, 1H), 7.47 (d, *J*=1.8 Hz, 1H), 7.52 (d, *J*=8.7 Hz, 1H), 8.32 (br s, 1H); MS *m/z* 241 (M⁺, 100). Anal. Calcd for C₁₅H₁₂ClN: C, 74.53; H, 5.00; N, 5.79. Found: C, 74.48; H, 5.24; N, 5.53.

3.3.7. 6-Chloro-2-(3-chlorophenyl)indole (**3g**)

Colorless crystals; mp 136–137 °C (hexane–Et₂O); IR (KBr) 3433, 1601 cm⁻¹; ¹H NMR δ 6.81 (d, *J*=1.8 Hz, 1H), 7.10 (dd, *J*=8.7, 1.8 Hz, 1H), 7.31 (dt, *J*=7.8, 0.9 Hz, 1H), 7.38 (t, *J*=7.8 Hz, 1H), 7.40 (s, 1H), 7.52 (d, *J*=7.8 Hz, 1H), 7.53 (d, *J*=8.7 Hz, 1H), 7.63 (d, *J*=1.8 Hz, 1H), 8.32 (br s, 1H); MS *m/z* 261 (M⁺, 100). Anal. Calcd for C₁₄H₉Cl₂N: C, 64.15; H, 3.46; N, 5.34. Found: C, 63.96; H, 3.49; N, 5.22.

3.3.8. 6-Chloro-2-(4-chlorophenyl)indole (**3h**)

Colorless crystals; mp 164–165 °C (hexane–Et₂O); IR (KBr) 3381 cm⁻¹; ¹H NMR δ 6.78 (d, *J*=1.4 Hz, 1H), 7.10 (dd, *J*=8.2, 1.8 Hz, 1H), 7.39 (s, 1H), 7.42 (d, *J*=8.7 Hz, 2H), 7.52 (d, *J*=8.2 Hz, 1H), 7.57 (d, *J*=8.7 Hz, 2H), 8.31 (br s, 1H); ¹³C NMR δ 100.39, 110.87, 121.23, 121.55, 126.31, 127.71, 128.34, 129.30, 130.43, 133.76, 137.20, 137.43; MS *m/z* 261 (M⁺, 100). Anal. Calcd for C₁₄H₉Cl₂N: C, 64.15; H, 3.46; N, 5.34. Found: C, 63.97; H, 3.56; N, 5.09.

3.3.9. 6-Chloro-2-(3-methoxyphenyl)indole (**3i**)

Colorless crystals; mp 105–107 °C (hexane–Et₂O); IR (KBr) 3451, 1605 cm⁻¹; ¹H NMR δ 3.88 (s, 3H), 6.79 (d, *J*=1.4 Hz, 1H), 6.89 (dd, *J*=8.7, 2.3 Hz, 1H), 7.09 (dd, *J*=8.2, 1.8 Hz, 1H), 7.17 (dd, *J*=2.3, 1.8 Hz, 1H), 7.23 (d, *J*=7.8 Hz, 1H), 7.37 (dd, *J*=8.2, 7.8 Hz, 1H), 7.39 (s, 1H), 7.52 (d, *J*=8.7 Hz, 1H), 8.33 (br s, 1H); MS *m/z* 257 (M⁺, 100). Anal. Calcd for C₁₅H₁₂ClNO: C, 69.91; H, 4.69; N, 5.43. Found: C, 69.65; H, 4.94; N, 5.44.

3.3.10. 6-Chloro-2-(naphthalene-1-yl)indole (**3j**)

A white solid; mp 130–131 °C (hexane–Et₂O); IR (KBr) 3346 cm⁻¹; ¹H NMR δ 6.78 (d, *J*=1.4 Hz, 1H), 7.15 (dd, *J*=8.2, 1.8 Hz, 1H), 7.45 (s, 1H), 7.52–7.56 (m, 3H), 7.61 (d, *J*=8.2 Hz, 1H), 7.63 (dd, *J*=7.3, 1.4 Hz, 1H), 7.91 (d, *J*=8.2 Hz, 1H), 7.93 (dd, *J*=6.9, 2.3 Hz, 1H), 8.27 (dd, *J*=6.9, 1.8 Hz, 1H), 8.33 (br s, 1H); ¹³C NMR δ 103.64, 110.78, 120.90, 121.41, 125.32, 125.44, 126.24, 126.80, 127.21, 127.42, 127.93, 128.54, 128.84, 130.56, 131.40, 133.91, 136.66, 137.44; MS *m/z* 277 (M⁺, 100). Anal. Calcd for C₁₈H₁₂ClN: C, 77.84; H, 4.35; N, 5.04. Found: C, 77.59; H, 4.51; N, 4.91.

3.3.11. 6-Chloro-2-(1,1-dimethylethyl)indole (**3k**)

Colorless crystals; mp 94–95 °C (hexane); IR (KBr) 3445, 1612 cm⁻¹; ¹H NMR δ 1.38 (s, 9H), 6.22 (dd, *J*=2.3, 0.9 Hz, 1H), 7.03 (dd, *J*=8.2, 1.8 Hz, 1H), 7.23 (d, *J*=1.8 Hz, 1H), 7.42 (d, *J*=8.2 Hz, 1H), 7.92 (br s, 1H); ¹³C NMR δ 30.18, 31.84, 97.03, 110.32, 120.19, 120.71, 126.73, 127.06, 136.10, 149.55; MS *m/z* 207 (M⁺, 100), 192 (100). Anal. Calcd for C₁₂H₁₄ClN: C, 69.39; H, 6.79; N, 6.74. Found: C, 69.49; H, 6.94; N, 6.67.

3.3.12. 5-Methyl-2-phenylindole (**3l**)¹⁸

Colorless needles; mp 217–218 °C (hexane) (lit.,¹⁹ 216–217 °C; lit.,²⁰ 210–211 °C). The spectral data (IR and ¹H NMR) were identical to those reported previously.^{19,20}

Acknowledgements

We thank Mrs. Miyuki Tanmatsu of this University for her assistance in determining the mass spectra and performing combustion analyses.

References and notes

- For a recent excellent review: (a) Cacchi, S.; Fabrizi, G. *Chem. Rev.* **2005**, *105*, 2873–2920; Recent general syntheses of indoles: (b) Tokuyama, H.; Makido, T.; Han-ya, Y.; Fukuyama, T. *Heterocycles* **2007**, *72*, 191–197; (c) Nielsen, S. D.; Ruhland, T.; Rasmussen, L. K. *Synlett* **2007**, 443–446; (d) Xu, D.-Q.; Yang, W.-L.; Luo, S.-P.; Wang, B.-T.; Wu, J.; Xu, Z.-Y. *Eur. J. Org. Chem.* **2007**, 1007–1021; (e) Schwarz, N.; Alex, K.; Ali Sayyed, I.; Khedkar, V.; Tillack, A.; Beller, M. *Synlett* **2007**, 1091–1095; (f) Della Rosa, C.; Kneeteman, M.; Mancini, P. *Tetrahedron Lett.* **2007**, *48*, 1435–1438; (g) Sayyed, I. A.; Alex, K.; Tillack, A.; Schwarz, N.; Michalik, D.; Beller, M. *Eur. J. Org. Chem.* **2008**, 4525–4528; (h) Fang, Y.-Q.; Lautens, M. *J. Org. Chem.* **2008**, *73*, 538–549; (i) Cucek, K.; Vercek, B. *Synthesis* **2008**, 1741–1746; (j) Sanz, R.; Guilarte, V.; Castroviejo, E. P. *Synlett* **2008**, 3006–3010; (k) Bondzic, B. P.; Farwick, A.; Liebich, J.; Elibracht, P. *Org. Biomol. Chem.* **2008**, *6*, 3723–3731; (l) Stuart, D. R.; Bertrand-Laperle, M.; Burgess, K. M. N.; Fagnou, K. *J. Am. Chem. Soc.* **2008**, *130*, 16474–16475; (m) Donaldo, J. R.; Taylor, R. J. *Synlett* **2009**, 59–62; (n) Varma, P. P.; Sherigara, B. S.; Mahadevan, K. M.; Hulikal, V. *Synth. Commun.* **2009**, *39*, 158–165; (o) Nakamura, I.; Nemoto, T.; Shiraiwa, N.; Terada, M. *Org. Lett.* **2009**, *11*, 1055–1058; (p) Lehmann, F.; Holm, M.; Laufer, S. *Tetrahedron Lett.* **2009**, *50*, 1708–1709.
- An excellent review: (a) Tokuyama, H.; Fukuyama, T. *Chem. Rec.* **2002**, *2*, 37–45; See also: (b) Kotani, M.; Yamago, S.; Satoh, A.; Tokuyama, H.; Fukuyama, T. *Synlett* **2005**, 1893–1896 and references in these papers.
- Suginome, M.; Fukuda, T.; Ito, Y. *Org. Lett.* **1999**, *1*, 1977–1979.
- (a) Ichikawa, J.; Wada, Y.; Miyazaki, H.; Mori, T.; Kuroki, H. *Org. Lett.* **2003**, *5*, 1455–1458; (b) Ichikawa, J.; Mori, T.; Miyazaki, H.; Wada, Y. *Synlett* **2004**, 1219–1222; (c) Mori, T.; Ichikawa, J. *Synlett* **2007**, 1169–1171.
- Contributions from our laboratory: (a) Kobayashi, K.; Kitamura, T.; Yoneda, K.; Morikawa, O.; Konishi, H. *Chem. Lett.* **2000**, 798–799; (b) Kobayashi, K.; Irisawa, S.; Matoba, T.; Matsumoto, T.; Yoneda, K.; Morikawa, O.; Konishi, H. *Bull. Chem. Soc. Jpn.* **2001**, *74*, 1109–1114; (c) Kobayashi, K.; Yoneda, K.; Mano, M.; Morikawa, O.; Konishi, H. *Chem. Lett.* **2003**, 32, 76–77; (d) Kobayashi, K.; Yoneda, K.; Mizumoto, T.; Umakoshi, H.; Morikawa, O.; Konishi, H. *Tetrahedron Lett.* **2003**, *44*, 4733–4736; (e) Kobayashi, K.; Takagoshi, K.; Kondo, S.; Morikawa, O.; Konishi, H. *Bull. Chem. Soc. Jpn.* **2004**, *77*, 553–559; (f) Kobayashi, K.; Yoneda, K.; Miyamoto, K.; Morikawa, O.; Konishi, H. *Tetrahedron* **2004**, *60*, 11639–11645; (g) Kobayashi, K.; Izumi, Y.; Hayashi, K.; Morikawa, O.; Konishi, H. *Bull. Chem. Soc. Jpn.* **2005**, *78*, 2171–2174; (h) Kobayashi, K.; Hayashi, K.; Iitsuka, D.; Morikawa, O.; Konishi, H. *Synthesis* **2006**, 1077–1082. See also pertinent references cited in these papers.
- Michelin, R. A.; Facchin, G.; Braga, D.; Sabatino, P. *Organometallics* **1986**, *5*, 2265–2274.
- (a) Ito, Y.; Kobayashi, K.; Seko, N.; Saegusa, T. *Bull. Chem. Soc. Jpn.* **1984**, *57*, 73–84; (b) Ugi, I.; Meyr, R. *Org. Synth.* **1973**, *Coll. Vol. V*, 1060–1063.
- Carling, R. W.; Leeson, P. D.; Moore, K. W.; Smith, J. D.; Moyes, C. R.; Mawer, I. M.; Thomas, S.; Chan, T.; Baker, R.; Foster, A. C.; Grimwood, S.; Kemp, J. A.; Marshall, G. R.; Tricklebank, M. D.; Saywell, K. L. *J. Med. Chem.* **1993**, *36*, 3397–3408.
- Foggasy, K.; Kovács, K.; Keseru, G. N.; Töke, L.; Faigl, F. *J. Chem. Soc., Perkin Trans. 1* **2001**, 1039–1043.
- Michelin, R. A.; Facchin, G.; Uguagliati, P. *Inorg. Chem.* **1984**, *23*, 961–969.
- Snyder, H. R.; Smith, C. W. *J. Am. Chem. Soc.* **1943**, *65*, 2452–2454.
- Brown, F.; Mann, F. G. *J. Chem. Soc., Chem. Commun.* **1948**, 847–858.
- Kabalka, G. W.; Wang, L.; Pagni, R. M. *Tetrahedron* **2001**, *57*, 8017–8028.
- Piozzi, F.; Langella, M. R. *Gazz. Chim. Ital.* **1963**, *40*, 1382–1391.
- Hegedus, L. S.; Winton, P. M.; Varaprath, S. J. *Org. Chem.* **1981**, *46*, 2215–2221.
- Bazile, Y.; Bras, J. P.; De Cointet, P.; Nanthavong, S.; Pigerol, C.; Broll, M.; Eymard, P.; Werbenec, J. P.; Fournier, J. *Eur. J. Med. Chem.* **1977**, *12*, 525–530.
- Suzuki, N.; Yasaki, S.; Yasuhara, A.; Sakamoto, T. *Chem. Pharm. Bull.* **2003**, *51*, 1170–1173.
- Crowther, A. F.; Mann, F. G.; Purdie, D. *J. Chem. Soc., Chem. Commun.* **1943**, 58–68.
- Ito, Y.; Kobayashi, K.; Saegusa, T. *J. Org. Chem.* **1979**, *44*, 2030–2032.
- Majumdar, K. C.; Samanta, S.; Chattopadhyay, B. *Tetrahedron Lett.* **2008**, *49*, 7213–7216.