



A simple and convenient approach to the Friedländer synthesis of pyrano[2,3-*b*]pyridines

Nagarajan Panneer Selvam, Thelagathoti Hari Babu, Paramasivan T. Perumal *

Organic Chemistry Division, Central Leather Research Institute, Adyar, Chennai 600 020, Tamilnadu, India

ARTICLE INFO

Article history:

Received 17 April 2009

Received in revised form 25 July 2009

Accepted 7 August 2009

Available online 13 August 2009

ABSTRACT

$\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ mediated efficient synthesis of pyrano[2,3-*b*]pyridines was achieved by the Friedländer reaction of 2-amino-3-cyano-4*H*-pyrans with cyclopentanone/cyclohexanone under solvent-free condition.

© 2009 Elsevier Ltd. All rights reserved.


1. Introduction

The presence of pyranopyridine scaffold in the frame work of several biologically active naturally occurring alkaloids of plant origin¹ has enthused researchers to synthesize and study their potential biological activities. They are known to possess anti-allergic, anti-inflammatory, and estrogenic properties.² In addition, benzopyrano[2,3-*b*]pyridines exhibit anti-proliferative,³ cancer chemopreventive,⁴ anti-bacterial (including anti-tubercular),⁵ anti-myopic,⁶ anti-histamic,⁷ hypotensive,⁸ anti-rheumatic,⁹ and anti-asthmatic activities.¹⁰

Despite numerous methods reported such as Skraup, Döbner-von Miller, and Combes,¹¹ the Friedländer annulation is the most simple and straightforward approach for the synthesis of poly-substituted pyridines and related azaheterocycles. Friedländer annulation is acid or base catalyzed annulation of *o*-aminoaryl ketones with carbonyl compounds containing a reactive α -methylene group. Among various Friedländer syntheses, cyclocondensation of *o*-aminobenzonitrile with various ketones has been less explored. Previous investigation^{12,13} on condensation of these derivatives required prolonged reaction times, use of hazardous reagents and limits the large scale application. In view of its immense biological applications, the development of simple and convenient protocol is of considerable interest.

Over the past few years, organic reactions under solvent-free conditions have gained much popularity.¹⁴ Solvent-free reactions are not only of interest from ecological point of view, but in many cases, also offer several synthetic advantages like higher yields, reduced pollution, low cost, improved selectivities, and simplicity in process and handling.¹⁵

Recently, $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ ¹⁶ has emerged both as the medium and catalyst in various organic transformations. In view of its inherent properties like reusability, greater selectivity, operational simplicity, non-corrosiveness, low cost, and ease of isolation, various synthetic transformations were accomplished. Previously, our group demonstrated the utility of $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ as an alternative to ionic liquid^{17d} and employed it as a reaction medium for carrying out various organic transformations. As part of our current studies on the design of new routes for the preparation of biologically active heterocyclic compounds,¹⁷ we herein disclose a simple and convenient method for the synthesis of pyrano[2,3-*b*]pyridines by Friedländer reaction mediated by $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ under solvent-free condition (Scheme 1).




Scheme 1.

2. Results and discussion

In our initial endeavor, we investigated the reaction of **1a** with cyclohexanone **2a** using various Lewis acid catalysts in different solvents (methanol, ethanol, and 1,2-dichloroethane) and neat at reflux or at 100 °C (Scheme 2 and Table 1). After systematic screening, we observed that tinchloride dihydrate gave moderate yield under solvent-free condition with equimolar amount of catalyst at 100 °C. Then, we optimized the reaction condition by increasing/decreasing the catalyst loading and temperature. The best result was obtained when the reaction was carried out with $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (also serving as reaction medium) under solvent-free condition at 110 °C. Further increase of temperature did not

* Corresponding author. Tel.: +91 44 24911386; fax: +91 44 24911589.

E-mail address: ptperumal@gmail.com (P.T. Perumal).

**Scheme 2.**

improve the product yield or increase the reaction rate. We also studied the recyclability of $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ as reaction medium. $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ is insoluble in dichloromethane. This property was utilized in recycling the reaction medium. Since $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ is insoluble in dichloromethane, it was used to bring out the organic part from the reaction mixture. Thus, dichloromethane leaves the $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ in reaction vessel, which was recycled for further reactions (three times). The yield did not vary much while recycling the reaction medium and the yields were found to be 56, 55, and 53, respectively.

Table 1
Catalyst and solvent screening

Entry	Catalyst	Yield ^a (%)			
		MeOH	EtOH	DCE	Neat
1	InCl_3	—	—	5	—
2	$\text{In}(\text{OTf})_3$	—	—	8	—
3	$\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$	—	—	45	54
4	$\text{NH}_2\text{SO}_3\text{H}$	—	—	—	Trace
5	CAN	—	—	—	—
6	BiCl_3	—	—	5	10
7	$\text{Bi}(\text{OTf})_3$	—	—	10	12
8	FeCl_3	—	—	6	12

^a All reactions were performed with 100 mol % catalyst at reflux or 100 °C for 5 h.

Table 2 summarizes our results on the Friedländer reaction of various 2-amino-3-cyano-4*H*-pyrans¹⁸ with cyclohexanone/cyclopentanone under optimized condition (**Scheme 3**). The reaction was amenable to wide variations in 2-amino-3-cyano-4*H*-pyrans and afforded the corresponding highly substituted pyranopyridines **3a–l** in moderate yields.

The structures of all products **3a–l** were confirmed by IR, ¹H NMR, ¹³C NMR, mass, and elemental analysis. In the IR spectrum of **3a**, disappearance of –CN group and a peak at 1638 cm^{–1} (–C=N–) confirmed the formation of pyridine ring. Absorptions at 3435, 3349, and 1209 cm^{–1} indicated the presence of –NH₂ and –C=O–C-groups, respectively. Signals at δ 150.2 (–C=N), 154.2 (–C–NH₂), 22.3–32.4 (–CH₂ groups) in the ¹³C NMR spectrum confirmed that cyclohexanone reacted with –NH₂ and –CN of 2-amino-3-cyano-4*H*-pyran. The ¹H NMR spectrum of **3a** displayed signals at δ 1.76, 2.20, and 2.71 (–CH₂ protons of cyclohexyl ring) and 4.03 (–NH₂ protons, D₂O exchangeable). Mass analysis also supported the structural assignment.

To further explore the potential of this protocol, we investigated the reaction of 2-amino-3-cyano chromenes with cyclohexanone/cyclopentanone and obtained the corresponding naphthopyranopyridines **5a–e** in moderate yields (**Scheme 4** and **Table 3**).

The structures of all products **5a–e** were confirmed by IR, ¹H NMR, ¹³C NMR, mass, and elemental analysis. The IR spectrum of **5e** showed absorptions at 3483, 3397, and 1109 cm^{–1} indicating the presence of –NH₂ and –C=O–C-groups, respectively. Disappearance of –CN group and a peak at 1628 cm^{–1} (–C=N–) confirmed the formation of pyridine ring. In the ¹H NMR spectrum, –NH₂ protons appeared as a broad singlet at δ 4.19 (D₂O exchangeable) and the methylene protons (–CH₂) of cyclohexyl ring were seen as

multiplets at δ 1.72, 2.26, and 2.83. Signals at δ 154.2 (–C=N), 160.4 (–C–NH₂), 22.3–32.4 (–CH₂ groups) in the ¹³C NMR spectrum confirmed that cyclohexanone reacted with –NH₂ and –CN of pyran. Mass analysis also supported the structural assignment.

A mechanistic rationalization for the reaction is given in **Scheme 5**.

3. Conclusion

In summary, we have developed a simple and convenient method for the synthesis of highly substituted pyranopyridines from 2-amino-3-cyano-4*H*-pyrans and cyclohexanone/cyclopentanone using inexpensive and commercially available $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ under solvent-free condition. The major advantage of this protocol is that $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ (as reaction medium) can be recycled upto three times without much change in yield. The generality of the reaction was also extended to the synthesis of naphthopyranopyridines in moderate yields.

4. Experimental

4.1. Materials and methods

$\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$, $\text{NH}_2\text{SO}_3\text{H}$, CAN, and FeCl_3 were obtained from S.D. Fine Chem. Ltd. $\text{Bi}(\text{OTf})_3$, $\text{In}(\text{OTf})_3$, InCl_3 , and BiCl_3 were purchased from Aldrich and used as received. All melting points were uncorrected. IR spectra were recorded on a Perkin Elmer FT-IR spectrophotometer. ¹H and ¹³C NMR spectra were recorded in CDCl_3 using TMS as an internal standard on JEOL spectrometer at 500 MHz and 125 MHz and Bruker spectrometer at 300 MHz and 75 MHz, respectively. Mass spectra were recorded by electrospray ionization method on Thermo Finnegan Mass spectrometer. Elemental analyses were recorded using a Thermo Finnegan FLASH EA 1112 CHN analyzer. Column chromatography was performed on silica gel (100–200 mesh, SRL, India). Analytical TLC was performed on precoated plastic sheets of silica gel G/UV-254 of 0.2 mm thickness (Macherey-Nagel, Germany).

4.2. General procedure for the synthesis of pyranopyridines

To a two-neck round-bottomed flask fitted with a mechanical stirrer and oil bath, 2 g of $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ was added and stirred at 50 °C. $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ became viscous liquid. To this viscous liquid, 2-amino-3-cyano-4*H*-pyran **1** (1 mmol) and cyclohexanone **2a** (1.1 mmol) were added and stirred at same temperature for about 15 min. Then stirring was continued at 110 °C until the disappearance of starting materials, as indicated by TLC. After the completion of the reaction, the reaction mixture was allowed to cool to 40 °C and triturated with dichloromethane (3×15 mL) and stirred for 10 min. The organic layer thus obtained was washed twice with water and with saturated sodium chloride solution, dried over anhydrous Na_2SO_4 , and concentrated under reduced pressure. The crude product was purified through column chromatography (40% ethyl acetate in petroleum ether). The round-bottom flask, which contains the $\text{SnCl}_2 \cdot 2\text{H}_2\text{O}$ was again utilized for recycling the reaction medium (successfully recycled three times). The same procedure was followed for the synthesis of naphthopyranopyridines **5a–e**.

4.2.1. Ethyl 5-amino-4-(4-chlorophenyl)-2-methyl-6,7,8,9-tetrahydro-4*H*-pyrano[2,3-*b*]quinoline-3-carboxylate (3a). Off white solid; mp 138–140 °C; *R*_f 0.65 (40% EtOAc/petroleum ether); IR ν_{max} (KBr): 3435, 3349, 2928, 1685, 1638, 1571, 1447, 1295, 1209, 1057, 833 cm^{–1}; ¹H NMR (500 MHz, CDCl_3) δ: 1.22 (t, 3H, *J*=6.9 Hz, $-\text{OCH}_2\text{CH}_3$), 1.76 (m, 4H, $-\text{CH}_2$), 2.20 (m, 2H, $-\text{CH}_2$), 2.42 (s, 3H, $-\text{CH}_3$), 2.71 (m, 2H, $-\text{CH}_2$), 4.03 (br s, 2H, D₂O exchangeable, –NH₂), 4.09 (m, 2H, $-\text{OCH}_2\text{CH}_3$), 4.80 (s, 1H, $-\text{CH}$), 7.18 (m, 4H, $-\text{ArH}$); ¹³C NMR (125 MHz, CDCl_3) δ: 14.2, 19.8, 22.3, 22.5, 22.9, 32.4, 37.9, 60.4, 99.2,

Table 2
Synthesis of pyrano[2,3-*b*]pyridines **3a–l**

Entry	2-Amino-3-cyano-4 <i>H</i> -pyrans (1)	Cyclic ketone (2)	Product (3) ^a	Time (h)	Yield ^b (%)
1				4.5	58
2				4.5	60
3				4.2	64
4				4.5	64
5				4.5	62
6				4.2	60
7				4.0	65

Table 2 (continued)

Entry	2-Amino-3-cyano-4H-pyrans (1)	Cyclic ketone (2)	Product (3) ^a	Time (h)	Yield ^b (%)
8				4.0	65
9				5.0	62
10				5.0	63
11				4.5	62
12				5.0	63

^a All products were characterized by IR, ¹H NMR, ¹³C NMR, mass spectral, and elemental analysis.^b Isolated yield.

106.4, 113.7, 128.7, 129.8, 132.8, 142.4, 150.2, 154.2, 154.3, 160.1, 166.7; MS (ESI LCQ-MS): *m/z* 399 [M+1]⁺. Anal. Calcd for C₂₂H₂₃N₂O₃Cl: C, 66.24; H, 5.81; N, 7.02%. Found: C, 66.41; H, 5.77; N, 7.04%.

4.2.2. Ethyl 5-amino-2-methyl-4-(3-nitrophenyl)-6,7,8,9-tetrahydro-4H-pyranol[2,3-*b*]quinoline-3-carboxylate (**3b**). Off white solid; mp

204–206 °C; *R*_f 0.57 (40% EtOAc/petroleum ether); IR ν_{max} (KBr): 3438, 3340, 2932, 1686, 1642, 1529, 1353, 1230, 1065, 801, 685 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ : 1.24 (t, 3H, *J*=7.7 Hz, -OCH₂CH₃), 1.78 (m, 4H, -CH₂), 2.33 (m, 2H, -CH₂), 2.46 (s, 3H, -CH₃), 2.74 (m, 2H, -CH₂), 4.11 (br s, 2H, D₂O exchangeable, -NH₂), 4.12 (m, 2H, -OCH₂CH₃), 4.95 (s, 1H, -CH), 7.42 (t, 1H, *J*=8.4 Hz, -Ar-






Table 3
Synthesis of naphthopyranopyridines **5a–e**

Entry	2-Amino-3-cyano chromenes (4)	Cyclic ketone (2)	Product (5) ^a	Time (h)	Yield ^b (%)
1				3.5	62
2				4.0	60
3				3.0	64
4				4.0	60
5				4.0	60

^a All products were characterized by IR, ¹H NMR, ¹³C NMR, mass spectral, and elemental analysis.

^b Isolated yield.



H), 7.58 (d, 1H, *J*=7.7 Hz, –Ar–H), 8.03 (d, 1H, *J*=8.4 Hz, –Ar–H), 8.19 (s, 1H, –Ar–H); ¹³C NMR (125 MHz, CDCl₃) δ: 14.3, 17.6, 23.7, 32.5, 34.7, 60.8, 98.4, 105.8, 114.0, 121.7, 122.9, 123.9, 129.4, 137.6, 146.1, 148.2, 150.1, 154.3, 154.8, 161.3, 166.5; MS (ESI LCQ-MS): *m/z* 410 [M+1]⁺. Anal. Calcd for C₂₂H₂₃N₃O₅: C, 64.54; H, 5.66; N, 10.26%. Found: C, 64.63; H, 5.69; N, 10.24%.

4.2.3. Methyl 5-amino-4-(3-fluorophenyl)-2-methyl-6,7,8,9-tetrahydro-4*H*-pyrano[2,3-*b*]quinoline-3-carboxylate (3c). Yellow solid; mp 222–224 °C; *R*_f 0.61 (40% EtOAc/petroleum ether); IR ν_{\max} (KBr): 3453, 3372, 2932, 1692, 1643, 1446, 1380, 1295, 1234, 1064, 774 cm⁻¹; ¹H NMR (500 MHz, CDCl₃) δ: 1.63 (m, 4H, –CH₂), 2.34 (m, 2H, –CH₂), 2.44 (s, 3H, –CH₃), 2.74 (m, 2H, –CH₂), 3.75 (s, 3H, –OCH₃), 4.02 (s, 2H, D₂O exchangeable, NH₂), 4.85 (s, 1H, –CH), 6.88 (t, 1H,

$J=8.4$ Hz, –Ar–H), 6.97 (d, 1H, $J=9.2$ Hz, –Ar–H), 7.08 (d, 1H, $J=6.9$ Hz, –Ar–H), 7.19 (s, 1H, –Ar–H); ^{13}C NMR (75 MHz, CDCl_3) δ : 19.8, 22.3, 22.5, 22.9, 29.7, 32.4, 38.2, 51.4, 99.0, 106.1, 113.7, 114.1, 115.3, 123.9, 130.1, 146.5, 150.3, 154.2, 160.4 ($J_{\text{C}-\text{F}}=240.2$ Hz), 163.8, 167.2; MS (ESI LCQ-MS): m/z 369 [M+1] $^+$. Anal. Calcd for $\text{C}_{21}\text{H}_{21}\text{N}_2\text{O}_3\text{F}$: C, 68.46; H, 5.75; N, 7.60%. Found: C, 68.62; H, 5.80; N, 7.67%.

4.2.4. Isopropyl 5-amino-4-(4-chlorophenyl)-2-methyl-6,7,8,9-tetrahydro-4H-pyranolo[2,3-b]quinoline-3-carboxylate (3d). Pale brown solid; mp 130–132 °C; R_f 0.52 (40% EtOAc/petroleum ether); IR ν_{max} (KBr): 3441, 3365, 2928, 1689, 1642, 1452, 1371, 1297, 1211, 1098, 1060, 834 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ : 1.16 (d, 3H, $J=6.1$ Hz, –CH₃), 1.22 (d, 3H, $J=6.1$ Hz, –CH₃), 1.76 (m, 4H, –CH₂), 2.29 (m, 2H, –CH₂), 2.41 (s, 3H, –CH₃), 2.71 (m, 2H, –CH₂), 4.03 (s, 2H, $D_2\text{O}$ exchangeable, –NH₂), 4.78 (s, 1H, –CH), 4.97 (m, 1H, –OCH(CH₃)₂), 7.20 (s, 4H, –Ar–H); ^{13}C NMR (75 MHz, CDCl_3) δ : 18.3, 22.4, 22.6, 22.9, 32.5, 37.5, 99.3, 106.6, 113.8, 128.3, 129.1, 129.5, 132.9, 142.4, 150.3, 154.3, 159.3, 166.3; MS (ESI LCQ-MS): m/z 412 [M+1] $^+$. Anal. Calcd for $\text{C}_{23}\text{H}_{25}\text{N}_2\text{O}_3\text{Cl}$: C, 66.90; H, 6.10; N, 6.78%. Found: C, 67.12; H, 6.14; N, 6.81%.

4.2.5. Methyl 5-amino-4-(2,4-dichlorophenyl)-2-methyl-6,7,8,9-tetrahydro-4H-pyranolo[2,3-b]quinoline-3-carboxylate (3e). Light brown solid; mp 222–224 °C; R_f 0.57 (40% EtOAc/petroleum ether); IR ν_{max} (KBr): 3483, 3396, 2934, 1713, 1642, 1451, 1397, 1224, 1064, 1099, 845, 655 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ : 1.65 (m, 2H, –CH₂), 1.83 (m, 2H, –CH₂), 2.31 (m, 2H, –CH₂), 2.48 (s, 3H, –CH₃), 2.69 (m, 2H, –CH₂), 3.63 (s, 3H, –OCH₃), 4.40 (s, 2H, $D_2\text{O}$ exchangeable, –NH₂), 5.21 (s, 1H, –CH), 7.14 (m, 2H, –Ar–H), 7.29 (s, 1H, –Ar–H); ^{13}C NMR (75 MHz, CDCl_3) δ : 22.4, 22.9, 32.0, 34.8, 50.8, 98.9, 105.8, 113.4, 129.2, 132.3, 140.7, 150.4, 154.1, 154.2, 161.7, 162.7, 167.0; MS (ESI LCQ-MS): m/z 419 [M+1] $^+$. Anal. Calcd for $\text{C}_{21}\text{H}_{20}\text{N}_2\text{O}_3\text{Cl}_2$: C, 60.15; H, 4.81; N, 6.68%. Found: C, 60.28; H, 4.83; N, 6.71%.

4.2.6. 1-(5-Amino-2-methyl-4-(3-nitrophenyl)-6,7,8,9-tetrahydro-4H-pyranolo[2,3-b]quinolin-3-yl)ethanone (3f). Pale brown solid; mp 238–240 °C; R_f 0.62 (40% EtOAc/petroleum ether); IR ν_{max} (KBr): 3433, 3357, 2925, 1706, 1631, 1528, 1351, 682 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ : 1.80 (m, 4H, –CH₂), 2.33 (m, 5H, –CH₃ and –CH₂), 2.49 (s, 3H, acetyl–CH₃), 2.75 (m, 2H, –CH₂), 4.10 (br s, 2H, $D_2\text{O}$ exchangeable, –NH₂), 5.06 (s, 1H, –CH), 7.41 (t, 1H, $J=7.7$ Hz, –Ar–H), 7.63 (d, 1H, $J=7.7$ Hz, –Ar–H), 8.02 (d, 1H, $J=8.4$ Hz, –Ar–H), 8.15 (s, 1H, –Ar–H); ^{13}C NMR (75 MHz, CDCl_3) δ : 18.3, 21.3, 22.8, 23.6, 32.3, 38.6, 110.4, 114.6, 115.3, 128.3, 130.1, 136.2, 139.4, 149.4, 153.4, 155.1, 156.3, 166.4, 198.6; MS (ESI LCQ-MS): m/z 380 [M+1] $^+$. Anal. Calcd for $\text{C}_{21}\text{H}_{21}\text{N}_3\text{O}_4$: C, 66.48; H, 5.58; N, 11.08%. Found: C, 66.72; H, 5.61; N, 11.04%.

4.2.7. 11-Amino-12-(3-nitrophenyl)-3,4,7,8,9,10-hexahydro-2H-chromeno[2,3-b]quinolin-1(12H)-one (3g). Pale yellow solid; mp 242–244 °C; R_f 0.50 (40% EtOAc/petroleum ether); IR ν_{max} (KBr): 3468, 3365, 2931, 1631, 1530, 1351, 1192, 1076, 716 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ : 1.22 (m, 2H, –CH₂), 2.02 (m, 4H, –CH₂), 2.35 (m, 4H, –CH₂), 2.75 (m, 4H, –CH₂), 4.15 (s, 2H, $D_2\text{O}$ exchangeable, –NH₂), 5.02 (s, 1H, –CH), 7.40 (t, 1H, $J=7.7$ Hz, –Ar–H), 7.69 (d, 1H, $J=7.7$ Hz, –Ar–H), 8.00 (d, 1H, $J=6.9$ Hz, –Ar–H), 8.17 (s, 1H, –Ar–H); ^{13}C NMR (75 MHz, CDCl_3) δ : 20.2, 22.8, 32.3, 33.6, 34.9, 36.9, 98.6, 114.1, 121.6, 122.8, 124.1, 128.8, 135.4, 145.5, 148.5, 150.7, 154.2, 154.8, 167.4, 197.0; MS (ESI LCQ-MS): m/z 392 [M+1] $^+$. Anal. Calcd for $\text{C}_{22}\text{H}_{21}\text{N}_3\text{O}_4$: C, 67.51; H, 5.41; N, 10.74%. Found: C, 67.64; H, 5.42; N, 10.78%.

4.2.8. 11-Amino-12-(4-nitrophenyl)-3,4,7,8,9,10-hexahydro-2H-chromeno[2,3-b]quinolin-1(12H)-one (3h). Off white solid; mp 242–244 °C; R_f 0.52 (40% EtOAc/petroleum ether); IR ν_{max} (KBr): 3410, 3368, 2930, 1637, 1564, 1520, 1348, 1190, 1074 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ : 1.23 (m, 2H, –CH₂), 1.81 (m, 6H, –CH₂), 2.03 (m,

2H, –CH₂), 2.33 (m, 2H, –CH₂), 2.75 (m, 2H, –CH₂), 4.06 (s, 2H, $D_2\text{O}$ exchangeable, –NH₂), 5.00 (s, 1H, –CH), 7.51 (d, 2H, $J=8.4$ Hz, –Ar–H), 8.09 (d, 2H, $J=8.4$ Hz, –Ar–H); ^{13}C NMR (75 MHz, CDCl_3) δ : 21.4, 21.9, 22.4, 32.1, 33.9, 34.8, 34.9, 33.1, 52.7, 111.3, 113.8, 114.6, 123.8, 130.1, 144.1, 153.4, 154.3, 160.2, 164.3, 198.6; MS (ESI LCQ-MS): m/z 392 [M+1] $^+$. Anal. Calcd for $\text{C}_{22}\text{H}_{21}\text{N}_3\text{O}_4$: C, 67.51; H, 5.41; N, 10.74%. Found: C, 67.83; H, 5.40; N, 10.69%.

4.2.9. 11-Amino-3,3-dimethyl-12-(3-nitrophenyl)-3,4,7,8,9,10-hexahydro-2H-chromeno[2,3-b]quinolin-1(12H)-one (3i). White solid; mp 290–292 °C [lit:¹⁹ 293 °C]; R_f 0.46 (40% EtOAc/petroleum ether); IR ν_{max} (KBr): 3424, 3387, 2935, 1637, 1564, 1535, 1421, 1370, 1232, 1032, 580 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ : 1.08 (s, 3H, –CH₃), 1.24 (s, 3H, –CH₃), 1.82 (m, 4H, –CH₂), 2.23 (m, 4H, –CH₂), 2.57 (m, 2H, –CH₂), 2.77 (m, 2H, –CH₂), 4.07 (br s, 2H, $D_2\text{O}$ exchangeable, –NH₂), 4.98 (s, 1H, –CH), 7.41 (t, 1H, $J=7.7$ Hz, –Ar–H), 7.70 (d, 1H, $J=7.7$ Hz, –Ar–H), 8.02 (d, 1H, $J=8.4$ Hz, –Ar–H), 8.18 (s, 1H, –Ar–H); ^{13}C NMR (75 MHz, CDCl_3) δ : 21.2, 21.4, 21.9, 26.1, 28.0, 31.1, 32.7, 40.2, 43.0, 49.5, 97.2, 112.1, 113.0, 120.5, 122.2, 144.9, 146.8, 150.5, 152.5, 153.0, 164.3, 195.2; MS (ESI LCQ-MS): m/z 420 [M+1] $^+$. Anal. Calcd for $\text{C}_{24}\text{H}_{25}\text{N}_3\text{O}_4$: C, 68.72; H, 6.01; N, 10.02%. Found: C, 69.01; H, 6.04; N, 10.09%.

4.2.10. 11-Amino-12-(2,4-dichlorophenyl)-3,3-dimethyl-3,4,7,8,9,10-hexahydro-2H-chromeno[2,3-b]quinolin-1(12H)-one (3j). Pale yellow solid; mp 282–284 °C; R_f 0.49 (40% EtOAc/petroleum ether); IR ν_{max} (KBr): 3404, 3348, 2933, 1649, 1369, 1230, 1168, 1039, 852, 567 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ : 0.97 (s, 3H, –CH₃), 1.09 (s, 3H, –CH₃), 1.78 (m, 4H, –CH₂), 2.32 (m, 4H, –CH₂), 2.56 (m, 2H, –CH₂), 2.72 (m, 2H, –CH₂), 4.38 (s, 2H, $D_2\text{O}$ exchangeable, –NH₂), 5.19 (s, 1H, –CH), 7.12 (d, 2H, $J=8.4$ Hz, –Ar–H), 7.31 (s, 1H, –Ar–H); ^{13}C NMR (75 MHz, CDCl_3) δ : 18.4, 21.8, 28.4, 32.3, 34.7, 47.1, 54.3, 61.2, 111.8, 113.4, 115.4, 126.8, 130.1, 133.1, 136.4, 136.7, 150.1, 154.9, 160.2, 164.8, 198.3; MS (ESI LCQ-MS): m/z 443 [M+1] $^+$. Anal. Calcd for $\text{C}_{24}\text{H}_{24}\text{N}_2\text{O}_2\text{Cl}_2$: C, 65.02; H, 5.46; N, 6.32%. Found: C, 65.28; H, 5.49; N, 6.31%.

4.2.11. 10-Amino-8-(4-chlorophenyl)-6-methyl-1,2,3,8-tetrahydro-5-oxa-4-aza-cyclopenta[b]naphthacene-7-carboxylic acid methyl ester (3k). Off white solid; mp 182–184 °C; R_f 0.59 (40% EtOAc/petroleum ether); IR ν_{max} (KBr): 3461, 3336, 2956, 1714, 1559, 1446, 1208, 1096, 820 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ : 1.72 (s, 3H, –CH₃), 1.96 (m, 2H, –CH₂), 2.58 (m, 2H, –CH₂), 2.93 (m, 2H, –CH₂), 3.86 (s, 3H, –OCH₃), 4.13 (br s, 2H, $D_2\text{O}$ exchangeable, –NH₂), 4.89 (s, 1H, –CH), 6.97 (d, 2H, $J=7.7$ Hz, –Ar–H), 7.15 (d, 2H, $J=7.7$ Hz, –Ar–H); ^{13}C NMR (75 MHz, CDCl_3) δ : 18.4, 24.2, 24.8, 34.7, 52.6, 96.4, 104.7, 105.8, 114.2, 130.1, 132.3, 136.3, 152.6, 154.8, 155.7, 163.9, 165.1; MS (ESI LCQ-MS): m/z 371 [M+1] $^+$. Anal. Calcd for $\text{C}_{20}\text{H}_{19}\text{N}_2\text{O}_3\text{Cl}$: C, 64.78; H, 5.16; N, 7.55%. Found: C, 64.93; H, 5.18; N, 7.49%.

4.2.12. 11-Amino-3,3-dimethyl-12-(4-nitrophenyl)-3,4,7,8,9,10-hexahydro-2H-chromeno[2,3-b]quinolin-1(12H)-one (3l). Off white solid; mp: 322–324 °C [lit:¹⁹ 323–325 °C]; R_f 0.49 (40% EtOAc/petroleum ether); IR ν_{max} (KBr): 3402, 3232, 2927, 1637, 1517, 1370, 1370, 1346, 1228, 1205, 1032, 802, 579 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ : 0.92 (s, 3H, –CH₃), 1.07 (s, 3H, –CH₃), 1.80 (m, 4H, –CH₂), 2.13 (m, 4H, –CH₂), 2.54 (m, 2H, –CH₂), 2.74 (m, 2H, –CH₂), 4.10 (br s, 2H, $D_2\text{O}$ exchangeable, –NH₂), 4.98 (s, 1H, –CH), 7.50 (d, 2H, $J=8.4$ Hz, –Ar–H), 8.08 (d, 2H, $J=8.4$ Hz, –Ar–H); ^{13}C NMR (75 MHz, CDCl_3) δ : 18.6, 22.3, 22.5, 23.0, 32.5, 34.6, 98.6, 114.7, 126.3, 127.2, 134.1, 136.2, 152.3, 154.4, 156.8, 162.0; MS (ESI LCQ-MS): m/z 420 [M+1] $^+$. Anal. Calcd for $\text{C}_{24}\text{H}_{25}\text{N}_3\text{O}_4$: C, 68.72; H, 6.01; N, 10.02%. Found: C, 68.99; H, 5.98; N, 10.09%.

4.2.13. 14-(3-Nitrophenyl)-9,11,12,14-tetrahydro-10H-7-oxa-8-azabenz[a]naphthacen-13-ylamine (5a). Pale yellow solid; mp 284–

286 °C [lit.²⁰ 287–289 °C]; R_f 0.53 (40% EtOAc/petroleum ether); IR ν_{max} (KBr): 3456, 3324, 2928, 1654, 1535, 1349, 1233, 1082, 816, 680 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ : 1.80 (m, 4H, $-\text{CH}_2$), 2.34 (m, 2H, $-\text{CH}_2$), 2.75 (m, 2H, $-\text{CH}_2$), 4.11 (br s, 2H, D_2O exchangeable, $-\text{NH}_2$), 5.32 (s, 1H, $-\text{CH}$), 7.81 (m, 5H, $-\text{Ar}-\text{H}$), 8.01 (m, 4H, $-\text{Ar}-\text{H}$), 8.44 (s, 1H, $-\text{Ar}-\text{H}$); ^{13}C NMR (125 MHz, CDCl_3) δ : 22.4, 22.8, 26.4, 32.1, 34.8, 98.8, 105.4, 113.4, 115.2, 117.5, 121.5, 122.4, 122.8, 123.9, 126.6, 132.1, 133.2, 134.5, 142.3, 151.0, 154.5, 159.9, 163.7; MS (ESI LCQ-MS): m/z 424 [$\text{M}+1$]⁺. Anal. Calcd for $\text{C}_{26}\text{H}_{21}\text{N}_3\text{O}_3$: C, 73.74; H, 5.00; N, 9.92%. Found: C, 73.99; H, 5.04; N, 9.98%.

4.2.14. 13-(4-Chlorophenyl)-9,10,11,13-tetrahydro-7-oxa-8-aza-benzo[*a*]-cyclopenta[*ll*]anthracene-12-ylamine (5b**).** Off white solid; mp 214–216 °C; R_f 0.61 (40% EtOAc/petroleum ether); IR ν_{max} (KBr): 3431, 3372, 2936, 1601, 1451, 1209, 1086, 823 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ : 1.96 (m, 2H, $-\text{CH}_2$), 2.48 (m, 2H, $-\text{CH}_2$), 2.68 (m, 2H, $-\text{CH}_2$), 4.21 (m, 2H, D_2O exchangeable, $-\text{NH}_2$), 5.41 (s, 1H, $-\text{CH}$), 7.10 (m, 4H, $-\text{Ar}-\text{H}$), 7.42 (m, 6H, $-\text{Ar}-\text{H}$); ^{13}C NMR (125 MHz, CDCl_3) δ : 22.1, 22.9, 25.7, 34.6, 97.3, 106.4, 114.9, 117.4, 119.2, 121.8, 122.3, 126.3, 127.4, 127.8, 128.3, 129.4, 130.2, 134.2, 141.8, 153.8, 157.8, 162.4, 164.3; MS (ESI LCQ-MS): m/z 399 [$\text{M}+1$]⁺. Anal. Calcd for $\text{C}_{25}\text{H}_{19}\text{N}_2\text{OCl}$: C, 75.28; H, 4.80; N, 7.02%. Found: C, 75.53; H, 4.84; N, 7.05%.

4.2.15. 13-(4-Nitrophenyl)-9,10,11,13-tetrahydro-7-oxa-8-aza-benzo[*a*]-cyclopenta[*ll*]anthracene-12-ylamine (5c**).** Pale brown solid; mp 202–204 °C; R_f 0.57 (40% EtOAc/petroleum ether); IR ν_{max} (KBr): 3451, 3341, 2940, 1611, 1534, 1358, 1440, 1198, 1091, 821 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ : 1.86 (m, 2H, $-\text{CH}_2$), 2.48 (m, 2H, $-\text{CH}_2$), 2.86 (m, 2H, $-\text{CH}_2$), 4.31 (br s, 2H, D_2O exchangeable, $-\text{NH}_2$), 5.36 (s, 1H, $-\text{CH}$), 6.86 (d, 1H, $J=8.4$ Hz, $-\text{Ar}-\text{H}$), 7.18 (m, 4H, $-\text{Ar}-\text{H}$), 7.40 (m, 3H, $-\text{Ar}-\text{H}$), 8.16 (d, 2H, $J=7.7$ Hz, $-\text{Ar}-\text{H}$); ^{13}C NMR (75 MHz, CDCl_3) δ : 23.5, 24.8, 25.3, 34.8, 96.4, 106.8, 115.1, 117.8, 118.3, 122.4, 123.7, 124.2, 125.9, 126.2, 127.4, 129.4, 134.3, 144.9, 152.1, 155.7, 161.0, 164.3; MS (ESI LCQ-MS): m/z 410 [$\text{M}+1$]⁺. Anal. Calcd for $\text{C}_{25}\text{H}_{19}\text{N}_3\text{O}_3$: C, 73.34; H, 4.68; N, 10.26%. Found: C, 73.57; H, 4.72; N, 10.31%.

4.2.16. 7-(3-Nitrophenyl)-9,10,11,12-tetrahydro-7*H*-14-oxa-13-aza-benzo[*a*]naphthacen-8-ylamine (5d**).** Light green solid; mp 154–156 °C; R_f 0.51 (40% EtOAc/petroleum ether); IR ν_{max} (KBr): 3454, 2930, 1609, 1529, 1380, 1347, 1103, 806, 679 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ : 1.71 (m, 4H, $-\text{CH}_2$), 2.48 (m, 2H, $-\text{CH}_2$), 2.88 (m, 2H, $-\text{CH}_2$), 4.12 (br s, 2H, D_2O exchangeable, $-\text{NH}_2$), 5.21 (s, 1H, $-\text{CH}$), 7.04 (d, 1H, $J=7.7$ Hz, $-\text{Ar}-\text{H}$), 7.41 (m, 5H, $-\text{Ar}-\text{H}$), 7.68 (d, 1H, $J=8.4$ Hz, $-\text{Ar}-\text{H}$), 7.90 (s, 1H, $-\text{Ar}-\text{H}$), 8.02 (m, 2H, $-\text{Ar}-\text{H}$); ^{13}C NMR (75 MHz, CDCl_3) δ : 22.2, 22.5, 23.2, 32.0, 34.8, 97.4, 105.4, 115.2, 115.8, 116.1, 121.2, 123.6, 124.8, 126.4, 127.4, 129.2, 132.3, 134.7, 144.7, 148.9, 150.4, 156.4, 160.9, 164.9; MS (ESI LCQ-MS): m/z 424 [$\text{M}+1$]⁺. Anal. Calcd for $\text{C}_{26}\text{H}_{21}\text{N}_3\text{O}_3$: C, 73.74; H, 5.00; N, 9.92%. Found: C, 73.98; H, 5.06; N, 9.97%.

4.2.17. 7-(2,4-Dichlorophenyl)-9,10,11,12-tetrahydro-7*H*-14-oxa-13-aza-benzo[*a*]naphthacen-8-ylamine (5e**).** Pale brown solid; mp 160–162 °C; R_f 0.37 (40% EtOAc/petroleum ether); IR ν_{max} (KBr): 3483, 3397, 3043, 1628, 1580, 1457, 1380, 1109, 866, 809 cm^{-1} ; ^1H NMR (500 MHz, CDCl_3) δ : 1.72 (m, 4H, $-\text{CH}_2$), 2.26 (m, 2H, $-\text{CH}_2$), 2.83 (m, 2H, $-\text{CH}_2$), 4.19 (s, 2H, D_2O exchangeable, $-\text{NH}_2$), 5.78 (s, 1H, $-\text{CH}$), 7.04 (d, 1H, $J=8.4$ Hz, $-\text{Ar}-\text{H}$), 7.11 (d, 1H, $J=8.4$ Hz, $-\text{Ar}-\text{H}$), 7.20 (d, 1H, $J=8.4$ Hz, $-\text{Ar}-\text{H}$), 7.36 (m, 2H, $-\text{Ar}-\text{H}$), 7.48 (m, 1H,

$-\text{Ar}-\text{H}$), 7.73 (d, 1H, $J=8.4$ Hz, $-\text{Ar}-\text{H}$), 8.12 (d, 1H, $J=8.4$ Hz, $-\text{Ar}-\text{H}$), 8.61 (d, 1H, $J=8.4$ Hz, $-\text{Ar}-\text{H}$); ^{13}C NMR (125 MHz, CDCl_3) δ : 22.3, 22.8, 29.7, 32.4, 38.2, 99.0, 106.1, 113.7, 114.2, 115.3, 121.3, 122.4, 123.9, 125.6, 126.9, 128.2, 130.1, 132.0, 133.7, 143.4, 147.8, 150.3, 154.2, 160.4, 164.0; MS (ESI LCQ-MS): m/z 447 [$\text{M}+1$]⁺. Anal. Calcd for $\text{C}_{26}\text{H}_{20}\text{N}_3\text{O}_2\text{Cl}_2$: C, 69.81; H, 4.51; N, 6.26%. Found: C, 70.13; H, 4.54; N, 6.30%.

Acknowledgements

One of the author, N.P.S. thanks the Council of Scientific and Industrial Research (CSIR), New Delhi for financial assistance.

References and notes

- (a) Ahmad, S. *J. Nat. Prod.* **1984**, *47*, 391; (b) Mitaka, S.; Skaltsounis, A.-L.; Till-equin, F.; Koch, M.; Pusset, J.; Chauviere, G. *J. Nat. Prod.* **1985**, *48*, 772; (c) Ulubelen, A. *Phytochemistry* **1984**, *23*, 2123; (d) Tantivatana, P.; Ruangrungsri, N.; Vaisiroiroj, V.; Lankin, D. C.; Bhacca, N. S.; Morris, R. P.; Cordell, G. A.; Johnson, L. F. *J. Org. Chem.* **1983**, *48*, 268; (e) Munoz, M. A.; Torres, R.; Cassels, B. K. *J. Nat. Prod.* **1982**, *45*, 367.
- (a) Faber, K.; Stueckler, H.; Kappe, T. *J. Heterocycl. Chem.* **1984**, *21*, 1171; (b) Johnson, J. V.; Rauckman, S.; Beccanari, P. D.; Roth, B. *J. Med. Chem.* **1989**, *32*, 1942; (c) Yamada, N.; Kadokawa, S.; Takahashi, K.; Umezawa, K. *Biochem. Pharmacol.* **1992**, *44*, 1211.
- Kolokythas, G.; Pouli, N.; Marakos, P.; Pratsinis, H.; Kletsas, D. *Eur. J. Med. Chem.* **2006**, *41*, 71.
- Azuine, M. A.; Tokuda, H.; Takayasu, J.; Enjyo, F.; Mukainaka, T.; Konoshima, T.; Nishino, H.; Kapadia, G. *J. Pharmacol. Res.* **2004**, *49*, 161.
- (a) Srivastava, S. K.; Tripathi, R. P.; Ramachandran, R. *J. Biol. Chem.* **2005**, *280*, 30273; (b) Brotz-Oesterheld, H.; Knezevic, I.; Bartel, S.; Lampe, T.; Warnecke-Eberz, U.; Ziegelbauer, K.; Habich, D.; Labischinski, H. *J. Biol. Chem.* **2003**, *278*, 39435.
- Toshiro, S.; Noriko, W. Eur. Pat. Appl. EP 647445 A1 19950412, 1995.
- Ito, Y.; Kato, H.; Yasuda, S.; Kato, N.; Iwasaki, N.; Nishino, H.; Takeshita, M. Jpn. Kokai Tokkyo Koho, JP 06107664 A2 19940419, 1994.
- Goto, K.; Yaoka, O.; Oe, T. PCT Int. Appl. WO 8401711 A1 19840510, 1984.
- Maruyama, Y.; Goto, K.; Terasawa, M. Ger. Offen. DE 3010751 19810806, 1981.
- Ukawa, K.; Ishiguro, T.; Kuriki, H.; Nohara, A. *Chem. Pharm. Bull.* **1985**, *33*, 4432.
- (a) *In Comprehensive Heterocyclic Chemistry*; Jones, G., Katritzky, A. R., Ress, C. W., Eds.; Pergamon: New York, NY, 1996; Vol. 5, p 167; (b) Cho, C. S.; Oh, B. H.; Kim, T. J.; Shim, S. C. *J. Chem. Soc., Chem. Commun.* **2000**, 1885; (c) Jiang, B.; Si, Y. C. *J. Org. Chem.* **2002**, *67*, 9449; (d) Kraupa, H. *Chem. Ber.* **1880**, *13*, 2086; (e) Friedländer, P. *Ber.* **1882**, *15*, 2572; (f) Mansake, R. H.; Kulka, M. *Org. React.* **1953**, *7*, 59; (g) Linderman, R. J.; Kirolos, S. K. *Tetrahedron Lett.* **1990**, *31*, 2689; (h) Theclitou, M. E.; Robinson, L. A. *Tetrahedron Lett.* **2002**, *43*, 3907.
- (a) Martinez-Grau, A.; Marco, J. *Bioorg. Med. Chem. Lett.* **1997**, *7*, 3165; (b) Li, J. R.; Zhang, L. J.; Yang, X. Q.; Li, Q.; Wang, D.; Wang, C. X.; Shi, D. X.; Zhang, Q. *Chin. Chem. Lett.* **2008**, *19*, 15; (c) Sheldon, R. A. *Green Chem.* **2005**, *7*, 267.
- (a) de los Rios, C.; Marco, J. L.; Carreiras, M. C.; Chinchon, P. M.; Garcia, A. G.; Villarroya, M. *Bioorg. Med. Chem.* **2002**, *10*, 2077; (b) Marco, J. L.; de los Rios, C.; Garcia, A. G.; Villarroya, M.; Carreiras, M. C.; Martins, C.; Eleuterio, A.; Morreal, A.; Orozco, M.; Luque, J. *Bioorg. Med. Chem.* **2004**, *12*, 2199; (c) Leon, R.; Marco-Contelles, J.; Garcia, A. G.; Villarroya, M. *Bioorg. Med. Chem.* **2005**, *13*, 1167; (d) Marco-Contelles, J.; Leon, R.; de los Rios, C.; Garcia, A. G.; Lopez, M. G.; Villarroya, M. *Bioorg. Med. Chem.* **2006**, *14*, 8176; (e) Marco, J. L.; de los Rios, C.; Carreiras, M. C.; Banos, J. E.; Badia, A.; Vivas, N. M. *Bioorg. Med. Chem.* **2001**, *9*, 727; (f) Marco-Contelles, J.; Leon, R.; Lopez, M. G.; Garcia, A. G.; Villarroya, M. *Eur. J. Med. Chem.* **2006**, *41*, 1464.
- (a) Loh, T. P.; Huang, J. M.; Goh, S. H.; Vittal, J. *J. Org. Lett.* **2000**, *29*, 1291; (b) Solvent-Free Organic Synthesis; Tanaka, K., Ed.; Wiley-VCH: Weinheim, 2003.
- (a) Cave, G. W. W.; Raston, C. L.; Scott, J. L. *Chem. Commun.* **2001**, 2159; (b) Tanaka, K.; Toda, F. *Chem. Rev.* **2000**, *100*, 1025.
- (a) Bez, G.; Gogoi, D. *Tetrahedron Lett.* **2006**, *47*, 5155; (b) Tang, L.; Ding, L.; Chang, W.-X.; Li, J. *Tetrahedron Lett.* **2006**, *47*, 303; (c) Yoo, C. L.; Fettinger, J. C.; Kurth, M. J. *J. Org. Chem.* **2005**, *70*, 6941.
- (a) Selvam, N. P.; Saravanan, C.; Muralidharan, D.; Perumal, P. T. *J. Heterocycl. Chem.* **2006**, *43*, 1379; (b) Selvam, N. P.; Perumal, P. T. *Tetrahedron* **2008**, *64*, 2972; (c) Selvam, N. P.; Saranya, S.; Perumal, P. T. *Can. J. Chem.* **2008**, *86*, 32; (d) Arumugam, P.; Perumal, P. T. *Chem. Lett.* **2006**, *35*, 635.
- Kuthan, J. *Adv. Heterocycl. Chem.* **1995**, *62*, 20.
- Rafael, G.; Antonio, G.; Marco-Contelles, J. *J. Chem. Res.* **2006**, *8*, 536.
- Jiarong, L.; Lijun, Z.; Dixin, S.; Qing, L.; Dong, W.; Chunxia, W.; Qi, Z.; Ling, Z.; Yanqiu, F. *Synlett* **2008**, 233.