



## 2,2'-Diamino-6,6'-dimethylbiphenyl as an efficient ligand in the palladium-catalyzed Suzuki–Miyaura and Mizoroki–Heck reactions

Jian-Mei Lu<sup>a</sup>, Hui Ma<sup>a</sup>, Sha-Sha Li<sup>b</sup>, Dan Ma<sup>a</sup>, Li-Xiong Shao<sup>a,\*</sup>

<sup>a</sup> College of Chemistry and Materials Engineering, Wenzhou University, Chashan University Town, Wenzhou, Zhejiang Province 325035, People's Republic of China

<sup>b</sup> Oujian College, Wenzhou University, Chashan University Town, Wenzhou, Zhejiang Province 325035, People's Republic of China

### ARTICLE INFO

#### Article history:

Received 28 February 2010

Received in revised form 20 April 2010

Accepted 23 April 2010

Available online 29 April 2010

#### Keywords:

2,2'-Diamino-6,6'-dimethylbiphenyl

Palladium

Suzuki–Miyaura reaction

Mizoroki–Heck reaction

Synthetic methods

### ABSTRACT

2,2'-Diamino-6,6'-dimethylbiphenyl was found to be an efficient ligand in the palladium-catalyzed Suzuki–Miyaura coupling reactions of aryl iodides, bromides, and chlorides and Mizoroki–Heck reactions of aryl iodides and bromides. Under appropriate conditions, all reactions gave the desired products in moderate to excellent yields. The ligand is inexpensive, air-stable and easy to available.

© 2010 Elsevier Ltd. All rights reserved.

### 1. Introduction

Palladium-catalyzed cross-coupling reactions are versatile and efficient methods for carbon–carbon bond formations.<sup>1</sup> Among them, the Suzuki–Miyaura and Mizoroki–Heck coupling reactions play important roles in modern synthetic chemistry.<sup>2,3</sup> Improvement of these reactions is greatly relied on the reactivity of the palladium catalyst by using increasing efficient supporting ligands. To date, many efforts are made to the search for more efficient ligands. During the past decades, the most common ligands used for these two coupling reactions are phosphine-based ones.<sup>4</sup> Since most of the phosphine-based ligands are air and/or moisture-sensitive, in recent years, phosphine-free ligand as *N*-heterocyclic carbenes (NHCs) have also been employed.<sup>5</sup> In addition, alkyl-substituted diamines, such as ethylenediamines and cyclohexane-1,2-diamines, acting as versatile ligands, have been widely used in the transition-metal catalyzed carbon–carbon bonds and carbon–heteroatom bonds formations during the last years.<sup>6,7</sup> As part of our ongoing research toward effective ligands for transition-metal catalyzed cross-coupling reactions for carbon–carbon bond formations, it was found that easily available 2,2'-diamino-6,6'-dimethylbiphenyl (L1),<sup>8</sup> is a useful ligand for the palladium-catalyzed Suzuki–Miyaura coupling reactions of aryl halides including

iodides, bromides, and chlorides and Mizoroki–Heck reactions of aryl iodides and bromides. Herein, we wish to report these results in detail.


### 2. Results and discussion

In initial investigations, we examined the Suzuki–Miyaura coupling reaction of the model substrate 4-methoxyphenyl iodide (**1a**) with 4-methylphenylboronic acid (**2a**), using 2,2'-diamino-6,6'-dimethylbiphenyl (L1) (10 mol %) (Scheme 1) as the ligand and Pd(OAc)<sub>2</sub> (5 mol %) as the catalyst in THF at room temperature for 48 h (Table 1). As can be seen from Table 1, of the bases screened, Cs<sub>2</sub>CO<sub>3</sub> showed the best result and the corresponding coupling product **3a** was obtained in 90% yield (Table 1, entry 5). For the other solvents as toluene, CH<sub>2</sub>Cl<sub>2</sub>, dioxane, DCE, and CH<sub>3</sub>CN, all showed inferior results compared to that of THF (Table 1, entries 8–12). When L1 was replaced by other analogues as L2,<sup>9</sup> L3,<sup>10</sup> L4,<sup>11</sup> L5,<sup>12</sup> and L6,<sup>12</sup> inferior results were found (Table 1, entries 13–17). Finally, it was found that if the reaction temperature was elevated to 40 °C, the corresponding reactions can be finished within 12 h with 96% yield using L1 as the ligand (Table 1, entry 18).

Thus, the optimal reaction conditions are using Pd(OAc)<sub>2</sub> (5 mol %) as the catalyst, L1 (10 mol %) as the ligand, Cs<sub>2</sub>CO<sub>3</sub> (2.0 equiv) as the base and THF (2.0 mL) as the solvent at the temperature of 40 °C.

With the optimal reaction conditions in hand, we firstly explored the scope and limitations of the reaction with a set of aryl


\* Corresponding author. Tel./fax: +86 577 86689300; e-mail address: [Shaolix@wzu.edu.cn](mailto:Shaolix@wzu.edu.cn) (L.-X. Shao).



Scheme 1. Ligands screened.

iodides and arylboronic acids. We were pleased to find out that all reactions afforded the desired coupling products **3** in high yields within 12 h and substituents, either electron-donating group as methoxy group or electron-withdrawing group as Cl atom on the phenyl ring of **1** have almost no significant effect on these reactions (Table 2, entries 1–3). 2-Thiophenyl iodide **1d** also led to the corresponding coupling product **3d** in high yield (Table 2, entry 4). Nevertheless, to our disappointment, when arylbromide as 4-methoxyphenyl bromide **1e** was used as the substrate, almost no desired product was obtained under the optimal reaction conditions (Table 2, entry 5). Gratifyingly, when the reaction temperature was elevated to 70 °C, consistent with the above studies, the reactions of aryl bromides **1** with arylboronic acids **2** also took place smoothly to give the corresponding products **3** in moderate to high yields (Table 2, entries 6–13). Substituents on the phenyl rings of

Table 1  
Optimization for the reaction conditions of **1a** with **2a**



Entry <sup>a</sup>	Ligand	Base	Solvent	Yield (%) <sup>b</sup>
1	L1	NaOH	THF	10
2	L1	KOH	THF	82
3	L1	CH <sub>3</sub> COOK	THF	9
4	L1	KHCO <sub>3</sub>	THF	3
5	L1	Cs <sub>2</sub> CO <sub>3</sub>	THF	90
6	L1	Na <sub>2</sub> CO <sub>3</sub>	THF	15
7	L1	K <sub>2</sub> CO <sub>3</sub>	THF	24
8	L1	Cs <sub>2</sub> CO <sub>3</sub>	Toluene	48
9	L1	Cs <sub>2</sub> CO <sub>3</sub>	CH <sub>2</sub> Cl <sub>2</sub>	36
10	L1	Cs <sub>2</sub> CO <sub>3</sub>	Dioxane	73
11	L1	Cs <sub>2</sub> CO <sub>3</sub>	DCE	24
12	L1	Cs <sub>2</sub> CO <sub>3</sub>	CH <sub>3</sub> CN	76
13	L2	Cs <sub>2</sub> CO <sub>3</sub>	THF	10
14	L3	Cs <sub>2</sub> CO <sub>3</sub>	THF	14
15	L4	Cs <sub>2</sub> CO <sub>3</sub>	THF	63
16	L5	Cs <sub>2</sub> CO <sub>3</sub>	THF	79
17	L6	Cs <sub>2</sub> CO <sub>3</sub>	THF	74
18 <sup>c</sup>	L1	Cs <sub>2</sub> CO <sub>3</sub>	THF	96


<sup>a</sup> Otherwise specified, all reactions were carried out using **1a** (0.5 mmol), **2a** (0.6 mmol), Pd(OAc)<sub>2</sub> (5 mol%), ligand (10 mol%), and base (2.0 equiv) in solvent (2.0 mL) at rt for 48 h.

<sup>b</sup> Isolated yields.

<sup>c</sup> The reaction was carried out at 40 °C for 12 h.

aryl bromides and arylboronic acids have little effect on the reaction. For instance, the reaction of 3-methoxyphenyl bromide **1f** with 4-chlorophenylboronic acid **2d** only gave the coupling product **3g** in 72% yield (Table 2, entry 11). It is worthy of noting that at this temperature (70 °C), Cl atom on the phenyl ring kept untouched (Table 2, entries 8 and 11).

Table 2  
Pd(OAc)<sub>2</sub>-catalyzed Suzuki–Miyaura coupling reaction of **1** with **2**



Entry <sup>a</sup>	<b>1</b> (R <sup>1</sup> /X)	<b>2</b> (R <sup>2</sup> )	Yield <sup>b</sup> (%)
1	<b>1a</b> (4-MeO/I)	<b>2b</b> (H)	<b>3b</b> (86)
2	<b>1b</b> (4-Cl/I)	<b>2c</b> (4-MeO)	<b>3c</b> (92)
3	<b>1c</b> (H/I)	<b>2c</b>	<b>3b</b> (92)
4	<b>1d</b>	<b>2c</b>	<b>3d</b> (95)
5 <sup>c</sup>	<b>1e</b> (4-MeO/Br)	<b>2b</b>	Trace
6	<b>1e</b>	<b>2b</b>	<b>3b</b> (95)
7	<b>1e</b>	<b>2a</b> (4-Me)	<b>3a</b> (91)
8	<b>1e</b>	<b>2d</b> (4-Cl)	<b>3c</b> (75)
9	<b>1e</b>	<b>2e</b> (3,5-Me, Me)	<b>3e</b> (94)
10	<b>1f</b> (3-MeO/Br)	<b>2b</b>	<b>3f</b> (94)
11	<b>1f</b>	<b>2d</b>	<b>3g</b> (72)
12	<b>1g</b> (4-F/Br)	<b>2c</b>	<b>3h</b> (90)
13	<b>1h</b> (H/Br)	<b>2c</b>	<b>3h</b> (94)


<sup>a</sup> Otherwise specified, all reactions were carried out using **1** (0.5 mmol), **2** (0.6 mmol), Pd(OAc)<sub>2</sub> (5 mol%), L1 (10 mol%), and Cs<sub>2</sub>CO<sub>3</sub> (1.0 mmol) in THF (2.0 mL) at 40 °C for 12 h (for iodides) or 70 °C for 24 h (for bromides).

<sup>b</sup> Isolated yields.

<sup>c</sup> The reaction was carried out at 40 °C.

It is known to all that despite the lower reactivity of aryl chlorides comparable to the common partners as organic bromides, iodides, and triflates, chlorides are arguably the most useful single class of substrates, because of their lower cost and the wider diversity of available compounds.<sup>13</sup> Encouraged by the success on the palladium-catalyzed Suzuki–Miyaura coupling reactions of aryl iodides and bromides with arylboronic acids using 2,2'-diamino-6,6'-dimethylbiphenyl (L1) as the ligand, we then turned our interests to the reactions of aryl chlorides. Optimization procedure was carried out using 4-methoxyphenyl chloride **1i** (0.5 mmol) and phenylboronic acid **2b** (0.75 mmol) as the substrates, Pd(OAc)<sub>2</sub> (5 mol%) as the catalyst, L1 (10 mol%) as the ligand, Cs<sub>2</sub>CO<sub>3</sub> (2.0 equiv) as the base and DMF as the solvent. The results are shown in Table 3. After several trials and errors, to our delight, it was found that the best result can be obtained using 1.0 mL DMF as the solvent at 110 °C and the corresponding coupling product **3b** was obtained in 75% yield (Table 3, entry 2). In other solvents as toluene and DMSO, only very lower yield or no product was obtained for the same reaction (Table 3, entries 5 and 6).


Under the reaction conditions using Pd(OAc)<sub>2</sub> (5 mol%) as the catalyst, L1 (10 mol%) as the ligand, Cs<sub>2</sub>CO<sub>3</sub> (2.0 equiv) as the base and DMF (1.0 mL) as the solvent at 110 °C, a variety of aryl chlorides and arylboronic acids were examined (Table 4). All the substituted aryl chlorides tested afforded the corresponding products **3** in moderate to good yields. Substituents on the phenyl ring of chlorides slightly affected the reaction and electron-withdrawing group on the phenyl ring of aryl chloride gave better result (Table 4, entries 1 vs 3).

**Table 3**  
Optimization for the Suzuki–Miyaura reaction of 4-methoxyphenyl chloride **1i**

Entry <sup>a</sup>	Solvent (mL)	Yield <sup>b</sup> (%)
1	DMF (2.0)	54
2	DMF (1.0)	75
3	DMF (0.5)	21
4	DMF (0.25)	13
5	Toluene (1.0)	5
6	DMSO (1.0)	—

<sup>a</sup> All reactions were carried out using **1** (0.5 mmol), **2** (0.75 mmol), Pd(OAc)<sub>2</sub> (5 mol %), L1 (10 mol %), and Cs<sub>2</sub>CO<sub>3</sub> (1.0 mmol) at 110 °C for 48 h.

<sup>b</sup> Isolated yields.


**Table 4**  
Suzuki–Miyaura reaction of chlorides **1**

Entry <sup>a</sup>	<b>1</b> (R <sup>1</sup> )	<b>2</b> (R <sup>2</sup> )	Yield <sup>b</sup> (%)
1	<b>1j</b> (4-COCH <sub>3</sub> )	<b>2b</b> (H)	<b>3i</b> (66)
2	<b>1k</b> (H)	<b>2c</b> (4-MeO)	<b>3b</b> (75)
3	<b>1l</b> (3-MeO)	<b>2b</b>	<b>3f</b> (54)
4	<b>1m</b> (2-Me)	<b>2c</b>	<b>3j</b> (59)
5	<b>1n</b> (4-Me)	<b>2c</b>	<b>3a</b> (62)


<sup>a</sup> All reactions were carried out using **1** (0.5 mmol), **2** (0.75 mmol), Pd(OAc)<sub>2</sub> (5 mol %), L1 (10 mol %), and Cs<sub>2</sub>CO<sub>3</sub> (1.0 mmol) in DMF (1.0 mL) at 110 °C for 48 h.

<sup>b</sup> Isolated yields.

Further, we investigated the Mizoroki–Heck coupling reaction using 2,2′-diamino-6,6′-dimethylbiphenyl (L1) as the ligand and Pd(OAc)<sub>2</sub> as the catalyst. For our initial investigation, we chose iodo-benzene **1c** (2.5 mmol) and methyl acrylate **4a** (3.0 mmol) as the model substrates. After several trials and errors, we were pleased to find out that in the presence of Pd(OAc)<sub>2</sub> (1 mol %), L1 (1 mol %), K<sub>2</sub>CO<sub>3</sub> (2 equiv) at 90 °C, the corresponding coupling product **5a** can be obtained in 94% yield (Scheme 2) (please see in the Supplementary data for the details).

**Scheme 2.** Optimal conditions for the coupling of **1c**.

A series of aryl halides and alkenes were then subjected to the reaction under the optimal conditions described above (Table 5). The results indicated that the combination of Pd(OAc)<sub>2</sub> and L1 is also efficient for the Mizoroki–Heck coupling reaction of aryl iodides. Both alkenes, such as methyl acrylate **4a** and styrene **4b** could be efficiently converted to the corresponding coupling products **5** in good to high yields (Table 5, entries 1–6). Besides aryl iodides, aryl bromides could also give products **5** in moderate to good yields under the similar reaction conditions (Table 5, entries 7–9). Nonetheless, the combination of Pd(OAc)<sub>2</sub> and L1 showed no activity for the Mizoroki–Heck coupling reaction of aryl chlorides.

**Table 5**  
Mizoroki–Heck reaction of aryl halides **1** with alkenes **4**

Entry <sup>a</sup>	<b>1</b> (R <sup>1</sup> /X)	<b>4</b> (R <sup>2</sup> )	Temp/Time	Yield <sup>b</sup> (%)
1	<b>1a</b> (4-MeO/I)	<b>4a</b> (CO <sub>2</sub> Me)	90 °C/36 h	<b>5b</b> , 99
2	<b>1b</b> (4-Cl/I)	<b>4a</b>	90 °C/24 h	<b>5c</b> , 82
3	<b>1c</b> (H/I)	<b>4b</b> (Ph)	110 °C/24 h	<b>5d</b> , 71
4	<b>1d</b> (2-I)	<b>4a</b>	90 °C/36 h	<b>5e</b> , 80
5	<b>1a</b>	<b>4b</b>	110 °C/24 h	<b>5f</b> , 64
6	<b>1b</b>	<b>4b</b>	110 °C/32 h	<b>5g</b> , 67
7 <sup>c</sup>	<b>1h</b> (H/Br)	<b>4a</b>	140 °C/24 h	<b>5a</b> , 57
8 <sup>c</sup>	<b>1o</b> (4-Me/Br)	<b>4a</b>	140 °C/24 h	<b>5h</b> , 68
9 <sup>c</sup>	<b>1p</b> (4-Cl/Br)	<b>4a</b>	110 °C/24 h	<b>5c</b> , 61

<sup>a</sup> Otherwise specified, all reactions were carried out using **1** (2.5 mmol), **4** (3.0 mmol), Pd(OAc)<sub>2</sub> (1 mol %), L1 (1 mol %), and K<sub>2</sub>CO<sub>3</sub> (5.0 mmol) in DMF (5.0 mL).

<sup>b</sup> Isolated yields.

<sup>c</sup> Pd(OAc)<sub>2</sub> (5 mol %) and L1 (5 mol %) were used.

### 3. Conclusion

In conclusion, we have developed a new combination of Pd(OAc)<sub>2</sub>-2,2′-diamino-6,6′-dimethylbiphenyl, which is proved to be efficient for Suzuki–Miyaura reactions of various aryl iodides, bromides, and chlorides with arylboronic acids and Mizoroki–Heck olefination of aryl iodides and bromides with various substituted alkenes under the appropriate reaction conditions. It also demonstrates great tolerance to a wide range of groups on all substrates of aryl halides, arylboronic acids and alkenes. The ligand is inexpensive, air-stable and easy to available.

## 4. Experimental section

### 4.1. General methods

<sup>1</sup>H and <sup>13</sup>C NMR spectra were recorded on a Bruker Avance-300 MHz spectrometer for solution in CDCl<sub>3</sub> with tetramethylsilane (TMS) as an internal standard; *J*-values are in hertz. THF, toluene, and dioxane were distilled from sodium (Na) under nitrogen (N<sub>2</sub>) atmosphere. DMF, CH<sub>3</sub>CN, CH<sub>2</sub>Cl<sub>2</sub>, and 1,2-dichloroethane (DCE) were distilled from CaH<sub>2</sub> under nitrogen (N<sub>2</sub>) atmosphere. Commercially obtained reagents were used without further purification. Flash column chromatography was carried out using Huanghai 300–400 mesh silica gel at increased pressure.

## 4.2. Experimental procedures

**4.2.1. General procedure for the palladium-catalyzed Suzuki–Miyaura cross-coupling reaction of arylboronic acids with aryl iodides and bromides.** (If Aryl halide is liquid). Under N<sub>2</sub> atmosphere, arylboronic acid **2** (0.6 mmol), Pd(OAc)<sub>2</sub> (5 mol %), ligand L1 (10 mol %), Cs<sub>2</sub>CO<sub>3</sub> (2.0 equiv), and THF (2.0 mL) were added into a Schlenk reaction tube, then aryl halide (0.5 mmol) was added. The mixture was stirred at 40 °C for 12 h (for iodide) or 70 °C for 24 h (for bromide). Then the solvent was removed under reduced pressure and the residue was purified by a flash column chromatography (SiO<sub>2</sub>).

(If Aryl halide is solid). Under N<sub>2</sub> atmosphere, aryl halide **1** (0.5 mmol), arylboronic acid **2** (0.6 mmol), Pd(OAc)<sub>2</sub> (5 mol %), ligand L1 (10 mol %), Cs<sub>2</sub>CO<sub>3</sub> (2.0 equiv), and THF (2.0 mL) were added into a Schlenk reaction tube. The mixture was stirred at 40 °C for 12 h (for iodide) or 70 °C for 24 h (for bromide). Then the solvent was removed under reduced pressure and the residue was purified by a flash column chromatography (SiO<sub>2</sub>).

**4.2.2. General procedure for the palladium-catalyzed Suzuki–Miyaura cross-coupling reaction of arylboronic acids with aryl chlorides.** Under N<sub>2</sub> atmosphere, arylboronic acid **2** (0.6 mmol), Pd(OAc)<sub>2</sub> (5 mol %), ligand L1 (10 mol %), Cs<sub>2</sub>CO<sub>3</sub> (2.0 equiv), and DMF (1.0 mL) were added into a Schlenk reaction tube, then aryl chloride **1** (0.5 mmol) was added. The mixture was stirred at 110 °C for 48 h. Then the solvent was diluted with EtOAc, washed with saturated brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and purified by a flash column chromatography (SiO<sub>2</sub>).

**4.2.2.1. Compound 3a<sup>14</sup>.** A white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz, TMS) δ 2.36 (s, 3H, CH<sub>3</sub>), 3.80 (s, 3H, OCH<sub>3</sub>), 6.94 (d, J=8.7 Hz, 2H, Ar), 7.20 (d, J=7.5 Hz, 2H, Ar), 7.43 (d, J=7.5 Hz, 2H, Ar), 7.49 (d, J=8.7 Hz, 2H, Ar). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz, TMS) δ 21.0, 55.2, 114.1, 126.5, 127.9, 129.4, 133.6, 136.3, 137.9, 158.9.

**4.2.2.2. Compound 3b<sup>14</sup>.** A white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz, TMS) δ 3.79 (s, 3H, OCH<sub>3</sub>), 6.94 (d, J=8.4 Hz, 2H, Ar), 7.27 (t, J=6.9 Hz, 1H, Ar), 7.39 (t, J=7.5 Hz, 2H, Ar), 7.49–7.54 (m, 4H, Ar). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz, TMS) δ 55.2, 114.1, 126.6, 126.7, 128.1, 128.7, 133.6, 140.7, 159.1.

**4.2.2.3. Compound 3c<sup>14</sup>.** A white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz, TMS) δ 3.84 (s, 3H, OCH<sub>3</sub>), 6.97 (d, J=9.0 Hz, 2H, Ar), 7.37 (d, J=9.0 Hz, 2H, Ar), 7.45–7.50 (m, 4H, Ar). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz, TMS) δ 55.3, 114.2, 127.9, 128.0, 128.8, 132.4, 132.6, 139.2, 159.3.

**4.2.2.4. Compound 3d<sup>15</sup>.** A white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz, TMS) δ 3.78 (s, 3H, OCH<sub>3</sub>), 6.88 (d, J=8.7 Hz, 2H, Ar), 7.00–7.03 (m, 1H, Ar), 7.16–7.18 (m, 2H, Ar), 7.51 (d, J=9.0 Hz, 2H, Ar). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz, TMS) δ 55.2, 114.2, 122.0, 123.7, 127.41, 127.3, 127.8, 144.3, 159.1.

**4.2.2.5. Compound 3e<sup>16</sup>.** A white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz, TMS) δ 2.37 (s, 3H, CH<sub>3</sub>), 3.84 (s, 3H, OCH<sub>3</sub>), 6.94–6.97 (m, 3H, Ar), 7.17 (s, 2H, Ar), 7.51 (d, J=9.0 Hz, 2H, Ar). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz, TMS) δ 21.4, 55.2, 114.0, 124.6, 128.1, 128.3, 133.9, 138.1, 140.8, 158.9.

**4.2.2.6. Compound 3f<sup>17</sup>.** A colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz, TMS) δ 3.83 (s, 3H, OCH<sub>3</sub>), 6.86–6.90 (m, 1H, Ar), 7.11–7.19 (m, 2H, Ar), 7.30–7.36 (m, 2H, Ar), 7.39–7.44 (m, 2H, Ar), 7.56–7.59 (m, 2H, Ar). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz, TMS) δ 55.2, 112.6, 112.8, 119.6, 127.1, 127.4, 128.7, 129.7, 141.0, 142.7, 159.9.

**4.2.2.7. Compound 3g<sup>18</sup>.** A colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz, TMS) δ 3.83 (s, 3H, OCH<sub>3</sub>), 6.87–6.91 (m, 1H, Ar), 7.05–7.13 (m, 2H, Ar), 7.30–7.39 (m, 3H, Ar), 7.47–7.51 (m, 2H, Ar). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz, TMS) δ 55.2, 112.7, 112.8, 119.4, 128.4, 128.8, 129.9, 133.4, 139.4, 141.4, 159.9.

**4.2.2.8. Compound 3h<sup>15</sup>.** A white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz, TMS) δ 3.85 (s, 3H, OCH<sub>3</sub>), 6.96–6.98 (m, 2H, Ar), 7.07–7.13 (m, 2H, Ar), 7.45–7.51 (m, 4H, Ar). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz, TMS) δ 55.2, 114.2, 115.5 (d, J<sub>C–F</sub>=21.4 Hz), 128.0, 128.1 (d, J<sub>C–F</sub>=7.7 Hz), 132.7, 136.9 (d, J<sub>C–F</sub>=3.2 Hz), 159.0, 162.0 (d, J<sub>C–F</sub>=244.3 Hz).

**4.2.2.9. Compound 3i<sup>19</sup>.** A white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz, TMS) δ 2.62 (s, 3H, CH<sub>3</sub>), 7.36–7.49 (m, 3H, Ar), 7.60–7.68 (m, 4H, Ar), 8.02 (d, J=8.1 Hz, 2H, Ar). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz, TMS) δ 26.6, 127.1, 127.2, 128.2, 128.8, 128.9, 135.8, 139.8, 145.7, 197.6.

**4.2.2.10. Compound 3j<sup>20</sup>.** A colorless oil. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz, TMS) δ 2.27 (s, 3H, CH<sub>3</sub>), 3.83 (s, 3H, OCH<sub>3</sub>), 6.94 (d, J=9.0 Hz, 2H, Ar), 7.20–7.25 (m, 6H, Ar). <sup>13</sup>C NMR (CDCl<sub>3</sub>, 75 MHz, TMS) δ 20.5, 55.2, 113.5, 125.7, 126.9, 129.9, 130.2, 130.3, 134.4, 135.4, 141.5, 158.5.

**4.2.3. General procedure for the palladium-catalyzed Heck–Mizoroki cross-coupling reaction of aryl halides with alkenes.** Under N<sub>2</sub> atmosphere, Pd(OAc)<sub>2</sub> (1 mol %), L1 (1 mol %), and K<sub>2</sub>CO<sub>3</sub> (2.0 equiv) and DMF (5.0 mL) were added into a Schlenk reaction tube successively, then aryl halides (2.5 mmol) and alkenes (3.0 mmol) were added. The mixture was stirred at the temperature and times listed in Scheme 2 and Table 5. Then the reaction solution was diluted with EtOAc, washed with saturated brine, dried over anhydrous Na<sub>2</sub>SO<sub>4</sub> and purified by a flash column chromatography (SiO<sub>2</sub>).

**4.2.3.1. Compound 5a<sup>21</sup>.** A white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz, TMS) δ 3.80 (s, 3H, OMe), 6.44 (d, J=16.2 Hz, 1H), 7.37–7.39 (m, 3H, Ar), 7.50–7.54 (m, 2H, Ar), 7.70 (d, J=16.2 Hz, 1H).

**4.2.3.2. Compound 5b<sup>22</sup>.** A white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz, TMS) δ 3.79 (s, 3H, OMe), 3.83 (s, 3H, OMe), 6.31 (d, J=15.9 Hz, 1H), 6.90 (d, J=8.7 Hz, 2H, Ar), 7.47 (d, J=8.7 Hz, 2H, Ar), 7.65 (d, J=15.9 Hz, 1H).

**4.2.3.3. Compound 5c<sup>22</sup>.** A white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz, TMS) δ 3.81 (s, 3H, OMe), 6.41 (d, J=15.9 Hz, 1H), 7.36 (d, J=8.7 Hz, 2H, Ar), 7.46 (d, J=8.7 Hz, 2H, Ar), 7.64 (d, J=15.9 Hz, 1H).

**4.2.3.4. Compound 5d<sup>22</sup>.** A white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz, TMS) δ 7.07 (s, 2H), 7.19–7.34 (m, 6H, Ar), 7.46–7.49 (m, 4H, Ar).

**4.2.3.5. Compound 5e<sup>23</sup>.** A white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz, TMS) δ 3.78 (s, 3H, OMe), 6.23 (d, J=15.6 Hz, 1H), 7.03 (dd, J=3.9, 5.1 Hz, 1H), 7.24 (d, J=3.9 Hz, 1H), 7.36 (d, J=5.1 Hz, 1H), 7.78 (d, J=15.6 Hz, 1H).

**4.2.3.6. Compound 5f<sup>24</sup>.** A white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz, TMS) δ 3.80 (s, 3H, OMe), 6.89 (d, J=7.8 Hz, 2H, Ar), 6.96 (d, J=16.8 Hz, 1H), 7.06 (d, J=16.8 Hz, 1H), 7.20–7.36 (m, 3H, Ar), 7.44 (d, J=8.1 Hz, 2H, Ar), 7.48 (d, J=7.8 Hz, 2H, Ar).

**4.2.3.7. Compound 5g<sup>24</sup>.** A white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz, TMS) δ 7.05 (s, 1H), 7.06 (s, 1H), 7.24–7.52 (m, 9H, Ar).



4.2.3.8. **Compound 5h**<sup>25</sup>. A white solid. <sup>1</sup>H NMR (CDCl<sub>3</sub>, 300 MHz, TMS)  $\delta$  2.35 (s, 3H, Me), 3.78 (s, 3H, OMe), 6.39 (d,  $J=16.8$  Hz, 1H), 7.17 (d,  $J=8.1$  Hz, 2H, Ar), 7.40 (d,  $J=8.1$  Hz, 2H, Ar), 7.66 (d,  $J=16.8$  Hz, 1H).

## Acknowledgements

Financial support from the National Natural Science Foundation of China (No. 20702013) and the start-up fund of Wenzhou University is greatly acknowledged. J.-M.L. thanks the Department of Education of Zhejiang Province for financial support (No. Y200907072). D.M. thanks Science and Technology Department of Zhejiang Province for financial support (No. 2008R40G2250090).

## Supplementary data

Supplementary data associated with this article can be found in online version at doi:10.1016/j.tet.2010.04.099. These data include MOL files and InChIkeys of the most important compounds described in the article.

## References and notes

- (a) Trost, B. M. *Science* **1991**, *234*, 1471–1477; (b) Tsuji, J. *Palladium Reagents and Catalysis: Innovations in Organic Synthesis*; Wiley: New York, NY, 1995; (c) *Handbook of Organopalladium Chemistry for Organic Synthesis*; Negishi, E., de Meijere, A., Eds.; John Wiley: New York, NY, 2002.
- (a) Miyaura, N.; Yamada, K.; Suzuki, A. *Tetrahedron Lett.* **1979**, *20*, 3437–3440; (b) Miyaura, N.; Suzuki, A. *Chem. Rev.* **1995**, *95*, 2457–2483; (c) Suzuki, A. *J. Organomet. Chem.* **1999**, *576*, 147–168; (d) Miyaura, N. *J. Organomet. Chem.* **2002**, *653*, 54–57; (e) Miyaura, N. *Topics in Current Chemistry*; Springer: Berlin, 2002; Vol. 219; (f) Miyaura, N. In *Metal-Catalyzed Cross-Coupling Reactions*, 2nd ed.; de Meijere, A., Diederich, F., Eds.; Wiley-VCH: Weinheim, 2004; Chapter 2, pp 41–124; (g) Suzuki, A. *Chem. Commun.* **2005**, 4759–4763; (h) Alonso, F.; Beletskaya, I. P.; Yus, M. *Tetrahedron* **2008**, *64*, 3047–3101.
- (a) Heck, R. F. *J. Am. Chem. Soc.* **1968**, *90*, 5518–5526; (b) Heck, R. F.; Nolley, J. P. *J. Org. Chem.* **1972**, *37*, 2320–2322; (c) Beletskaya, I. P.; Cheprakov, A. V. *Chem. Rev.* **2000**, *100*, 3009–3066; (d) Yi, L.; Liebscher, J. *Chem. Rev.* **2007**, *107*, 133–173; (e) Ruvk, R. T.; Huffman, M. A.; Kim, M. M.; Shevlin, M.; Kandur, W. V.; Davies, W. *Angew. Chem., Int. Ed.* **2008**, *47*, 4711–4714; (f) Henriksen, S. T.; Norrby, P. O.; Päivi Kaukoranta, P.; Pher, G. *J. Am. Chem. Soc.* **2008**, *130*, 10414–10421.
- (a) Phan, N. T. S.; Van Der Sluys, M.; Jones, C. W. *Adv. Synth. Catal.* **2006**, *348*, 609–679; (b) Andersen, N. G.; Keay, B. A. *Chem. Rev.* **2001**, *101*, 997–1030; (c) *Applied Homogeneous Catalysis with Organometallic Compounds*; Cornils, B., Herrmann, W. A., Eds.; VCH: New York, NY, 1996; Vols. 1 and 2; (d) Parshall, G. W.; Ittel, S. D. *Homogeneous Catalysis*, 2nd ed.; John Wiley: New York, NY, 1992, Chapter 8; (e) Martin, R.; Buchwald, S. L. *Acc. Chem. Res.* **2008**, *41*, 1461–1473.
- For recent selected papers, see: (a) Zhang, X.-Q.; Qiu, Y.-P.; Rao, B.; Luo, M.-M. *Organometallics* **2009**, *28*, 3093–3099; (b) Morgan, B. P.; Galdamez, G. A.; Gilliard, R. J., Jr.; Smith, R. C. *Dalton Trans.* **2009**, 2020–2028; (c) Zhou, Y.-B.; Xi, Z.-X.; Chen, W.-Z.; Wang, D.-Q. *Organometallics* **2008**, *27*, 5911–5920; (d) Broggi, J.; Clavier, H.; Nolan, S. P. *Organometallics* **2008**, *27*, 5525–5531; (e) Diebolt, O.; Braunstein, P.; Nolan, S. P.; Cazin, C. S. J. *Chem. Commun.* **2008**, 3190–3192.
- For some selected papers, see: (a) Lu, S.-M.; Bolm, C. *Adv. Synth. Catal.* **2008**, *350*, 1101–1105; (b) Lee, M.-r.; Stahl, S. S.; Gellman, S. H. *Org. Lett.* **2008**, *10*, 5317–5319; (c) Correa, A.; Carril, M.; Bolm, C. *Chem.—Eur. J.* **2008**, *14*, 10919–10922; (d) Carril, M.; Correa, A.; Bolm, C. *Angew. Chem., Int. Ed.* **2008**, *47*, 4862–4865; (e) Correa, A.; Elmore, S.; Bolm, C. *Chem.—Eur. J.* **2008**, *14*, 3527–3529; (f) Correa, A.; Carril, M.; Bolm, C. *Angew. Chem., Int. Ed.* **2008**, *47*, 2880–2883.
- For selected papers using other ligands, please see: (a) Borhade, S. R.; Waghmode, S. B. *Tetrahedron Lett.* **2008**, *49*, 3423–3429; (b) Mino, T.; Shindo, H.; Kaned, T.; Koizumi, T.; Kasashima, Y.; Sakamoto, M.; Fujita, T. *Tetrahedron Lett.* **2009**, *50*, 5358–5360; (c) Firouzabadi, H.; Iranpoor, N.; Gholinejad, M. *Tetrahedron* **2009**, *65*, 7079–7084; (d) Lakshmi Kantam, M.; Srinivas, P.; Yadav, J.; Likhari, P. R.; Bhargava, S. *J. Org. Chem.* **2009**, *74*, 4882–4885; (e) Srinivas, P.; Likhari, P. R.; Maheswaran, M.; Sridhar, B.; Ravikumar, K.; Kantam, M. L. *Chem.—Eur. J.* **2009**, *15*, 1578–1581.
- (a) Liang, Y.-X.; Gao, S.; Wan, H.-H.; Wang, J.-W.; Chen, H.-L.; Zhang, Z.; Hu, X.-Q. *Tetrahedron: Asymmetry* **2003**, *14*, 1267–1273; (b) Gillespie, K. M.; Sanders, C. J.; Westmoreland, L.; Thickett, C. P.; Scott, P. J. *Org. Chem.* **2002**, *67*, 3450–3458.
- Amirnasr, M.; Kickelbick, G.; Dehghanpour, S. *Helv. Chim. Acta* **2006**, *89*, 274–284.
- Chen, Y.-X.; Chan, A. S. C.; Li, Y.-M.; Lam, K.-H. *Chin. J. Chem.* **2001**, *19*, 794–799.
- Miyano, S.; Nawa, M.; Mori, A.; Harukichi, H. *Bull. Chem. Soc. Jpn.* **1984**, *57*, 2171–2176.
- Greencia, E. C.; Horaguchi, T. *J. Heterocycl. Chem.* **2006**, *43*, 1441–1446.
- (a) Grushin, V. V.; Alper, H. In *Activation of Unreactive Bonds and Organic Synthesis*; Murai, S., Ed.; Springer: Berlin, 1999; pp 193–226; (b) Grushin, V. V.; Alper, H. *Chem. Rev.* **1994**, *94*, 1047–1062.
- Wang, J.-W.; Meng, F.-H.; Zhang, L.-F. *Organometallics* **2009**, *28*, 2334–2337.
- Hatakeyama, T.; Hashimoto, S.; Ishizuka, K.; Nakamura, M. *J. Am. Chem. Soc.* **2009**, *131*, 11949–11963.
- So, C. M.; Lee, H. W.; Lau, C. P.; Kwong, F. Y. *Org. Lett.* **2009**, *11*, 317–320.
- Scheuermann, G. M.; Rumi, L.; Steurer, P.; Bannwarth, W.; Mühlaupt, R. *J. Am. Chem. Soc.* **2009**, *131*, 8262–8270.
- Denmark, S. E.; Smith, R. C.; Chang, W.-T.; Muhuhi, J. M. *J. Am. Chem. Soc.* **2009**, *131*, 3104–3118.
- Han, Y.; Lee, L. J.; Huynh, H. V. *Organometallics* **2009**, *28*, 2778–2786.
- Yoshikai, N.; Matsuda, H.; Nakamura, E. *J. Am. Chem. Soc.* **2009**, *131*, 9590–9599.
- Lee, E. Y.; Kim, Y.; Lee, J. S.; Park, J. *Eur. J. Org. Chem.* **2009**, 2943–2946.
- Ahmed-Omer, B.; Barrow, D. A.; Wirth, T. *Tetrahedron Lett.* **2009**, *50*, 3352–3355.
- Toguem, S.-M. T.; Hussain, M.; Malik, I.; Villinger, A.; Langer, P. *Tetrahedron Lett.* **2009**, *50*, 4962–4964.
- McNulty, J.; Das, P. *Eur. J. Org. Chem.* **2009**, 4031–4035.
- Iwasawa, T.; Kamei, T.; Watanabe, S.; Nishiuchi, M.; Kawamura, Y. *Tetrahedron Lett.* **2008**, *49*, 7430–7433.