

## Report

# Congenital Dyserythropoietic Anemia Type I Is Caused by Mutations in Codanin-1

Orly Dgany,<sup>1,4,\*</sup> Nili Avidan,<sup>5,\*</sup> Jean Delaunay,<sup>6</sup> Tatyana Krasnov,<sup>1,4</sup> Lea Shalmon,<sup>1,4</sup> Hanna Shalev,<sup>7</sup> Tal Eidelitz-Markus,<sup>2</sup> Joseph Kapelushnik,<sup>7</sup> Daniel Cattan,<sup>8</sup> Alexandre Pariente,<sup>9</sup> Michel Tulliez,<sup>10</sup> Aurore Créten,<sup>6</sup> Pierre-Olivier Schischmanoff,<sup>6</sup> Achille Iolascon,<sup>11</sup> Eithan Fibach,<sup>12</sup> Ariel Koren,<sup>13</sup> Jochen Rössler,<sup>14</sup> Martine Le Merrer,<sup>15</sup> Isaac Yaniv,<sup>3,4</sup> Rina Zaizov,<sup>3,4</sup> Edna Ben-Asher,<sup>5</sup> Tsvyia Olender,<sup>5</sup> Doron Lancet,<sup>5</sup> Jacques S. Beckmann,<sup>5</sup> and Hannah Tamary<sup>1,3,4</sup>

<sup>1</sup>Pediatric Hematology Laboratory, Felsenstein Research Center, Rabin Medical Center, Departments of <sup>2</sup>Pediatrics E and <sup>3</sup>Pediatric Hematology Oncology, Schneider Children's Medical Center of Israel, Petah Tikva; <sup>4</sup>Sackler School of Medicine, Tel Aviv University, Tel Aviv; <sup>5</sup>Department of Molecular Genetics and Crown Human Genome Center, The Weizmann Institute of Science, Rehovot, Israel; <sup>6</sup>Service d'Hématologie, Hôpital de Bicêtre, Faculté de Médecine Paris-Sud, and INSERM U 473, Kremlin-Bicêtre, France; <sup>7</sup>Pediatrics Department, Soroka Medical Center, Beersheba, Israel; <sup>8</sup>Service d'Hépatogastroentérologie, Centre Hospitalier, Villeneuve-Saint-Georges, France; <sup>9</sup>Unité d'Hépatogastroentérologie, Centre Hospitalier, Pau, France; <sup>10</sup>Service d'Hématologie Biologique, Hôpital Henri Mondor, Créteil, France; <sup>11</sup>Pediatric Department, University of Foggia and CEINGE, University Federico II, Naples; <sup>12</sup>Department of Hematology, Hadassah University Hospital, Ein-Kerem, Jerusalem; <sup>13</sup>Pediatric Department B, Ha'Emek Medical Center, Afula, Israel; <sup>14</sup>Paediatric Haematology, Oncology, Children's University Hospital, Essen, Germany; and <sup>15</sup>Hôpital des Enfants Malades, Paris

Congenital dyserythropoietic anemias (CDAs) constitute a rare group of inherited red-blood-cell disorders associated with dysplastic changes in late erythroid precursors. CDA type I (CDAI [MIM 224120], gene symbol *CDAN1*) is characterized by erythroid pathological features such as internuclear chromatin bridges, spongy heterochromatin, and invagination of the nuclear membrane, carrying cytoplasmic organelles into the nucleus. A cluster of 45 highly inbred Israeli Bedouin with CDAI enabled the mapping of the *CDAN1* disease gene to a 2-Mb interval, now refined to 1.2 Mb, containing 15 candidate genes on human chromosome 15q15 (Tamary et al. 1998). After the characterization and exclusion of 13 of these genes, we identified the *CDAN1* gene through 12 different mutations in 9 families with CDAI. This 28-exon gene, which is transcribed ubiquitously into 4738 nt mRNA, was reconstructed on the basis of gene prediction and homology searches. It encodes codanin-1, a putative o-glycosylated protein of 1,226 amino acids, with no obvious transmembrane domains. Codanin-1 has a 150-residue amino-terminal domain with sequence similarity to collagens and two shorter segments that show weak similarities to the microtubule-associated proteins, MAP1B (neuraxin) and synapsin. These findings, and the cellular phenotype, suggest that codanin-1 may be involved in nuclear envelope integrity, conceivably related to microtubule attachments. The specific mechanisms by which codanin-1 underlies normal erythropoiesis remain to be elucidated.

Congenital dyserythropoietic anemias (CDAs), a group of inherited disorders associated with morphological and functional abnormalities of erythropoiesis, have been classified into three types (I–III), with some patients still

unassigned (Wickramasinghe 1997; Delaunay and Iolascon 1999). The autosomal recessive CDA type II (CDAIL [MIM 224100]), with more than 250 cases described to date (Iolascon et al. 2001), is the most common form. The disease gene maps to 20q11.2 in most studied families (Gasparini et al. 1997). The least common of the CDAs, the autosomal dominant CDA type III (CDAL [MIM 105600]), was localized to chromosome 15q22 upon linkage analysis of a large Swedish family (Lind et al. 1995).


CDA type I (CDAI [MIM 224120]) is an autosomal recessive disease. Patients with CDAI present with mod-

Received August 14, 2002; accepted for publication September 16, 2002; electronically published November 14, 2002.

Address for correspondence and reprints: Dr. Hannah Tamary, Pediatric Hematology Oncology, Schneider's Children Medical Center of Israel, Petah Tikva, 49202 Israel. E-mail: htamary@post.tau.ac.il

\* These authors contributed equally to this work.

© 2002 by The American Society of Human Genetics. All rights reserved. 0002-9297/2002/7106-0024\$15.00



**Figure 1** Physical map and genomic organization of the *CDAN1* locus. *a*, Relative positions of landmark microsatellite markers are indicated in the CDAI candidate interval defined by markers D15S779 and D15S778. The CDAI interval is indicated by green and brown arrows. An informative SNP (*pink*) within the newly defined transcript LOC146050 enabled us to restrict the linkage interval. The genes and transcripts identified are boxed: known genes in green, unknown transcripts in blue, inferred genes in purple, *CDAN1* in red. The expression patterns of these genes were tested by RT-PCR on RNA from erythroid precursor cells (Pope et al. 2000) and are marked by a plus sign (+) or a minus sign (–) in the Express lane. Resequenced genes are marked by a plus sign (+) or a minus sign (–) in the Reseq lane. *b*, Expanded view of the 15-kb segment showing the genomic organization of *CDAN1* and of two alternative transcripts isolated and sequenced from erythroid and fibroblast cells. Coding and noncoding exons are depicted as filled and open boxes, respectively. Red and blue exons are based on two partial transcripts, DKFZp434G2127 and BI855138, respectively. Green and pink exons are based on gene prediction and RT-PCR, respectively. The lengths of the corresponding transcripts and inferred proteins are indicated to the right.

erate-to-severe macrocytic anemia. Bone marrow aspirates reveal binuclear intermediate- and late-erythroid precursors as well as internuclear chromatin bridges. Ultrastructural erythroid features include spongy heterochromatin and invagination of the nuclear membrane, carrying cytoplasm and cytoplasmic organelles into the nucleus. Dysmorphic features, mainly syndactyly and the absence or hypoplasia of phalanges and nails, have also been observed in several patients (Wickramasinghe 1997). Arrest of DNA synthesis (Wickramasinghe 1997) and apoptotic features in erythroid precursors have been described (Tamary et al. 1996). Interferon  $\alpha 2$  was incidentally found to ameliorate the anemia and to partially reverse the morphological abnormalities of the erythroid precursors by an unknown mechanism (Lavabre-Bertrand et al. 1995; Wickramasinghe et al. 1997;

Parez et al. 2000). A cluster of 45 Israeli Bedouin with CDAI enabled mapping of the disease gene to a region between markers D15S779 and D15S778 (Tamary et al. 1998). All patients showed a similar clinical picture, and in all the subjects, the diagnosis was confirmed by bone marrow electron microscopy. DNA was extracted from whole blood, and RNA was extracted from the diagnostic bone marrow aspiration. All studies were approved by the institutional review board of Rabin Medical Center. Using new polymorphic markers from genomic clones (AC019011, AC018924) and an informative SNP within the defined transcript for the putative transcription factor LOC146050, we further refined the critical CDAI interval to 1.2 Mb (fig. 1a).

In an attempt to identify the underlying mutations, we proceeded with systematic *in silico* analyses through

**Table 1****CODANIN-1 Mutations in Patients with CDAI**

ID, Family Origin, Genotype, and Exon <sup>a</sup>	Nucleotide Position <sup>b</sup>	Amino Acid Change	Frequency	Detection Method <sup>c</sup>
I, Israeli Bedouin, H: 24	3238C→T	Arg→Trp (R1040W)	3/376	+NcoI
II, French Polynesian, H: 26	3503C→T	Pro→Leu (P1129L)	0/142	-AciI
III, European, H: 12	1910A→G	Asn→Ser (N598S)	0/196	+DdeI
IV, European, CH: 14	2129C→T	Pro→Leu (P671L)	1/196	-MspI
19	2716T→A	Phe→Ile (F866I)	0/170	MALDI-TOF
V, European, CH: 14	2158C→T	Arg→stop (R680X)	0/200	+BtsI
14	2254C→T	Arg→Trp (R712W)	1/164	+BsrI
VI, European, CH: 23	3106C→T	Arg→stop (R996X)	0/196	-TaqI
24	3242A→T	Asp→Val (D1041V)	0/180	MALDI-TOF
VII, European, CH: 14	2206G→A	Glu→Lys (E696K)	0/160	MALDI-TOF
VIII, European, CH: 24	3221C→A	Ser→Phe (S1034F)	0/180	+StyI
IX, Arab, CH: 19	2719G→A	Val→Met (V867M)	1/180	+NlaIII

NOTE.—A total of 184 chromosomes from healthy nonrelated Bedouin individuals were analyzed as controls for the Bedouin mutation. For each of the additional mutations, close to 200 chromosomes from unrelated control individuals were analyzed. Three carriers, each for one mutation, were found. This is in accordance with the estimated frequency of CDAI heterozygotes in the European population.

<sup>a</sup> ID I included five families; all other IDs included one family each. H = homozygote; CH = compound heterozygote.

<sup>b</sup> Nucleotide position in *CODANIN-1* cDNA accession number AF525398.


<sup>c</sup> The mutation abolishes (-) or creates (+) a restriction site.

gene prediction and homology searches of the critical CDAI interval and found 17 transcripts or putative genes. Fifteen genes prevailed as potential candidates, based on their expression as determined by RT-PCR in erythroid cells grown in liquid culture (Pope et al. 2000), including two redefined and two newly characterized transcripts, namely UBR1 (GenBank accession number AF525401), TTBK (GenBank accession number AF525400), FLJ008 (GenBank accession number AF525397) and LOC146050 (GenBank accession number AF525399). The remaining two genes, *Calpain3* (*CAPN3*) and *TGM7*, were not considered, because they are not expressed in erythroid cells. The predicted exons were amplified from genomic DNA or from mRNA of erythroid precursors of one Bedouin patient with CDAI and of his healthy brother. These were subjected to sequence verification and mutation detection. Such systematic and comprehensive PCR and sequencing analyses were carried out for all coding regions of 14 of the 15 candidate genes (excluding KIAA1300, fig. 1a). All PCR reactions were performed according to standard procedures, and the products were subsequently sequenced using dye terminators of Big-Dyes kits on an ABI 3100 sequencer (Perkin-Elmer/Applied Biosystems). Sequence

comparisons were performed using the STADEN package (Bonfield et al. 1995) and Sequencher software (gene codes cooperation).

In the first 13 genes tested, no segregating mutation was identified; however, we eventually identified a homozygous mutation in the Bedouin patient in the fourteenth gene scrutinized (table 1). This gene was reconstructed on the basis of prediction programs and homology searches using two partial transcripts and two EST sequences (fig. 1b). The gene, thus identified as *CDAN1*, has 28 exons spanning 15 kb of genomic DNA (fig. 1b). It encompasses a putative 4,738-nucleotide mRNA encoding a protein of 1,226 amino acids, which we propose to name Codanin-1, for CDA type I. Subsequently, 11 additional mutations in eight other patient groups with CDAI were identified in this gene (table 1).

A single homozygous C→T substitution in exon 24, converting arginine to tryptophan at codon 1040 and creating an *NcoI* restriction site, was found in all 45 analyzed Bedouin patients with CDAI (fig. 2C). One of 184 Israeli Arab and 2 of 192 independent Bedouin control chromosomes were found to be carriers of this mutation (table 1). Subsequently, two other patients with CDAI from two unrelated families were found to be



**Figure 2** The mutation in a Bedouin family. *A*, Pedigree of a Bedouin family, showing the segregation of the founder chromosome with the disease. *B*, Sequence chromatogram from a wild-type (WT) and diseased (M) genotype showing a C→T substitution in exon 25, converting arginine to tryptophan at codon 863. *C*, Segregation of the Bedouin mutation, as assayed by *Nco*I restriction. A 412-bp PCR product is generated between the F and R primers. The mutation results in a gain of restriction site and, therefore, the restriction digest produced fragments of 175 and 237 bp, instead of the 412-bp band observed in healthy individuals.

homozygous for two additional *CDAN1* missense mutations: a proline→leucine substitution at codon 1129, found in a French Polynesian patient, and an asparagine→serine substitution at codon 598, present in three affected brothers from a family of French origin (table 1). It is noteworthy that the three sibs also exhibit sensorineural deafness and lack of motile sperm cells. These symptoms can be accounted for by a large 70-kb deletion 1 Mb distal to the *CDAN1* gene (N. Avidan, H. Tamary, O. Dgany, D. Cattan, A. Pariente, M. Thulliez, N. Borot, L. Moati, A. Barthelme, L. Shalmon, T. Krasnov, E. Ben-Asher, T. Olender, M. Khen, I. Yaniv, R. Zaizov, H. Shalev, J. Delaunay, M. Fellous, D. Lancet, and J. S. Beckmann, unpublished data). Hence, discovery of the A→G substitution in exon 12 indicates that these brothers carry, in addition to the 70-kb deletion, a missense mutation in *CDAN1* responsible for the CDAI phenotype. It is surprising to find tightly linked independent mutations on the same haplotype. However, cases of contiguous gene syndromes have been reported (Bitner-Glindzicz et al. 2000; Parvari et al. 2001; Shanske et al. 2001); given the growing number of mapped genetic disorders, such cases may be frequent enough to be encountered.


Three sporadic European patients with CDAI were found to be compound heterozygotes for *CDAN1* mutations. One patient has two distinct missense mutations, whereas the other two have one null and one missense mutation each (table 1). Segregation analysis of each mutation in respective families and genotyping of control chromosomes was done by restriction fragment analysis (NEBiolabs) or by mass spectrometry SNP scoring (Sauer et al. 2002) (primers available upon request). Segregation analysis demonstrated that all patients inherited one mutation from each parent, thus confirming that the patients are true compound heterozygotes. In addition, two patients of European descent and one of Arab descent were shown to carry one distinct missense mutation each (table 1). The unidentified mutations may be located in the promoter or introns of *CDAN1* or in some as-yet-unidentified exons. It is noteworthy that one additional patient (diagnosed by A.I.), who presented the clinical characteristics of CDAI, was found to be merely a compound heterozygote for two synonymous nucleotide changes (data not shown). For this patient with CDAI, pathogenic mutations remain to be identified either in *CDAN1* or in a second locus involved in the etiology of this disease. In any event, our data imply a considerable molecular homogeneity for CDAI.

In total, 12 different mutations in nine families with CDAI were identified: 4 in exon 14, 3 in exon 24, 2 in exon 19, and 1 each in exons 12, 23, and 26. The mutation-clustering pattern suggests either that these exons are more prone to mutation or that they encode essential functional domains (table 1). Furthermore, although

nonsense mutations were found in two compound heterozygotes, no homozygote for null-type mutations has been identified. This suggests that codanin-1 may have a unique and essential function, with very little redundancy. Future knockout studies and further screening of patients with CDAI should indicate whether the lack of this protein is lethal. In addition, the diversity of the severity of the disease among homozygous Bedouin patients with CDAI suggests that modifier genes may modulate the phenotype. Hence, at this stage, it is impossible to establish precise phenotype-genotype correlation.

Northern blot analysis was performed with a cDNA probe for exons 26–28 on RNA from eight different human tissues (Clontech). All tissues expressed the same 4.7-kb band (fig. 3A), suggesting that the gene is ubiquitously expressed and that the inferred mRNA is close to full length. A 369-bp shorter, alternatively spliced variant was found upon RT-PCR (data not shown) to be coexpressed in fibroblast cell line cultures from both patients with CDAI and healthy control individuals but not in erythroid cells. It is generated by in-exon splicing of exons 11–14 (fig. 1b), which preserves the original reading frame. This mRNA isoform encodes a putative protein of 1,103 amino acids. This major *CDAN1* mRNA appears to be expressed at a level at least tenfold lower than  $\beta$ -actin control as evidenced by their relative exposure times (fig. 3B). This observation is in agreement with the results of expression array experiments (data not shown) and the low number of *CDAN1* ESTs present in the databases.

BLAST homology searches (Altschul et al. 1990) against several genomic sequence databases showed no obvious human codanin-1 paralog but revealed two putative orthologs: one in the mouse syntenic region (84% identity) and one in *Fugu rubripes* (44% identity) (fig. 4). We also identified a putative ortholog in *Drosophila melanogaster* (GenBank accession number AF487678S2), which shares 23% identity with codanin-1. This protein, *vanaso*, has been proposed to be involved in the fly's olfactory behavior and is unlikely to be a functional *CDAN1* ortholog. Codanin-1 has no clear intracellular location-specific domains, such as a transmembrane or a signal peptide segments; it contains, however, numerous o-glycosylation consensus sites. In addition, the amino-terminal domain shares a significant homology with fibrillar collagens as revealed by FASTA (Pearson and Lipman 1988) (fig. 4). A multiple alignment was performed using the CLUSTALX program with the default parameters (Higgins et al. 1996). Further domain analysis was performed by BLOCKS (Henikoff et al. 2000), defining 13 interspecies sequence conservation regions (fig. 4). Subsequently, these BLOCKS were compared with all BLOCK database motifs, using the Local Alignment of Multiple Alignment method (Petrokovski 1996). Similarities to two domains of the microtubule-




**Figure 3** Northern blot analysis. Panels A and B show *CDAN1* and  $\beta$ -actin mRNA exposed for 1 week and overnight, respectively. The membrane was purchased from Clontech and was hybridized according to the manufacturer's instructions. The membrane was first probed with codanin-1 (exons 26–28 [fig. 1]) and subsequently re-probed with  $\beta$ -actin cDNA as an internal control.

associated protein MAP1B (Noble et al. 1989) (fig. 4, *green shaded blocks C and E*) and to two domains of fibrillar collagen triple-helix repeats (Silver et al. 2002) (fig. 4, *pink shaded blocks A and G*) were unveiled. Using CYRCA, more remote, indirect similarities were further detected between the triple-helix repeats blocks and synapsins blocks, a protein family known to be involved in binding diverse substrates including actin, spectrin, and microtubules (Angers et al. 2002). CYRCA is a method developed to identify blocks of potentially similar function and structure appearing in different contexts (Kunin et al. 2001). On the basis of the disease phenotype, which includes spongy heterochromatin and invagination of the nuclear membrane, and the protein similarity results, we speculate that codanin-1 might be involved in nuclear membrane integrity by connecting the nuclear membrane and microtubules (Dreger et al. 2001; Lippincott-Schwartz 2002).

Further analysis of codanin-1, using antibodies and animal models, should help to unravel the mechanism by which this protein ensures nuclear integrity during erythropoiesis, to clarify the pathogenesis of the disease, and possibly to facilitate the development of novel strategies for its management. Furthermore, identification of the gene involved in CDAI may facilitate the cloning of the genes involved in other types of CDA.





**Figure 4** Multiple alignment of human codanin-1. Multiple alignment with its putative mouse and fugu orthologs. Identical and similar residues in the multiple-alignment chart are represented by asterisks (\*) and colons (:), respectively; weak conservation is represented by periods (.). Conserved BLOCKS are confined within blue boxes. Shaded boxes are used for known block similarity: green, fibrillar collagens (IPB00885) (A, G); pink box, MAP1B (IPB000102) (C, E). Arrows indicate positions of mutated residues; green and blue arrows specify missense mutations within families I–VI and VII–IX, respectively, and red arrows specify null mutations. Note the cluster of mutations around aa 1034–41 and 866–867. The thick black arrow indicates the region with shared similarity to fibrillar collagens. The thin black line represents a split between exons.

## Acknowledgments

This work was supported in part by the Israel Ministry of Health Chief Scientist Grant (to H.T.), the Arc-en-Ceil/Keshet exchange program (to H.T. and J.D.), and by an Israel Ministry of Science Culture and Sports grant for the National Laboratory for Genome Infrastructure (to D.L.). It was also supported by the Crown Human Genome Center, the Alfred Krupp Foundation, the German-Israeli Foundation for Scientific Research and Development, and the Weizmann Institute Glasberg, Levy, Nathan Brunschwig and Levine funds (all to D.L.). R.Z. is incumbent of the J. Maus and G. Ceaseman-Maus Chair in Pediatric Hematology; D.L. holds the Ralph and Lois Silver Chair in Human Genomics; and J.S.B. is a recipient of the Hermann-Mayer chair. We also thank Profs. J.-P. Dommergues and G. Tchernia and Drs. F. Bernaudin, L. Roda, and H. Chambost for referring their patients to us; and Dr. M. Meunier-Rotival and Ms. C. Driancourt and M. A. Proust for their invaluable help.

## Electronic-Database Information

Accession numbers and URLs for data presented herein are as follows:

BLAST, <http://www.ncbi.nlm.nih.gov/BLAST/> (for codanin-1 homologs and orthologs, as well as genetic markers)

Blocks, <http://bioinformatics.weizmann.ac.il/blocks/> (for biological sequence analysis) (Henikoff et al. 2000)

GenBank, <http://www.ncbi.nlm.nih.gov/Genbank/> (for KIAA0770 [AB018313], DKFZP564G2022 [NM\_015497], ZFP106 [NM\_022473], SNAP23 [NM\_003825], FLJ10460 [NM\_018097], FLJ23375 [NM\_024956], CCNDBP1 [NM\_012142], EPB42 [NM\_000119], TGM5 [NM\_004245], UBR1 [AF525401], TTBK [AF525400], FLJ008 [AF525397], CDANI [AF525398], LOC146050 [AF525399], and vanaso [AF487678S2])

Local Alignment of Multiple Alignment (LAMA), [http://bioinformatics.weizmann.ac.il/blocks-bin/LAMA\\_search.sh](http://bioinformatics.weizmann.ac.il/blocks-bin/LAMA_search.sh) (for sequence motifs search) (Petrokovski 1996)

Online Mendelian Inheritance in Man (OMIM), <http://www.ncbi.nlm.nih.gov/Omim/> (for CDAI [MIM 224120], CDAII [MIM 224100], and CDAIII [MIM 105600])

## References

- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215:403–410
- Angers A, Fioravante D, Chin J, Cleary LJ, Bean AJ, Byrne JH (2002) Serotonin stimulates phosphorylation of alypsia synapsin and alters its subcellular distribution in sensory neurons. *J Neurosci* 22:5412–5422
- Bitner-Glindzicz M, Lindley KJ, Rutland P, Blaydon D, Smith VV, Milla PJ, Hussain K, Furth-Lavi J, Cosgrove KE, Shepherd RM, Barnes PD, O'Brien RE, Farndon PA, Sowden J, Liu XZ, Scanlan MJ, Malcolm S, Dunne MJ, Aynsley-Green A, Glaser B (2000) A recessive contiguous gene deletion causing infantile hyperinsulinism, enteropathy and deafness identifies the Usher type 1C gene. *Nat Genet* 26:56–60
- Bonfield JK, Smith K, Staden R (1995) A new DNA sequence assembly program. *Nucleic Acids Res* 23:4992–4999
- Delaunay J, Iolascon A (1999) The congenital dyserythropoietic anaemias. *Baillieres Best Pract Res Clin Haematol* 12:691–705
- Dreger M, Bengtsson L, Schoneberg T, Otto H, Hucho F (2001) Nuclear envelope proteomics: novel integral membrane proteins of the inner nuclear membrane. *Proc Natl Acad Sci USA* 98:11943–11948
- Gasparini P, del Giudice EM, Delaunay J, Totaro A, Granatiero M, Melchionda S, Zelante L, Iolascon A (1997) Localization of the congenital dyserythropoietic anemia II locus to chromosome 20q11.2 by genomewide search. *Am J Hum Genet* 61:1112–1116
- Henikoff J, Greene E, Petrokovski S, Henikoff S (2000) Increased coverage of protein families with the blocks database servers. *Nucleic Acids Res* 28:228–230
- Higgins DG, Thompson JD, Gibson TJ (1996) Using CLUSTAL for multiple sequence alignments. *Methods Enzymol* 266:383–402
- Iolascon A, Delaunay J, Wickramasinghe SN, Perrotta S, Gigante M, Camaschella C (2001) Natural history of congenital dyserythropoietic anemia type II. *Blood* 98:1258–1260
- Kunin V, Chan B, Sitbon E, Lithwick G, Petrokovski S (2001) Consistency analysis of similarity between multiple alignments: prediction of protein function and fold structure from analysis of local sequence motifs. *J Mol Biol* 307:939–949
- Lavabre-Bertrand T, Blanc P, Navarro R, Saghroun M, Vanereau H, Braun M, Wagner A, Taib J, Lavabre-Bertrand C, Navarro M (1995) Alpha-interferon therapy for congenital dyserythropoiesis type I. *Br J Haematol* 89:929–932
- Lind L, Sandstrom H, Wahlin A, Eriksson M, Nilsson-Sojka B, Sikstrom C, Holmgren G (1995) Localization of the gene for congenital dyserythropoietic anemia type III, CDAN3, to chromosome 15q21-q25. *Hum Mol Genet* 4:109–112
- Lippincott-Schwartz J (2002) Cell biology: ripping up the nuclear envelope. *Nature* 416:31–32
- Noble M, Lewis SA, Cowan NJ (1989) The microtubule binding domain of microtubule-associated protein MAP1B contains a repeated sequence motif unrelated to that of MAP2 and tau. *J Cell Biol* 109:3367–3376
- Parez N, Dommergues M, Zupan V, Chambost H, Fieschi JB, Delaunay J, Miélot F, Cramer EM, Dommergues JP, Wickramasinghe SN, Tchernia G (2000) Severe congenital dyserythropoietic anaemia type 1: prenatal management, transfusion support and alpha-interferon therapy. *Br J Haematol* 110:420–423
- Parvari R, Brodyansky I, Elpeleg O, Moses S, Landau D, Hershkovitz E (2001) A recessive contiguous gene deletion of chromosome 2p16 associated with cystinuria and a mitochondrial disease. *Am J Hum Genet* 69:869–75
- Pearson WR, Lipman DJ (1988) Improved tools for biological sequence comparison. *Proc Natl Acad Sci USA* 85:2444–2448
- Petrokovski S (1996) Searching databases of conserved sequence regions by aligning protein multiple-alignments. *Nucleic Acids Res* 24:3836–3845
- Pope SH, Fibach E, Sun J, Chin K, Rodgers GP (2000) Two-phase liquid culture system models normal human adult

- erythropoiesis at the molecular level. *Eur J Haematol* 64: 292-303
- Sauer S, Gelfand DH, Boussicault F, Bauer K, Reichert F, Gut IG (2002) Facile method for automated genotyping of single nucleotide polymorphisms by mass spectrometry. *Nucleic Acids Res* 30:e22
- Shanske A, Ferreira JC, Leonard JC, Fuller P, Marion RW (2001) Hirschsprung disease in an infant with a contiguous gene syndrome of chromosome 13. *Am J Med Genet* 102: 231-236
- Silver F, Horvath I, Foran D (2002) Mechanical implications of the domain structure of fiber-forming collagens: comparison of the molecular and fibrillar flexibilities of the alpha1-chains found in types I-III collagen. *J Theor Biol* 216: 243-254
- Tamary H, Shalev H, Luria D, Shaft D, Zoldan M, Shalmon L, Gruinspan A, Stark B, Chaison M, Shinar E, Resnitzky P, Zaizov R (1996) Clinical features and studies of erythropoiesis in Israeli Bedouins with congenital dyserythropoietic anemia type I. *Blood* 87:1763-1770
- Tamary H, Shalmon L, Shalev H, Halil A, Dobrushin D, Ashkenazi N, Zoldan M, Resnitzky P, Korostishevsky M, Bonne-Tamir B, Zaizov R (1998) Localization of the gene for congenital dyserythropoietic anemia type I to a <1-cM interval on chromosome 15q15.1-15.3. *Am J Hum Genet* 62:1062-1069
- Wickramasinghe SN (1997) Dyserythropoiesis and congenital dyserythropoietic anemias. *Br J Haematol* 98:785-797
- Wickramasinghe SN, Hasan R, Smythe J (1997) Reduced interferon-alpha production by Epstein-Barr virus transformed B-lymphoblastoid cell lines and lectin-stimulated lymphocytes in congenital dyserythropoietic anaemia type I. *Br J Haematol* 98:295-298