

Mutations in Glucose Transporter 9 Gene *SLC2A9* Cause Renal Hypouricemia

Hirotsuka Matsuo,* Toshinori Chiba, Shushi Nagamori, Akiyoshi Nakayama, Hideharu Domoto, Kanokporn Phetdee, Pattama Wiriyasermkul, Yuichi Kikuchi, Takashi Oda, Junichiro Nishiyama, Takahiro Nakamura, Yuji Morimoto, Keiko Kamakura, Yutaka Sakurai, Shigeaki Nonoyama, Yoshikatsu Kanai, and Nariyoshi Shinomiya

(American Journal of Human Genetics 83, 744–751; December 12, 2008)

In the version of Table 1 published online ahead of the print issue, the final entry in the first column, “Frequency,” was incorrectly edited. The correct entry is 21,060 and appears in the table here. The table has now been corrected both in print and online. The journal regrets this error.

Table 1. Frequency of Hypouricemia of the Japan Maritime Self-Defense Force

Urate (mg/dl)	Frequency	Cumulative Frequency	Relative Frequency (%)	Cumulative Relative Frequency (%)
0.0–0.5	3	3	0.01	0.01
0.6–1.0	27	30	0.13	0.14
1.1–1.5	2	32	0.01	0.15
1.6–2.0	7	39	0.03	0.18
2.1–2.5	29	68	0.14	0.32
2.6–3.0	132	200	0.62	0.94
3.1–	21060	21260	99.06	100.00

*Correspondence: hmatsuo@ndmc.ac.jp

DOI 10.1016/j.ajhg.2008.11.013. ©2008 by The American Society of Human Genetics. All rights reserved.