

Weakly nonextensive thermostatics and the Ising model with long-range interactions

 R. Salazar^{1,a}, A.R. Plastino^{1,2,3}, and R. Toral^{1,4}
¹ Departament de Física, Universitat de les Illes Balears, 07071 Palma de Mallorca, Spain

² Faculty of Astronomy and Geophysics, National University La Plata, Casilla de Correo 727, La Plata 1900, Argentina

³ CONICET, Casilla de Correo 727, La Plata 1900, Argentina

⁴ Instituto Mediterraneo de Estudios Avanzados (IMEDEA), Campus UIB, 07071 Palma de Mallorca, Spain

Received 24 May 2000

Abstract. We introduce a new nonextensive entropic measure S_χ that grows like N^χ , where N is the size of the system under consideration. This kind of nonextensivity arises in a natural way in some N -body systems endowed with long-range interactions described by $r^{-\alpha}$ interparticle potentials. The power law (weakly nonextensive) behavior exhibited by S_χ is intermediate between (1) the linear (extensive) regime characterizing the standard Boltzmann-Gibbs entropy and (2) the exponential law (strongly nonextensive) behavior associated with the Tsallis generalized q -entropies. The functional S_χ is parametrized by the real number $\chi \in [1, 2]$ in such a way that the standard logarithmic entropy is recovered when $\chi = 1$. We study the mathematical properties of the new entropy, showing that the basic requirements for a well behaved entropy functional are verified, *i.e.*, S_χ possesses the usual properties of positivity, equiprobability, concavity and irreversibility and verifies Khinchin axioms except the one related to additivity since S_χ is nonextensive. For $1 < \chi < 2$, the entropy S_χ becomes superadditive in the thermodynamic limit. The present formalism is illustrated by a numerical study of the thermodynamic scaling laws of a ferromagnetic Ising model with long-range interactions.

PACS. 05.20.-y Classical statistical mechanics – 05.20.Gg Classical ensemble theory – 02.70.Lq Monte Carlo and statistical methods 75.10.Hk Classical spin models

1 Introduction

There is nowadays an intense research activity on the mathematical properties and physical applications of new versions of the Maximum Entropy Principle based on generalized or alternative entropic functionals [1–18]. This line of inquiry has been greatly stimulated by the work of Tsallis, who showed that it is possible to build up a mathematically consistent and physically meaningful generalization of the standard Boltzmann-Gibbs-Jaynes thermostatical formalism on the basis of a nonextensive entropic measure [1]. The main motivation behind Tsallis proposal is that there are important physical scenarios, such as self-gravitating systems [19,20], electron-plasma two dimensional turbulence [21], among many others, which are characterized by a nonextensive behavior: due to the long range of the relevant interactions some of the thermodynamic variables usually regarded as additive, such as the internal energy, lose their extensive character. This suggests that a *nonextensive (non-additive)* entropy functional might be appropriate for their thermostatical description. The Jaynes' maximum entropy

approach to statistical mechanics [22,23] suggests in a natural way the possibility of incorporating *alternative entropy functionals* to the variational principle. The new entropy functional introduced by Tsallis has the form [1]

$$S_q = \frac{1}{q-1} \left(1 - \sum_{i=1}^w p_i^q \right), \quad (1)$$

where $(p_i, i = 1, \dots, w)$ are the microstate probabilities describing the system under consideration and the entropic index q is any real number. The standard Boltzmann-Gibbs entropy $S = -\sum_{i=1}^w p_i \ln p_i$ is recovered in the limit $q \rightarrow 1$. The measure S_q is *nonextensive*. The entropy of a composite system $A \oplus B$ constituted by two subsystems A and B , independent in the sense that $p_{ij}^{(A \oplus B)} = p_i^{(A)} p_j^{(B)}$, verifies the Tsallis' q -additive relation

$$S_q(A \oplus B) = S_q(A) + S_q(B) + (1-q)S_q(A)S_q(B). \quad (2)$$

We see from the above equation that the Tsallis' parameter q can be regarded as a measure of the degree of nonextensivity. Many relevant mathematical properties of the standard thermostatics are verified by the Tsallis'

^a e-mail: rafael@imedea.uib.es

generalized formalism or can be appropriately generalized [1, 18]. Self-gravitating systems constituted the first physical problem discussed within Tsallis' nonextensive thermostatics [20] and Tsallis' theory has recently been applied to other physical problems [24–29]. One of the main consequences of the intensive effort devoted in recent years to the study of Tsallis theory is that there is now a growing consensus that there are many problems in statistical physics, biology, economics, and other areas, where a generalization of the standard approach based on Boltzmann-Gibbs-Jaynes extensive entropy might be useful. A comprehensive bibliography on the current research literature on Tsallis' theory and the statistical physics of nonextensive systems can be found in [30, 31].

Inspired on Tsallis' pioneering proposal, various nonextensive entropic measures endowed with interesting properties have been recently advanced [6–12]. Moreover, it has been proven that some physically relevant mathematical properties are shared by large families of entropic measures [3, 4]. The aim of the present work is to explore the possibility of developing a new thermostatical formalism different from the Tsallis one. Our proposal is based on a nonadditive entropic functional characterized by a degree of nonextensivity weaker than the one exhibited by Tsallis measure. As we shall see, Tsallis entropy varies exponentially with the size of the system under consideration, while the new measure introduced here only scales as a power of the size of the system. That is to say, our proposal is tantamount of considering a nonextensive regime intermediate between (1) the standard extensive one associated with Boltzmann-Gibbs entropy, and (2) the exponentially nonextensive one described by Tsallis formalism.

There is an important physical motivation for introducing an entropy endowed with power law nonextensivity. Systems with long range interactions constitute the physical scenarios where the need of a generalization of the standard thermostatical formalism can be more clearly appreciated. Consider a system of N particles in a d -dimensional (one particle) configuration space. If the dependence of the interparticle potential with the interparticle distance r is given by $r^{-\alpha}$, it can be shown [32] that the system's energy levels scale as $N\tilde{N}$, where

$$\tilde{N} = \frac{N^{1-\alpha/d} - \alpha/d}{1 - \alpha/d}. \quad (3)$$

Hence, for large N , the internal energy scales as a power of the size of the system. That is,

$$E \sim N^{2-\alpha/d} \quad (4)$$

In the case of extensive systems governed by short-range interactions, the internal energy E and the entropy S scale in the same way: they both grow linearly with N . On the other hand, the temperature T is an intensive variable and does not change with N . How can these scaling laws be generalized to the non-extensive setting? A possible path towards the alluded generalization starts with the

Helmholtz free energy,

$$F = E - TS. \quad (5)$$

From the above expression it seems reasonable to expect both E and TS to scale in the same fashion [32]. The only way to fulfill this expectation, if the standard extensive entropy is used, is to require that the temperature scales as [32, 33]

$$T \sim \tilde{N} \sim N^{1-\alpha/d}, \quad (6)$$

losing its intensive character.

It would be very appealing to have, within the nonextensive scenario, an entropy S_χ endowed with the same scaling law as the one exhibited by the energy. If, as it occurs for extensive systems, the entropy and the energy behave in the same way, the temperature would still be an intensive quantity. What we need in order to have a thermostatical formalism with these nice properties is an entropy functional with power law nonextensivity, scaling as

$$S_\chi \sim N^\chi \sim N^{2-\alpha/d}. \quad (7)$$

From the above discussion it is clear that we are going to assume that the exponent χ appearing in the entropic scaling law is related with d and α by

$$\chi = 2 - \alpha/d, \quad (8)$$

so that the physically motivated range of values for χ is

$$1 \leq \chi \leq 2. \quad (9)$$

The purpose of this work is to study the basic properties of a possible candidate for a weakly-nonextensive thermostatics, based on a new non-additive entropy different from the Tsallis q -entropy, and to consider its application to a magnetic system with long-range interactions. The paper is organized as follows. In Section 2 the exponential behavior of Tsallis entropy is analyzed and a power-law weakly nonextensive entropy is introduced. The basic mathematical properties of the new measure are studied in Section 3. Two-state systems are considered in Section 4. In Section 5 the weakly nonextensive entropy is applied to a ferromagnetic Ising model with long-range interactions. Finally, some conclusions are drawn in Section 6.

2 Exponential vs. power law nonextensivity

2.1 Tsallis entropy and exponential nonextensivity

The nonextensive behavior of Tsallis functional, encapsulated in equation (2), is the most important single property distinguishing Tsallis measure from the standard additive logarithmic entropy. An important consequence of relation (2) that has not been fully appreciated in the literature is that Tsallis entropy may vary exponentially with

the size of the system. In order to clarify the above assertion let us consider the Tsallis entropy S_q of a composite system $A = \oplus_{j=1}^N A^{(j)}$ consisting on N identical and independent subsystems $\{A^{(j)}, j = 1, \dots, N\}$. Let us assume that each subsystem has M microstates and is characterized by the same set of probabilities $(\tilde{p}_i, i = 1, \dots, M)$. The entropy $S_q(\oplus_{j=1}^N A^{(j)}) \equiv S_q(N)$ associated with the composite system is then given by

$$S_q(N) = \frac{1}{q-1} \left\{ 1 - \sum_{i_1 \dots i_N} (\tilde{p}_{i_1} \dots \tilde{p}_{i_N})^q \right\}, \quad (10)$$

where each subindex i_k under the summation sign runs from 1 to M . Since each subsystem is characterized by the same probabilities, the above expression can be cast under the guise

$$S_q(N) = \frac{1}{q-1} \left\{ 1 - \left(\sum_{i=1}^M \tilde{p}_i^q \right)^N \right\}, \quad (11)$$

which leads to

$$S_q(N) = \frac{1}{q-1} \{ 1 - [1 + (1-q)S_q(1)]^N \}, \quad (12)$$

where

$$S_q(1) = \frac{1}{q-1} \left(1 - \sum_{i=1}^M \tilde{p}_i^q \right) \quad (13)$$

is the Tsallis entropy associated with each individual subsystem.

The expression (12) is valid for any probability distribution $(\tilde{p}_1, \dots, \tilde{p}_M)$ characterizing the individual subsystems. Let us consider the simplest case corresponding to subsystems with only two states ($M = 2$) characterized by an equiprobability distribution ($\tilde{p}_i = 1/M, i = 1, \dots, M$). A system of N noninteracting spins constitutes an example of this situation. The explicit expression for the entropy of each subsystem is then $S_q(1) = (2^{1-q} - 1)/(1 - q)$, and the Tsallis measure corresponding to the composite system adopts the value (for $q < 1$ and large N),

$$S_q \sim \frac{1}{1-q} 2^{(1-q)N}. \quad (14)$$

If $q > 1$ the q -entropy tends to the constant limit value $\frac{1}{q-1}$ as $N \rightarrow \infty$. We are going to consider only the $q < 1$ regime. In that case the Tsallis entropy exhibits an exponential behavior as a function of the size N of the system. We have shown in [36] that Tsallis entropy in Ising models with long-range and short-range interactions follows the same exponential behavior with the size N of the system.

2.2 Power law nonextensive entropy

We believe that it is worth exploring the possibility of a thermostatical formalism based on a nonextensive

entropy S_χ growing as a power N^χ of the size of the system, instead of growing exponentially. In accordance with the physical arguments discussed in the Introduction (see Eqs. (8, 9)), we shall restrict our considerations to values of $\chi \in [1, 2]$. Let us assume that the functional S_χ associated with a given discrete probability distribution $\{p_i, i = 1, \dots, w\}$ is given by

$$S_\chi(p_1, \dots, p_w) = \sum_{i=1}^w \phi_\chi(p_i), \quad (15)$$

$\phi_\chi(p_i)$ being an appropriate function of the individual microstate probabilities p_i . Notice that the entropy given by equation (15) is a symmetric function of the probabilities p_i . In order to find a suitable expression for the function $\phi_\chi(z)$ it is enough to consider again the equiprobability distribution associated with a collection of N identical independent two-state subsystems. In that case we have $w = 2^N$ and

$$\begin{aligned} S_\chi &= w \phi_\chi(1/w) \\ &= 2^N \phi_\chi(2^{-N}) \\ &\sim N^\chi, \end{aligned} \quad (16)$$

which implies

$$\phi_\chi(2^{-N}) \sim 2^{-N} N^\chi. \quad (17)$$

The simplest choice for a function $\phi_\chi(z)$ complying with the above relation is

$$\phi_\chi(z) = z(-\ln z)^\chi. \quad (18)$$

Unfortunately, this function is not adequate for our purposes. Its second derivative $d^2\phi_\chi/dz^2$ lacks a definite sign within the relevant range of values of z and χ , leading thus to an entropy functional without a definite concavity. However, since we are only interested in the large N asymptotic regime, any function $\phi_\chi(z)$ behaving like (18) in the limit $z \rightarrow 0$ would do. As we shall see, the function

$$\phi_\chi(z) = \frac{1}{2} \left[z(-\ln z)^\chi + z(-\ln z)^{1/\chi} \right] \quad (19)$$

leads to the measure

$$S_\chi = \frac{1}{2} \sum_{i=1}^w \left[p_i(-\ln p_i)^\chi + p_i(-\ln p_i)^{1/\chi} \right], \quad (20)$$

which complies with all the basic properties of a well behaved entropy. It is clear that in the case $\chi = 1$ the standard entropy $S_1 = -\sum_{i=1}^w p_i \ln p_i$ is recovered.

The optimization of S_χ under the constraints imposed by normalization,

$$\sum_{i=1}^w p_i = 1, \quad (21)$$




Fig. 1. The real roots of $\phi''_\chi(z) = 0$ as a function of χ . There are no real roots within the interval $[1/\chi_c, \chi_c]$, where $\chi_c \approx 2.1762$.




Fig. 2. The function $F(z)$ defined by the equations (25, 26) for $\chi = 1.0, 1.25, 1.5, 1.75, 2.0$.

and the mean values

$$\langle \mathcal{A}^{(r)} \rangle \equiv \sum_{i=1}^w p_i \mathcal{A}_i^{(r)} = \mathcal{A}_\chi^{(r)} \quad (r = 1, \dots, R) \tag{22}$$

of a given set $\{\mathcal{A}^{(r)}\}$ of observable leads to the variational problem

$$\delta \left\{ S_\chi - \sum_{r=1}^R \beta_r \langle \mathcal{A}^{(r)} \rangle - \alpha \sum_{i=1}^w p_i \right\} = 0, \tag{23}$$

whose solution is of the form

$$p_i = F \left(\alpha + \sum_{r=1}^R \beta_r \mathcal{A}_i^{(r)} \right) \quad i = 1, \dots, w. \tag{24}$$

Here α and $\{\beta_r\}$ are appropriate Lagrange multipliers and $F(z)$ is the inverse function of

$$\phi'_\chi(z) = \frac{1}{2} \left\{ (-\ln z)^\chi + (-\ln z)^{\frac{1}{\chi}} - \chi(-\ln z)^{\chi-1} - \frac{1}{\chi}(-\ln z)^{\frac{1}{\chi}-1} \right\}, \tag{25}$$

which verifies the relation

$$F[\phi'_\chi(z)] = \phi'_\chi[F(z)] = z. \tag{26}$$

The function $F(z)$ is well defined because the second derivative of $\phi_\chi(z)$ is negative,

$$\begin{aligned} \frac{d^2 \phi_\chi}{dz^2} &= \frac{1}{2z} \left[\chi(\chi - 1)(-\ln z)^{\chi-2} \right. \\ &\quad \left. + \frac{1}{\chi} \left(\frac{1}{\chi} - 1 \right) (-\ln z)^{\frac{1}{\chi}-2} \right. \\ &\quad \left. - \chi(-\ln z)^{\chi-1} + \frac{1}{\chi}(-\ln z)^{\frac{1}{\chi}-1} \right] \\ &< 0, \end{aligned} \tag{27}$$

for $0 < z < 1$, and $1 < \chi < 2$. Actually, $\phi''_\chi(z)$ has a definite (negative) sign within the larger interval $[1/\chi_c, \chi_c]$, where $\chi_c \approx 2.1762$. This can be appreciated in Figure 1, where the real roots of $\phi''_\chi(z) = 0$ are depicted as a function of χ . We see that there are no real roots $\chi \in [1/\chi_c, \chi_c]$.

It is important to realize that the lack of a simple analytical expression for the function $F(z)$ does not constitute a serious conceptual problem, nor does it pose any practical difficulty for the numerical treatment of problems involving the Jaynes' maximum entropy distributions (24). The form of the function $F(z)$ is shown in Figure 2.

The canonical ensemble probability distribution associated with the entropy S_χ ,

$$p_i = F(\alpha + \beta \epsilon_i), \tag{28}$$

is obtained when the S_χ is extremalized under the constraints of normalization and the mean value of the energy,

$$\langle U \rangle = \sum_i p_i \epsilon_i, \tag{29}$$

where ϵ_i stands for the energy of the microstate i . The Lagrange multiplier β appearing in (28) is the one associated with the energy constraint and corresponds, within the present thermostatical formalism, to the inverse temperature. That is, taking Boltzmann constant $k = 1$, we have $\beta = 1/T$.

3 Main properties of the new entropy

3.1 Khinchin axioms

Khinchin proposed a set of four axioms [34], which are usually regarded as reasonable requirements for a well behaved information measure. Our entropy measure S_χ verifies the first three of them:

- (i) $S_\chi = S_\chi(p_1, \dots, p_w)$,
i.e., the entropy is a function of the probabilities p_i only.

- (ii) $S_\chi(p_1, \dots, p_w) \leq S_\chi(\frac{1}{w}, \dots, \frac{1}{w}) \equiv S_\chi^{\text{equipr.}}(w)$, *i.e.*, S_χ adopts its extreme at equiprobability (this property will be proved in Sect. 3.2).
- (iii) $S_\chi(p_1, \dots, p_w) = S_\chi(p_1, \dots, p_w, p_{w+1} = 0)$, that is, the entropy of a system does not change if we incorporate a new microstate with zero probability. This property, known as *expansibility*, is clearly verified since $\phi_\chi(0) = 0$.
- (iv) The fourth Khinchin axiom concerns the behavior of the entropy of a composite system in connection to the entropies of the subsystems. We will comment on this axiom later.

3.2 General mathematical properties

Let us now consider the properties related to positivity, certainty, concavity, equiprobability, additivity and irreversibility, which are some of the most important features characterizing an information or entropic measure [1, 8, 12, 13].

3.2.1 Positivity

It is plain from equation (19) that $\phi_\chi(0) = \phi_\chi(1) = 0$ and also that $\phi_\chi(p) \geq 0$ for $p \in [0, 1]$. This implies the *positivity condition*

$$S_\chi \geq 0. \tag{30}$$

3.2.2 Certainty

The equality symbol in equation (30) holds only at *certainty*, that is, when one microstate has probability one and all the other microstates have zero probability,

$$S_\chi(1, 0, \dots, 0) = S_\chi(0, 1, \dots, 0) = \dots = 0. \tag{31}$$

Indeed, S_χ vanishes *if and only if* we have certainty.

3.2.3 Concavity

Considering (p_1, \dots, p_w) as independent variables, the second partial derivatives of S_χ are

$$\frac{\partial^2 S_\chi}{\partial p_j \partial p_k} = \frac{\partial^2 S_\chi}{\partial^2 p_j} \delta_{jk} < 0 \quad \text{for } 0 < p_j < 1. \tag{32}$$

Expression (32) guarantees definite *concavity* over probability space (see, for instance, [23]).

3.2.4 Equiprobability

The probability distribution that extremizes S_χ under the normalization constraint is

$$p_i = F(\alpha) = 1/w \tag{33}$$

Therefore, since S_χ has negative concavity, it is *maximal at equiprobability*. A well behaved entropy should also be, at equiprobability, a monotonically increasing function of the number of states w . For large w we have $S_\chi(w) \sim (\ln w)^\chi$. Therefore, for large values of w , $S_\chi^{\text{equipr.}}(w)$ is an *increasing function of w* .

3.2.5 Nonextensivity

The nonextensive behavior of the entropy S_χ is determined by the relation of the entropy of a composite system with the individual entropies of its constituent subsystems. Let us consider systems A and B with associated probabilities $\{a_i, i = 1, \dots, w_A\}$ and $\{b_j, j = 1, \dots, w_B\}$, respectively. If systems A and B are independent, *i.e.*, the composite system $A \oplus B$ has associated probabilities $\{a_i b_j; i = 1, \dots, w_A; j = 1, \dots, w_B\}$, then the entropy $S_\chi(A \oplus B)$ of the composite system minus the sum of the entropies of its subsystems,

$$\Delta S_\chi(A, B) \equiv S_\chi(A \oplus B) - S_\chi(A) - S_\chi(B) \tag{34}$$

is the quantity characterizing the nonextensive features of the measure S_χ . When $\Delta S_\chi(A, B) > 0$ ($\Delta S_\chi(A, B) < 0$) we have superadditivity (subadditivity). From the examination of particular examples we conclude that $\Delta S_\chi(A, B)$ does not have always the same sign. However, the region of probability space where $\Delta S_\chi(A, B)$ is *positive* is much larger than the region where that quantity is negative. Furthermore, if we consider N identical subsystems (instead of just two of them), the region of probability space corresponding to subadditive behavior tends to vanish as N grows. Consequently, the entropy S_χ becomes superextensive in the thermodynamic limit. Particular examples illustrating these features of the measure S_χ are provided in Section 4. Moreover, in Section 5 we are going to consider the nonextensivity of S_χ in connection with the thermodynamic properties of an Ising model endowed with long-range interactions.

3.2.6 Irreversibility

One of the most important roles played by entropic functionals within theoretical physics is to characterize the “arrow of time”. When they verify an H -theorem, they provide a quantitative measure of macroscopic irreversibility. We will now show, for some simple systems, that the present measure S_χ satisfies an H -theorem, *i.e.*, its time derivative has a definite sign.

Let us calculate the time derivative of S_χ

$$\frac{dS_\chi}{dt} = \sum_{i=1}^w \frac{dp_i}{dt} \phi'_\chi(p_i), \tag{35}$$

for a system whose probabilities p_i evolve according to the master equation

$$\frac{dp_i}{dt} = \sum_{j=1}^w [P_{ji} p_j - P_{ij} p_i], \tag{36}$$




Fig. 3. Entropy S_χ for a two-state system ($w = 2$) as a function of p for different values of χ indicated on the figure.

where P_{ij} is the transition probability per unit time between microscopic configurations i and j . Assuming a system with a uniform equilibrium distribution and detailed balance, *i.e.*, $P_{ij} = P_{ji}$, we obtain from (35)

$$\frac{dS_\chi}{dt} = \frac{1}{2} \sum_{i=1}^w \sum_{j=1}^w P_{ij} (p_i - p_j) \left(\phi'_\chi(p_j) - \phi'_\chi(p_i) \right). \quad (37)$$

Since $\phi''_\chi(p_i) < 0$, the quantities $(p_i - p_j)$ and $(\phi'_\chi(p_j) - \phi'_\chi(p_i))$ have the same sign. Then we obtain

$$\frac{dS_\chi}{dt} \geq 0. \quad (38)$$

The equality holds for equiprobability, *i.e.*, at equilibrium, while in any other cases the entropy S_χ increases with time. Therefore, S_χ exhibits *irreversibility*.

3.2.7 Jaynes thermodynamic relations

It is noteworthy that, within the present Jaynes' maximum entropy formalism, the usual thermodynamical relations involving the entropy, the relevant mean values, and the associated Lagrange multipliers, *i.e.*,

$$\frac{\partial S_\chi}{\partial \langle \mathcal{A}^{(r)} \rangle} = \beta_r \quad (39)$$

are verified. Hence, our formalism exhibits the usual thermodynamical Legendre transform structure. Actually, this property is verified by a wide family of entropy functionals [3,4]. A particular important example of (39) is furnished by the canonical ensemble thermodynamic relation

$$\frac{\partial S_\chi}{\partial \langle U \rangle} = \beta = \frac{1}{T}. \quad (40)$$




Fig. 4. The subadditivity and superadditivity of S_χ . The quantity $(\Delta S_\chi)_{\text{rel.}}$ as a function of p and q , for a composite system constituted by the independent two-state subsystems A (with associated probabilities $\{q, 1 - q\}$), and B (with associated probabilities $\{p, 1 - p\}$).

4 Two-state systems

In order to illustrate some of the above properties, we consider a two-state system (with associated probabilities $\{p, 1 - p\}$). In this case, S_χ only depends on the variable p . In fact, from its definition, we have

$$S_\chi(p) = \frac{1}{2} \left[p(-\ln p)^\chi + p(-\ln p)^{\frac{1}{\chi}} + (1-p)(-\ln(1-p))^\chi + (1-p)(-\ln(1-p))^{\frac{1}{\chi}} \right], \quad (41)$$

The shape of $S_\chi(p)$ for different values of χ is shown in Figure 3, which exhibits the positivity and concavity of S_χ . In fact, from expression (41), the first derivative of $S_\chi(p)$ vanishes at $p = 1/2$ and $d^2 S_\chi / dp^2 < 0 \quad \forall p \in [0, 1]$. Since the second derivative is always negative, $S_\chi(p)$ is maximal at equiprobability. Moreover, as shown in the general case, taking into account the concavity of S_χ and that S_χ vanishes at the certainty, then S_χ is positive for all p .

The non-additivity of S_χ is illustrated in Figure 4 for two independent two-state systems A (with probabilities p and $1 - p$) and B (with probabilities q and $1 - q$), through the plot of the relative difference

$$(\Delta S_\chi)_{\text{rel.}} = [S_\chi(A \oplus B) - (S_\chi(A) + S_\chi(B))] / S_\chi(A \oplus B) \quad (42)$$

as a function of p and q . For most values of p and q the nonextensive measure S_χ behaves in a superadditive fashion. Only for values lying in a small region near the edges of the (p, q) -square does S_χ become subextensive.

The nonextensive behavior of S_χ in the thermodynamic limit can be illustrated by recourse to a system




Fig. 5. The superadditivity of S_χ in the thermodynamic limit. (a) The quantity $(\Delta S_\chi)_{\text{rel.}}$ associated with a system constituted by N identical two-state subsystems with probabilities p and $(1-p)$, as a function of p and for different values of N . (b) $(\Delta S_\chi)_{\text{rel.}}$ as a function of p for a system constituted by $N-1$ identical two-state subsystems with probabilities $\{\frac{1}{2}, \frac{1}{2}\}$ along with one extra two-state subsystem with probabilities p and $(1-p)$.

constituted by N two-state subsystem. Let us first assume that each one of the subsystems is described by the same probabilities p and $(1-p)$. Consequently, the entropy associated with the composite system is

$$S_\chi(N) = \frac{1}{2^N} \sum_{k=0}^N \binom{N}{k} p^k (1-p)^{N-k} \times \left[(-\ln(p^k (1-p)^{N-k}))^\chi + (-\ln(p^k (1-p)^{N-k}))^{\frac{1}{\chi}} \right] \quad (43)$$

It can be shown after some algebra that, for $0 < p < 1$,

$$[-\ln(1-p)]^\chi N^\chi < S_\chi(N) < [-\ln p]^\chi N^\chi + [-\ln p]^{1/\chi} N^{1/\chi}. \quad (44)$$

Hence, for any given value of $p \in (0, 1/2)$ there exist an M such that

$$N > M \implies S_\chi(N) > N S_\chi(1), \quad (45)$$

which means that S_χ becomes superadditive for large enough values of N . This is shown in Figure 5a, where the quantity

$$(\Delta S_\chi)_{\text{rel.}} = [S_\chi(N) - N S_\chi(1)] / S_\chi(N) \quad (46)$$

is depicted for different values of N . In a similar way, Figure 5b shows the behavior of $(\Delta S_\chi)_{\text{rel.}}$ as a function of p for a composite system consisting of $N-1$ two-state subsystems each with probabilities $\{\frac{1}{2}, \frac{1}{2}\}$ along with one extra subsystem with probabilities p and $(1-p)$.

5 Ferromagnetic Ising model with long-range interactions

In this section we apply the canonical thermostatics associated with the entropy S_χ to an N -body system described by an $r^{-\alpha}$ interparticle potential. The energy of this kind of systems scales as a power of the number of particles [32]. The consideration of systems exhibiting this power-law nonextensivity constituted our main motivation for introducing the measure S_χ . As argued in Section 1, an entropy endowed with the same nonextensive behavior as the one associated with the energy may lead to a thermostatical description preserving the intensive character of the temperature. In order to illustrate these ideas we are going to study a long-range Ising model described by the Hamiltonian

$$\mathcal{H} = J \sum_{i,j=1}^N \frac{1 - S_i S_j}{r_{ij}^\alpha}, \quad (S_i = \pm 1, \forall i), \quad (47)$$

where J is an appropriate coupling constant, and the sum runs over all the pairs of sites on a d -dimensional lattice with periodic boundary conditions and r_{ij} stands for the distance between the sites i and j . It is clear that the range of the interaction is determined by the value of the exponent α . In particular, the standard (short-range) first-neighbors interaction is recovered in the limit $\alpha \rightarrow \infty$, while the mean field approximation is obtained when $\alpha = 0$ (replacing J by J/N). These extreme cases illustrate, respectively, the two possible thermodynamic behaviors that, according to the value of α , are admitted by the system (47). On the one hand we have the extensive regime corresponding to $\alpha > d$. On the other hand we have the non-extensive regime associated with $\alpha < d$. In order to clarify this let us estimate the internal energy per particle at zero temperature. We have [32]

$$\frac{E}{N} \sim \int_1^\infty dr r^{d-1} r^{-\alpha}. \quad (48)$$

It is easy to see that the above integral converges if $\alpha > d$ and diverges if $0 \leq \alpha \leq d$. The following is a standard notation used in previous works:

$$\tilde{N} \equiv 1 + \frac{1}{d} \int_1^{N^{1/d}} dr r^{d-1} r^{-\alpha} = \frac{N^{1-\alpha/d} - \alpha/d}{1 - \alpha/d}, \quad (49)$$

Here we consider the non-extensive regime for the one-dimensional case $d = 1$ and $\alpha = 0.8$. We have performed numerical simulations by using a novel approach recently introduced [2] to study systems with long range interactions governed by generalized entropies as the one considered here. The method relies upon the calculation of the number of states, $\Omega(\epsilon_k)$, with a given energy ϵ_k .

Note that, the number of possible configurations and associated probabilities p_i is in general very large, $W = 2^N$ for Ising models for example. However the number of permitted energies or energy levels K is not so large, because there is a large number of states $\Omega(\epsilon_k)$ sharing the same energy ϵ_k . We can rewrite the sums in equations (20, 29) taking into account the K energy levels weighted by $\Omega(\epsilon_k)$:

$$S_\chi = \frac{1}{2} \sum_{k=1}^K \Omega(\epsilon_k) p(\epsilon_k) \quad (50)$$

$$\times [(-\ln p(\epsilon_k))^\chi + (-\ln p(\epsilon_k))^{1/\chi}], \quad (51)$$

$$\langle O \rangle = \sum_{k=1}^K \Omega(\epsilon_k) p(\epsilon_k) O(\epsilon_k).$$

Hence, the knowledge of the number of states $\Omega(\epsilon_k)$ allows, by the use of these expressions, the calculation of the entropy and any averages of interest.

To calculate the probabilities $p(\epsilon_k)$ we use the definition $p(\epsilon_k) = F(\alpha + \beta\epsilon_k)$, where the function F is computed numerically as the inverse function of $\phi'_\chi(z)$ (see Fig. 2).

To compute the histogram $\Omega(\epsilon_k)$, the ‘‘Histogram by Overlapping Windows (HOW)’’ method [2, 35, 36] is used. A naive way of computing the histogram consists in generating different system configurations randomly and counting how many times a configuration with energy ϵ_k appears. However, since the $\Omega(\epsilon_k)$ values span too many orders of magnitude it is not possible to find in this way a histogram over all the energy levels. The HOW method avoids this problem by generating system configurations only in a restricted energy interval and estimating the relative weights $\Omega(\epsilon_k)/\Omega(\epsilon_l)$ of these energy levels from the number of times they appear in the sample. From the overlap between energy intervals, one gets the complete $\Omega(\epsilon_k)$ function, apart from an irrelevant normalization factor. Details about the HOW method can be found in [35, 36].

Finally we note a particularity of this new formalism, which is that the determination of the α constant from the normalization condition $\sum_{i=1}^w p_i = 1$, requires to solve the following equation

$$\sum_{k=1}^K \Omega(\epsilon_k) F(\alpha + \beta\epsilon_k) = 1, \quad (52)$$

where ϵ_k , β , $\Omega(\epsilon_k)$ are input data for the equation. This equation for α was solved numerically using a dichotomic searching method.

Using the described procedure we have calculated the dependence over the temperature T of the internal energy $E(N, T)$, spontaneous magnetization $M(N, T) = |\sum_{i=1}^N s_i|$, entropy $S(N, T)$ and free energy $F(N, T) =$

$E - TS$ for the one-dimensional long-range Ising model in the non-extensive regime $\alpha = 0.8$ and the corresponding value $\chi = 1.2$ in the equation (8).

This system has been recently studied within the standard Boltzmann-Gibbs thermostatics [33], as well as within Tsallis non-extensive q -formalism [2]. In [33] it was numerically verified that the scaling laws of the main thermodynamical quantities associated with the Gibbs canonical ensemble are

$$\begin{aligned} E(N, T)/(N\tilde{N}) &= e(T/\tilde{N}), \\ M(N, T)/N &= m(T/\tilde{N}), \\ S(N, T)/N &= s(T/\tilde{N}), \\ F(N, T)/(N\tilde{N}) &= f(T/\tilde{N}). \end{aligned} \quad (53)$$

Since the energy and the entropy scale in different ways, the temperature T has to be scaled as T/\tilde{N} . A similar situation arises within Tsallis q -generalized formalism, the concomitant scaling laws being [2]

$$\begin{aligned} E_q(N, T)/(N\tilde{N}) &= e(TA_q^E(N)/(N\tilde{N})), \\ M_q(N, T)/N &= m(TA_q^E(N)/(N\tilde{N})), \\ S_q(N, T)/(A_q(N)) &= s(TA_q^S(N)/(N\tilde{N})), \\ F_q(N, T)/(N\tilde{N}) &= f(TA_q(N)/(N\tilde{N})), \end{aligned} \quad (54)$$

where [2]

$$\begin{aligned} A_q(N) &= (2^{N(1-q)} - 1)/(1 - q), \\ A_q^S(N) &= (2^{N|1-q|} - 1)/|1 - q|, \\ A_q^E(N) &= A_q(N)^2/A_q^S(N). \end{aligned} \quad (55)$$

The scaling laws corresponding to the generalized χ -canonical ensemble associated to the weakly nonextensive entropy S_χ are,

$$\begin{aligned} E(N, T)/(N\tilde{N}) &= e(TR_\chi(N)), \\ M(N, T)/N &= m(TR_\chi(N)), \\ S_\chi(N, T)/(N^\chi + N^{1/\chi}) &= s(TR_\chi(N)), \\ F(N, T)/(N\tilde{N}) &= f(TR_\chi(N)), \end{aligned} \quad (56)$$

where

$$R_\chi(N) = \frac{N^\chi + N^{1/\chi}}{N^\chi - (2 - \chi)N}. \quad (57)$$

It is plain from the above equation that

$$\lim_{N \rightarrow \infty} R_\chi(N) = 1, \quad (58)$$

so that for large enough values of N the scaling laws (56) become

$$\begin{aligned} E(N, T)/(N^\chi) &= e(T), \\ M(N, T)/N &= m(T), \\ S(N, T)_\chi/(N^\chi) &= s(T), \\ F(N, T)/(N^\chi) &= f(T). \end{aligned} \quad (59)$$




Fig. 6. Plot of (a) the internal energy and (b) the magnetization as a function of the scaled temperature. In the insert of (b) we plot with solid line the behavior of the scaling factor $R_\chi(N)$ for all the N values and by points for the actual simulated sizes $N = 100, 400, 1000, 2000, 3000$. We use here $\chi = 1.2$.

According to the above equations, the thermodynamic curves of the Ising models (47) computed with increasing N -values must collapse without the need of a temperature rescaling. Numerical evidence of this scaling behavior is provided by Figures 6, 7. It must be realized, however, that the N -values used are not large enough to see a complete collapse of the curves depicted. In order to reach complete collapse we need $R_\chi(N) \approx 1$ (see the insert of Fig. 6b).

We plot in Figure 6 the scaled curves for the internal energy and magnetization using the equations (56). In the insert of that plot we see the asymptotical behavior for the scaling factor $R_\chi(N)$, and with points we indicate the sizes plotted $N = 100, 400, 1000, 2000, 3000$. Unfortunately we would need very large sizes $N \sim 10^8$ to observe an actual scaling without any scale factor, that is $R_\chi \sim 1$. However from the Figure 6 one can see that there is an actual tendency to the collapse. In fact we expect that the final universal curves will resemble to those corresponding to $N = 2000, 3000$ and we see in the Figure 6a tendency towards a complete collapse.

Also in Figure 7 we show the collapse for the entropy in the main plot with the scaled relations equations (56) and in the insert without scaling. In addition we can see




Fig. 7. Plot of the entropy S_χ as a function of the scaled temperature for the indicated system sizes. In the insert we plot the expected no scale temperature behavior.

in this plot the actual superadditive behavior of the new entropy in an interacting system instead of the simple N independent two states systems of the last section.

6 Conclusions

We have shown that the entropy functional S_χ verifies the main properties usually regarded as essential requirements for physically meaningful entropy functionals [1,8,12,13]. The entropy S_χ verifies the first three Khinchin axioms. The measure S_χ satisfies the requirements of positivity, equiprobability, concavity and irreversibility. The associated Jaynes' maximum entropy scheme also complies with Jaynes' thermodynamic relations. The properties verified by S_χ suggest that it might be a useful measure for describing nonextensive physical phenomena as well as for other practical applications.

The entropic functional S_χ grows as a power N^χ of the size N of the system and is thus characterized by a weak nonextensive behavior (as opposed to the strong, exponential nonextensivity exhibited by Tsallis measure). An attentive reader might think that a power law nonextensive measure can be obtained in a trivial way by just defining a new "entropy" equal to $(S_{\text{Boltzmann}})^\chi$. However, this simple proposal would not lead to anything new. The probability distributions maximizing $(S_{\text{Boltzmann}})^\chi$ are the same exponential distributions that maximize $S_{\text{Boltzmann}}$. This means that the standard (extensive) thermostatics would be obtained again. On the contrary, the entropic functional S_χ furnishes a nontrivial new thermostatics that might be appropriate for dealing with some nonextensive systems. As an application of the thermostatical formalism associated with S_χ we considered a ferromagnetic Ising model with long-range interactions. We studied numerically the concomitant scaling laws.

Tsallis thermostatics, which has attracted a great deal of attention recently [24–31], arises from a nonextensive entropic measure S_q growing exponentially with the size of the system. Now, if there are in nature

extensive systems (that is, those successfully studied within standard statistical mechanics and described by Gibbs exponential distributions) as well as Tsallis' strong (exponentially) nonextensive systems, it is not unreasonable to expect also the existence of systems exhibiting an intermediate (weak) power-law kind of nonextensivity.

Let us consider a family of dynamical systems characterized by a set of parameters λ . We assume that there is a certain region Σ in λ -space such that the system's thermodynamic behavior is extensive for $\lambda \in \Sigma$ and nonextensive otherwise. It is possible that, as a continuous change in the parameters λ is considered, the system abruptly jumps from an extensive regime to an exponentially nonextensive one. Notice, however, that this kind of discontinuity would be more drastic than the one associated with a change of exponents (that is, the jump from one power-law to a new power-law characterized by a different exponent). We think that it would be physically more sensible if there is an "onset of nonextensivity" boundary between the extensive and nonextensive regions in λ -space, characterized by a weak, power-law nonextensive regime. A suggestive analogy can be made here with the onset of chaos in nonlinear dynamics. Consider, for instance, the well known Logistic map. For values of the map's parameter corresponding to chaotic behavior the concomitant Lyapunov exponent is positive and near trajectories diverge exponentially in time. However, critical values of the map parameter associated with a vanishing Lyapunov exponent still do exhibit a form of "weak chaos". In those critical situations, corresponding to the onset chaos, near trajectories do not diverge exponentially in time: they diverge in a power-law way [37,38].

It would be important to gain a further clarification of the physical meaning of the parameter χ , and to understand under what circumstances might nature choose to maximize the functional S_χ . The only way to attain such an understanding is by a detailed study of the dynamics of particular nonextensive systems endowed with long range interactions. We hope that our present contribution may stimulate further work within this line of inquiry.

A.R.P. acknowledges the AECI Scientific Cooperation Program (Spain), and the CONICET (Argentine Agency) for partial financial support. R. Toral and R. Salazar acknowledge the financial support from DGES, grants PB94-1167 and PB97-0141-C02-01.

References

1. C. Tsallis, J. Stat. Phys. **52**, 479 (1988); C. Tsallis, R.S. Mendes, A.R. Plastino, Physica A **261**, 534 (1998); E.M.F. Curado, C. Tsallis, J. Phys. A **24**, L69 (1991).
2. R. Salazar, R. Toral, Phys. Rev. Lett. **83**, 4233 (1999).
3. A. Plastino, A.R. Plastino, Phys. Lett. A **226**, 257 (1997).
4. R.S. Mendes, Physica A **242**, 299 (1997); E.M.F. Curado, Braz. J. Phys. **29**, 36 (1999).
5. P.T. Landsberg, Braz. J. Phys. **29**, 46 (1999).
6. P.T. Landsberg, V. Vedral, Phys. Lett. A **247**, 211 (1998).
7. S. Abe, Phys. Lett. A **224**, 326 (1997).
8. A.R.R. Papa J. Phys. A **31**, 1 (1998).
9. E.P. Borges, I. Roditi, Phys. Lett. A **246**, 399 (1998).
10. C. Anteneodo, A.R. Plastino, J. Phys. A **32**, 1089 (1999).
11. A.K. Rajagopal, S. Abe, Phys. Rev. Lett. **83**, 1711 (1999).
12. R. Rossignoli, N. Canosa Phys. Lett. A **264**, 148 (1999).
13. K.S. Fa, J. Phys. A **31**, 8159 (1998).
14. D. Holste, I. Grosse, H. Herzog, J. Phys. A **31**, 2551 (1998).
15. E.P. Borges, J. Phys. A **31**, 5281 (1998); E.K. Lenzi, E.P. Borges, R.S. Mendes, J. Phys. A **32**, 8551 (1999).
16. A. Compte, D. Jou, J. Phys. A **29**, 4321 (1996); A. Compte, D. Jou, Y. Katayama, J. Phys. A **30**, 1023 (1997).
17. L. Borland, Phys. Rev. E **57**, 6634 (1998).
18. E.K. Lenzi, L.C. Malacarne, R.S. Mendes, Phys. Rev. Lett. **80**, 218 (1998).
19. W.C. Saslaw, *Gravitational Physics of Stellar and Galactic Systems* (Cambridge University Press, Cambridge, 1985).
20. A.R. Plastino, A. Plastino, Phys. Lett. A **174**, 384 (1993).
21. B.M.R. Boghosian, Phys. Rev. E **53**, 4754 (1995).
22. E.T. Jaynes, in *Statistical Physics*, edited by W.K. Ford (Benjamin, New York, 1963); A. Katz, *Statistical Mechanics* (Freeman, San Francisco, 1967).
23. C. Beck, F. Schlögl, *Thermodynamics of Chaotic Systems: An Introduction* (Cambridge University Press, Cambridge, 1993).
24. C. Tsallis, S.V.F. Levy, A.M.C. Souza, R. Maynard, Phys. Rev. Lett. **75**, 3589 (1995); Erratum: **77**, 5442 (1996).
25. M.L. Lyra, C. Tsallis, Phys. Rev. Lett. **80**, 53 (1998); C. Anteneodo, C. Tsallis, Phys. Rev. Lett. **80**, 5313 (1998).
26. M. Buiatti, P. Grigolini, A. Montagnini, Phys. Rev. Lett. **82**, 3383 (1999).
27. A.K. Rajagopal, R.S. Mendes, E.K. Lenzi, Phys. Rev. Lett. **80**, 3907 (1998).
28. D.B. Ion, M.L.D. Ion, Phys. Rev. Lett. **81**, 5714 (1998).
29. M.L.D. Ion, D.B. Ion, Phys. Rev. Lett. **83**, 463 (1999).
30. C. Tsallis, *Physics World* **10**, 42 (1997); an updated bibliography on Tsallis' theory and its applications is available at <http://tsallis.cat.cbpf.br/biblio.htm>.
31. Several review articles on Tsallis' thermostatistics and its applications appeared in Braz. J. Phys. **29** (1999), Special Issue, *Nonextensive Statistical Mechanics and Thermodynamics*, edited by S.R.A. Salinas, C. Tsallis, which can be seen at http://sbf.if.usp.br/WWW_pages/Journals/BJP/Vol29/Num1/index.htm.
32. C. Tsallis, Fractals **3**, 541 (1995).
33. S.A. Cannas, F.A. Tamarit, Phys. Rev. B **54**, R12661 (1996).
34. A.I. Khinchin, *Mathematical Foundations of Information Theory* (Dover, New York, 1957).
35. G. Bhanot, R. Salvador, S. Black, P. Carter, R. Toral. Phys. Rev. Lett., **59**, 803 (1987).
36. R. Salazar, R. Toral, Physica A (to be published, 2000).
37. C. Tsallis, A.R. Plastino, W.-M. Zheng. Chaos, Solitons and Fractals **8**, 885 (1997).
38. U.M.S. Costa, M.L. Lyra, A.R. Plastino, C. Tsallis. Phys. Rev. E, **56**, 245 (1997).