

Dilute limit of a strongly-interacting model of spinless fermions and hardcore bosons on the square lattice

N.G. Zhang^a and C.L. Henley

Department of Physics, Cornell University, Ithaca, New York 14853-2501, USA

Received 8 May 2003

Published online 28 May 2004 – © EDP Sciences, Società Italiana di Fisica, Springer-Verlag 2004

Abstract. In our model, spinless fermions (or hardcore bosons) on a square lattice hop to nearest neighbor sites, and also experience a hard-core repulsion at the nearest neighbor separation. This is the simplest model of correlated electrons and is more tractable for exact diagonalization than the Hubbard model. We study systematically the dilute limit of this model by a combination of analytical and several numerical approaches: the two-particle problem using lattice Green functions and the t -matrix, the few-fermion problem using a modified t -matrix (demonstrating that the interaction energy is well captured by pairwise terms), and for bosons the fitting of the energy as a function of density to Schick's analytical result for dilute hard disks. We present the first systematic study for a strongly-interacting lattice model of the t -matrix, which appears as the central object in older theories of the existence of a two-dimensional Fermi liquid for dilute fermions with strong interactions. For our model, we can (Lanczos) diagonalize the 7×7 system at all fillings and the 20×20 system with four particles, thus going far beyond previous diagonalization works on the Hubbard model.

PACS. 71.10.Fd Lattice fermion models (Hubbard model, etc.) – 71.10.Pm Fermions in reduced dimensions (anyons, composite fermions, Luttinger liquid, etc.) – 05.30.Jp Boson systems – 74.20.Mn Nonconventional mechanisms (spin fluctuations, polarons and bipolarons, resonating valence bond model, anyon mechanism, marginal Fermi liquid, Luttinger liquid, etc.)

1 Introduction

Since the discovery of the high-temperature superconductors in 1986, there has been intense study of a number of two-dimensional models that are believed to model the electronic properties of the CuO_2 plane of the cuprate superconductors, for example, the Hubbard model, the $t - J$ model, and the Heisenberg model [1, 2]. Two-dimensional quantum models with short-range kinetic and interaction terms are difficult to study. In one dimension, there are exact solutions using the Bethe ansatz and a host of related analytical techniques [3], and there is a very accurate numerical method, the density-matrix renormalization group (DMRG) [4], that can be applied to large systems relatively easily. In two dimensions, on the other hand, there are few exact solutions (one famous nontrivial case is the Hubbard model with one hole in a half-filled background, the Nagaoka state [5]), and current numerical methods are not satisfactory (quantum Monte Carlo is plagued by the negative sign problem [1] at low temperatures and at many fillings of interest and the DMRG in two dimensions [6] is still in early development stage).

The most reliable method for studying complicated quantum systems is exact diagonalization, which means enumerating all basis states and diagonalizing the resulting Hamiltonian matrix. Of course, this method is computationally limited by the growth of the Hilbert space which is in general exponential in the number of particles and the lattice size. The 4×4 Hubbard model with 16 electrons, 8 spin-up and 8 spin-down, after reduction by particle conservation, translation, and the symmetries of the square, has 1,310,242 states in the largest matrix block [7], and can be diagonalized using the well-known Lanczos method. The Hubbard model has been diagonalized for the 4×4 lattice (see e.g., Ref. [8]), and at low filling (four electrons) for 6×6 [9] with extensive employment of symmetries.

In this paper we present a model of spinless fermions (or hardcore bosons) which contains the basic ingredients of short-range hopping on a lattice and interaction but is simpler than the Hubbard model. As the interaction is replaced by a hard-core, the Hilbert space is reduced and we can diagonalize the 7×7 system at all fillings, or the 20×20 system with four particles, which is many lattice constants larger than the tractable size for diagonalizations of the Hubbard model. We have systematically studied this system at the dilute limit, and the main results of this study

^a *Present address:* Dept. of Physics, George Washington University, Washington, DC 20052
e-mail: ngzhang@gwu.edu

are the following. First, using lattice Green functions we solve analytically the problems of two and a few particles and check the results for large lattices using diagonalization (Sect. 2). We find that the few-particle energies can largely be approximated by two-particle energies when the lattice size is large. Second, we derive the t-matrix formalism for particles with nearest-neighbor hopping and use it to solve the problems of two particles (Sect. 3) and a few fermions (Sect. 4). Comparing our analytical results with exact diagonalization data, we confirm that in the dilute limit, almost all of the interaction correction is accounted for by the two-body terms of the t-matrix approximation. This result contributes to the understanding of the validity of Fermi liquid theory in a finite-system context. Third, we study the energy per particle curve for two-dimensional dilute bosons and check Schick's well-known analytical results [10] using exact diagonalization (Sect. 5). Our findings here complement Monte Carlo studies [11,12]. To our best knowledge, this work appears to be the first systematic study of the dilute limit of this spinless fermion model.

1.1 The spinless fermion model

We have asked the question: *Is there a model that contains the basic ingredient of short-range hopping and interaction but is simpler, in the exact diagonalization sense, than the Hubbard model?* The answer is yes: we can neglect the spin. We obtain the following Hamiltonian for *spinless* fermions,

$$H = -t \sum_{\langle i,j \rangle} (c_i^\dagger c_j + c_j^\dagger c_i) + V \sum_{\langle i,j \rangle} \hat{n}_i \hat{n}_j, \quad (1.1)$$

where c_i^\dagger and c_i are spinless fermion creation and annihilation operators at site i , $\hat{n}_i = c_i^\dagger c_i$ the number operator, t the nearest-neighbor hopping amplitude, and V the nearest-neighbor interaction. Note that with spinless fermions, there can be at the most one particle per site; no on-site interaction (as that in the Hubbard model) is possible, and we have included in our Hamiltonian nearest-neighbor repulsion.

The spinless fermion model, equation (1.1), is a two-state model, and the number of basis states for a N -site system is 2^N , which is a significant reduction from the 4^N of the Hubbard model. We can further reduce the number of basis states by taking the nearest-neighbor interaction $V = +\infty$, i.e., infinite repulsion, which excludes nearest neighbors, giving roughly $2^{N/2}$ states.

The spinless fermion model with infinite repulsion equation (1.1) contains a significant reduction of the Hilbert space. After using particle conservation and translation symmetry (but not point group symmetry), the largest matrix for the 7×7 system has 1,906,532 states (for 11 particles), and we can therefore compute for all fillings the 7×7 system whereas for the Hubbard model 4×4 is basically the limit [13]. This of course means that for certain limits we can also go much further than the Hubbard model, for example, we can handle four particles on a 20×20 lattice where the number of basis states is

2,472,147. This extended capability with our model has enabled us to obtain a number of results that are difficult to obtain with the Hubbard model.

An added feature of our model is that the basis set for the spinless fermion problem is identical to that for the hardcore boson problem, because with hardcore repulsion, there can be at the most one boson at one site also. Therefore, without computational difficulty, we can study numerically both the spinless fermion and hardcore boson problem.

Spinless fermions can also be realized in experiments, for example, the spin polarized ^3He due to a strong magnetic field, or ferro or ferri-magnetic electronic systems where one spin-band is filled [16].

Considering the tremendous effort that has been devoted to the Hubbard model and the close resemblance of our model, equation (1.1), to the Hubbard model, it is surprising that works on this spinless model have been rather sparse, though it has been commented that the spinless model offers considerable simplifications [17]. The one-dimensional spinless fermion model with finite repulsion is solved exactly using Bethe ansatz [18]. The infinite-dimensional problem is studied in reference [19]. A very different approach using the renormalization group for fermions is done in reference [20]. A Monte Carlo study of the two-dimensional model at half-filling only and low temperatures is in reference [21], which, dating back to 1985, is one of the earliest quantum Monte Carlo simulations for fermions. (It is no coincidence that they chose the model with the smallest Hilbert space.) The prior work most comparable to ours may be the studies of four spinless electrons in a 6×6 lattice, with Coulomb repulsion, by Pichard et al. [22]; their motivation was the Wigner crystal melting and the competition of Coulomb interactions with Anderson localization when a disorder potential is turned on.

In two separate publications, we have studied the dense limit of this two-dimensional model of spinless fermions (or hardcore bosons) with infinite nearest-neighbor repulsion [14,23]. There, near half-filled [24], stripes (that are holes lining up across the lattice) are natural objects. The present paper focuses on the dilute limit, treating the problem of a few particles. One of the goals of these papers is to advertise this model of spinless fermions to the strongly-correlated electron community, as we believe that it is the simplest model of correlated fermions and deserves more research effort and better understanding.

1.2 The t-matrix

At the dilute limit of our model, the scattering t-matrix is of fundamental importance. For two particles, we expect that, at least when the potential V is small, we can write a perturbative equation for energy,

$$E = \mathcal{E}(\mathbf{q}_1) + \mathcal{E}(\mathbf{q}_2) + \Delta E(\mathbf{q}_1, \mathbf{q}_2), \quad (1.2)$$

which is to say that the exact interacting energy of two particles is the noninteracting energy $\mathcal{E}(\mathbf{q}_1) + \mathcal{E}(\mathbf{q}_2)$, for a

pair of momenta \mathbf{q}_1 and \mathbf{q}_2 , plus a correction term ΔE due to the interaction V . And with more than two particles, at least when the particle density is low, we expect to have

$$E = \sum_{\mathbf{q}} \mathcal{E}(\mathbf{q}) + \frac{1}{2} \sum_{\mathbf{q}, \mathbf{q}'} \Delta E(\mathbf{q}, \mathbf{q}'). \quad (1.3)$$

Equation (1.3) is central in Fermi liquid theory, where it is justified by the so-called ‘‘adiabatic continuation’’ idea, which says that interacting fermion states correspond one-to-one to noninteracting ones as we slowly switch on a potential.

In the boson case, because many bosons can occupy one quantum mechanical state and form a condensate, equation (1.3) should be modified, but with only two bosons, we expect equation (1.2) should be valid (in that the correction vanishes in the dilute limit). Equations (1.2) and (1.3) are used when we look at a list of noninteracting energies and draw correspondences with the interacting energies, the energy shift being packaged in the term ΔE : the t-matrix term.

We are not aware of a systematic study of the t-matrix for a lattice model. In this paper, we present the first such study for the two-particle problem in Section 3 (for bosons and fermions) and the few-fermion problem in Section 4. We check the t-matrix results with exact diagonalization data and show that our t-matrix on a lattice is the sum of the two-body scattering terms to infinite order.

2 The two-particle problem

The two-particle problem has appeared in many different contexts. The most familiar one is the hydrogen atom problem in introductory quantum mechanics textbooks. The two-magnon problem is closely related mathematically to our two-particle problem, and it has been solved in arbitrary dimensions for ferromagnets (see e.g., Ref. [25]). Another important two-particle problem is the Cooper problem, with two electrons in the presence of a Fermi sea (see e.g., Ref. [26]). And motivated by the possibility of Cooper pair formation in high-temperature superconductors, there have also been a number of studies on bound states on a two-dimensional lattice [27–32]. The two-electron problem in the plain two-dimensional repulsive Hubbard model is studied in reference [33], and ground state energy in the large-lattice limit is obtained analytically.

In this section, we present a rather complete calculation for the two-particle problem in our model, treating both bosons and fermions. We calculate eigenenergies for all states for a finite-size lattice, and our calculation is more complicated than the Hubbard model [33] case because of nearest-neighbor (in place of on-site) interaction. Where the Green function in the Hubbard case was a scalar object, in our case it is replaced by a 4×4 matrix, corresponding to the four nearest neighbor sites where the potential acts. This Green function study of the two-particle problem is closely related to the treatment of

the two-electron problem in the Hubbard model [33] and that in an extended Hubbard model [31]. We will show the use of lattice symmetry and recursion relations to simplify the problem with nearest-neighbor interactions. Note that bound states are not of interest, since they do not exist (with two particles) under the infinite repulsive interaction in our model (see Sect. 3.5).

2.1 Preliminary

In this two-particle calculation, we will work in momentum space, and we will start with a Hamiltonian more general than equation (1.1) [31],

$$H = T + U, \quad (2.1)$$

$$T = \sum_{\mathbf{r}_1 \mathbf{r}_2} t(\mathbf{r}_2 - \mathbf{r}_1) c_{\mathbf{r}_1}^\dagger c_{\mathbf{r}_2}, \quad (2.2)$$

$$U = \sum_{\mathbf{r}_1 \mathbf{r}_2} V(\mathbf{r}_2 - \mathbf{r}_1) c_{\mathbf{r}_1}^\dagger c_{\mathbf{r}_1} c_{\mathbf{r}_2}^\dagger c_{\mathbf{r}_2}. \quad (2.3)$$

Here we have allowed hopping and interaction between any two lattice sites, but we require that both depend only on the separation between the two vectors and both have inversion symmetry. That is $t(\mathbf{r}_1, \mathbf{r}_2) = t(\mathbf{r}_2 - \mathbf{r}_1)$, $t(-\mathbf{r}) = t(\mathbf{r})$, $V(\mathbf{r}_1, \mathbf{r}_2) = V(\mathbf{r}_2 - \mathbf{r}_1)$, and $V(-\mathbf{r}) = V(\mathbf{r})$. In momentum space, equations (2.2) and (2.3) become,

$$T = \sum_{\mathbf{p}} \mathcal{E}(\mathbf{p}) c_{\mathbf{p}}^\dagger c_{\mathbf{p}}, \quad (2.4)$$

$$U = \frac{1}{2N} \sum_{\mathbf{p} \mathbf{p}' \mathbf{k}} V(\mathbf{k}) c_{\mathbf{p}}^\dagger c_{\mathbf{p}'}^\dagger c_{\mathbf{p}'+\mathbf{k}} c_{\mathbf{p}-\mathbf{k}}, \quad (2.5)$$

where

$$\mathcal{E}(\mathbf{p}) = \sum_{\mathbf{r}} t(\mathbf{r}) e^{i\mathbf{p}\mathbf{r}}, \quad (2.6)$$

$$V(\mathbf{k}) = \sum_{\mathbf{r}} V(\mathbf{r}) e^{i\mathbf{k}\mathbf{r}}, \quad (2.7)$$

with $\mathcal{E}(-\mathbf{p}) = \mathcal{E}(\mathbf{p})$ and $V(-\mathbf{k}) = V(\mathbf{k})$. Equations (2.2) and (2.3) reduce to our nearest-neighbor Hamiltonian equation (1.1) if we take,

$$t(\mathbf{r}) = \begin{cases} -t, & \mathbf{r} = (\pm 1, 0)(0, \pm 1), \\ 0, & \text{otherwise,} \end{cases} \quad (2.8)$$

$$V(\mathbf{r}) = \begin{cases} V, & \mathbf{r} = (\pm 1, 0)(0, \pm 1), \\ 0, & \text{otherwise,} \end{cases} \quad (2.9)$$

where we have taken the lattice constant to be unity [34], and the nearest-neighbor vectors will be called

$$\mathbf{R}_1 = (1, 0), \mathbf{R}_2 = (-1, 0), \mathbf{R}_3 = (0, 1), \mathbf{R}_4 = (0, -1). \quad (2.10)$$

Then we have,

$$\mathcal{E}(\mathbf{p}) = -2t(\cos p_x + \cos p_y), \quad (2.11)$$

$$V(\mathbf{k}) = 2V(\cos k_x + \cos k_y). \quad (2.12)$$

Note that the structure of later equations depends sensitively on having four sites in equation (2.9) where $V(\mathbf{r}) \neq 0$, but does not depend much on the form of equation (2.8) and the resulting dispersion equation (2.11).

Using momentum conservation of equation (2.1), the two-particle wave function that we will use is,

$$|\psi\rangle = \sum_{\mathbf{q}} g(\mathbf{q})|\mathbf{q}, \mathbf{P} - \mathbf{q}\rangle, \quad (2.13)$$

where the sum is over the whole Brillouin zone, and the coefficient $g(\mathbf{q})$ satisfies,

$$g(\mathbf{P} - \mathbf{q}) = s_{bf}g(\mathbf{q}), \quad (2.14)$$

where $s_{bf} = 1$ for bosons and -1 for fermions.

2.2 Green function equations

Applying the more general form of the Hamiltonian operator equations (2.4) and (2.5) to the state equation (2.13), the Schrodinger equation $(E - T)|\psi\rangle = U|\psi\rangle$ becomes

$$(E - \mathcal{E}(\mathbf{q}) - \mathcal{E}(\mathbf{P} - \mathbf{q}))g(\mathbf{q}) = \frac{1}{N} \sum_{\mathbf{k}} V(\mathbf{q} - \mathbf{k})g(\mathbf{k}). \quad (2.15)$$

Equation (2.15) is a matrix equation $\mathcal{A}g = Eg$ where $\mathcal{A}_{\mathbf{q}\mathbf{k}} = (\mathcal{E}(\mathbf{q}) + \mathcal{E}(\mathbf{P} - \mathbf{q}))\delta_{\mathbf{q}\mathbf{k}} + V(\mathbf{q} - \mathbf{k})/N$. If V is not infinity, this $N \times N$ matrix \mathcal{A} can be diagonalized, and E and $g(\mathbf{q})$ are respectively the eigenvalue and eigenvector. To deal with $V = +\infty$, we need some further manipulations.

We consider the case when $E \neq \mathcal{E}(\mathbf{q}) + \mathcal{E}(\mathbf{P} - \mathbf{q})$, for any \mathbf{q} , which is to say, the energy E is not the energy of a noninteracting pair. The (lattice) Fourier transform of the coefficients $g(\mathbf{q})$ is

$$\tilde{g}(\mathbf{r}) = \sum_{\mathbf{q}} e^{-i\mathbf{q}\cdot\mathbf{r}}g(\mathbf{q}); \quad (2.16)$$

this is just the real-space wavefunction in terms of the relative coordinate \mathbf{r} . Define the lattice Green function,

$$G(E, \mathbf{P}; \mathbf{r}, \mathbf{r}') = \frac{1}{N} \sum_{\mathbf{q}} \frac{e^{i\mathbf{q}\cdot(\mathbf{r}' - \mathbf{r})}}{E - \mathcal{E}(\mathbf{q}) - \mathcal{E}(\mathbf{P} - \mathbf{q})}, \quad (2.17)$$

then after dividing the first factor from both sides of our Schrodinger equation, equation (2.15), and Fourier transforming, we obtain

$$\tilde{g}(\mathbf{r}) = \sum_{\mathbf{r}'} G(E, \mathbf{P}; \mathbf{r}, \mathbf{r}')V(\mathbf{r}')\tilde{g}(\mathbf{r}'). \quad (2.18)$$

In the following we return to the nearest-neighbor potential $V(\mathbf{r})$ in equation (2.9). The Green function sum in equation (2.18) then has only four terms,

$$\tilde{g}(\mathbf{r}) = \sum_j G(E, \mathbf{P}; \mathbf{r}, \mathbf{R}_j)(V\tilde{g}(\mathbf{R}_j)), \quad (2.19)$$

summed over the separations in equation (2.10). If we also restrict \mathbf{r} to the four nearest-neighbor vectors [35], then equation (2.19) becomes,

$$\tilde{g}(\mathbf{R}_i) = \sum_j G(E, \mathbf{P}; \mathbf{R}_i, \mathbf{R}_j)(V\tilde{g}(\mathbf{R}_j)). \quad (2.20)$$

If we define the 4×4 matrix,

$$\mathcal{G}_{ij}(E, \mathbf{P}) = G(E, \mathbf{P}; \mathbf{R}_i, \mathbf{R}_j), \quad (2.21)$$

and a 4×1 vector $\phi_j = \tilde{g}(\mathbf{R}_j)$, then we obtain a simple matrix equation,

$$(I - \mathcal{G}(E, \mathbf{P})V)\phi = 0. \quad (2.22)$$

We can also rewrite this equation as an equation for energy using the determinant,

$$\det(I - \mathcal{G}(E, \mathbf{P})V) = 0. \quad (2.23)$$

With $V = +\infty$, we have even simpler equations

$$\mathcal{G}(E, \mathbf{P})(V\phi) = 0, \quad (2.24)$$

and

$$\det \mathcal{G}(E, \mathbf{P}) = 0. \quad (2.25)$$

Notice we write $V\phi$ to denote the limit as $V \rightarrow \infty$; it would not do to write simply ϕ in equations (2.22) and (2.24), since $\phi \rightarrow 0$ as $V \rightarrow \infty$ (being the amplitude of the relative wavefunction at the forbidden separations $\{\mathbf{R}_i\}$).

For the Hubbard model, there is only on-site interaction, so $V(\mathbf{r})$ is nonzero only when $\mathbf{r} = 0$, and the sum in equation (2.18) has only one term. Equation (2.20) is simply a scalar equation, which, after \tilde{g} cancels from both sides of the equation and using equation (2.17), gives,

$$1 = \frac{V}{N} \sum_{\mathbf{q}} \frac{1}{E - \mathcal{E}(\mathbf{q}) - \mathcal{E}(\mathbf{P} - \mathbf{q})}, \quad (2.26)$$

which is exactly the result in reference [33].

2.2.1 Simplifications for rectangular boundaries

We specialize to the case of total momentum $\mathbf{P} = 0$ and rectangular-boundary lattices. We have from equations (2.17) and (2.21),

$$\mathcal{G}_{ij}(E) = \frac{1}{N} \sum_{\mathbf{q}} \frac{\cos(q_x(R_{jx} - R_{ix})) \cos(q_y(R_{jy} - R_{iy}))}{E - 2\mathcal{E}(\mathbf{q})}, \quad (2.27)$$

where the potential is nonzero on the sites $\{\mathbf{R}_i\}$ given by equation (2.10) and in the last step we have used the symmetry properties of the dispersion relation $\mathcal{E}(q_x, q_y) = \mathcal{E}(q_x, -q_y) = \mathcal{E}(-q_x, q_y)$. Obviously equation (2.27) is a function of displacements $\mathbf{R}_j - \mathbf{R}_i$, which (in view of Eq. (2.10) can be (0,0), (1,1), (2,0), or any vector related by square symmetry. It is convenient for this and later

sections to define a new notation for the Green function \mathcal{G}_{ij} , emphasizing its dependence on $\mathbf{R}_j - \mathbf{R}_i = (m, n)$,

$$\Gamma(E, m, n) = \frac{1}{N} \sum_{\mathbf{q}} \frac{\cos(mq_x) \cos(nq_y)}{E + 4 \cos q_x + 4 \cos q_y}, \quad (2.28)$$

where the sum is over the N wavevectors $\mathbf{q} = (2\pi l_x/L_x, 2\pi l_y/L_y)$ with $0 \leq l_x < L_x$ and $0 \leq l_y < L_y$ (for one Brillouin zone), and we have used the expression for $\mathcal{E}(\mathbf{q})$ from equation (2.11) (and taken $t = 1$).

This Green function for rectangular-boundary lattices satisfies the following reflection properties,

$$\begin{aligned} \Gamma(E, m, n) &= \Gamma(E, -m, n) \\ &= \Gamma(E, m, -n) = \Gamma(E, -m, -n). \end{aligned} \quad (2.29)$$

And if we have a square lattice ($L_x = L_y$) we also have

$$\Gamma(E, m, n) = \Gamma(E, n, m), \quad (2.30)$$

and equation (2.27) can be written as,

$$\mathcal{G}_{ij}(E) = \Gamma(E, R_{jx} - R_{ix}, R_{jy} - R_{iy}). \quad (2.31)$$

Using the reflection properties of $\Gamma(E, m, n)$, equation (2.29), and the definition equation (2.31), our Green function matrix becomes,

$$\mathcal{G}_{ij}(E) = \begin{pmatrix} a & c & b & b \\ c & a & b & b \\ b & b & a & d \\ b & b & d & a \end{pmatrix}, \quad (2.32)$$

where $a = \Gamma(E, 0, 0)$, $b = \Gamma(E, 1, 1)$, $c = \Gamma(E, 2, 0)$, and $d = \Gamma(E, 0, 2)$. The eigenvalues and eigenvectors of this matrix are,

$$\begin{aligned} \lambda_{f1} &= a - c, & (1, -1, 0, 0) \\ \lambda_{f2} &= a - d, & (0, 0, 1, -1) \\ \lambda_{b1, b2} &= a + \frac{c+d}{2} \pm \frac{\sqrt{16b^2 + (c-d)^2}}{2}, \\ & & (v_{1,2}, v_{1,2}, 1, 1) \end{aligned} \quad (2.33)$$

where v_1 and v_2 are complicated functions of a, b, c , and d .

The exact energy E makes the matrix $\mathcal{G}_{ij}(E)$ singular, which means that one of the eigenvalues has to be zero. From equation (2.24), the null eigenvector of \mathcal{G} is $V\phi = V(\tilde{g}(\mathbf{R}_1), \tilde{g}(\mathbf{R}_2), \tilde{g}(\mathbf{R}_3), \tilde{g}(\mathbf{R}_4))$, in terms of $\{\mathbf{R}_i\}$ as in equation (2.10). The relative wavefunction should be odd or even under inversion, depending on statistics, i.e. $\tilde{g}(-\mathbf{r}) = s_{bf} e^{i\mathbf{P}\cdot\mathbf{r}} \tilde{g}(\mathbf{r})$ which follows immediately from equations (2.16) and (2.14). Inversion, acting on nearest-neighbor vectors equation (2.10), induces $\mathbf{R}_1 \leftrightarrow \mathbf{R}_2$ and $\mathbf{R}_3 \leftrightarrow \mathbf{R}_4$; thus with $\mathbf{P} = 0$, we should have $V\phi_1 = -V\phi_2$ and $V\phi_3 = -V\phi_4$ for fermions, and $V\phi_1 = V\phi_2$ and $V\phi_3 = V\phi_4$ for bosons. Inspecting the eigenvectors we obtained in equation (2.33), we see that those corresponding to $\lambda_{f1, f2}$ are antisymmetric under inversion – corresponding to a “p-wave-like” (relative angular momentum 1) state for fermions. Setting $\lambda_{f1} = 0$, we get $a = c$,

or setting $\lambda_{f2} = 0$ $a = d$, which respectively mean

$$\Gamma(E, 0, 0) - \Gamma(E, 2, 0) = 0, \text{ or} \quad (2.34)$$

$$\Gamma(E, 0, 0) - \Gamma(E, 0, 2) = 0. \quad (2.35)$$

Associated with the even eigenvectors are $\lambda_{b1, b2}$ which are identified as boson eigenvalues.

2.2.2 Simplifications for square boundaries

The boson eigenvalues, equation (2.33), are rather complicated for general rectangular-boundary lattices. For a square-boundary lattice, using equation (2.30), we get $c = d$ in the matrix equation (2.32), which makes the fermion eigenvalues $\lambda_{f1, f2}$ degenerate. The boson eigenvalues in equation (2.33) simplify greatly to $\lambda_{b1} = a + 2b + c$ and $\lambda_{b2} = a - 2b + c$, which means that the boson energy equations are,

$$\Gamma(E, 0, 0) + 2\Gamma(E, 1, 1) + \Gamma(E, 2, 0) = 0, \quad (2.36)$$

$$\Gamma(E, 0, 0) - 2\Gamma(E, 1, 1) + \Gamma(E, 2, 0) = 0. \quad (2.37)$$

The corresponding eigenvectors simplify too, to $(1, 1, 1, 1)$ and $(1, 1, -1, -1)$ respectively, which may be described as “s-wave-like” and “d-wave-like”, i.e. relative angular momentum 0 and 2.

2.3 Large-L asymptotics for two-boson energy

Equations (2.34, 2.35, 2.36) and (2.37) are much better starting points for analytical calculations than the original determinant equation (2.25). In the center of the problem is the lattice Green function $\Gamma(E, m, n)$ defined in equation (2.28). Many of the lattice calculations come down to evaluating these lattice Green functions [28–32, 36]. In this section, we derive the large-lattice two-boson energy using the recursion and symmetry relations of the Green function $\Gamma(E, m, n)$.

The Green function $\Gamma(E, m, n)$ for general m and n and finite lattice are difficult to evaluate. The good thing is that there are a number of recursion relations connecting the Green functions at different m and n [37, 38]. These are trivial to derive after noting that equation (2.17) (for $\mathbf{P} = 0$) can be written

$$[E - (4 + \Delta_{\mathbf{r}}^2) - (4 + \Delta_{\mathbf{r}'}^2)]G(E, 0; \mathbf{r}, \mathbf{r}') = \delta_{\mathbf{r}=0} \delta_{\mathbf{r}'=0} \quad (2.38)$$

where $\Delta_{\mathbf{r}}^2$ is the discrete Laplacian, $(\Delta_{\mathbf{r}}^2 + 4)f(\mathbf{r}) \equiv \sum_i f(\mathbf{r} + \mathbf{R}_i)$ for any function $f(\mathbf{r})$, where the sum is over neighbor vectors equation (2.10). The two recursion relations that we will use are

$$E\Gamma(E, 0, 0) + 4\Gamma(E, 1, 0) + 4\Gamma(E, 0, 1) = 1, \quad (2.39)$$

$$\begin{aligned} \Gamma(E, 0, 0) + 2\Gamma(E, 1, 1) + \Gamma(E, 2, 0) \\ + \frac{1}{2}E\Gamma(E, 1, 0) = 0. \end{aligned} \quad (2.40)$$

Using equations (2.30, 2.39) and (2.40), the boson equation (2.36) for square-boundary lattices simplifies to

$$\Gamma(E, 0, 0) = \frac{1}{E}, \quad (2.41)$$

with eigenvector $(1, 1, 1, 1)$.

Next we compute the leading form of $\Gamma(E, 0, 0)$ for large L of a square-boundary lattice. The calculation is close to that in reference [33] for the Hubbard model. We define $E = -8 + \Delta E$. Because the lowest energy of an independent particle is $\mathcal{E}(\mathbf{0}) = -4$, ΔE is the energy correction to two independent particle energy at zero momentum. Then we have, from equation (2.28),

$$\begin{aligned} \Gamma(E, 0, 0) &= \frac{1}{N} \sum_{\mathbf{q}} \frac{1}{E + 4 \cos q_x + 4 \cos q_y}, \\ &= -\frac{1}{4N} \sum_{\mathbf{q}} \frac{1}{2 - \cos q_x - \cos q_y - \Delta E/4}, \\ &\approx \frac{1}{L^2 \Delta E} - \frac{1}{4\pi} \int_{2\pi/L}^{\pi} \frac{dq}{q} \\ &= \frac{1}{L^2 \Delta E} - \frac{\ln L}{4\pi} + \text{const.} \end{aligned} \quad (2.42)$$

We should discuss the number of approximations we have made to extract this leading dependence in L . First except in the $\mathbf{q} = 0$ term we have ignored the ΔE term, assuming it is small as compared to \mathbf{q}^2 (with $\mathbf{q} \neq 0$). This is justified as we only want the leading term in the large- L limit. Using an integral for a lattice sum is another approximation. We choose the lower limit of integration to be $2\pi/L$ corresponding to the first wavevectors after $(0, 0)$ is taken out of the sum. We also used the quadratic approximation for the energy dispersion $\mathcal{E}(\mathbf{q})$ appearing in the denominator.

Using the boson energy equation (2.41) and the large- L limit of the Green function equation (2.42), we get,

$$\frac{1}{-8 + \Delta E} \approx \frac{1}{L^2 \Delta E} - \frac{\ln L}{4\pi} + \text{const.} \quad (2.43)$$

In the large- L limit, $\Delta E \rightarrow 0$ (as it is the interaction correction to the noninteracting energy), so we get, to the leading order of L ,

$$\Delta E = \frac{4\pi}{L^2 \ln L}. \quad (2.44)$$

We will check equation (2.44) in Section 2.4.

2.4 Large- L asymptotics for few-particle energy

The procedure used in Section 2.3 for two bosons can also be applied to problems with a few particles. For a few particles on a large lattice with short-range (here nearest-neighbor) interaction, two-particle interaction is the main contribution to energy. We write for two particles,

$$E(2, L) = E_0(2, L) + \Delta E(L). \quad (2.45)$$




Fig. 1. Boson and fermion $2(E(M, L) - E_0(M, L)) / (M(M-1))$ versus L for $M = 2, 3, 4, 5$. All curves appear to converge at large L . The fermion (p-wave) result is much less than the boson result (s-wave). The $M = 4$ plot goes to $L = 20$ and the $M = 5$ plot to $L = 10$. The boson $M = 5$ curve is too high to be included in this plot.

Here in this section we use the notation $E(M, L)$ and $E_0(M, L)$ to denote the M -particle exact and noninteracting ground state energies respectively and emphasize the dependence of ΔE on L by using $\Delta E(L)$. It is reasonable to expect that the energy for M particles is the noninteracting energy plus interaction corrections from the $M(M-1)/2$ pairs of particles. We then have,

$$E(M, L) \approx E_0(M, L) + \frac{M(M-1)}{2} \Delta E(L). \quad (2.46)$$

For bosons, $E_0(M, L) = -4M$, because in the ground state, all bosons occupy the zero-momentum state. On the other hand, for fermions, because of Pauli exclusion, no two fermions can occupy the same state, the noninteracting ground state is obtained from filling the M fermions from the lowest state ($\mathbf{k} = 0$) up.

Equation (2.46) implies that plotting $2(E(M, L) - E_0(M, L)) / (M(M-1))$ versus L for different M should all asymptotically at large L approach $\Delta E(L)$. In Figure 1, we do such plots, for bosons and fermions with $M = 2, 3, 4, 5$. The fermion results, from p-wave scattering (as our spinless fermion wave function has to be antisymmetric), are much smaller than the boson results (bold curves) from s-wave scattering.

Note that in our calculation for $\Gamma(E, 0, 0)$ equation (2.42), we have neglected the contribution of ΔE in the denominator except for the first term ($\mathbf{q} = 0$). Now with the leading form of ΔE equation (2.44), we can obviously plug $E \approx -8 + \Delta E$ into equation (2.42) to get the form of the next term,

$$\Delta E = \frac{4\pi}{L^2 \ln L} \left(A + \frac{B}{\ln L} + \frac{C}{(\ln L)^2} \right). \quad (2.47)$$




Fig. 2. Boson $(E(M, L) + 4M)L^2 \ln L / (2\pi M(M-1))$ versus $1/\ln L$ for $M = 2, 3, 4, 5$. Quadratic polynomial fitting is done for $M = 2$ and $M = 3$. The fitted constant coefficients are approximately one, and the other coefficients from $M = 2$ and $M = 3$ are comparable.

Using equations (2.46) and (2.47), we get, for a few bosons ($E_0(M, L) = -4M$),

$$\frac{(E(M, L) + 4M)L^2 \ln L}{2\pi M(M-1)} = A + B \frac{1}{\ln L} + C \left(\frac{1}{\ln L} \right)^2. \quad (2.48)$$

In Figure 2, we plot $(E(M, L) + 4M)L^2 \ln L / (2\pi M(M-1))$ versus $1/\ln L$ for $M = 2, 3, 4, 5$, using the boson data in Figure 1. Quadratic polynomial fitting is done for $M = 2, 3$, where we have more data than $M = 4, 5$. The coefficient $A \approx 1$ for both fits, implying, from equation (2.47), the leading-order term in $\Delta E(L)$ is indeed $4\pi / (L^2 \ln L)$. B and C from two fits are also comparable.

To summarize, from equations (2.46) and (2.47) and fitting in Figure 2, we find that in our model the energy of a small number M of bosons on a large $L \times L$ lattice is to the leading order of L ,

$$E(M, L) \approx -4M + \frac{M(M-1)}{2} \frac{4\pi}{L^2 \ln L}. \quad (2.49)$$

For two *fermions* on a large $L \times L$ lattice, the non-interacting energy – the lead term in equation (2.49) – is obviously lower for $\mathbf{P} = (0, 1)$ than for $\mathbf{P} = (0, 0)$. We have not worked out the asymptotic behavior for $\mathbf{P} \neq (0, 0)$.

3 The two-particle t-matrix

In Section 2.3 and Section 2.4, we studied the ground state energy of a few particles on a large lattice, and we showed that the energy of M particles can be approximated by

summing the energy of the $M(M-1)/2$ pairs. In this section, we reformulate the equations for two particles and derive a scattering matrix, the t-matrix. The t-matrix gives us equations of the form equations (1.2) and (1.3) which are more precise statements of the ideas presented in Sections 2.3 and 2.4. They apply to small lattices and to excited states.

One possible objection to the formulas (1.2) and (1.3) is that they appear to be perturbative, yet the interaction potential in our problem is infinitely strong, so the first-order (first Born approximation) scattering amplitude, being proportional to the potential, is infinite too. However, this singular potential scattering problem (e.g., hard-sphere interaction in 3D) has been solved (see Ref. [39]) by replacing the potential with the so-called scattering length, which is finite even when the potential is infinite. As we review in Appendix A, a perturbation series (Born series) can be written down (that corresponds to a series of the so-called ladder diagrams) and even though each term is proportional to the potential, the sum of all terms (the t-matrix, ΔE in Eqs. (1.2) and (1.3)) is finite.

Because the t-matrix captures two-body interaction effects, it is the centerpiece of dilute fermion and boson calculations with strong interactions. Field-theoretical calculations in both three and two dimensions are based on the ladder diagrams and the t-matrix. See Fetter and Walecka [40] for the 3D problem, Schick [10] for the 2D boson problem and Bloom [41] the 2D fermion problem. For lattice fermion problems, Kanamori [42] derived the t-matrix for a tight-binding model that is essentially a Hubbard model (this work is also described in Yosida [43]). And in reference [25], the t-matrix is worked out explicitly for the Hubbard model, and Kanamori's result is obtained. Reference [25] also evaluated the t-matrix for the dilute limit in three dimensions and obtained a functional dependence on particle density.

Rudin and Mattis [44] used the t-matrix expression derived in references [25, 42] and found upper and lower bounds of the fermion t-matrix in two dimensions in terms of particle density. Rudin and Mattis's result for the low-density limit of the two-dimensional Hubbard model is of the same functional form as Bloom's diagrammatical calculation for the two-dimensional fermion hard disks [41]. Since the discovery of high-temperature superconductors, Bloom's calculation has received a lot of attention because of its relevance to the validity of the Fermi liquid description of dilute fermions in two dimensions. There have been a number of works on the 2D dilute Fermi gas [45–47] and on the dilute limit of 2D Hubbard model [48], all using the t-matrix, but these results have not been checked by numerical calculations.

3.1 Setup and symmetry

To have an equation in the form of equation (1.2), we start with any pair of momentum vectors \mathbf{q}_1 and \mathbf{q}_2 and write noninteracting energy of the pair $E_0 = \mathcal{E}(\mathbf{q}_1) + \mathcal{E}(\mathbf{q}_2)$ and total momentum $\mathbf{P} = \mathbf{q}_1 + \mathbf{q}_2$. Because our Hamiltonian, equations (2.4) and (2.5), conserves total momentum, we

can restrict our basis states to $|\mathbf{q}, \mathbf{P} - \mathbf{q}\rangle$. It is tempting to take $|\mathbf{q}_1, \mathbf{P} - \mathbf{q}_1\rangle$ and $|\mathbf{q}_2, \mathbf{P} - \mathbf{q}_2\rangle$ as our nonperturbed states, but there can be other two-particle states with the same total momentum \mathbf{P} and energy E_0 .

In fact, using our energy dispersion function equation (2.11), if we write $\mathbf{q}_1 = (q_{1x}, q_{1y})$ and $\mathbf{q}_2 = (q_{2x}, q_{2y})$, and define $\mathbf{q}_3 = (q_{1x}, q_{2y})$ and $\mathbf{q}_4 = (q_{2x}, q_{1y})$ then we have, $\mathbf{q}_1 + \mathbf{q}_2 = \mathbf{q}_3 + \mathbf{q}_4$ and $\mathcal{E}(\mathbf{q}_1) + \mathcal{E}(\mathbf{q}_2) = \mathcal{E}(\mathbf{q}_3) + \mathcal{E}(\mathbf{q}_4)$. We call this fact, that component exchanges in the pair \mathbf{q}_1 and \mathbf{q}_2 give a pair \mathbf{q}_3 and \mathbf{q}_4 that have the same total momentum and energy, the *pair component exchange symmetry* of our energy dispersion function $\mathcal{E}(\mathbf{q})$. This symmetry is due to the fact that our $\mathcal{E}(\mathbf{q})$ is separable into a x part and a y part (i.e., $\mathcal{E}(\mathbf{q}) = \mathcal{E}_x(q_x) + \mathcal{E}_y(q_y)$ where $\mathcal{E}_x(q) = -2t \cos q = \mathcal{E}_y(q)$ in our model) [49].

The pair component exchange symmetry says that if $q_{1x} \neq q_{2x}$ and $q_{1y} \neq q_{2y}$, then the state $|\mathbf{q}_3, \mathbf{q}_4\rangle$, with \mathbf{q}_3 and \mathbf{q}_4 defined above using component exchange, has the same total momentum and energy as $|\mathbf{q}_1, \mathbf{q}_2\rangle$. The degenerate perturbation theory requires $|\mathbf{q}_3, \mathbf{q}_4\rangle$ should be included in the set of nonperturbed states with $|\mathbf{q}_1, \mathbf{q}_2\rangle$.

With a noninteracting two-particle energy E_0 and total momentum \mathbf{P} , we divide the N wavevectors into two disjoint sets,

$$Q_0 = \{\mathbf{q} | \mathcal{E}(\mathbf{q}) + \mathcal{E}(\mathbf{P} - \mathbf{q}) = E_0\}, \bar{Q} = \{\mathbf{q} | \mathbf{q} \notin Q_0\}. \quad (3.1)$$

Note that if $\mathbf{q} \in Q_0$ then $\mathbf{P} - \mathbf{q} \in Q_0$. Denote N_0 the number of elements in Q_0 and $\bar{N} = N - N_0$ the number of elements in \bar{Q} . With this separation of \mathbf{q} , our eigenstate equation (2.13) becomes

$$|\psi\rangle = \sum_{\mathbf{q} \in Q_0} g(\mathbf{q}) |\mathbf{q}, \mathbf{P} - \mathbf{q}\rangle + \sum_{\mathbf{q} \in \bar{Q}} g(\mathbf{q}) |\mathbf{q}, \mathbf{P} - \mathbf{q}\rangle, \quad (3.2)$$

where the first sum contains all states whose noninteracting energy is degenerate. Using the idea of degenerate perturbation theory, we expect to be able to find a secular matrix \mathcal{T} , $N_0 \times N_0$, for the degenerate states in Q_0 only, and \mathcal{T} will eventually be our momentum space t-matrix, which we will derive now.

Note that using equation (2.14), the number of independent states in the first sum of equation (3.2) is less than N_0 . We include both $|\mathbf{q}, \mathbf{P} - \mathbf{q}\rangle$ and $|\mathbf{P} - \mathbf{q}, \mathbf{q}\rangle$ in our calculation because we are considering boson and fermion problems at the same time: the symmetric solution $g(\mathbf{q}) = g(\mathbf{P} - \mathbf{q})$ is a boson solution and the antisymmetric solution $g(\mathbf{q}) = -g(\mathbf{P} - \mathbf{q})$ is a fermion solution (see Eq. (2.14)).

3.2 Derivation of the t-matrix

Our purpose is to derive a set of closed equations for $g(\mathbf{q})$, the coefficient in our two-particle state equation (3.2), with $\mathbf{q} \in Q_0$.

The Schrodinger equation for the two-particle state $|\psi\rangle$, equation (2.15), can now be written as,

$$(E - \mathcal{E}(\mathbf{q}) - \mathcal{E}(\mathbf{P} - \mathbf{q}))g(\mathbf{q}) = \frac{1}{N} \sum_{\mathbf{r}'} e^{i\mathbf{q}\cdot\mathbf{r}'} V(\mathbf{r}') \tilde{g}(\mathbf{r}'), \quad (3.3)$$

where $\tilde{g}(\mathbf{r})$ is the Fourier transform of $g(\mathbf{q})$ as defined in equation (2.16).

For $\mathbf{q} \in \bar{Q}$, if we assume that $E \neq \mathcal{E}(\mathbf{q}) + \mathcal{E}(\mathbf{P} - \mathbf{q})$, equation (3.3) becomes

$$\tilde{g}^0(\mathbf{r}) = \tilde{g}(\mathbf{r}) - \sum_{\mathbf{r}'} \bar{G}(E, \mathbf{P}; \mathbf{r}, \mathbf{r}') V(\mathbf{r}') \tilde{g}(\mathbf{r}'), \quad (3.4)$$

where we have defined a Green function for the set \bar{Q} ,

$$\bar{G}(E, \mathbf{P}; \mathbf{r}, \mathbf{r}') = \frac{1}{\bar{N}} \sum_{\mathbf{q} \in \bar{Q}} \frac{e^{i\mathbf{q}\cdot(\mathbf{r}' - \mathbf{r})}}{E - \mathcal{E}(\mathbf{q}) - \mathcal{E}(\mathbf{P} - \mathbf{q})}, \quad (3.5)$$

and a Fourier transform with vectors in Q_0 ,

$$\tilde{g}^0(\mathbf{r}) = \sum_{\mathbf{q} \in Q_0} e^{-i\mathbf{q}\cdot\mathbf{r}} g(\mathbf{q}). \quad (3.6)$$

By restricting to the nearest-neighbor repulsion potential equation (2.9), equation (3.4) becomes,

$$\tilde{g}^0(\mathbf{r}) = \tilde{g}(\mathbf{r}) - \sum_j \bar{G}(E, \mathbf{P}; \mathbf{r}, \mathbf{R}_j) V \tilde{g}(\mathbf{R}_j) \quad (3.7)$$

summed over neighbor vector equation (2.10). Now restricting $\mathbf{r} = \mathbf{R}_i$ in equation (3.7), we get a set of four equations,

$$\phi_{0i} = \phi_i - \sum_j \bar{G}_{ij}(E, \mathbf{P}) (V \phi_j), \quad (3.8)$$

where we have written

$$\bar{G}_{ij}(E, \mathbf{P}) = \bar{G}(E, \mathbf{P}; \mathbf{R}_i, \mathbf{R}_j) \quad (3.9)$$

and $\phi_i = \tilde{g}(\mathbf{R}_i)$ and $\phi_{0i} = \tilde{g}^0(\mathbf{R}_i)$. Equation (3.8) is a matrix equation,

$$\phi_0 = (I - \bar{\mathcal{G}}(E, \mathbf{P})V)\phi, \quad (3.10)$$

where $\bar{\mathcal{G}}$ is 4×4 , ϕ and ϕ_0 are 4×1 , and V is a scalar (strength of potential). And we can invert the matrix to get,

$$\phi = (I - \bar{\mathcal{G}}(E, \mathbf{P})V)^{-1} \phi_0. \quad (3.11)$$

This is a key result, as we have expressed the desired function \tilde{g} , a Fourier transform of $g(\mathbf{q})$ including all \mathbf{q} , in terms of \tilde{g}^0 which includes only $\mathbf{q} \in Q_0$; the information about other $\mathbf{q} \in \bar{Q}$ was packaged into the Green function $\bar{\mathcal{G}}(E, \mathbf{P})$.

Now we go back to equation (3.3), restrict the summation to \mathbf{R}_i , and substitute in $V \tilde{g}_i$ from equation (3.11), and we get,

$$(E - \mathcal{E}(\mathbf{q}) - \mathcal{E}(\mathbf{P} - \mathbf{q}))g(\mathbf{q}) = \sum_{\mathbf{q}' \in Q_0} T(E, \mathbf{P}; \mathbf{q}, \mathbf{q}')g(\mathbf{q}'). \quad (3.12)$$

where in the last step we have used the Fourier transform of $\tilde{g}^0(\mathbf{R}_j)$ equation (3.6) and defined,

$$T(E, \mathbf{P}; \mathbf{q}, \mathbf{q}') = \frac{1}{N} \sum_{ij} e^{i\mathbf{q}\mathbf{R}_i} e^{-i\mathbf{q}'\mathbf{R}_j} (V(I - \bar{\mathcal{G}}(E)V)^{-1})_{ij}. \quad (3.13)$$

If we restrict $\mathbf{q} \in Q_0$ in equation (3.12), then we have,

$$(E - E_0)g(\mathbf{q}) = \sum_{\mathbf{q}' \in Q_0} T(E, \mathbf{P}; \mathbf{q}, \mathbf{q}')g(\mathbf{q}'), \quad (3.14)$$

which means,

$$E = E_0 + \text{Eigenvalue}(\mathcal{T}(E)), \quad (3.15)$$

where we have written

$$\mathcal{T}_{\mathbf{q}, \mathbf{q}'}(E) = T(E, \mathbf{P}; \mathbf{q}, \mathbf{q}') \quad (3.16)$$

and left out the dependence on \mathbf{P} . $\mathcal{T}_{\mathbf{q}, \mathbf{q}'}$ is the t-matrix in momentum space. Both \mathbf{q} and \mathbf{q}' in equation (3.14) are in Q_0 , which means that if there are N_0 elements in Q_0 then the matrix $\mathcal{T}(E)$ is $N_0 \times N_0$.

Equation (3.15) is our desired equation that shows explicitly the interaction correction to the noninteracting energy E_0 . In Appendix A, we show the physical meaning of $T(E, \mathbf{P}; \mathbf{q}, \mathbf{q}')$ in the language of diagrammatic perturbation theory, namely it is the sum total of all the terms with repeated scattering of the same two particles. This t-matrix formalism for the two-particle problem is therefore exact, and it is exactly equivalent to the Schrodinger equation and the Green function formalism in Section 2. The resulting equation is an implicit equation on E , of the form $E = E_0 + \Delta E(E)$ of equation (1.2), and we will show in a later section that for fermions the approximation $E \approx E_0 + \Delta E(E_0)$ is often very good.

Note also that for our case $V = +\infty$, the t-matrix expression equation (3.13) becomes

$$T(E, \mathbf{P}; \mathbf{q}, \mathbf{q}') = \frac{1}{N} \sum_{ij} e^{i\mathbf{q}\mathbf{R}_i} e^{-i\mathbf{q}'\mathbf{R}_j} (-\bar{\mathcal{G}}(E)^{-1})_{ij}, \quad (3.17)$$

where the potential V cancels out, giving a finite value. This is one of the advantages of the t-matrix formalism that it can deal with infinite (singular) potential, for which straightforward perturbation theory would diverge.

The definition of $T(E, \mathbf{P}; \mathbf{q}, \mathbf{q}')$ in equation (3.13) is a Fourier transform of the real space quantity $V(I - \bar{\mathcal{G}}(E)V)^{-1}$. Here $\bar{\mathcal{G}}$ is 4×4 because we have nearest-neighbor interaction. When there is only on-site interaction, as is in the usual Hubbard model case, $\bar{\mathcal{G}}(E) = \bar{G}(E, \mathbf{P}, (0, 0), (0, 0))$, equation (3.5), is a scalar. Then, we can simply use the scalar quantity $V/(I - \bar{G}V)$, which is the t-matrix that has appeared in Kanamori [42], Mattis [25], Rudin and Mattis [44], and Yosida [43]. Our expression, equation (3.13), is more complicated because we have nearest-neighbor interaction (and thus the relevance of \mathbf{R}_j).

3.3 Symmetry considerations

In Section 2.2.1, after deriving the general Green function equation using $\mathcal{G}(E)$, we specialized to rectangular-boundary lattices and used lattice reflection symmetries to diagonalize the 4×4 matrix $\mathcal{G}(E)$ and obtained scalar

equations. Here our t-matrix equation (3.15) requires us to find the eigenvalues of the t-matrix \mathcal{T} . In this section, we use particle permutation symmetry and pair component exchange symmetry to diagonalize the $N_0 \times N_0$ t-matrix $\mathcal{T}(E)$ for a few special cases.

3.3.1 $N_0 = 1$

There is only one momentum vector in Q_0 . Let us write $Q_0 = \{\mathbf{q}_1\}$ (this implies that $\mathbf{P} - \mathbf{q}_1 = \mathbf{q}_1$). Then there is only one unperturbed two-particle basis state $|\mathbf{q}_1, \mathbf{q}_1\rangle$ (see Eq. (3.2)). This must be a boson state, and $\mathcal{T}(E)$ is a number. We write the resulting scalar equation as,

$$E = E_0 + T_1(E). \quad (3.18)$$

3.3.2 $N_0 = 2$

Here $Q_0 = \{\mathbf{q}_1, \mathbf{q}_2\}$ with $\mathbf{q}_1 + \mathbf{q}_2 = \mathbf{P}$. The basis states are $|\mathbf{q}_1, \mathbf{q}_2\rangle$ and $|\mathbf{q}_2, \mathbf{q}_1\rangle$. The symmetric (boson) combination is $(|\mathbf{q}_1, \mathbf{q}_2\rangle + |\mathbf{q}_2, \mathbf{q}_1\rangle)/\sqrt{2}$, and the antisymmetric (fermion) combination is $(|\mathbf{q}_1, \mathbf{q}_2\rangle - |\mathbf{q}_2, \mathbf{q}_1\rangle)/\sqrt{2}$. These have to be the eigenvectors of $\mathcal{T}(E)$. And that is to say that if we define

$$\mathcal{S}_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ 1 & -1 \end{pmatrix}, \quad (3.19)$$

then we have $\mathcal{S}_2 = \mathcal{S}_2^t$, $\mathcal{S}_2^2 = I$, and

$$\mathcal{S}_2 \mathcal{T}(E) \mathcal{S}_2 = \begin{pmatrix} T_{1,1}(E) & 0 \\ 0 & T_{1,-1}(E) \end{pmatrix}. \quad (3.20)$$

Here $T_{1,1}(E)$ and $T_{1,-1}(E)$ are scalars that correspond to boson and fermion symmetries respectively. And our t-matrix equation (3.15) is reduced to two scalar equations,

$$E = E_0 + T_{1,1}(E), \quad E = E_0 + T_{1,-1}(E), \quad (3.21)$$

for bosons and fermions respectively. Our notation for the eigenvalues of $\mathcal{T}(E)$ is always to write T with subscripts that are the coefficients (in order) of the N_0 two-particle basis vectors.

3.3.3 $N_0 = 4$

Here $Q_0 = \{\mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3, \mathbf{q}_4\}$ with $\mathbf{q}_1 + \mathbf{q}_2 = \mathbf{q}_3 + \mathbf{q}_4 = \mathbf{P}$. The basis states are $|\mathbf{q}_1, \mathbf{q}_2\rangle$, $|\mathbf{q}_2, \mathbf{q}_1\rangle$, $|\mathbf{q}_3, \mathbf{q}_4\rangle$, and $|\mathbf{q}_4, \mathbf{q}_3\rangle$. Using particle permutation symmetry, we get two states with even symmetries appropriate for bosons, which generically would be

$$a(|\mathbf{q}_1, \mathbf{q}_2\rangle + |\mathbf{q}_2, \mathbf{q}_1\rangle) + b(|\mathbf{q}_3, \mathbf{q}_4\rangle + |\mathbf{q}_4, \mathbf{q}_3\rangle), \\ -b(|\mathbf{q}_1, \mathbf{q}_2\rangle + |\mathbf{q}_2, \mathbf{q}_1\rangle) + a(|\mathbf{q}_3, \mathbf{q}_4\rangle + |\mathbf{q}_4, \mathbf{q}_3\rangle), \quad (3.22)$$

and two odd (fermion-type) states,

$$a(|\mathbf{q}_1, \mathbf{q}_2\rangle - |\mathbf{q}_2, \mathbf{q}_1\rangle) + b(|\mathbf{q}_3, \mathbf{q}_4\rangle - |\mathbf{q}_4, \mathbf{q}_3\rangle),$$

$$-b(|\mathbf{q}_1, \mathbf{q}_2\rangle - |\mathbf{q}_2, \mathbf{q}_1\rangle) + a(|\mathbf{q}_3, \mathbf{q}_4\rangle - |\mathbf{q}_4, \mathbf{q}_3\rangle) \quad (3.23)$$

where a and b are arbitrary coefficients to be determined.

Recall $N_0 = 4$ means the pair $(\mathbf{q}_1, \mathbf{q}_2)$ has the same total momentum and energy as $(\mathbf{q}_3, \mathbf{q}_4)$, which may happen for various reasons. When the reason is the pair component exchange symmetry (of Sect. 3.1), i.e. $\mathbf{q}_3 = (q_{1x}, q_{2y})$ and $\mathbf{q}_4 = (q_{2x}, q_{1y})$, then $a = b = 1/2$, due to a hidden symmetry under the permutation $1 \leftrightarrow 3, 2 \leftrightarrow 4$. The only effect this permutation has on the momentum transfers $\mathbf{q}_i - \mathbf{q}_j$ is to change the sign of one or both components; but the potential $V(\mathbf{r})$ is symmetric under reflection through either coordinate axis, hence $V(\mathbf{q}_i - \mathbf{q}_j)$ is invariant under the permutation. Since the t -matrix depends only on $V(\mathbf{q}_i - \mathbf{q}_j)$, it inherits this symmetry. Next, if we define

$$\mathcal{S}_4 = \frac{1}{2} \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & -1 & -1 \\ 1 & -1 & 1 & -1 \\ 1 & -1 & -1 & 1 \end{pmatrix}, \quad (3.24)$$

then we have $\mathcal{S}_4 = \mathcal{S}_4^t$, $\mathcal{S}_4^2 = I$, and $\mathcal{S}_4 \mathcal{T}(E) \mathcal{S}_4$ becomes diagonal with four eigenvalues of $\mathcal{T}(E)$: $T_{1,1,1,1}(E)$, $T_{1,1,-1,-1}(E)$, $T_{1,-1,1,-1}(E)$, and $T_{1,-1,-1,1}(E)$. And our t -matrix equation (3.15) is reduced to

$$E = E_0 + T_{1,1,1,1}(E), \quad E = E_0 + T_{1,1,-1,-1}(E), \quad (3.25)$$

for bosons and

$$E = E_0 + T_{1,-1,1,-1}(E), \quad E = E_0 + T_{1,-1,-1,1}(E), \quad (3.26)$$

for fermions.

The three cases $N_0 = 1, 2$, and $N_0 = 4$ with pair component exchange symmetry are three special cases in which we know the eigenvectors of \mathcal{T} and can therefore diagonalize \mathcal{T} from symmetry considerations easily.

Different or larger values of N_0 are possible when \mathbf{P} has a special symmetry, e.g. when $P_x = P_y$, $N_0 = 8$ generically since Q_0 includes pairs such as $(q_{1y}, q_{1x}), (q_{2y}, q_{2x})$. For these general cases, we return to equation (3.15) and diagonalize \mathcal{T} numerically. For example, on a $L \times L$ lattice, the pairs $(0,1)(0,-1)$ and $(1,0)(-1,0)$ have the same total energy and momentum, but this is not due to the pair component exchange symmetry. In this case, we numerically diagonalize the 4×4 matrix $\mathcal{T}(E)$, and we find that in the fermion eigenvectors, equation (3.23), $a \neq b$.

3.4 Solving for energy

The example system that we will study here is 10×11 with $\mathbf{P} = (0, 0)$. The noninteracting and interacting energies of the system are in Table 1. It can be seen that all of the energies listed in Table 1 are of the three cases discussed in Section 3.3: $N_0 = 1$, $N_0 = 2$, and $N_0 = 4$ due to pair component exchange symmetry.




Fig. 3. $f(E) = E_0 + T_1(E)$ versus E for 10×11 lattice with $\mathbf{P} = (0, 0)$ and $E_0 = -8.0$ (i.e., $E_0 = \mathcal{E}(0) + \mathcal{E}(0)$.) The intersections with the line $y = E$ are the exact two-particle energies.

We solve for energy E in the implicit equation, $E = E_0 + T(E)$, where $T(E)$ represents the eigenvalues of $\mathcal{T}(E)$, e.g., $T_{1,-1}(E)$. We plot $f(E) = E_0 + T(E)$ along with a line $y = E$. Their intersections are the desired energies E .

3.4.1 $N_0 = 1$ case

In Figure 3, we plot $f(E)$ versus E for the 10×11 lattice with $\mathbf{P} = (0, 0)$ and the noninteracting energy $E_0 = -8.0 = \mathcal{E}(0) + \mathcal{E}(0)$. Here $Q_0 = \{(0, 0)\}$, and the nonperturbed state is $|\mathbf{q}_1 = (0, 0), \mathbf{P} - \mathbf{q}_1 = (0, 0)\rangle$ which can only be a boson state. The energy intersections from Figure 3 are -7.906 , -7.299 , -6.971 , -6.022 , and so on. Looking into Table 1, we see that these are all boson energies.

In Figure 3, note also that the energy -6.601 , which is an exact eigenenergy from exact diagonalization, does not appear as an intersection in Figure 3. This is a special energy, being also a noninteracting energy. Earlier, as mentioned at the beginning of Section 3.2, we assumed that our $E \neq \mathcal{E}(\mathbf{q}) + \mathcal{E}(\mathbf{P} - \mathbf{q})$ for any $\mathbf{q} \in \bar{Q}$, so this energy is excluded from our t -matrix formulation. We will address later in Section 3.4.3 this kind of exact solutions that are also noninteracting energies.

Note that our equation $E = E_0 + T(E)$ is a reformulation of the Schrodinger equation with certain symmetry considerations, and it should be satisfied by all energies E with the same symmetry. Building $T(E)$ from E_0 and \mathbf{P} does not automatically give us a unique interacting energy E that corresponds to the noninteracting energy E_0 . However, we can see clearly from Figure 3, if we perturb the exact solutions by a small amount $E \rightarrow E + \delta$, then $f(E)$ changes drastically except for the lowest energy $E = -7.906$. That is to say that these other energies, for example $E = -6.971$, are exact solutions of the equation $f(E) = E$, but they are not stable solutions. From the plot, only $E = -7.906$ comes close to being stable.

Table 1. The 12 low-lying noninteracting and exact two-particle energies of the 10×11 lattice with total momentum $\mathbf{P} = (0, 0)$. \mathbf{q}_1 and $\mathbf{q}_2 = \mathbf{P} - \mathbf{q}_1$ are the momentum vectors.

\mathbf{q}_1	\mathbf{q}_2	$\mathcal{E}(\mathbf{q}_1) + \mathcal{E}(\mathbf{q}_2)$	boson	fermion
(0, 0)	(0, 0)	-8.0000000000	-7.9068150537	-7.3117803781
(0, 1)	(0, -1)	-7.3650141313	-7.2998922545	-7.1770594424
(1, 0)	(-1, 0)	-7.2360679774	-6.9713379459	-6.4994071102
(1, -1)	(-1, 1)	-6.6010821088	-6.6010821088	-6.4700873024
(1, 1)	(-1, -1)	-6.6010821088	-6.0227385416	-5.5449437453
(0, 2)	(0, -2)	-5.6616600520	-5.4277094111	-5.1475674826
(2, 0)	(-2, 0)	-5.2360679774	-5.0769765528	-4.8309218202

We can be more precise about this notion of stability. If we have an iteration $x_{n+1} = f(x_n)$, and x^* is a fix point (i.e., $f(x^*) = x^*$), then the iteration is linear stable at x^* if and only if $|f'(x^*)| < 1$. In our plots, we have included a line $y = E$ with slope one, which can be used as a stability guide. An intersection (fix point) is linearly stable when the function $f(E)$ at the intersection is not as steep as the straight line.

3.4.2 $N_0 = 2$ case

In Figure 4 we plot for $E_0 = -7.365$ and $\mathbf{P} = (0, 0)$ with $Q_0 = \{(0, 1), (0, -1)\}$. The boson function $f(E) = E_0 + T_{1,1}(E)$ is the dotted line in the top graph, and the fermion function $f(E) = E_0 + T_{1,-1}(E)$ is the solid line in the bottom graph.

The intersections closest to $E_0 = -7.365$ are -7.299 , the first excited boson energy (see Tab. 1), and -7.311 , the lowest fermion energy. Note that the curve on which the fermion intersection (-7.311) lies is very flat. In other words for this fermion energy $E \approx E_0 + T(E_0)$, i.e., the first iteration using the noninteracting energy gives an energy very close to the exact value. More precisely, we find with $E_0 = -7.365014$, $f(E_0) = E_0 + T(E_0) = -7.310584$, which is very close to $E = -7.31178$. Many t-matrix calculations [25, 42–44], use the first iteration $E \approx E_0 + T(E_0)$ as an approximation to the exact energy, and we see in this case this approximation is very good. (We will come back to this point later in Sect. 3.5.)

3.4.3 $N_0 = 4$ case

In Figures 5 and 6, we plot $f(E)$ for $E_0 = \mathcal{E}(1, -1) + \mathcal{E}(-1, 1) = \mathcal{E}(1, 1) + \mathcal{E}(-1, -1) = -6.601$ and $\mathbf{P} = (0, 0)$. For this $N_0 = 4$ case we have two boson functions, plotted in Figure 5, $f(E) = E_0 + T_{1,1,1,1}(E)$ (dotted line) and $f(E) = E_0 + T_{1,1,-1,-1}(E)$ (dot-dashed line), and we have two fermion functions, plotted in Figure 6, $f(E) = E_0 + T_{1,-1,1,-1}(E)$ (solid line) and $f(E) = E_0 + T_{1,-1,-1,1}(E)$ (dashed line). The fermion intersections closest to E_0 are -6.499 and -6.470 . Here again the two fermion curves are very flat. The two boson intersections closest to E_0 are -6.022 and -6.601 . Note that the latter is also a noninteracting energy, and it is the intersection of the horizontal line $E = E_0$ with $y = E$.




Fig. 4. $f(E)$ versus E for 10×11 lattice with $\mathbf{P} = (0, 0)$ and $E_0 = -7.365$ (i.e., $E_0 = \mathcal{E}(0, 1) + \mathcal{E}(0, -1)$.) The top graph (dotted line) is for boson $f(E) = E_0 + T_{1,1}(E)$, and the top graph (solid line) for fermion $f(E) = E_0 + T_{1,-1}(E)$. The fermion curve is essentially flat near $E = E_0$.

One interesting observation of the fermion plot in Figure 6 is that pairs of closely spaced energies (for example -7.311 and -7.177) lie on different symmetry curves. We know that if we have a square lattice (for example 10×10) then the noninteracting fermion energies come in pairs. Here, we have chosen a 10×11 lattice that is close to a square but does not have exact degeneracies. We see




Fig. 5. Boson $f(E)$ versus E for 10×11 lattice with $\mathbf{P} = (0, 0)$ and $E_0 = -6.601$. The dotted line is for $T_{1,1,1,1}$ and the horizontal dot-dashed line for $T_{1,1,-1,-1}$ (which corresponds to a noninteracting state, see text at the end of this section).




Fig. 6. Fermion $f(E)$ versus E for 10×11 lattice with $\mathbf{P} = (0, 0)$ and $E_0 = -6.601$. The solid line is for $T_{1,-1,1,-1}$ and the long-dashed line for $T_{1,-1,-1,1}$. Note that closely spaced fermion energy pairs are separated by symmetry.

that the resulting closely spaced pairs are separated by symmetry considerations.

Another interesting observation from Figure 5 for bosons is that we have a horizontal line that corresponds to $T_{1,1,-1,-1}(E) = 0$. For this case the noninteracting energy is an exact energy. That is to say, $(1, 1, -1, -1)$ is a null vector of $\mathcal{T}(E)$ (see Sect. 3.3.3), or the eigenstate,

$$|\mathbf{q}_1, \mathbf{q}_2\rangle + |\mathbf{q}_2, \mathbf{q}_1\rangle - |\mathbf{q}_3, \mathbf{q}_4\rangle - |\mathbf{q}_4, \mathbf{q}_3\rangle, \quad (3.27)$$

with $\mathbf{q}_3 = (q_{1x}, q_{2y})$ and $\mathbf{q}_4 = (q_{2x}, q_{1y})$ is an exact eigenstate of the Hamiltonian. This can be shown easily using the Schrodinger equation (2.15). We have $g(\mathbf{q}_1) = g(\mathbf{q}_2) = 1$, $g(\mathbf{q}_3) = g(\mathbf{q}_4) = -1$, and $g(\mathbf{q}) = 0$ for all other \mathbf{q} , and

we can easily show $V(\mathbf{q} - \mathbf{q}_1) + V(\mathbf{q} - \mathbf{q}_2) - V(\mathbf{q} - \mathbf{q}_3) - V(\mathbf{q} - \mathbf{q}_4) = 0$ (because $V(\mathbf{k})$ can be separated into a sum of two terms that involve the x and y components separately).

Transforming to the real space, without worrying about normalization, we can have

$$\begin{aligned} \tilde{g}(\mathbf{r}) &= \sum_{\mathbf{q}} e^{-i\mathbf{q}\cdot\mathbf{r}} g(\mathbf{q}) \\ &\sim (e^{-iq_{1x}x} - e^{-iq_{2x}x}) (e^{-iq_{1y}y} - e^{-iq_{2y}y}), \end{aligned} \quad (3.28)$$

where we have used the fact mentioned above that $g(\mathbf{q})$ is not zero for only four \mathbf{q} 's which are related by pair component exchange symmetry. It is clear from equation (3.28) that $\tilde{g}(0, y) = 0 = \tilde{g}(x, 0)$, which means that the wave function in relative position space is “d-wave” like, having nodes along x and y axes (thus happens to have nodes at every relative position where the potential would be nonzero).

3.5 Fermion: noninteracting to interacting

In this section we use the t-matrix techniques developed in the preceding sections to study the relationship between the noninteracting energies and the interacting energies. We start with the table of energies in Table 1 for the 10×11 lattice with $\mathbf{P} = (0, 0)$. We have asked in the introduction to this section whether we can go from the noninteracting to the interacting energies and now we know that we have an equation $E = E_0 + T(E)$ where $T(E)$ is the symmetry reduced scalar t-matrix function. From our graphs (Figs. 4 and 6) we have commented that for fermions the curve of $T(E)$ around E_0 is quite flat (which is not the case for bosons). And we mentioned that this implies that the approximation $E \approx E_0 + T(E_0)$ is close to the exact energy. Now in this section, we study the t-matrix approach for a specific system. We will denote $E_1 = E_0 + T(E_0)$, the first iteration result, and $E_{n+1} = E_0 + T(E_n)$, the n th iteration result.

In Table 2 we show the t-matrix calculation for the 10×11 lattice. We show for the lowest few states the noninteracting energy E_0 , the first t-matrix iteration E_1 , the fifth t-matrix iteration E_5 , and the exact energy E_{exact} . In Figure 7 these energy levels are plotted graphically. From the table, it is clear that the first t-matrix iteration result E_1 is quite close to the exact energy, and the fifth iteration result E_5 gives a value that is practically indistinguishable from the exact value.

In principle, the t-matrix energy for a pair of particles might be shifted out of the continuum of two independent particle energies, creating a bound state. The potential has s-like symmetry so this is possible only for an s-like relative wavefunction, i.e. for bosons since spinless fermions must have a spatially antisymmetric wavefunction. Our constraint affects low-energy states like a repulsive potential, invariably shifting their energies upwards compared to the noninteracting case (as evident in the exact energies, e.g. Tabs. 1 and 2). Thus bound state formation never occurs for the low-energy states which we studied.

Table 2. Fermion energies for 10×11 lattice with $\mathbf{P} = (0, 0)$. $E_0 = \mathcal{E}(\mathbf{q}_1) + \mathcal{E}(\mathbf{q}_2)$ is the noninteracting energy. $E_n = E_0 + T(E_{n-1})$ where $T(E)$ is the symmetry reduced t-matrix. Here only fermion energies (from $T_{1,-1}$ or $T_{1,-1,1,-1}$ and $T_{1,-1,-1,1}$) are included.

\mathbf{q}_1	\mathbf{q}_2	E_0	E_1	E_5	E_{exact}
(0,1)	(0,-1)	-7.365014	-7.310598893	-7.311780378	-7.311780378
(1,0)	(-1,0)	-7.236067	-7.175212790	-7.177059440	-7.177059442
(1,-1)	(-1,1)	-6.601082	-6.493807907	-6.499407060	-6.499407110
(1,1)	(-1,-1)	-6.601082	-6.460962404	-6.470087137	-6.470087302
(0,2)	(0,-2)	-5.661660	-5.532751985	-5.544942250	-5.544943745
(2,0)	(-2,0)	-5.236067	-5.134290466	-5.147558003	-5.147567483




Fig. 7. Two-fermion energy levels for the 10×11 lattice with $\mathbf{P} = (0, 0)$. From left to right, the lowest few noninteracting energies E_0 , first t-matrix iteration E_1 , fifth t-matrix iteration E_5 , and the exact energy E_{exact} are plotted. Note that the third noninteracting energy from the bottom is doubly degenerate (see Tab. 2).

Furthermore, nearest-neighbor hopping on a bipartite lattice admits an exact symmetry relating states of energy $-E \rightarrow +E$, so (in the limit of infinite V) the pair states near the band maximum have their energies *reduced* by the hard-core exclusion and are not bound either.

4 A few fermions: shell effect and t-matrix

In Section 2, we used lattice Green function to study the problem of two particles (bosons and fermions), and at the end of that section, in Section 2.4, we obtained the ground state energy of a few particles on a large lattice by summing up the energy of each pair of particles. This section contains a much more detailed study of the few-fermion problem: we will consider first the fermion shell effect and then we will study the interaction correction to energy (ground state and excited states) for a few fermions (three, four, and five) using the t-matrix.

Our results – summarized in Section 4.6 – confirm that, in the dilute limit, almost all of the interaction correction

is accounted for by the *two-body* terms of the t-matrix approximation, equation (4.1). But (recall Eq. (1.3)) that is a hallmark of a Fermi liquid picture; i.e., our numerical results suggest its validity at low densities. This is a nontrivial result, in that firstly, the validity of Fermi liquid theory in a finite-system context has rarely been considered. Standard t-matrix theory depends on a Fermi surface which (at $T = 0$) is completely sharp in momentum space, and every pair's t-matrix excludes scattering into the same set of occupied states. In a finite system, however, the allowed \mathbf{q} vectors fall on a discrete grid, and since the total number of particles is finite, the t-matrices of different pairs see a somewhat different set of excluded states (since they do not exclude themselves, and one particle is a non-negligible fraction of the total).

Secondly, and more essentially, the analytic justifications of Fermi liquid theory exist only in the cases of spinful fermions (in a continuum). That case is dominated by s-wave scattering, so that the t-matrix approaches a constant in the limit of small momenta (and hence in the dilute limit). Our spinless case is rather different, as will be elaborated in Section 5, because the t-matrix is dominated by the p-wave channel, which vanishes at small momenta. Thus the \mathbf{q} dependence is crucial in our case, and the numerical agreement is less trivial than it would be for s-wave scattering.

In this section, after an exhibition of the shell effect (Sect. 4.1), we present a general recipe for the multi-fermion t-matrix calculation. This is developed by the simplest cases, chosen to clarify when degeneracies do or do not arise.

4.1 Fermion shell effect

At zero temperature, the ground state of noninteracting fermions is formed by filling the one-particle states one by one from the lowest to higher energies. For our model of spinless fermions on a square lattice, we have the two ingredients for the shell effect: fermionic exclusion and degeneracies of one-particle states due to the form of our energy function and lattice symmetry. Shell effects have been noted previously in interacting models [50]; our code, permitting non-rectangular boundary conditions, allows us to see even more cases of them.

In Figure 8 we show the exact and for comparison the noninteracting ground state energies for the 5×8 lattice for




Fig. 8. Shell effect for 5×8 lattice (fermion, top graph; boson, bottom graph). Exact, interacting groundstate energies are compared with noninteracting energies for up to seven particles. Energy increment $E(M) - E(M - 1)$ is shown.

up to seven particles. The energy increment curve $E(M) - E(M - 1)$ is plotted and shows clearly the shell effect.

With fermions, the filled shells for the 5×8 lattice are at $M = 3$ (with momentum vectors $(0, 0)(0, \pm 1)$ occupied) and $M = 5$ (with $(0, 0)(0, \pm 1)(\pm 1, 0)$ occupied). For comparison, because bosons can all be at the zero-momentum state, where energy is -4 , the total noninteracting energy is $-4M$, so the exact energy curve shows smooth changes when M increases. There is no shell effect.

4.2 General multi-fermion theory

The key notion for generalizing our two-fermion approach to M fermions is that the set Q_0 now consists of every M -tuple α of wavevectors that gives the same total momentum and noninteracting energy. This defines a reduced Hilbert space, with the corresponding basis states $|\phi_\alpha\rangle$. We can construct an approximate, effective Hamiltonian $H_0 + H_{\text{tm}}$ acting within Q_0 -space, where H_{tm} is a sum of pairwise t-matrix terms, each of which changes just two

fermion occupancies:

$$H_{\text{tm}} = \sum'_{\alpha\beta} \mathbf{T}^{\alpha\beta}. \quad (4.1)$$

The notation $\sum'_{\alpha\beta}$ means the sum only includes the pair (α, β) when $|\phi_\alpha\rangle$ differs from $|\phi_\beta\rangle$ by a change of two fermions.

Thus, each term in equation (4.1) is associated with a particular fermion pair $(\mathbf{q}_i, \mathbf{q}_j)$. Each such pairwise t-matrix can be viewed as a sum of all possible repeated scatterings of those two fermions through intermediate states, except that intermediate states which are already included in Q_0 are excluded. (The most important omission of this approximation would be the processes in which three or more fermions are permuted before the system returns to the Q_0 Hilbert space.) Each term is a two-fermion t-matrix calculated according to the recipe of Sections 3.2 and 3.4 – thus each term has its own two-fermion wavevector set Q_0^{ij} and complementary set \bar{Q}^{ij} , as defined in equation (3.1). The only change in the recipe is to augment the set \bar{Q}^{ij} of wavevectors forbidden in the intermediate scatterings of the two fermions, since they cannot scatter into states already occupied by the other fermions in states α and β . (See Eq. (4.2) for an example.)

The t-matrix treatment is a form of perturbation expansion, for which the small parameter is obviously not V (which is large) but instead $1/L^2$, as is evident from equation (2.44). That is, as the lattice size is increased (with a fixed set of fermions), the approximation captures a larger and larger fraction of the difference $E_{\text{exact}} - E_0$.

4.3 A three-fermion t-matrix calculation

We first compute the energy of three fermions ($M = 3$) for the 8×9 lattice with $\mathbf{P} = 0$. For this example calculation, we have chosen $L_x \neq L_y$ to reduce the number of degeneracies in the noninteracting spectrum. In Figure 9 we show the lowest five noninteracting levels and the corresponding states in momentum space.

Let us consider the lowest noninteracting state in the 8×9 , $\mathbf{P} = (0, 0)$, and $M = 3$ system, with three momentum vectors: $\mathbf{q}_1 = (0, 1)$, $\mathbf{q}_2 = (0, 0)$, and $\mathbf{q}_3 = (0, -1)$ (see Fig. 9). And let us first consider the interaction of the pair \mathbf{q}_1 and \mathbf{q}_2 . The noninteracting energy of the pair is $E_0^{12} = \mathcal{E}(\mathbf{q}_1) + \mathcal{E}(\mathbf{q}_2) = -7.682507$ and the total momentum is $\mathbf{P}_{12} = \mathbf{q}_1 + \mathbf{q}_2 = (0, 1)$. As usual, we use E_0^{12} and \mathbf{P}_{12} to form the set Q_0^{12} (Eq. (3.1)). Here there are no other degenerate vectors so $Q_0^{12} = \{(0, 0), (0, 1)\}$. The three-particle problem is different from the two-particle case in the choice of \bar{Q}^{12} , the set of momentum vectors that the two particles can scatter into. Due to the presence of the third particle and Pauli exclusion, the two particles at $\mathbf{q}_1 = (0, 1)$ and $\mathbf{q}_2 = (0, 0)$ cannot be scattered into the momentum vector $\mathbf{q}_3 = (0, -1)$, so we must exclude \mathbf{q}_3 from \bar{Q}^{12} . Furthermore, even though there is no particle at $\mathbf{P}_{12} - \mathbf{q}_3 = (0, 1) - (0, -1) = (0, 2)$, this momentum cannot be scattered into, because otherwise the other particle would be scattered into the occupied \mathbf{q}_3 . That is to




Fig. 9. Lowest five noninteracting energy levels for the 8×9 lattice with $M = 3$ fermions and total momentum $\mathbf{P} = (0, 0)$. States in momentum space are drawn.

say, the momentum vectors that can be scattered into are

$$\bar{Q}^{12} = \{\mathbf{q} | \mathbf{q} \neq \mathbf{q}_1, \mathbf{q}_2, \mathbf{q}_3, \mathbf{P}_{12} - \mathbf{q}_3\} \quad (4.2)$$

This exclusion is shown graphically in Figure 10.

The t-matrix formalism can then be applied using Q_0^{12} and \bar{Q}^{12} to compute energy correction $\tilde{T}^{12}(E)$ for the interaction of the \mathbf{q}_1 and \mathbf{q}_2 pair. Here $\tilde{T}^{12}(E)$ is the “fermion” function $T_{1,-1}(E)$ (Eq. (3.21)), corresponding to the antisymmetric eigenvector of the t-matrix $\tilde{T}(E)$; the tilde denotes the modification due to exclusion of the set \bar{Q}^{12} . When the t-matrix contributes a small correction, it is accurate to use the bare values, $E_0^{ij} \equiv \mathcal{E}(\mathbf{q}_i) + \mathcal{E}(\mathbf{q}_j)$, and this approximation was used for all tables and figures in this section.

The total energy within this approximation is a sum of the t-matrix corrections for all possible pairs in the system, which are $(\mathbf{q}_2, \mathbf{q}_3)$, and $(\mathbf{q}_1, \mathbf{q}_3)$ in the present case:

$$E_{\text{tm}} = \mathcal{E}(\mathbf{q}_1) + \mathcal{E}(\mathbf{q}_2) + \mathcal{E}(\mathbf{q}_3) + \tilde{T}^{12}(E^{12}) + \tilde{T}^{13}(E^{13}) + \tilde{T}^{23}(E^{23}). \quad (4.3)$$

This is a special case of the effective Hamiltonian equation (4.1), which reduces to a 1×1 matrix in the nondegenerate case. (That is, whenever the set Q_0 of multi-fermion occupations has just one member.) The momentum space exclusions due to the presence of other particles are depicted in Figure 10, and the numerical values of this calculation are given in Table 3.

A more accurate approximation is to enforce a self-consistency,

$$E^{ij} \equiv E_0^{ij} + \tilde{T}^{ij}(E^{ij}) \quad (4.4)$$

where as defined above $E_0^{ij} \equiv \mathcal{E}(\mathbf{q}_i) + \mathcal{E}(\mathbf{q}_j)$. It should be cautioned that the physical justification is imperfect:




Fig. 10. Momentum space exclusions in t-matrix $M = 3$ calculation for the state $(0,0)(0,1)(0,-1)$. The crosses indicate exclusions when calculating pair energy for $(0,0)(0,1)$ (left figure), $(0,0)(0,-1)$ (middle), and $(0,1)(0,-1)$ (right) respectively.

Table 3. T-matrix calculation for the 8×9 lattice with $M = 3$ noninteracting particles $\mathbf{q}_1 = (0, 0)$, $\mathbf{q}_2 = (0, 1)$, and $\mathbf{q}_3 = (0, -1)$. The total noninteracting energy is $E_0 = \mathcal{E}(\mathbf{q}_1) + \mathcal{E}(\mathbf{q}_2) + \mathcal{E}(\mathbf{q}_3)$ and the total t-matrix correction is $\tilde{T} = \tilde{T}^{12} + \tilde{T}^{13} + \tilde{T}^{23}$. The energy calculated using the t-matrix is then $E_{\text{tm}} = E_0 + \tilde{T}$ and the exact energy from diagonalization is E_{exact} . $E_0^{ij} = \mathcal{E}(\mathbf{q}_i) + \mathcal{E}(\mathbf{q}_j)$, is the noninteracting energy of the (i, j) pair.

Q_0^{ij}	\mathbf{P}_{ij}	E_0^{ij}	\tilde{T}^{ij}
$(0,0)(0,1)$	$(0,1)$	-7.532088886	0.041949215
$(0,0)(0,-1)$	$(0,-1)$	-7.532088886	0.041949215
$(0,1)(0,-1)$	$(0,0)$	-7.064177772	0.118684581
Column sum			$\tilde{T} = 0.202583012$
Noninteracting total		$E_0 = -11.064177772$	
T-matrix total		$E_{\text{tm}} = -10.861594761$	
Exact total		$E_{\text{exact}} = -10.871031687$	

if we visualize this approximation via a path integral or a Feynman diagram, the self-consistent formula would mean that other pairs may be scattering simultaneously with pair (ij) , yet we did not take into account that the other pairs’ fluctuations would modify the set of sites \bar{Q}^{ij} accessible to this pair. In any case, analogous to the two-particle t-matrix (Sect. 3), we could solve equation (4.4) iteratively setting $E_{n+1}^{ij} = E_0^{ij} + \tilde{T}^{ij}(E_n^{ij})$, until successive iterates agree within a tolerance that we chose to be 10^{-15} , which happened after some tens iterations.

Using the same procedure, we can also calculate the t-matrix energies for the nondegenerate excited states of the $M = 3$ system in Figure 9: the $(-1, 0)(0, 0)(1, 0)$ and $(0, 2)(0, 0)(0, -2)$ states. The results are shown in Table 4. Figure 11 shows graphically the noninteracting energy levels, the t-matrix energies for the three nondegenerate states, and the exact energies from diagonalization, and the arrows link the noninteracting energies E_0 with the t-matrix results $E_{\text{tm}} = E_0 + \tilde{T}$. The agreement between E_{tm} and E_{exact} is good.

4.4 A five-fermion t-matrix calculation

We now consider briefly a $M = 5$ calculation, again for the 8×9 lattice. The noninteracting ground state is unique, with momentum vectors $\mathbf{q}_1 = (0, 0)$, $\mathbf{q}_2 = (0, 1)$, $\mathbf{q}_3 = (0, -1)$, $\mathbf{q}_4 = (1, 0)$, and $\mathbf{q}_5 = (-1, 0)$. In Figure 12 we show the excluded set \bar{Q}^{24} of the t-matrix computation

Table 4. Lowest 15 noninteracting, exact, and t-matrix energies for 8×9 lattice with $M = 3$ and $\mathbf{P} = (0, 0)$.

E_0	E_{exact}	E_{tm}
-11.064178	-10.871031687	-10.861594761
-10.828427	-10.608797838	-10.595561613
-9.892605	-9.672121352	
-9.892605	-9.519017636	
-9.892605	-9.497189108	
-9.892605	-9.462304364	
-9.892605	-9.398345108	
-9.892605	-9.345976806	
-8.694593	-8.252919763	-8.210179503
-8.239901	-8.015024904	
-8.239901	-7.946278078	
-8.239901	-7.809576487	
-8.239901	-7.800570818	
-8.239901	-7.690625772	
-8.239901	-7.615399722	




Fig. 11. Noninteracting, t-matrix, and exact energies of the three-particle fermion system on the 8×9 lattice with $\mathbf{P} = (0, 0)$. The bracketed numbers refer to the degeneracies of the level (see Fig. 9). The arrows associate the noninteracting states with the t-matrix results. We have worked on nondegenerate noninteracting states so far.

for the pair $(\mathbf{q}_2, \mathbf{q}_4)$. The momentum vectors $(\mathbf{q}_1, \mathbf{q}_3, \mathbf{q}_5)$ filled with other fermions are excluded, of course; three more wavevectors are excluded since the other member of the pair would have to occupy one of $\mathbf{q}_1, \mathbf{q}_3,$ or \mathbf{q}_5 , due to conservation of the total momentum $\mathbf{P} = (1, 1)$. The t-matrix results for all 10 pairs are presented in Table 5.

One might think that the pair, $\mathbf{q}_2 = (0, 1)$ and $\mathbf{q}_4 = (1, 0)$, exhibits pair-exchange symmetry with $(0, 0)(1, 1)$, so that $N_0 = 4$ as in Sections 3.3.3 and 3.4.3. However, since




Fig. 12. Momentum space exclusions in t-matrix $M = 5$ calculation for momentum vectors $\mathbf{q}_2 = (0, 1)$ and $\mathbf{q}_4 = (1, 0)$ (dots without crosses). These two fermions are excluded from scattering into momenta from the set $\bar{Q}^{2,4}$ (marked by crosses). The ground state is shown, with occupied momenta $(0, 0)$, $(0, 1)$, $(0, -1)$, $(1, 0)$, and $(-1, 0)$ (solid dots).

Table 5. T-matrix calculation for the 8×9 lattice with five particles $(0, 0)$, $(0, 1)$, $(0, -1)$, $(1, 0)$, and $(-1, 0)$. The exclusions in \bar{Q} for the pair $(0, 1)(1, 0)$ are depicted in Figure 12.

Q_0^{ij}	E_0^{ij}	\tilde{T}^{ij}
$(0, 0) (0, 1)$	-7.532088886	0.045184994
$(0, 0) (0, -1)$	-7.532088886	0.045184994
$(0, 0) (1, 0)$	-7.414213562	0.056898969
$(0, 0) (-1, 0)$	-7.414213562	0.056898969
$(0, 1) (0, -1)$	-7.064177772	0.118684581
$(0, 1) (1, 0)$	-6.946302449	0.081095408
$(0, 1) (-1, 0)$	-6.946302449	0.081095408
$(0, -1) (1, 0)$	-6.946302449	0.081095408
$(0, -1) (-1, 0)$	-6.946302449	0.081095408
$(1, 0) (-1, 0)$	-6.828427125	0.131405343
$E_0 = -17.892604897$		$\tilde{T} = 0.778639481$
$E_{\text{exact}} = -17.145715214$		$E_{\text{tm}} = -17.113965417$

$(0, 0)$ is occupied, the $(0, 1)(1, 0)$ pair cannot scatter into the $(0, 0)(1, 1)$ pair: hence $(0, 1)(1, 0)$ is a generic pair with $N_0^{2,4} = 2$. In general, if a pair is ever free to scatter into a degenerate pair state with a different occupation, that must be part of a many-particle state degenerate with the original one. Thus, the complicated t-matrix pairs with $N_0^{ij} > 2$ can arise in a many-fermion calculation only when the noninteracting many-fermion states are themselves degenerate.

4.5 Degenerate states

In the ground state examples considered up to now (Sects. 4.3 and 4.4), the noninteracting states were all nondegenerate. Let us now study a *degenerate* state in the third lowest level (six-fold degenerate) of $M = 3$ fermions on the 8×9 lattice: $\mathbf{q}_2 = (0, 1)$, $\mathbf{q}_3 = (1, 0)$, $\mathbf{q}_4 = (-1, -1)$. (See Fig. 9, row 3.) In this state, the pair $[\mathbf{q}_2 = (0, 1), \mathbf{q}_3 = (1, 0)]$ has the same total energy and momentum as the pair $[\mathbf{q}_1 = (0, 0), \mathbf{q}_5 = (1, 1)]$, on account of the pair component exchange symmetry (see Sect. 3.1); consequently $[\mathbf{q}_2, \mathbf{q}_3]$ can be scattered into $[\mathbf{q}_1, \mathbf{q}_5]$ contrary to the previous example in Section 4.4. Indeed, each of the six basis states in row 3 of Figure 9

is connected to the next one by a two-body component exchange symmetry.

Following the two-fermion calculation with $N_0 = 4$ pairs (see Sect. 3.3.3), the degenerate pairs $[\mathbf{q}_2, \mathbf{q}_3]$ and $[\mathbf{q}_1, \mathbf{q}_5]$ must be handled in the same set Q_0^{23} . The results equations (3.23) and (3.26) imply

$$\begin{aligned} \mathbf{T}^{23}(E^{23})|\mathbf{q}_2\mathbf{q}_3\rangle &= \frac{1}{2}(T_{1,-1,1,-1} + T_{1,-1,-1,1})|\mathbf{q}_2\mathbf{q}_3\rangle \\ &+ \frac{1}{2}(T_{1,-1,1,-1} - T_{1,-1,-1,1})|\mathbf{q}_1\mathbf{q}_5\rangle. \end{aligned} \quad (4.5)$$

Here $T_{1,-1,1,-1}$ and $T_{1,-1,-1,1}$ depend implicitly on $\mathbf{P} = (0,0)$, on the momenta, and on the energy E^{23} , as well as on \bar{Q}^{23} which depends on the occupation (\mathbf{q}_4) of the third fermion. In this notation, each \mathbf{T}^{ij} acting on any state produces two terms as in equation (4.5). The total t-matrix correction Hamiltonian is $\sum_{ij} \mathbf{T}^{ij}$, summed over all 18 possible pairs appearing in the degenerate noninteracting states. When we apply this to each state in the third row of Figure 9, we finally obtain a 6×6 matrix mixing these states. Diagonalization of this matrix would give the correct t-matrix corrections (and eigenstates) for this “multiplet” of six states. We have not carried out such a calculation.

It is amusing to briefly consider the states in row 5 of Figure 9, a different sixfold degenerate set. Unlike the row 3 case, these states separate into two subsets of three states, of which one subset has $q_y = \{-2, +1, +1\}$ and the other subset has the opposite $\{q_y\}$ components. Scatterings cannot mix these subsets, so the 6×6 matrix breaks up into two identical 3×3 blocks. Hence the t-matrix energies from row 5 consist of three twofold degenerate levels. By comparison, the *exact* interacting energies derived from these noninteracting states come in three *nearly* degenerate pairs, such that the intra-pair splitting is much smaller than the (already small) splitting due to the t-matrix.

4.6 Errors of the t-matrix

How good are the t-matrix results? From our example calculations on the 8×9 lattice, in Tables 3, 4, and 5, we see that E_{tm} and E_{exact} are close.

In Figure 13 we plot the noninteracting, t-matrix, and exact energies for $M = 3$, $\mathbf{P} = (0,0)$ ground state on a series of near square lattices $L \times (L+1)$. The noninteracting ground state momentum vectors are $(0,0)(0,1)(0,-1)$ for this series of lattices. We do not plot for $L > 12$, because, as can be seen in the bottom graph, the t-matrix energy E_{tm} approaches the exact energy E_{exact} rapidly. The error of the t-matrix result, $E_{\text{tm}} - E_{\text{exact}}$, decays very fast as the size of the lattice increases – very roughly as the L^{-6} power. Even at $L = 6$, i.e. at a density $n \approx 0.05$, the t-matrix approximation captures 95% of the interaction energy $E_{\text{exact}} - E_0$. These figures are based on using the bare energies in $\tilde{T}^{ij}(E_0^{ij})$ in equation (4.3). If we carried out the self-consistent calculation described in Section 4.3, the error $E_{\text{tm}} - E_{\text{exact}}$ would be smaller by a factor of roughly 2.5.




Fig. 13. Noninteracting, t-matrix, and exact energies for $M = 3$, $\mathbf{P} = (0,0)$ ground state $(0,0)(0,1)(0,-1)$ on a series of $L \times (L+1)$ lattices as a function of L .

5 The dilute limit: energy curves

In this section, we are interested in the functional form of the energy as a function of particle density for both bosons and fermions in the dilute limit. In the three-dimensional case, the problem of dilute quantum gases with strong, repulsive, short-range interactions was first addressed in the language of diagrammatic field theory by Galitskii [51] for fermions and Beliaev [52] for bosons. At that time, the ground state energy as an expansion in the particle density was also obtained for hard-sphere fermion and boson gases by Yang and collaborators [53] using a pseudopotential method. The field theoretical methods were later adapted to two dimensions in particular by Schick [10] for hard-disk bosons and by Bloom [41] for hard-disk fermions. Some other relevant analytic papers using a t-matrix approach for the Hubbard model were discussed in Section 1.2: Kanamori [42] and Mattis [25] in $d = 3$ and Rudin and Mattis [44] for $d = 2$.

For both hard-disk fermions and bosons in two dimensions, the leading-order correction to the noninteracting energy is found to be in the form of $n/\ln n$, where n is particle density. Expansions with second-order coefficients different from the results of Schick and Bloom were found in references [54,55] for the boson case and in references [46,58] for the fermion case. There is no consensus at this time on the correct second-order coefficient for both the boson and fermion problems.

Recently, reference [56] has proved rigorously the leading-order expansion of the two-dimensional dilute boson gas found by Schick [10]. Numerically, the dilute boson problem on a two-dimensional lattice has been studied using quantum Monte Carlo in references [11,12], and they obtain good fit with Schick’s result. As we mentioned in Section 1.2, more recently, because of a question regarding the validity of the Fermi liquid theory in two dimensions Bloom’s calculation [41] has received renewed attention [46,48], but this result has not been checked by numerical studies.

5.1 Dilute bosons

For two-dimensional hard disk bosons, the energy per particle E/M at the low-density limit from diagrammatic calculations is obtained (in the spirit of Ref. [52]) by Schick [10]

$$\frac{E}{M} = \frac{2\pi\hbar^2}{m} \frac{n}{|\ln(na^2)|} \left(1 + \mathcal{O}\left(\frac{1}{\ln(na^2)}\right) \right), \quad (5.1)$$

where $n = M/N$ is particle density, m the mass of the boson, and a the two-dimensional scattering length. As mentioned above, the coefficient of the second-order term, has not been settled (a recent result is Ref. [57]).

This hard-disk calculation was carried out using the kinetic energy $\hbar^2 k^2/2m$. In our lattice model, our hopping energy dispersion is (Eq. (2.11))

$$\mathcal{E}(\mathbf{k}) = -2t(\cos(k_x) + \cos(k_y)) \approx -4t + tk^2, \quad (5.2)$$

where we have Taylor-expanded the dispersion function near $\mathbf{k} = 0$ because in the dilute limit, at the ground state, the particles occupy momentum vectors close to zero. Therefore if we use $t = \hbar = 1$ and the effective mass m^* such that we have the form $\hbar^2 k^2/2m^*$, then $m^* = 1/2$ for our system. So for our model, Schick's expansion equation (5.1) should become,

$$\frac{E}{M} = -4 + \frac{4\pi n}{|\ln(na^{*2})|} \left(1 + \mathcal{O}\left(\frac{1}{\ln(na^{*2})}\right) \right), \quad (5.3)$$

where we have used a^* to denote the scattering length in our lattice system. There is no straightforward correspondence between Schick's scattering length a in the continuum and our a^* on the lattice. With infinite nearest-neighbor repulsion, the closest distance that our particles can come to is $\sqrt{2}$. We expect roughly $1 < a^* < \sqrt{2}$, and will determine a more precise value from curve fitting.

In Figure 14 we show the boson energy per particle (E/M) versus particle per site (M/N) curve for ten lattices, ranging from 25 sites to 42 sites, with three or more particles ($M \geq 3$). The data from all these lattices collapse onto one curve, especially in the low-density limit.

Equation (5.3), Schick's result applied to our model, suggests the following leading order fitting form for E/M versus n at the low-density limit,

$$\frac{E/M + 4}{4\pi n} = A + \frac{B}{|\ln(na^{*2})|}. \quad (5.4)$$

That is to say, if we plot $(E/M + 4)/(4\pi n)$ versus $1/|\ln(na^{*2})|$, then, if Schick is correct, we should get a straight line, with intercept $A = 0$ and slope $B = 1$, with one adjustable parameter a^* .

In Figure 15, we plot $(E/M + 4)/(4\pi n)$ versus $1/|\ln(na^{*2})|$ for the low-density limit ($n \leq 0.15$) for three choices of $a^* = 1.0, 1.36, \sqrt{2}$. The data points appear to lie on straight lines. For $a^* = 1.36$ the fitted intercept is $A = -0.016$ and the slope $B = 0.959$. In Table 6 we show the fitted slope and intercept for a number of a^*




Fig. 14. Boson energy per particle E/M versus particle density M/N data for ten lattices and $M \geq 3$. Data from different lattices collapse onto one curve. The solid line corresponds to the fitting function $-4 + 4\pi n(A + B/|\ln(na^{*2})|)$ with $a^* = 1.36$, $A = -0.016$, and $B = 0.959$, which is equation (5.4) with parameters from Table 6.




Fig. 15. $(E/M + 4)/(4\pi n)$ versus $1/|\ln(na^{*2})|$ plot to check Schick's formula for two-dimensional dilute bosons (Eq. (5.3)). The data points are for $M \geq 3$ and $n \leq 0.15$ from those in Figure 14, for lattices from 5×5 to 6×7 . For the three a^* values, the $a^* = 1.36$ choice gives $A = -0.016 \approx 0$ and $B = 0.959 \approx 1$.

choices. The slope is zero close to $a^* = 1.34$ and the intercept is zero close to $a^* = 1.39$. Our data thus suggest $a^* = 1.36 \pm 0.03$.

In Figure 14, the solid line is the function $-4 + 4\pi n(A + B/|\ln(na^{*2})|)$ using $a^* = 1.36$, $A = -0.016$, and $B = 0.959$, and we obtain a good fit up to $n = 0.15$.

For bosons, quantum Monte Carlo can be used to obtain zero temperature energies for reasonably large systems. For a dilute boson gas on a square lattice with on-site hardcore but not nearest-neighbor interaction,

Table 6. Intercept A and slope B in linear fitting $(E/M + 4)/(4\pi n)$ versus $1/|\ln(na^*)|$ for bosons, using equation (5.4). The slope is one close to $a^* = 1.33$ and the intercept is zero close to $a^* = 1.39$. So we get $a^* = 1.36 \pm 0.03$. The fitting for three choices of a^* is plotted in Figure 15.

a^*	B	A	a^*	B	A	a^*	B	A
1.00	1.855	-0.251	1.32	1.033	-0.039	1.37	0.941	-0.0099
1.10	1.547	-0.178	1.33	1.014	-0.033	1.38	0.923	-0.0043
1.20	1.289	-0.112	1.34	0.995	-0.027	1.39	0.906	0.0013
1.30	1.072	-0.050	1.35	0.977	-0.021	1.40	0.889	0.0069
1.31	1.053	-0.044	1.36	0.959	-0.016	1.414	0.865	0.015

reference [11] has fitted the first term of Schick's formula equation (5.1), and reference [12] has used higher-order terms and included the fitting of the chemical potential also. The agreement is good in both studies.

5.2 Dilute fermions

For fermions, it customary to write the energy per particle expansion in terms of the Fermi wavevector k_F . For two-dimensional dilute hard disk fermions with a general spin s , the energy per particle from diagrammatic calculations, is obtained by Bloom [41]

$$\frac{E}{M} = \frac{\hbar^2 k_F^2}{4m} \left(1 + 2s \frac{1}{|\ln(k_F a)|} + \mathcal{O} \left(\frac{1}{\ln(k_F a)} \right)^2 \right), \quad (5.5)$$

(see Ref. [41] for the spin-1/2 calculation and Ref. [58] for general s).

Equation (5.5) means that for our spinless fermions ($s = 0$), the leading order correction to the noninteracting energy in equation (5.5) is zero, which is due to the fact that equation (5.5) is derived for s-wave scattering. In our model, without spin, only antisymmetric spatial wavefunctions are allowed for fermions, and therefore the leading-order correction to the noninteracting energy should be from p-wave scattering. Reference [40] contains a formula for p-wave scattering in three dimensions where the leading-order correction to $E - E_0$ is proportional to $(k_F a)^3$ while the s-wave correction is proportional to $k_F a$. We are not aware of a two-dimensional p-wave calculation in the literature [59], and we have not worked out this p-wave problem in two dimensions. We expect that the p-wave contribution to energy should be considerably smaller than that from the s-wave term. In Section 2.4, we have considered the case of a few fermions on a large $L \times L$ lattice, and in Figure 1 we have studied the interaction correction to the noninteracting energy ΔE . It was shown there that ΔE for our spinless fermions is much smaller than that for bosons.

Using $k_F^2 = 4\pi n/(2s + 1)$, we can rewrite equation (5.5) as

$$\frac{E}{M} = \frac{\pi \hbar^2 n}{(2s + 1)m} \left(1 + 4s \frac{1}{|\ln(na^2)|} + \mathcal{O} \left(\frac{1}{\ln(na^2)} \right)^2 \right). \quad (5.6)$$

In this form, it is revealed that the second term of equation (5.6) is identical to the first term of the boson

expression equation (5.1), apart from the replacement $n \rightarrow 2sn/(2s + 1)$. In other words, the dominant interaction term for spinfull fermions is *identical* to the term for bosons, provided we replace n by the density of all spin species but one, i.e. of the spin species which can s-wave scatter off a given test particle.

6 Conclusion

We have studied a two-dimensional model of strongly-interacting fermions and bosons. This model is the simplest model of correlated electrons. It is very difficult to study two-dimensional quantum models with short-range kinetic and potential terms and strong interaction. There are very few reliable analytical methods, and many numerical methods are not satisfactory. With our simplified model of spinless fermions and infinite nearest-neighbor repulsion, we can use exact diagonalization to study systems much larger (in lattice size) than that can be done with the Hubbard model, and we have checked our numerical results with analytical results obtained from using lattice Green function, t-matrix, and field theory (Schick's result [10]).

It is somewhat puzzling that with the essential role the t-matrix plays in almost every calculation in the dilute limit with strong interactions, no systematic study of the t-matrix for a lattice model has been made, as far as we know. We believe that our work on the two-particle t-matrix and the few-fermion t-matrix is the first such study. Some approximations that are routinely made in t-matrix calculations are graphically presented, especially the use of first t-matrix iteration in calculating fermion energy. And we demonstrate the qualitative difference between the boson and fermion t-matrices. We believe that this study is a solid step in understanding dilute fermions in two dimensions, and is of close relevance to the 2D Fermi liquid question.

The dilute boson and fermion energy per particle curves were studied in Section 5. The boson curve was fitted nicely with a previous diagrammatic calculation, and our work on dilute bosons complements quantum Monte Carlo results [12]. For the fermion problem in our model, the leading order contribution to energy is from p-wave scattering; therefore, the series of results based on s-wave calculations by Bloom [41], Bruch [58], and Engelbrecht et al. [46] are not directly applicable.

Our model of spinless fermions and hardcore bosons with infinite nearest-neighbor repulsion involves a

$$T_3(E, \mathbf{P}; \mathbf{q}, \mathbf{q}') = \frac{1}{N^2} \sum_{\mathbf{q}'', \mathbf{q}''' \in \bar{Q}} \frac{V(\mathbf{q} - \mathbf{q}'')V(\mathbf{q}'' - \mathbf{q}''')V(\mathbf{q}''' - \mathbf{q}')}{(E - \mathcal{E}(\mathbf{q}'') - \mathcal{E}(\mathbf{P} - \mathbf{q}''))(E - \mathcal{E}(\mathbf{q}''') - \mathcal{E}(\mathbf{P} - \mathbf{q}'''))} \quad (\text{A.8})$$




Fig. 16. The three figures represent perturbative terms involving $V(\mathbf{q} - \mathbf{q}')$, $T_2(E, \mathbf{P}; \mathbf{q}, \mathbf{q}')$ and $T_3(E, \mathbf{P}; \mathbf{q}, \mathbf{q}')$. The t-matrix, $T(E, \mathbf{P}; \mathbf{q}, \mathbf{q}')$, is the sum of all these terms, i.e., it is the sum of the ladder diagrams to infinite order.

significant reduction of the size of the Hilbert space as compared to the Hubbard model. This enables us to obtain exact diagonalization results for much larger lattices than that can be done with the Hubbard model, and this also enables us to check the various analytical results (Green function, t-matrix, diagrammatics) in the dilute limit with diagonalization for much larger systems than that has been done in previous works. This paper and a companion paper [14] on the dense limit are the first systematic study of the spinless fermion model in two dimensions. We hope that the comprehensiveness of this paper can not only draw more attention to this so far basically overlooked model but also serve as a guide for diagonalization and analytical studies in the dilute limit.

This work was supported by the National Science Foundation under grant DMR-0240953. We used the computer facilities of the Cornell Center for Materials Research (CCMR) under the NSF Materials Research Science and Engineering Centers (MRSEC) grant DMR-0079992. We thank G.S. Atwal for helpful discussions.

Appendix A: Physical meaning of $\mathcal{T}(E)$

In this section we give yet another derivation of the t-matrix which makes more explicit the physical meaning of $T(E, \mathbf{P}; \mathbf{q}, \mathbf{q}')$ equation (3.13).

Before we get into a lot of algebra, let us describe the physical idea. In scattering theory we know that the Born series is a perturbation series of the scattering amplitude in terms of the potential. In Figure 16 we show the first three terms graphically, where the first term, the first Born approximation, is particularly simple—it is the Fourier transform of the potential. We also know that when the potential is weak the first few terms are a good approximation to the scattering amplitude, but when the potential is strong, we need all terms. In this section, we will show that our t-matrix $T(E, \mathbf{P}, \mathbf{q}, \mathbf{q}')$ is the sum of all such two-body scattering terms.

We start with equation (2.15) which we copy here for convenience,

$$(E - \mathcal{E}(\mathbf{q}) - \mathcal{E}(\mathbf{P} - \mathbf{q}))g(\mathbf{q}) = \frac{1}{N} \sum_{\mathbf{q}'} V(\mathbf{q} - \mathbf{q}')g(\mathbf{q}'). \quad (\text{A.1})$$

For $\mathbf{q} \in Q_0$ we break up the sum over \mathbf{q}' into two terms and get,

$$(E - E_0)g(\mathbf{q}) = \frac{1}{N} \sum_{\mathbf{q}' \in Q_0} V(\mathbf{q} - \mathbf{q}')g(\mathbf{q}') + \frac{1}{N} \sum_{\mathbf{q}' \in \bar{Q}} V(\mathbf{q} - \mathbf{q}')g(\mathbf{q}'). \quad (\text{A.2})$$

For $\mathbf{q} \in \bar{Q}$ we can rewrite equation (A.1) to get,

$$g(\mathbf{q}') = \frac{1}{N} \sum_{\mathbf{q}''} \frac{V(\mathbf{q}' - \mathbf{q}'')}{E - \mathcal{E}(\mathbf{q}') - \mathcal{E}(\mathbf{P} - \mathbf{q}'')} g(\mathbf{q}''). \quad (\text{A.3})$$

Plug equation (A.3) into the second sum in equation (A.2) and rearrange terms, we get,

$$(E - E_0)g(\mathbf{q}) = \frac{1}{N} \sum_{\mathbf{q}' \in Q_0} V(\mathbf{q} - \mathbf{q}')g(\mathbf{q}') + \frac{1}{N} \sum_{\mathbf{q}''} T_2(E, \mathbf{P}; \mathbf{q}, \mathbf{q}'')g(\mathbf{q}''), \quad (\text{A.4})$$

where we have defined,

$$T_2(E, \mathbf{P}; \mathbf{q}, \mathbf{q}'') = \frac{1}{N} \sum_{\mathbf{q}' \in \bar{Q}} \frac{V(\mathbf{q} - \mathbf{q}')V(\mathbf{q}' - \mathbf{q}'')}{E - \mathcal{E}(\mathbf{q}') - \mathcal{E}(\mathbf{P} - \mathbf{q}')}. \quad (\text{A.5})$$

Now break the sum over \mathbf{q}'' in equation (A.4) into two parts, and we get

$$(E - E_0)g(\mathbf{q}) = \frac{1}{N} \sum_{\mathbf{q}' \in Q_0} V(\mathbf{q} - \mathbf{q}')g(\mathbf{q}') + \frac{1}{N} \sum_{\mathbf{q}' \in Q_0} T_2(E, \mathbf{P}; \mathbf{q}, \mathbf{q}')g(\mathbf{q}') + \frac{1}{N} \sum_{\mathbf{q}' \in \bar{Q}} T_2(E, \mathbf{P}; \mathbf{q}, \mathbf{q}')g(\mathbf{q}'). \quad (\text{A.6})$$

Plug equation (A.3) into the last term of equation (A.6) and we get

$$(E - E_0)g(\mathbf{q}) = \frac{1}{N} \sum_{\mathbf{q}' \in Q_0} V(\mathbf{q} - \mathbf{q}')g(\mathbf{q}') + \frac{1}{N} \sum_{\mathbf{q}' \in Q_0} T_2(E, \mathbf{P}; \mathbf{q}, \mathbf{q}')g(\mathbf{q}') + \frac{1}{N} \sum_{\mathbf{q}'} T_3(E, \mathbf{P}; \mathbf{q}, \mathbf{q}')g(\mathbf{q}'), \quad (\text{A.7})$$

where we have defined

See equation (A.8) above

$$(E - E_0)g(\mathbf{q}) = \sum_{\mathbf{q}' \in Q_0} \left[\frac{1}{N} \sum_{ij} e^{i\mathbf{q} \cdot \mathbf{R}_i} e^{-i\mathbf{q}' \cdot \mathbf{R}_j} V (\delta_{ij} + V \bar{G}_{ij} + V^2 (\bar{G}_{ij})^2 + \dots) \right]. \quad (\text{A.13})$$

Continue this process, we obtain

$$(E - E_0)g(\mathbf{q}) = \frac{1}{N} \sum_{\mathbf{q}' \in Q_0} (V(\mathbf{q} - \mathbf{q}') + T_2(E, \mathbf{P}; \mathbf{q}, \mathbf{q}') + T_3(E, \mathbf{P}; \mathbf{q}, \mathbf{q}') + \dots)g(\mathbf{q}'). \quad (\text{A.9})$$

What we have done here is the traditional perturbation theory using iteration. Equation (A.9) is the Born series for scattering amplitude. The first term $V(\mathbf{q} - \mathbf{q}')$, the Fourier transform of the potential $V(\mathbf{r})$, is the first Born approximation. The content of higher order terms T_2, T_3, \dots can be obtained from their definition. Equation (A.5) says that T_2 involves two scatterings under V , and equation (A.8) says that T_3 involves three scatterings. Thus the Born series equation (A.9) can be graphically depicted as the ladders in Figure 16 [60], and it involves multiple scatterings to all orders. Note that each term in the Born series is infinite for infinite potential V . Next we will show that summing all the terms in the series gives the t-matrix and the potential V cancels out, giving a finite value.

It is easy to check that

$$T_2(E, \mathbf{P}; \mathbf{q}, \mathbf{q}') = V^2 \sum_{ij} e^{i\mathbf{q} \cdot \mathbf{R}_i} e^{-i\mathbf{q}' \cdot \mathbf{R}_j} \bar{G}_{ij}(E, \mathbf{P}), \quad (\text{A.10})$$

where $\bar{G}_{ij}(E, \mathbf{P})$ is our good old Green function equation (3.5),

$$T_3(E, \mathbf{P}; \mathbf{q}, \mathbf{q}') = V^3 \sum_{ij} e^{i\mathbf{q} \cdot \mathbf{R}_i} e^{-i\mathbf{q}' \cdot \mathbf{R}_j} (\bar{G}(E, \mathbf{P}))_{ij}^2, \quad (\text{A.11})$$

and

$$T_n(E, \mathbf{P}; \mathbf{q}, \mathbf{q}') = V^n \sum_{ij} e^{i\mathbf{q} \cdot \mathbf{R}_i} e^{-i\mathbf{q}' \cdot \mathbf{R}_j} (\bar{G}(E, \mathbf{P}))_{ij}^{n-1}. \quad (\text{A.12})$$

Plug these results into equation (A.9), we get

See equation (A.13) above

Now we come to a formal step,

$$V(I + V\bar{G} + V^2(\bar{G})^2 + \dots) = V(I - V\bar{G}(E))^{-1}, \quad (\text{A.14})$$

and the interesting result is that the infinite potential V cancels out, giving a finite value $-\bar{G}(E)^{-1}$.

If we can do this formal sum, then we get from equation (A.13),

$$(E - E_0)g(\mathbf{q}) = \sum_{\mathbf{q}' \in Q_0} T(E, \mathbf{P}; \mathbf{q}, \mathbf{q}')g(\mathbf{q}'),$$

which is exactly our momentum space T-matrix equation (3.12) and $T(E, \mathbf{P}; \mathbf{q}, \mathbf{q}')$ is exactly our t-matrix equation (3.13).

References

1. E. Dagotto, Rev. Mod. Phys. **66**, 763 (1994)
2. E. Manousakis, Rev. Mod. Phys. **63**, 1 (1991)
3. E.H. Lieb, D.C. Mattis, *Mathematical Physics in One Dimension* (Academic, New York, 1966)
4. S.R. White, Phys. Rev. Lett. **69**, 2863 (1992); S.R. White, D.J. Scalapino, Phys. Rev. Lett. **80**, 1272 (1998)
5. Y. Nagaoka, Phys. Rev. **147**, 392 (1966)
6. R.J. Bursill, Phys. Rev. B **60**, 1643 (1999); C.J. Bolech, S.S. Kancharla, G. Kotliar, Phys. Rev. B **67**, 075110 (2003)
7. H.Q. Lin, J.E. Gubernatis, Comp. Phys. **7**, 400 (1993)
8. G. Fano, F. Ortolani, A. Parola, Phys. Rev. B **46**, 1048 (1992)
9. H. Bruus, J.-C. Angles d'Auriac, Phys. Rev. B **55**, 9142 (1997)
10. M. Schick, Phys. Rev. A **3**, 1067 (1971)
11. F. Becca, L. Capriotti, S. Sorella, A. Parola, Phys. Rev. B **62**, 15277 (2000)
12. G. Batrouni (private communication); K. Bernardet, G.G. Batrouni, J.-L. Meunier, G. Schmid, M. Troyer, A. Dorneich, Phys. Rev. B **65**, 104519 (2002)
13. Our exact diagonalization computer program can handle arbitrary periodic boundaries specified by two vectors on the square lattice and uses translation symmetry to reduce the matrix size. It is described in references [14, 15]
14. N.G. Zhang, C.L. Henley, Phys. Rev. B **68**, 014506 (2003)
15. N.G. Zhang, Ph.D. thesis, Cornell University (2002)
16. G. Zala, B.N. Narozhny, I.L. Aleiner, V.I. Fal'ko, Phys. Rev. B **69**, 075306 (2004)
17. D. Vollhardt, in *Proceedings of the Enrico Fermi School, Course CXXI*, edited by Broglia, Schrieffer (North-Holland, Amsterdam, 1994)
18. C.N. Yang, C.P. Yang, Phys. Rev. **150**, 321 and 327 (1966)
19. G.S. Uhrig, R. Vlaming, Phys. Rev. Lett. **71**, 271 (1993); G.S. Uhrig, R. Vlaming, Physica B **194-196**, 451 (1994); G.S. Uhrig, R. Vlaming, Physica B **206-207**, 694 (1995); G.S. Uhrig, R. Vlaming, Ann. Physik **4**, 778 (1995)
20. R. Shankar Rev. Mod. Phys. **66**, 129 (1994)
21. J.E. Gubernatis, D.J. Scalapino, R.L. Sugar, W.D. Toussaint, Phys. Rev. B **32**, 103 (1985)
22. J.-L. Pichard, G. Benenti, G. Katomeris, F. Selva, X. Waintal, *Exotic States in Quantum Nanostructures*, edited by S. Sarkar (Kluwer, Dordrecht, 2002). One could speculate that, if their model included our infinitely strong nearest-neighbor repulsion, the Hilbert space (identical to ours) would be substantially reduced, without very much error in the energies
23. C.L. Henley, N.G. Zhang, Phys. Rev. B **63**, 233107 (2001)
24. Note that with spinless fermions and hardcore bosons and infinite nearest-neighbor repulsion, the filling (particle per lattice site) in our model goes from 0 to 1/2 only
25. D.C. Mattis, *The Theory of Magnetism*, Vol. I (Springer, Berlin, 1981), p. 252
26. G. Baym, *Lectures on Quantum Mechanics*, Chap. 8, *Cooper Pairs* (Benjamin-Cummings, Reading, 1981)
27. S.-J. Dong, C.N. Yang, Rev. Math. Phys. **1**, 139 (1989)
28. F. Marsiglio, J.E. Hirsch, Physica C **171**, 554 (1990)

29. H.Q. Lin, Phys. Rev. B **44**, 4674 (1991)
30. A.G. Petukhov, J. Galan, J.A. Verges, Phys. Rev. B **46**, 6212 (1992)
31. A.S. Blaer, H.C. Ren, O. Tchernyshyov, Phys. Rev. B **55**, 6035 (1997)
32. S. Basu, R.J. Gooding, P.W. Leung, Phys. Rev. B **63**, 100506 (2001)
33. M. Fabrizio, A. Parola, E. Tosatti, Phys. Rev. B **44**, 1033 (1991)
34. We are slightly abusing the notation V . $V(\mathbf{r})$ is the potential function for all space. V is the value of the potential at four points ($V = V(\mathbf{r} = (1, 0))$ for example). And $V(\mathbf{k})$ the Fourier transform of $V(\mathbf{r})$. We will differentiate these meanings by including explicitly the argument, \mathbf{r} or \mathbf{k} . Without argument, we then mean $V = V(\mathbf{r} = (1, 0))$
35. The following derivation is indebted to reference [31] which solved a similar problem
36. Incidentally the interesting problem of computing the resistance between two points on a lattice resistor network turns out to be an application of the lattice Green functions. See J. Cserti, Am. J. Phys. **68**, 896 (2000)
37. S. Katsura, S. Inawashiro, J. Math. Phys. **12**, 1622 (1971)
38. T. Morita, J. Math. Phys. **12**, 1744 (1971)
39. L.D. Landau, E.M. Lifshitz, *Statistical Physics* Vol. I (Addison, Reading, 1969), p. 234
40. A.L. Fetter, J.D. Walecka, *Quantum Theory of Many-Particle Systems* (McGraw, New York, 1971)
41. P. Bloom, Phys. Rev. B **12**, 125 (1975)
42. J. Kanamori, Prog. Theor. Phys. **30**, 275 (1963)
43. K. Yosida, *Theory of Magnetism* (Springer, Berlin, 1998), p. 191
44. S. Rudin, D.C. Mattis, Phys. Lett. **110A**, 273 (1985)
45. M. Randeria, J.-M. Duan, L.-Y. Shieh, Phys. Rev. B **41**, 327 (1990)
46. J.R. Engelbrecht, M. Randeria, L. Zhang, Phys. Rev. B **45**, 10135 (1992)
47. J.R. Engelbrecht, M. Randeria, Phys. Rev. B **45**, 12419 (1992)
48. H. Fukuyama, O. Narikiyo, Y. Hasegawa, J. Phys. Soc. Jpn **60**, 372 (1991); H. Fukuyama, O. Narikiyo, Y. Hasegawa, J. Phys. Soc. Jpn **60**, 2013 (1991)
49. Note that, anisotropic hopping dispersion $\mathcal{E}(\mathbf{q}) = -2t_x \cos q_x - 2t_y \cos q_y$, where $\mathcal{E}_x(q) = -2t_x \cos q$ and $\mathcal{E}_y(q) = -2t_y \cos q$, also has this pair component exchange symmetry
50. N. Furukawa, M. Imada, J. Phys. Soc. Jpn **61**, 3331 (1992); M. Guerrero, G. Ortiz, J.E. Gubernatis, Phys. Rev. B **62**, 600 (2000)
51. V. Galitskii, Sov. Phys. JETP **7**, 104 (1958)
52. S. Beliaev, Sov. Phys. JETP **7**, 289 (1958)
53. K. Huang, C.N. Yang, Phys. Rev. **105**, 767 (1957); K. Huang, C.N. Yang, J.M. Luttinger, Phys. Rev. **105**, 776 (1957); T.D. Lee, C.N. Yang, Phys. Rev. **105**, 1119 (1957); T.D. Lee, K. Huang, C.N. Yang, Phys. Rev. **106**, 1135 (1957)
54. L.W. Bruch, Physica **93A**, 95 (1978)
55. D.F. Hines, N.E. Frankel, D.J. Mitchell, Phys. Lett. **68A**, 12 (1978)
56. E.H. Lieb, J. Yngvason, J. Stat. Phys. **103**, 509 (2001)
57. J.O. Andersen, Eur. Phys. J. B **28**, 389 (2002)
58. L.W. Bruch, Physica **94A**, 586 (1978); G.S. Atwal (personal communication) has also derived equation (5.5) for general s
59. Reference [58] discusses spin degeneracy, but does not contain a formula for p-wave terms
60. B.R. Holstein, *Topics in Advanced Quantum Mechanics* (Addison, Redwood, 1992), p. 74