

The heavy quark potential from Wilson's exact renormalization group

Ulrich Ellwanger¹, Manfred Hirsch^{2,3,a}, Axel Weber^{2,3,b}

¹ Laboratoire de Physique Théorique et Hautes Energies^c Université de Paris XI, bâtiment 211, F-91405 Orsay Cedex, France (e-mail: ellwange@qcd.th.u-psud.fr)

² Institut für Theoretische Physik, Universität Heidelberg, Philosophenweg 16, D-69120 Heidelberg, Germany

³ Laboratoire de Physique Théorique et Hautes Energies Université de Paris XI, bâtiment 211, F-91405 Orsay Cedex, France

Received: 28 February 1997 / Revised version: 12 May 1997

Abstract. We perform a calculation of the full momentum dependence of the gluon and ghost propagators in pure SU(3) Yang-Mills theory by integrating Wilson's exact renormalization group equations with respect to an infrared cutoff k . The heavy quark potential in the quenched approximation can be expressed in terms of these propagators. Our results strongly indicate a $1/p^4$ -behaviour of the heavy quark potential for $p^2 \rightarrow 0$. We show in general, that effective actions which satisfy Schwinger-Dyson equations, correspond to (quasi-) fixed points of Wilson's exact renormalization group equations.

1 Introduction

The computation of reliable phenomenological numbers on the basis of non-perturbative QCD is still an unsolved problem. Numerical methods seem to be required, as the simulation of the theory on a space-time lattice. A semi-phenomenological tool is provided by the Schwinger-Dyson equations (SDEs), which are reviewed in this respect in [1]. In the present paper we discuss an approach to non-perturbative phenomena in non-abelian gauge theories, which is based on the integration of exact renormalization group equations (ERGEs) [2] in continuum quantum field theory [3].

ERGEs describe the continuous evolution of effective Lagrangians, or effective actions, with a scale (or infrared cutoff) k , and allow to obtain the full quantum effective action for $k \rightarrow 0$ from a bare (microscopic) effective action at a cutoff scale $k = \Lambda$. In contrast to standard renormalization group equations they describe this evolution including all irrelevant couplings, or higher dimensional operators, and are exact in spite of the appearance of only one-loop diagrams. The price to pay is the fact that one has to deal with an infinite system of coupled differential

equations which for practical purposes requires some truncation. The types of approximations or systematic expansions, which can be employed, depend on the phenomena under consideration:

One can expand the effective Lagrangian (or Hamiltonian or action) in powers of momenta or derivatives, but keep all powers of the involved field. This kind of expansion within the ERGEs is appropriate for the calculation of effective potentials, which are required to study the existence and nature of phase transitions (see [4] for some early literature).

Alternatively, one can expand the effective action in powers of fields, keeping all powers of the momenta [5]. This allows, e.g., to study the formation of bound states by using ERGEs to look for the appearance of poles in four-point functions [6]. Finally, in order to describe dynamical symmetry breaking, composite fields can be introduced [7, 8] and different expansions can be applied to the parts of the effective action involving fundamental and composite fields, respectively.

The ansatz of the present paper is based on the expansion of the effective action in powers of fields, which allows to study the full momentum dependence of the (gluon and ghost) propagators. Within this expansion the knowledge of the running 3- and 4-point functions is required in order to integrate the ERGE for the running of the 2-point function, the knowledge of the 4- and 5-point functions is required for the ERGE of the 3-point function etc. [5, 6].

In the case of gauge theories this expansion gets to some extent reorganized: due to gauge invariance (or, more precisely, BRST invariance) parts of higher N-point functions (with certain Lorentz and gauge index structures) are completely determined by lower N-point functions.

^a Supported by a DAAD-fellowship HSP II financed by the German Federal Ministry for Research and Technology; e-mail: m.hirsch@thphys.uni-heidelberg.de

^b Supported by a LGFG-fellowship of the Land Baden-Württemberg and a DAAD-fellowship; current address: Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Circuito Exterior C.U., A.P. 70-543, 04510 México, D.F., Mexico; e-mail: axel@nuclecu.unam.mx

^c Laboratoire associé au Centre National de la Recherche Scientifique – URA D0063

The usual procedure consists in imposing Slavnov-Taylor identities (STIs) on the effective action, which can be solved for “dependent” parts of N-point functions in terms of “independent” ones. For instance, in the present case of a pure Yang-Mills theory a gauge invariant term in the effective action of the form

$$F_{\mu\nu}^a(f(D^2)F_{\mu\nu})^a \quad (1.1)$$

(where D denotes the covariant derivative and f an arbitrary function) describes simultaneously the “independent” effective gluon propagator or 2-point function, and certain “dependent” Lorentz and gauge index structures of the 3-, 4- and higher gluonic N-point functions. Additional “independent” structures of the gluonic 3- and higher N-point functions are described in terms of gauge invariant expressions involving at least three powers of the field strength $F_{\mu\nu}$.

The application of ERGEs to gauge theories faces some technical problems, since ERGEs are based on an intermediate infrared momentum cutoff k , and such a cutoff generally breaks gauge or BRST invariance [9–14]. One way to deal with these problems is the modification of the STIs [11, 13, 14]: the modified STIs impose “fine tuning conditions” on those couplings in the effective action at $k \neq 0$, which break gauge or BRST invariance. These “fine tuning conditions” are such that they guarantee the BRST invariance of the effective action for $k \rightarrow 0$.

As a consequence of the (modified) STIs, at any value of k , the “dependent” parts of the effective action can thus be obtained either via the modified STIs in terms of the “independent” parts at the scale k , or directly from the integration of the ERGEs. Below we will employ approximations which consist in neglecting contributions to the r.h. sides of the ERGEs. In the absence of approximations, both methods to obtain the dependent parts lead to identical results. In the presence of approximations one should determine the dependent parts of the effective action from the modified STIs. Approximations on the r.h. sides of ERGEs for independent parts of the effective action do thus not imply a violation of the modified STIs.

In the presence of approximations one can nevertheless study the ERGEs for the dependent parts of the effective action and see, whether these ERGEs deviate strongly from the corresponding equations obtained from the k -derivatives of the modified STIs. As emphasized in [14], this method allows for a numerical self-consistency check of the approximation and will also be employed in the present paper.

As described in Sect. 3, our approximation consists in the neglect of the contribution of certain parts of the 3- and 4-point functions to the ERGEs of the 2-point functions. We take those parts of the 3- and 4-point functions into account, however, which are determined in terms of the 2-point functions through the standard part of the STIs. In the purely gluonic sector these are those obtained by an expansion of (1.1) in powers of the gauge field A_μ^a .

No approximation is imposed on the momentum dependence of the 2-point functions (the gluon and ghost propagators). These propagators by themselves are gauge

dependent and not directly observable quantities. A certain combination thereof, however, has a physical meaning: The potential between infinitely heavy quarks can be written as a product of the dressed quark-gluon vertices and the full gluon propagator, and the quark-gluon vertex function is given by the ghost propagator function. As a result the heavy quark potential depends on both propagator functions, which we calculate here in the quenched approximation.

The aim of this paper is to see, how much information on the heavy quark potential in quenched QCD can be obtained within the approximation we employ, which is seen to be self-consistent for a range of momenta p^2 down to a certain value $p^2 \sim k_0^2$ (see Sect. 5). We find strong evidence for a $1/p^4$ -behaviour within the trustworthy range of p^2 .

Both the method and the results can be compared with the Schwinger-Dyson formalism. To this end we derive in Sect. 2 a new and general relation between ERGEs and SDEs: we show that effective actions, which satisfy SDEs in the presence of an infrared cutoff k , are (quasi-) fixed points of the ERGEs. Thus a search for fixed points of the ERGEs for $k \rightarrow 0$ is equivalent to the search for solutions of SDEs.

Investigations of the gluon propagator have been performed in the context of SDEs both in the axial [15, 16] and Landau [17] gauge. In most cases solutions with a singular behaviour like $1/p^4$ have been found to be consistent. (A behaviour less singular than $1/p^2$ has been put forward on phenomenological grounds in [18] and claimed to exist in the axial gauge in [16], but recently such a form of the gluon propagator in the infrared has been argued to be inconsistent [19].) In the Landau gauge the ghost sector has been completely neglected; thus it is much less straightforward to relate these results to physical observables. In Sect. 6 we will discuss the comparison between the ERGE and SDE approach in more detail.

The paper is organized as follows: in Sect. 2 we discuss the derivation of the ERGEs, and their relation to SDEs. In Sect. 3 we present our truncated ansatz for the pure Yang-Mills action. In Sect. 4 we discuss the technical procedure, and in Sect. 5 our numerical results. Section 6 is devoted to conclusions and an outlook.

2 ERGEs and SDEs

The form of the ERGEs can most conveniently be derived assuming a path integral representation for the generating functional of the theory. To this end we restrict ourselves, for the time being, to the case of a single scalar field. It is straightforward to supplement the following equations with the appropriate minus signs, Lorentz and internal indices in the cases of fermions, vector fields and internal symmetries. To start with, the path integral formulation of the theory requires for its proper definition a regularization of the ultraviolet divergences. This will be implemented through a modification of the propagator, i.e. the term quadratic in the fields in the bare action. In addition

we introduce an infrared cutoff k ; the ERGEs will describe the variation of the generating functional with k , where the UV cutoff Λ will be held fixed. (In the framework of the ERGEs, in particular in the application to gauge theories, one can consistently set $\Lambda = \infty$, see below). The UV and IR regularized generating functional of connected Green functions $\mathcal{G}_k^A(J)$ can then be represented as

$$e^{-\mathcal{G}_k^A(J)} = \mathcal{N} \int \mathcal{D}\varphi e^{-\frac{1}{2}(\varphi, P_0^{-1}\varphi) - \Delta S_k^A - S_{int}(\varphi) + (J, \varphi)}, \quad (2.1)$$

where $S_{int}(\varphi)$ is independent of k . Here (J, φ) etc. is a short-hand notation for

$$(J, \varphi) \equiv \int \frac{d^4p}{(2\pi)^4} J(p)\varphi(-p) \quad . \quad (2.2)$$

P_0^{-1} denotes the bare inverse propagator (e.g. $P_0^{-1}(p^2) = p^2$ in the case of a massless scalar field), S_{int} the bare interaction, and

$$\Delta S_k^A = \frac{1}{2}(\varphi, R_k^A \varphi) \quad (2.3)$$

implements the UV and IR cutoffs. In the case of a massless field a convenient choice for R_k^A is

$$R_k^A(p^2) = p^2 \frac{1 - e^{-\frac{p^2}{\Lambda^2}} + e^{-\frac{p^2}{k^2}}}{e^{-\frac{p^2}{\Lambda^2}} - e^{-\frac{p^2}{k^2}}}, \quad (2.4)$$

such that the full propagator

$$P_k^A = (P_0^{-1} + R_k^A)^{-1} = \frac{e^{-\frac{p^2}{\Lambda^2}} - e^{-\frac{p^2}{k^2}}}{p^2} \quad (2.5)$$

has the desired properties: exponential decay in the UV, and finiteness in the IR.

The ERGE for \mathcal{G}_k^A describes its variation with k , at fixed Λ . It can be obtained by differentiating (2.1) with respect to k , and replacing φ under the path integral by variations with respect to the sources. One finds

$$\begin{aligned} \partial_k \mathcal{G}_k^A(J) &= -\frac{1}{2} \int \frac{d^4q}{(2\pi)^4} \partial_k R_k^A(q^2) \\ &\cdot \left\{ \frac{\delta^2 \mathcal{G}_k^A(J)}{\delta J(q) \delta J(-q)} - \frac{\delta \mathcal{G}_k^A(J)}{\delta J(q)} \frac{\delta \mathcal{G}_k^A(J)}{\delta J(-q)} \right\}. \end{aligned} \quad (2.6)$$

The effective action $\Gamma_k^A(\varphi)$ is defined by the Legendre transform of $\mathcal{G}_k^A(J)$,

$$\Gamma_k^A(\varphi) = \mathcal{G}_k^A(J) + (J, \varphi) \quad , \quad (2.7)$$

and it is convenient to define an effective action $\widehat{\Gamma}_k^A$ with the IR cutoff terms subtracted:

$$\widehat{\Gamma}_k^A = \Gamma_k^A - \Delta S_k^A \quad (2.8)$$

Inserting the Legendre transformation into (2.6) one obtains the ERGE for $\widehat{\Gamma}_k^A$:

$$\begin{aligned} \partial_k \widehat{\Gamma}_k^A(\varphi) &= \frac{1}{2} \int \frac{d^4q}{(2\pi)^4} \partial_k R_k^A(q^2) \\ &\cdot \left(\frac{\delta^2 \widehat{\Gamma}_k^A(\varphi)}{\delta \varphi(q) \delta \varphi(-q)} + R_k^A(q^2) \right)^{-1}. \end{aligned} \quad (2.9)$$

The boundary condition of $\widehat{\Gamma}_k^A(\varphi)$ for $k \rightarrow \Lambda$ can be obtained from a careful consideration of the limit $P_k^A \rightarrow 0$ for $k \rightarrow \Lambda$ and reads

$$\widehat{\Gamma}_\Lambda^A(\varphi) = \frac{1}{2}(\varphi, (P_0^A)^{-1}\varphi) + S_{int}(\varphi) \quad . \quad (2.10)$$

On the other hand, for $k \rightarrow 0$ $\widehat{\Gamma}_k^A(\varphi)$ becomes the full quantum effective action of a theory, which is defined by a bare interaction $S_{int}(\varphi)$ and a fixed UV cutoff Λ . Since the integration of the ERGEs allows us to relate effective actions $\widehat{\Gamma}_k^A$ with different k , it can be used to compute $\widehat{\Gamma}_0^A$ in terms of $\widehat{\Gamma}_\Lambda^A$ related to S_{int} by (2.10).

For most applications in particle physics this is actually all we want: there always exists a scale Λ (at least the Planck scale) beyond which we expect unknown new interactions or particles with masses larger than Λ . Physics beyond Λ , including quantum fluctuations involving momenta p^2 with $p^2 \geq \Lambda^2$, will certainly affect the form of S_{int} or the effective action $\widehat{\Gamma}_\Lambda^A$. Given an ansatz for $\widehat{\Gamma}_\Lambda^A$, however, we are generally interested in the effect of quantum fluctuations involving momenta with $0 \leq p^2 \leq \Lambda^2$. If we integrate the ERGE for $\widehat{\Gamma}_k^A$ from $k = \Lambda$ down to $k = 0$, these quantum fluctuations have been fully taken into account.

Next we turn to the relation between ERGEs and SDEs. To this end it is first convenient to write the ERGEs in terms of the generating functional (or partition function) $Z_k^A(J)$, where $Z_k^A(J)$ is given by

$$Z_k^A(J) = e^{-\mathcal{G}_k^A(J)} \quad (2.11)$$

and the r.h. side of (2.11) by the path integral (2.1). The ERGE for $Z_k^A(J)$ reads

$$\partial_k Z_k^A(J) = -\frac{1}{2} \int \frac{d^4q}{(2\pi)^4} \partial_k R_k^A(q^2) \frac{\delta^2 Z_k^A(J)}{\delta J(q) \delta J(-q)} \quad (2.12)$$

The full set of SDEs can also most conveniently be expressed in terms of $Z(J)$ [20]. They follow from the vanishing of the path integral over a total derivative with respect to φ . Subsequently we will consider the SDEs in the presence of an UV cutoff Λ and an IR cutoff k as in (2.1). In terms of $Z_k^A(J)$ they read

$$\begin{aligned} &\left(J(p) - (P_0^{-1}(p^2) + R_k^A(p^2)) \frac{\delta}{\delta J(-p)} \right. \\ &\quad \left. - \frac{\delta S_{int}(\varphi)}{\delta \varphi(-p)} \Big|_{\varphi = \frac{\delta}{\delta J}} \right) Z_k^A(J) = 0 \quad . \end{aligned} \quad (2.13)$$

After replacing Z_k^A by \mathcal{G}_k^A according to (2.11), and \mathcal{G}_k^A by Γ_k^A according to (2.7), one obtains the SDEs for the one-particle irreducible vertices, which are generally considered in the literature for $k = 0$ and $\Lambda = \infty$.

Let us now denote the entire l.h.side of (2.13) by $\Sigma_k^A(J)$. Next we evaluate the k -derivative of Σ_k^A . The k -derivative hits R_k^A and Z_k^A , and we have to use the ERGE (2.12). The result can be written in the form

$$\partial_k \Sigma_k^A = -\frac{1}{2} \int \frac{d^4 q}{(2\pi)^4} \partial_k R_k^A(q^2) \frac{\delta^2 \Sigma_k^A}{\delta J(q) \delta J(-q)}. \quad (2.14)$$

Thus we can make the following important statement: suppose that we managed to find a functional $Z_k^A(J)$, which satisfies the SDEs (2.13) for some value of k , identically in the sources J . Thus the corresponding quantity $\Sigma_k^A(J)$ vanishes. Now let us start to vary k . Since the r.h. side of (2.14) vanishes, $\Sigma_{k'}^A$ will still vanish for $k' \neq k$. (Here, of course, it has been used that Z_k^A varies with k according to the ERGE (2.12).) Thus we find that $Z_{k'}^A$ still satisfies the SDE (2.13), with k replaced by k' . In other words, if we found a functional Z_k^A which satisfies the SDE (2.13) for some value of k , and compute $Z_{k'}^A$ by integrating the ERGE (2.12), $Z_{k'}^A$ is guaranteed to satisfy the SDEs for all values for k' . The same statement will hold for the other generating functionals \mathcal{G}_k^A and Γ_k^A . Thus generating functionals which satisfy k -dependent SDEs can be named quasi-fixed points of the ERGEs ("quasi" because they still depend on k). In particular, for $k \rightarrow 0$, \mathcal{G}_0^A and Γ_0^A will satisfy the standard SDEs (without an infrared cutoff, but still an ultraviolet cutoff Λ) if such a quasi-fixed point is approached in the infrared.

Note that this property is independent of the form of S_{int} in (2.13): Any SDE with any form of S_{int} in (2.13) plays the role of a quasi-fixed point. (In particular, since we work in the presence of an UV cutoff Λ , S_{int} can contain arbitrary higher dimensional interactions suppressed by powers of Λ .)

These formal arguments let us suspect, that results obtained by the integration of the ERGEs can be quite similar to results obtained by solving SDEs, although the formal arguments no longer hold in a strict sense as soon as the effective actions are approximated or truncated in some way.

Let us note, for later use, that on the r.h. sides of the ERGEs (2.6) and (2.9) we can replace $R_k^A(p^2)$ by $R_k(p^2)$, where $R_k(p^2)$ is defined by

$$R_k(p^2) = R_k^{\Lambda=\infty}(p^2) \quad . \quad (2.15)$$

In the limit $\Lambda \rightarrow \infty$ the momentum integrations on the r.h. sides of (2.6) and (2.9) are still exponentially damped with the choice (2.4) for R_k^A . Accordingly we denote by \mathcal{G}_k and $\widehat{\Gamma}_k$ the generating functionals, which satisfy the ERGEs with R_k^A replaced by R_k . Of course we are still able to compute $\widehat{\Gamma}_{k=0}$ in terms of an ansatz for $\widehat{\Gamma}_{k=\Lambda}$. In physical terms UV divergences do not appear, since $\widehat{\Gamma}_\Lambda$ is already supposed to contain all quantum effects involving

momenta with $p^2 \geq \Lambda^2$. We just loose the possibility to formally equate $\widehat{\Gamma}_\Lambda$ with S_{int} appearing in a path integral of the form of (2.1), although a slightly different, more involved path integral representation for $\widehat{\Gamma}_k$ still exists [21]. From universality we expect in any case, that $\widehat{\Gamma}_0$ is independent of minor modifications of $\widehat{\Gamma}_\Lambda$, so one choice is a priori as good as the other. Thus, subsequently, we define a theory by the ERGEs (2.6) or (2.9), with R_k^A replaced by R_k , and a finite ansatz for $\widehat{\Gamma}_\Lambda$ (or \mathcal{G}_Λ) at a fixed scale Λ .

3 Yang-Mills theories

The application of ERGEs to gauge theories requires some treatment of the breaking of gauge invariance, which is introduced through the intermediate IR cutoff k . Here we use the method of imposing modified STIs [11, 13, 14] in the formulation presented in [13, 14]. In [14], in particular, ERGEs for SU(3) Yang-Mills theory have already been integrated numerically within an approximation, where only marginal and relevant couplings in the effective action have been taken into account.

In the present paper we extend the form of the effective action considerably, namely we allow for an arbitrary momentum dependence of the gluon and ghost propagators. Again we consider the SU(3) Yang-Mills theory in four-dimensional Euclidean space-time, where the classical action reads

$$S = \int d^4 x \left\{ \frac{1}{4} F_{\mu\nu}^a F_{\mu\nu}^a + \frac{1}{2\alpha} \partial_\mu A_\mu^a \partial_\nu A_\nu^a + \partial_\mu \bar{c}^a (\partial_\mu c^a + g f^a_{bc} A_\mu^b c^c) - K_\mu^a (\partial_\mu c^a + g f^a_{bc} A_\mu^b c^c) - L^a \frac{1}{2} g f^a_{bc} c^b c^c + \bar{L}^a \frac{1}{\alpha} \partial_\mu A_\mu^a \right\} \quad (3.1)$$

with

$$F_{\mu\nu}^a = \partial_\mu A_\nu^a - \partial_\nu A_\mu^a + g f^a_{bc} A_\mu^b A_\nu^c. \quad (3.2)$$

We have included the usual gauge fixing and ghost parts and coupled external sources to the BRST variations

$$\begin{aligned} \delta A_\mu^a &= (\partial_\mu c^a + g f^a_{bc} A_\mu^b c^c) \zeta \\ \delta c^a &= \frac{1}{2} g f^a_{bc} c^b c^c \zeta \\ \delta \bar{c}^a &= -\frac{1}{\alpha} \partial_\mu A_\mu^a \zeta, \end{aligned} \quad (3.3)$$

where ζ is a Grassmann parameter. The invariance of S at $\bar{L} = 0$ under BRST transformations can then be expressed as

$$\begin{aligned} 0 &= \int d^4 x \left\{ \delta A_\mu^a \frac{\delta S}{\delta A_\mu^a} + \delta c^a \frac{\delta S}{\delta c^a} + \delta \bar{c}^a \frac{\delta S}{\delta \bar{c}^a} \right\} \Bigg|_{\bar{L}=0} \\ &= \zeta \int d^4 x \left\{ \frac{\delta S}{\delta K_\mu^a} \frac{\delta S}{\delta A_\mu^a} - \frac{\delta S}{\delta L^a} \frac{\delta S}{\delta c^a} - \frac{\delta S}{\delta \bar{L}^a} \frac{\delta S}{\delta \bar{c}^a} \right\} \Bigg|_{\bar{L}=0}. \end{aligned} \quad (3.4)$$

IR cutoffs in the gluon and ghost propagators will be introduced by adding a term of the form

$$\Delta S_k = \frac{1}{2} (A_\mu^a, R_{k,\mu\nu} A_\nu^a) + (\bar{c}^a, \tilde{R}_k c^a) \quad (3.5)$$

to the action. Explicit expressions for $R_{k,\mu\nu}$ and \tilde{R}_k will be given below. The effective action

$$\Gamma_k(A, c, \bar{c}, K, L, \bar{L}) \equiv \hat{\Gamma}_k + \Delta S_k \quad (3.6)$$

satisfies an ERGE of the form [13, 14]

$$\partial_k \hat{\Gamma}_k = \int \frac{d^4 p}{(2\pi)^4} \left\{ \frac{1}{2} \partial_k R_{k,\mu\nu}(p^2) \left(\Gamma_k^{(2)} \right)_{A_\nu^a(-p), A_\mu^a(p)}^{-1} - \partial_k \tilde{R}_k(p^2) \left(\Gamma_k^{(2)} \right)_{-\bar{c}^a(-p), c^a(p)}^{-1} \right\}. \quad (3.7)$$

Here $\left(\Gamma_k^{(2)} \right)_{A_\mu^a(-p), A_\nu^b(q)}^{-1}$ is the $(A_\mu^a(-p), A_\nu^b(q))$ -component of the inverse of the matrix $\Gamma_k^{(2)} = \delta^2 \Gamma_k / \delta \bar{\varphi} \delta \varphi$ of second derivatives of Γ_k with respect to the fields $\bar{\varphi}^B = \{A_\mu^a(-p), -\bar{c}^a(-p), c^a(-p)\}$ and $\varphi^B = \{A_\mu^a(p), c^a(p), \bar{c}^a(p)\}$, where the index of $\bar{\varphi}$ and φ runs over the different fields, momenta, and Lorentz and gauge indices. For $k \rightarrow 0$, $\hat{\Gamma}_0$ has to satisfy the STI (3.4). To this end $\hat{\Gamma}_k$ has to satisfy, in particular at the starting point $k = \Lambda$, a modified STI of the form [13, 14]

$$\begin{aligned} & \int \frac{d^4 p}{(2\pi)^4} \left\{ \frac{\delta \hat{\Gamma}_k}{\delta K_\mu^a(-p)} \frac{\delta \hat{\Gamma}_k}{\delta A_\mu^a(p)} - \frac{\delta \hat{\Gamma}_k}{\delta L^a(-p)} \frac{\delta \hat{\Gamma}_k}{\delta c^a(p)} \right. \\ & \left. - \frac{\delta \hat{\Gamma}_k}{\delta \bar{L}^a(-p)} \frac{\delta \hat{\Gamma}_k}{\delta \bar{c}^a(p)} \right\} \Bigg|_{\bar{L}=0} \\ & = \int \frac{d^4 p}{(2\pi)^4} \sum_B \left[R_{k,\mu\nu}(p^2) \frac{\delta^2 \hat{\Gamma}_k}{\delta K_\nu^a(-p) \delta \varphi^B} \left(\Gamma_k^{(2)} \right)_{\bar{\varphi}^B, A_\mu^a(p)}^{-1} \right. \\ & \quad - \tilde{R}_k(p^2) \frac{\delta^2 \hat{\Gamma}_k}{\delta L^a(-p) \delta \varphi^B} \left(\Gamma_k^{(2)} \right)_{\bar{\varphi}^B, c^a(p)}^{-1} \\ & \quad \left. - \tilde{R}_k(p^2) \frac{\delta^2 \hat{\Gamma}_k}{\delta \bar{L}^a(-p) \delta \varphi^B} \left(\Gamma_k^{(2)} \right)_{\bar{\varphi}^B, \bar{c}^a(p)}^{-1} \right] \Bigg|_{\bar{L}=0}. \quad (3.8) \end{aligned}$$

The crucial point is that once (3.8) is satisfied for $k = \Lambda$, it will be satisfied by $\hat{\Gamma}_k$ for any $k < \Lambda$ provided $\hat{\Gamma}_k$ is obtained from $\hat{\Gamma}_\Lambda$ by integrating the ERGE. In particular $\hat{\Gamma}_{k=0}$ will satisfy the standard STI (3.4), if the IR cutoff functions $R_{k,\mu\nu}$ and \tilde{R}_k vanish identically for $k \rightarrow 0$.

Most importantly, $\hat{\Gamma}_k$ at the starting point $k = \Lambda$ can thus not be identified with the classical action (3.1), but it has to contain symmetry breaking terms specified by the need to satisfy (3.8). The origin of these symmetry breaking terms can be understood from the interpretation of $\hat{\Gamma}_\Lambda$ as a quantum effective action, where momenta p^2 with $p^2 \geq \Lambda^2$ have already been integrated out; the symmetry breaking induced by this ‘‘IR’’ cutoff Λ then generates the symmetry breaking terms in $\hat{\Gamma}_\Lambda$.

As stated above, our approach towards the integration of the ERGE (3.7) is based on an expansion in powers of the gluon and ghost fields. We are interested in the form of the gluon and ghost propagators, since a certain combination thereof (see below) determines the heavy quark potential in the quenched approximation. It is thus essential to allow for an arbitrary momentum dependence of the 2-point functions in our ansatz for the effective action.

Expanding the ERGE (3.7) in powers of fields one easily finds that the full 3-gluon, 4-gluon, ghost-gluon and 4-ghost vertices appear on the r.h. sides of the ERGEs for the gluon and ghost propagators. In the sequel we will refer to parts of the vertices with certain Lorentz and gauge index structures as ‘‘operators’’. One can distinguish two different kinds of such operators:

- a) ‘‘Independent’’ operators, which are not fixed in terms of the propagators or other operators by the STI. An example is the contribution to the 4-gluon vertex obtained from a term involving four powers of the field strength $F_{\mu\nu}^a$ in the effective action. (The r.h. side of the corresponding ERGE depends on the full 2-, 3-, 4-, 5- and 6-point functions.)
- b) ‘‘Dependent’’ operators, which are fixed by the modified STI in terms of the propagators and operators of type a). In general (3.8) yields a system of coupled nonlinear integral equations for the dependent operators. Examples are the operators corresponding to those parts of the 3- and 4-gluon vertices, which are obtained by an expansion of the expression $F_{\mu\nu}^a (f(D^2) F_{\mu\nu})^a$ in powers of the gauge field (see (1.1)), and a gluonic mass term (which we do not consider as part of the gluon propagator in this context).

Since our goal is the determination of the propagators, the only important operators for our purpose are those, that contribute significantly to the ERGEs for the 2-point functions. In a first approximation, we will neglect the contributions of the operators of type a), which are the ones not determined by the STI. Such an approximation is only expected to hold for a limited range of scales k , and can to some extent be checked a posteriori (see below). In any case, it turns the ERGE (3.7) into a closed system of integro-differential equations for the propagators, which can be integrated numerically.

Furthermore, with the simpler ansatz used in [14] it has proved to be permissible in a quantitative sense to determine the operators of type b) through the standard form of the STI (3.8) with a vanishing r.h. side instead of using the modified STI, with the important exception of the gluon mass term. Again, this may only be true within a certain range of k . We will therefore proceed in the same way here, i.e. we assume that we can approximate the operators of type b) by the corresponding ones obtained from the standard STI, as far as their quantitative contribution to the flow of the 2-point functions is concerned.

In practical terms, this assumption implies a tremendous technical simplification in the determination of these operators. For example, for the 3- and 4-gluon vertices we can use the expressions resulting from an expansion of $F_{\mu\nu}^a (f(D^2) F_{\mu\nu})^a$ in powers of A_μ^a (apart from a momen-

tum-dependent field renormalization, cf. the explicit expressions in the appendix). We consider it to be essential, however, to use the modified form of the STI to fix the gluon mass term, which is a truly relevant parameter of the effective action.

Let us stress that the approximation we employ for the operators of type b) does not imply that we are working with an effective action, which violates the modified STI. In principle we could, for every value of $k \neq 0$, determine the operators of type b) by solving the system of coupled non-linear integral equations given by (3.8), so that the resulting effective action would be in accord with the modified STI. What we actually do is to approximate the contributions of the operators of type b) to the flow of the 2-point functions by the contributions of the corresponding operators obtained from the standard STI, in the same way as we neglect the contributions of the operators of type a).

It is important to note, that there exists a non-trivial quantitative check for the self-consistency of the approximations employed. It is based on the consistency of the ERGE with the k -derivative of the modified STI in the absence of approximations, and will be discussed in more detail in the next section. In particular, this self-consistency condition will be used to determine the range of scales k , where we consider the approximations to be permissible.

The approximations employed here can be improved systematically by taking more and more operators of type a) into account, as well as by using the operators of type b) corresponding to a (partial or complete) solution of (3.8). The purpose of the present paper is to see, which results for the heavy quark potential will be obtained within the "minimal" approach described above.

Let us now write down those terms in the k -dependent effective action, which we take into account on the r.h. sides of the ERGEs for the 2-point functions. In a slight abuse of notation, we will denote this part of the effective action by $\widehat{\Gamma}_k$. Firstly, we include the gluon and ghost 2-point functions themselves, which are given by the momentum-dependent functions $f_{1,k}$ and $f_{2,k}$, respectively, and where we take a mass term for the gluons into account. Since we will consider the Landau gauge later on, it is not necessary to introduce a further momentum-dependent function in the longitudinal part of the gluon 2-point function here. Secondly, we describe the 3-gluon, 4-gluon and ghost-gluon vertices of the type b) above (determined from the standard STI), which we take into account on the r.h. sides of the ERGEs for the 2-point functions. Finally we include the external source terms containing K_μ^a , L^a and \bar{L}^a , which are required to formulate the STI. These latter terms are constrained by two further identities satisfied by $\widehat{\Gamma}_k$, which can be shown to be invariant under the RG flow [11, 13, 14]:

$$\partial_\mu \frac{\delta \widehat{\Gamma}_k}{\delta K_\mu^a} = \frac{\delta \widehat{\Gamma}_k}{\delta \bar{c}^a} \quad , \quad (3.9a)$$

$$\frac{\delta \widehat{\Gamma}_k}{\delta \bar{L}^a} = \frac{1}{\alpha} \partial_\mu A_\mu^a \quad . \quad (3.9b)$$

Explicitly, then, our ansatz for $\widehat{\Gamma}_k$ reads

$$\begin{aligned} \widehat{\Gamma}_k(A, c, \bar{c}, K, L, \bar{L}) &= \frac{1}{2} \int_{p_1, p_2} A_\mu^a(p_1) \left\{ (p_1^\mu \delta_{\mu\nu} - p_{1\mu} p_{1\nu}) f_{1,k}(p_1^2) \right. \\ &\quad \left. + \frac{p_{1\mu} p_{1\nu}}{\alpha} + m_k^2 \delta_{\mu\nu} \right\} A_\nu^a(p_2) \\ &\quad + i g_k f_{bc}^a \int_{p_1, p_2, p_3} f_{3, \mu\nu\rho}(p_1, p_2, p_3) \\ &\quad \times A_\mu^a(p_1) A_\nu^b(p_2) A_\rho^c(p_3) \\ &\quad + g_k^2 f_{bc}^a f_{de}^a \int_{p_1, \dots, p_4} f_{4, \mu\nu\rho\sigma}(p_1, p_2, p_3, p_4) \\ &\quad \times A_\mu^b(p_1) A_\nu^c(p_2) A_\rho^d(p_3) A_\sigma^e(p_4) \\ &\quad + \int_{p_1, p_2} \bar{c}^a(p_1) p_1^\mu f_{2,k}(p_1^2) c^a(p_2) \\ &\quad + i g_k f_{bc}^a \int_{p_1, p_2, p_3} \bar{c}^a(p_1) p_{1,\mu} \\ &\quad \times f_{2,k}(p_1^2) f_{2,k}^{-1}(p_2^2) A_\mu^b(p_2) c^c(p_3) \\ &\quad + i \int_{p_1, p_2} K_\mu^a(p_1) p_{1\mu} f_{2,k}(p_1^2) c^a(p_2) \\ &\quad - g_k f_{bc}^a \int_{p_1, p_2, p_3} K_\mu^a(p_1) f_{2,k}(p_1^2) f_{2,k}^{-1}(p_2^2) A_\mu^b(p_2) c^c(p_3) \\ &\quad - \frac{g_k}{2} f_{bc}^a \int_{p_1, p_2, p_3} L^a(p_1) c^b(p_2) c^c(p_3) \\ &\quad - \frac{i}{\alpha} \int_{p_1, p_2} \bar{L}^a(p_1) p_{1\mu} A_\mu^a(p_2) \quad , \end{aligned} \quad (3.10)$$

where

$$\int_{p_1, \dots, p_n} \equiv \int \prod_{i=1}^n \left(\frac{d^4 p_i}{(2\pi)^4} \right) \cdot (2\pi)^4 \delta^4 \left(\sum_{i=1}^n p_i \right). \quad (3.11)$$

The k -dependent 3-gluon and 4-gluon vertex functions $f_{3, \mu\nu\rho}$ and $f_{4, \mu\nu\rho\sigma}$ are given in terms of the gluon propagator function $f_{1,k}$ and the ghost propagator function $f_{2,k}$ in the appendix.

In order to establish the connection with the heavy quark potential, we introduce quark fields in the effective action. In the limit where the quark mass tends to infinity, all quantum corrections corresponding to diagrams which contain inner quark lines, are naively suppressed by powers of the quark mass. On the other hand, the STI requires a momentum dependent quark-gluon vertex function, where the momentum dependence involves the function $f_{2,k}$ above. (Within the standard approach, the origin of this non-trivial contribution is due to UV divergent diagrams, which are not suppressed by the heavy quark mass.) After an eventual redefinition of the quark fields, such that their kinetic term is properly normalized, the standard STI constrains the quark-gluon coupling in the effective action to be of the form

$$- g_k \lambda_{AB}^a \int_{p_1, p_2, p_3} \bar{\psi}^A(p_1) \gamma_\mu f_{2,k}^{-1}(p_2^2) A_\mu^a(p_2) \psi^B(p_3) \quad , \quad (3.12)$$

where λ denotes the SU(3) generators in the fundamental representation. Here we have implicitly used the Kc -coupling from (3.10) and the fact that the quantum corrections to the vertices, which couple external sources to the BRST-variations of the quark fields, are suppressed by powers of the quark mass in the Landau gauge.

Let us now consider the 2-point function V_k of the quark currents J_μ^a , given by

$$J_\mu^a = \lambda_{AB}^a \bar{\psi}^A \gamma_\mu \psi^B \quad . \quad (3.13)$$

In the heavy quark limit, the only contribution to V_k comes from the dressed one-gluon exchange diagram, so from (3.12) and the form of the gluon propagator we obtain the current 2-point function

$$V_k(p^2) = \frac{g_k^2}{p^2 f_{1,k}(p^2) f_{2,k}^2(p^2)} \quad . \quad (3.14)$$

We have omitted the gluon mass term m_k^2 in our definition of $V_k(p^2)$. In principle we could have replaced the factor $p^2 f_{1,k}(p^2)$ in the denominator of $V_k(p^2)$, which is due to the gluon propagator, by the factor involving m_k^2 and the infrared cutoff term. However, once the integration of the ERGE is pursued down to $k = 0$, the modified STI (3.8) turns into the standard STI and enforces $m_{k=0}^2 = 0$, and the infrared cutoff term vanishes identically. Using (3.14) to define $V_k(p^2)$, these effects have been anticipated, and $V_k(p^2)$ corresponds more closely to the physical function $V_0(p^2)$ already at finite k .

In the heavy quark limit $V_k(p^2)$ can be identified, up to trivial factors, with the Fourier transform of the potential between quarks in quenched QCD. Although we will compute the gluon and ghost propagator functions $f_{1,k}$ and $f_{2,k}$ individually by integrating the ERGEs, it is only the combination appearing in (3.14), which has a physical meaning.

A priori it may seem that the functions $f_{1,k}$ and $f_{2,k}$ can be changed at will by redefining the gluon, ghost and anti-ghost fields in the form of multiplication with momentum dependent functions. We have checked explicitly, however, that the STI restricts the relations between the vertices and the two-point functions such that finally all possible field redefinitions cancel in the expression for $V_k(p^2)$.

As noted above and in [11, 13, 14], m_k^2 in $\widehat{\Gamma}_k$ is fixed in terms of the other parameters in $\widehat{\Gamma}_k$ by the non-vanishing r.h. side of (3.8), where we use our ansatz (3.10) for $\widehat{\Gamma}_k$. After a consideration of terms $\sim A \cdot c$ of (3.8), one arrives at a non-linear relation of the form

$$m_k^2 = \frac{g_k^2}{(4\pi)^2} k^2 \text{ST}(k^2, m_k^2, f_{1,k}, f_{2,k}) \quad , \quad (3.15)$$

where we have anticipated the Landau gauge and omitted the gauge parameter α on the r.h. side. The expression $\text{ST}(k^2, \dots)$ involves momentum integrations over a number of one-loop diagrams, in which the gluon and ghost propagators and hence the functions $f_{1,k}$ and $f_{2,k}$ appear, and will be given in the appendix. Generally (3.15) can be

solved for m_k^2 only numerically, but in any case m_k^2 is not an independent parameter of $\widehat{\Gamma}_k$.

To determine the flow of the coupling parameter g_k , we will use the ERGE for the KcA -vertex with vanishing momenta. Then it turns out, that in the Landau gauge g_k actually does not run at all with k ; it remains a constant, which will be specified once and for all at the starting point $k = \Lambda$. (It is straightforward, on the other hand, to define a ‘‘physical’’ running coupling in terms of the dressed one-gluon exchange diagram or the function V_k of (3.14), which runs with k due to the running of $f_{1,k}$ and $f_{2,k}$. Within our approximation this running agrees to one-loop order with the usual perturbative running, as we have checked explicitly.) In the Landau gauge the independent running parameters in $\widehat{\Gamma}_k$ are thus only the two functions $f_{1,k}$ and $f_{2,k}$. The aim will be the determination of these quantities by integrating the ERGEs with suitable boundary conditions.

Finally we have to introduce and specify the IR cutoff term ΔS_k in (3.5). If we demand a reasonable behaviour of the full propagators including the IR cutoff terms which appear on the r.h. side of the ERGE (3.7), we are led to more sophisticated choices than an expression of the form of (2.4) with $\Lambda \rightarrow \infty$. (Note that $R_{k,\mu\nu}$ and \tilde{R}_k have to vanish identically for $k \rightarrow 0$ in order that the modified STI (3.8) turns into the standard one (3.4). To this end Λ has to be put equal to ∞ inside $R_{k,\mu\nu}$ and \tilde{R}_k , which is perfectly consistent as discussed in Sect. 2.) First, we do not want to generate any poles on the real axis through the introduction of ΔS_k . This enforces a dependence of $R_{k,\mu\nu}$ and \tilde{R}_k on the functions $f_{1,k}$ and $f_{2,k}$ such that they are proportional to $f_{1,k}$ and $f_{2,k}$, respectively. Now, however, it is no longer automatically guaranteed, that ΔS_k serves as an IR cutoff if, e.g., $f_{1,k}(p^2)$ vanishes for $p^2 \rightarrow 0$ (as we might possibly expect). This problem is overcome by writing the functions $f_{1,k}$ and $f_{2,k}$ also in the exponents. Taking the presence of the gluon mass term m_k^2 into account, our choice of the cutoff functions thus reads

$$\begin{aligned} R_{k,\mu\nu}(p^2) &= (p^2 f_{1,k}(p^2) + m_k^2) \frac{e^{-(p^2 f_{1,k}(p^2) + m_k^2)/k^2}}{1 - e^{-(p^2 f_{1,k}(p^2) + m_k^2)/k^2}} \\ &\times \left(\delta_{\mu\nu} - \frac{p_\mu p_\nu}{p^2} \right) \\ &+ \left(\frac{p^2}{\alpha} + m_k^2 \right) \frac{e^{-(p^2/\alpha + m_k^2)/k^2}}{1 - e^{-(p^2/\alpha + m_k^2)/k^2}} \frac{p_\mu p_\nu}{p^2} \quad , \quad (3.16a) \end{aligned}$$

$$\tilde{R}_k(p^2) = p^2 f_{2,k}(p^2) \frac{e^{-p^2 f_{2,k}(p^2)/k^2}}{1 - e^{-p^2 f_{2,k}(p^2)/k^2}} \quad . \quad (3.16b)$$

In order to obtain the full propagators, these terms have to be added to the quadratic terms $\sim A \cdot A$ or $\sim \bar{c} \cdot c$ in the action $\widehat{\Gamma}_k$. Let us at this point remark a subtlety which has already been noted in [14]: the modified STI (3.8) determines, after its expansion to $\mathcal{O}(A \cdot c)$, only the longitudinal part of the gluon propagator and requires the presence of a term of the form $m_k^2 p_\mu p_\nu / p^2$. Then it is the




Fig. 1. Diagrammatic form of the ERGEs for the ghost propagator function $f_{2,k}$ and the gluon propagator function $f_{1,k}$. *Internal curly lines* denote the full gluon propagator, *internal dotted lines* the full ghost propagator and full points the full vertices from the ansatz (3.10) for $\widehat{\Gamma}_k$. The *crossed circles* denote insertions of $\partial_k R_{k,\mu\nu}$ resp. $\partial_k \widehat{R}_k$

condition of locality of the effective action, i.e. the need to cancel terms of the form $p_\mu p_\nu / p^2$, which requires the presence of the same mass term m_k^2 in the transverse part of the gluon propagator. Only after this consideration has been completed, we are allowed to approach the Landau gauge $\alpha = 0$. In [14] we had checked that $\alpha = 0$ is preserved by the RG flow (and IR stable), and that the r.h. sides of the ERGEs remain finite. Since this choice simplifies the calculations considerably, we will work in the Landau gauge throughout the rest of this paper. The full gluon and ghost propagators, as derived from $\Gamma_k = \widehat{\Gamma}_k + \Delta S_k$, then read

$$\frac{1 - e^{-(p^2 f_{1,k}(p^2) + m_k^2)/k^2}}{p^2 f_{1,k}(p^2) + m_k^2} \left(\delta_{\mu\nu} - \frac{p_\mu p_\nu}{p^2} \right) \quad (3.17a)$$

and

$$\frac{1 - e^{-p^2 f_{2,k}(p^2)/k^2}}{p^2 f_{2,k}(p^2)}, \quad (3.17b)$$

respectively.

4 Procedure

Let us now turn to the computation of $f_{1,k}$ and $f_{2,k}$ by integrating the ERGEs. The ERGE for $f_{1,k}$ can most easily be obtained by studying the terms quadratic in A of the functional ERGE (3.7). Its diagrammatic form is shown in Fig. 1. Note that the 3- and 4-gluon vertices are more complicated than the ones of a classical Yang-Mills action and are given in the appendix. The form of the 3-gluon vertex coincides with the one used in the framework of SDEs [17] because it corresponds to a special solution of the standard STIs [22] (up to a momentum-dependent field redefinition); we have not been able to find the corresponding 4-gluon vertex elsewhere in the literature. The ERGE for $f_{2,k}$ is obtained from the terms $\sim \bar{c}c$ of (3.7), and is also shown in diagrammatic form in Fig. 1.

The crossed circles in Fig. 1 denote insertions of $\partial_k R_{k,\mu\nu}$ resp. $\partial_k \widehat{R}_k$ according to the ERGE (3.7). Due to

the complicated expressions (3.16) for these cutoff functions, i.e. their dependence on m_k^2 , $f_{1,k}$ and $f_{2,k}$, the k -derivatives ∂_k acting on $R_{k,\mu\nu}$ and \widehat{R}_k have to be decomposed into partial derivatives $\partial/\partial k$, $\partial f_{1,k}/\partial k$, $\partial/\partial f_{1,k}$, etc.

The system of ERGEs for $f_{1,k}$ and $f_{2,k}$ (in the Landau gauge), which is obtained from the diagrams of Fig. 1, is thus of the form

$$\begin{aligned} \partial_k f_{1,k} = & \frac{g_\Lambda^2}{(4\pi)^2} \left(h_1^0(k^2, m_k^2, f_{1,k}, f_{2,k}) \right. \\ & + \partial_k f_{1,k} * h_1^1(k^2, m_k^2, f_{1,k}, f_{2,k}) \\ & + \partial_k f_{2,k} * h_1^2(k^2, m_k^2, f_{1,k}, f_{2,k}) \\ & \left. + \partial_k m_k^2 * h_1^3(k^2, m_k^2, f_{1,k}, f_{2,k}) \right), \quad (4.1a) \end{aligned}$$

$$\begin{aligned} \partial_k f_{2,k} = & \frac{g_\Lambda^2}{(4\pi)^2} \left(h_2^0(k^2, m_k^2, f_{1,k}, f_{2,k}) \right. \\ & + \partial_k f_{1,k} * h_2^1(k^2, m_k^2, f_{1,k}, f_{2,k}) \\ & + \partial_k f_{2,k} * h_2^2(k^2, m_k^2, f_{1,k}, f_{2,k}) \\ & \left. + \partial_k m_k^2 * h_2^3(k^2, m_k^2, f_{1,k}, f_{2,k}) \right). \quad (4.1b) \end{aligned}$$

The evaluation of the expressions h_i^j requires a numerical computation of the one-loop integrals appearing in Fig. 1. The $*$ denotes corresponding convolutions with respect to the loop momentum. It is evident from (4.1), that we still need the knowledge of $\partial_k m_k^2$ in order to determine $\partial_k f_{1,k}$ and $\partial_k f_{2,k}$. $\partial_k m_k^2$ can be obtained from the same diagrams in Fig. 1, which are relevant for $\partial_k f_{1,k}$, in the limit of vanishing external momentum. Again this gives us an expression of the form

$$\begin{aligned} \partial_k m_k^2 = & \frac{g_\Lambda^2}{(4\pi)^2} \left(h_3^0(k^2, m_k^2, f_{1,k}, f_{2,k}) \right. \\ & + \partial_k f_{1,k} * h_3^1(k^2, m_k^2, f_{1,k}, f_{2,k}) \\ & + \partial_k f_{2,k} * h_3^2(k^2, m_k^2, f_{1,k}, f_{2,k}) \\ & \left. + \partial_k m_k^2 * h_3^3(k^2, m_k^2, f_{1,k}, f_{2,k}) \right). \quad (4.2) \end{aligned}$$

Alternatively, we can obtain a similar expression by differentiating (3.15) with respect to k . We thus have two different equations at our disposal to complete the system (4.1). In the absence of approximations, the two different completed systems would yield exactly the same results, as follows from the compatibility of the full (3.7) and (3.8). Since the r.h. sides of the (4.2) and (3.15) involve our ansatz (3.10) for $\widehat{\Gamma}_k$ rather than the full effective action, however, we obtain different results. (Nevertheless, they agree on the one-loop level [14].) The use of $\widehat{\Gamma}_k$ instead of the full effective action involves two different kinds of approximations related to the operators of type a) and b), respectively (see the previous section).

The compatibility of the two completed systems of equations considered above turns out to be sensitive to both kinds of approximations. As in [14], we will use the difference between the corresponding results as a quantitative measure of the inadequacy of the approximations (for the respective scale k). Here we concentrate on the

effect of the approximations on the gluon and ghost propagator functions: We will take a certain discrepancy (in fact 10%) in the results for $\partial_k f_{1,k}$ or $\partial_k f_{2,k}$ at some value of the momentum as an indication that the approximations have a considerable influence on the contributions to the flow of $f_{1,k}$ or $f_{2,k}$.

We are of course aware of the fact that the described consistency check merely yields a necessary condition for the validity of the approximation, not a sufficient one. There exists, however, another a-posteriori justification for our procedure: The results we obtain will show a remarkable degree of universality in the sense of [23] (see the next section) as long as $k > k_0$, where the scale k_0 is determined from the above consistency condition. Since universality is only expected to hold when gauge symmetry is taken into account properly, and no significantly contributing operators are neglected, this property provides a strong indication of the consistency of our procedure.

In practice we will use the form of (4.2) as derived from the ERGE (3.7) in order to complete the system of linear equations for $\partial_k f_{1,k}$ and $\partial_k f_{2,k}$. The other system of equations, with (4.2) taken from the derivative of (3.15), will be employed for the consistency check at every scale k .

Given the ERGEs for $f_{1,k}$ and $f_{2,k}$, we still need suitable boundary conditions in order to proceed with the numerical integration. These boundary conditions can be chosen such that $\hat{\Gamma}_\Lambda$ resembles as close as possible the classical (bare) Yang-Mills action (3.1). One would thus choose, at $k = \Lambda$, $f_{1,\Lambda}(p^2) = f_{2,\Lambda}(p^2) = 1$ and g_Λ not too large in order to start in the perturbative regime. It is possible, however, to improve the form of the effective action at the starting point $k = \Lambda$. By definition, $\hat{\Gamma}_\Lambda$ should include all quantum effects involving internal momenta $p^2 \geq \Lambda^2$. On the one hand these are certainly small for a small coupling g_Λ , on the other hand it is fairly straightforward to include them to one-loop order. To this end one has to calculate the one-loop diagrams, which contribute to the gluon and ghost propagator, with IR cutoff functions R_k as in (3.16) and $k = \Lambda$. (Inside these cutoff functions we can set, to one-loop order, $f_{1,k}(p^2) = f_{2,k}(p^2) = 1$ and $m_k^2 = 0$). The diagrams have to be regularized in the UV (e.g. dimensionally) and renormalized such that, e.g., the renormalization conditions

$$f_{1,\Lambda}(0) = 1 \quad , \quad f_{2,\Lambda}(0) = 1 \quad (4.3)$$

are satisfied. As a result one obtains

$$\begin{aligned} f_{1,\Lambda}(p^2) &= 1 + \delta f_1^{1\text{-loop}}(p^2) \quad , \\ f_{2,\Lambda}(p^2) &= 1 + \delta f_2^{1\text{-loop}}(p^2) \end{aligned} \quad (4.4)$$

where the one-loop contributions $\delta f_i^{1\text{-loop}}$ are given in the appendix. For large p^2 the functions $\delta f_i^{1\text{-loop}}$ behave asymptotically as

$$\begin{aligned} \delta f_1^{1\text{-loop}}(p^2) &\sim \frac{g_\Lambda^2}{(4\pi)^2} \cdot \frac{13}{2} \ln\left(\frac{p^2}{\Lambda^2}\right) \quad , \\ \delta f_2^{1\text{-loop}}(p^2) &\sim \frac{g_\Lambda^2}{(4\pi)^2} \cdot \frac{9}{4} \ln\left(\frac{p^2}{\Lambda^2}\right) \quad . \end{aligned} \quad (4.5)$$

What we have achieved now is an appropriate behaviour of these functions even for momenta $p^2 \gg \Lambda^2$, far above the starting scale, provided $g_\Lambda^2/(4\pi)^2 \cdot \ln(p^2/\Lambda^2) \ll 1$. Now $V_k(p^2)$ of (3.14) has the appropriate logarithmic p^2 -dependence even for $p^2 \gg \Lambda^2$, corresponding to a one-loop RG improvement of the ‘‘physical’’ coupling constant:

$$V_{k=\Lambda}(p^2 \gg \Lambda^2) \sim \frac{g_\Lambda^2}{p^2 \left(1 + \frac{11g_\Lambda^2}{(4\pi)^2} \ln\left(\frac{p^2}{\Lambda^2}\right)\right)} \quad . \quad (4.6)$$

As boundary conditions for the integration of the ERGEs we thus use propagator functions $f_{i,\Lambda}$ according to (4.4), the numerical solution of (3.15) for m_Λ^2 , and values for g_Λ (as the only free parameter) from $g_\Lambda = 2.0$ down to $g_\Lambda = 1.4$.

It is clear that, in order to integrate the ERGEs (4.1) for the functions $f_{1,k}$ and $f_{2,k}$, numerical methods have to be employed. To this end we need a parametrization of these functions, which allows for a reliable fit for all values of the scale k . Since these functions always behave like $f_{i,k}(p^2 \rightarrow \infty) \rightarrow 1 + \delta f_i^{1\text{-loop}}$, the following parametrization turns out to be convenient:

$$f_{i,k}(p^2) = 1 - \sum_j \frac{\alpha_j^i}{1 + \gamma_j^i p^2} + \delta f_i^{1\text{-loop}}(p^2) \quad (4.7)$$

with up to 6 pairs of parameters α_j^i , γ_j^i for $i = 1, 2$, respectively.

Next we have to diagonalize (4.1) in the space of momenta p^2 . To this end we evaluate the functions h_i^j in (4.1) and (4.2) for a number of momenta $p^2 = p_j^2$ with up to 35 different values for p_j^2 in the range $10^{-1}k_0^2$ to $10\Lambda^2$, where k_0^2 is determined dynamically (see below). (It turns out to be convenient to space the momenta p_j^2 equally on a logarithmic scale.) By solving the system of linear algebraic equations we then obtain expressions for $\partial_k f_{i,k}(p_j^2)$ and $\partial_k m_k^2$. One step in the integration of the ERGE then amounts to a computation of

$$f_{i,k+\Delta k}(p_j^2) = f_{i,k}(p_j^2) + \Delta k \cdot \partial_k f_{i,k}(p_j^2) \quad (4.8)$$

with $|\Delta k/k| < 3 \cdot 10^{-2}$, and a subsequent determination of the new fit parameters α_j^i , γ_j^i of (4.7).

The new value of the parameter m_k^2 gets determined by a new numerical solution of (3.15) rather than by the numerical integration of $\partial_k m_k^2$; since it is a relevant parameter (the only one), for $k^2 \ll \Lambda^2$ accumulated theoretical and numerical errors would easily generate a m_k^2 of $\mathcal{O}(\Lambda^2)$ instead of $\mathcal{O}(k^2)$, as it should be according to (3.15).

Of course we have varied many details of the numerical procedure in order to test the robustness of the final results. An upper limit of ~ 35 different values for p_j^2 is dictated by the finite amount of available computing time, and this also keeps us from choosing arbitrarily small values for the gauge coupling g_Λ at the starting point $k^2 = \Lambda^2$, since then the ERGEs would have to be integrated over arbitrarily many orders of magnitude of the scale k^2 in order to arrive at the physically interesting non-perturbative regime.

5 Results

Our results can most easily be represented in terms of a function $F_k(p^2)$,

$$F_k(p^2) = \frac{1}{g_A^2} f_{1,k}(p^2) f_{2,k}^2(p^2) \quad , \quad (5.1)$$

which is simply related to the potential $V_k(p^2)$ in (3.14):

$$V_k(p^2) = \frac{1}{p^2 F_k(p^2)} \quad . \quad (5.2)$$

In Figs. 2–4 we show our results for $F_k(p^2)$ for a bare coupling $g_A = 1.4$. We plotted this function for different values of k^2 , $k^2 = \Lambda^2$, $10^{-1}\Lambda^2$, $10^{-2}\Lambda^2$, $10^{-3}\Lambda^2$, $k^2 = k_0^2 = 7.43 \cdot 10^{-5}\Lambda^2$, as full lines, and for some still smaller value $k^2 = \hat{k}^2 = 3.68 \cdot 10^{-5}\Lambda^2$ as a dashed line. p^2 is given in units of Λ^2 ; the maximal value of p^2 is Λ^2 , $10^{-1}\Lambda^2$ and, in order to resolve the small p^2 region, $10^{-2}\Lambda^2$ in Figs. 2 to 4, respectively. The curves with decreasing values at $p^2 = 0$ correspond to decreasing values of k^2 . We see that, for small enough k^2 and small enough p^2 , $F_k(p^2)$ approaches a form $\sim p^2$ near the origin, which corresponds to a $1/p^4$ -behaviour of the potential $V_k(p^2)$.

Just before $F_k(p^2)$ becomes 0 for $p^2 \rightarrow 0$, however, the two different methods of evaluating $\partial_k m_k^2$ (cf. (4.2) and the discussion below) lead to different results for $\partial_k f_{1,k}$ or $\partial_k f_{2,k}$, with a relative difference of ~ 0.1 . We denoted the corresponding scale by k_0 . For slightly smaller values of k the equation for the gluon mass term m_k^2 (3.15) possesses no longer a solution for m_k^2 .

The incompatibility of the RG flow with the modified STIs (3.8) below k_0 indicates, that our approximation becomes unreliable in this regime: Numerically different results for $\partial_k f_{1,k}$ or $\partial_k f_{2,k}$ show, that the neglected contributions play an important role on the r.h. side of the ERGE for the gluon or ghost propagator, respectively, and that our ansatz is too restrictive in this regime.

The gluon mass term m_k^2 itself, which was determined by (3.15) to be of $\mathcal{O}(\Lambda^2)$ at the starting point $k = \Lambda$, does not show an unusual behaviour for $k \sim k_0$. m_k^2 is always of $\mathcal{O}(k^2)$ or, more precisely, $m_k^2 \sim -0.6 \cdot k^2$ for $k \rightarrow k_0$.

Thus we cannot have confidence in the form of the RG flow for $k < k_0$. However, generally the RG flow modifies $F_k(p^2)$ only for $p^2 \lesssim k^2$ (cf. Figs. 2–4), hence our result for F_{k_0} is nevertheless trustworthy for $p^2 \gtrsim k_0^2$. Our results also indicate that, within a more general parametrization of $\hat{\Gamma}_k$, $F_k(0)$ approaches 0 for $k^2 \rightarrow 0$: In Figs. 2–4 we have shown the function $F_k(p^2)$ for some value $k^2 = \hat{k}^2 < k_0^2$ as a dashed line, and we see that the decrease of $F_k(p^2)$ at $p^2 = 0$ continues. For still smaller values of k^2 , however, the integration of the system of differential equations becomes numerically unstable and does no longer allow to obtain reliable results.

In any case, already our results for $F_{k_0}(p^2)$ for $p^2 \gtrsim k_0^2$ strongly indicate a form $\sim p^2$ near the origin, and the trustworthy range of p^2 may be sufficient for some phenomenological investigations involving, e.g., heavy quarks.

It might be interesting to get some feeling for the orders of magnitude of the different scales involved. Up to now all scales are only defined relative to the starting scale Λ . In order to relate this scale to a physical scale one can try to compare our result for $V_{k_0}(p^2)$, for $p^2 \gtrsim k_0^2$, with a phenomenological parametrization of this potential, which contains dimensionful parameters known in units of MeV. A convenient form of such a parametrization is given by Richardson [24], which reads in our convention

$$V_R(p^2) = \frac{48\pi^2}{(33 - 2N_f)p^2 \ln(1 + p^2/\Lambda_R^2)} \quad (5.3)$$

with $\Lambda_R \cong 400$ MeV, and for a rough comparison with our results (in pure SU(3) Yang-Mills theory) we may set $N_f = 0$. In terms of our function $F_k(p^2)$ of (5.1) this ansatz reads

$$F_R(p^2) = \frac{11 \ln(1 + p^2/\Lambda_R^2)}{16\pi^2} \quad . \quad (5.4)$$

We performed a fit of (5.4) to our function $F_{k_0}(p^2)$ for the region of momenta $k_0^2 \ll p^2 \ll \Lambda^2$ to determine the ratio Λ/Λ_R . The result is shown in Figs. 5 and 6, where we plotted $F_{k_0}(p^2)$ as a full line and $F_R(p^2)$, (5.4), as a dashed line, against p^2 in GeV². The optimized relation between Λ and Λ_R is such that $\Lambda = 12.1$ GeV, which implies that the scale k_0 , where our approximation ceases to be appropriate, is ~ 104 MeV. One can see that our potential is surprisingly close to the phenomenological one of Richardson, both in the perturbative (Fig. 5) and in the nonperturbative (Fig. 6) regime. Significant deviations show up only for momenta $p^2 \lesssim k_0^2$.

Let us now look at the dependence of our results on the bare coupling g_A . We find that also for larger values of g_A , $F_k(0)$ becomes small for small k , until our consistency condition indicates the inadequacy of the approximation at some scale k_0 , with $k_0^2/\Lambda^2 = 6.13 \cdot 10^{-4}$, $2.69 \cdot 10^{-3}$, and $7.98 \cdot 10^{-3}$ for $g_A = 1.6$, 1.8 and 2.0, respectively. In Fig. 7 we compare the different results for $F_{k_0}(p^2)$ with $g_A = 1.4$ (full line), $g_A = 1.6$ (long-dashed line), $g_A = 1.8$ (short-dashed line) and $g_A = 2.0$ (dotted line). In each case we determined the physical value of the starting scale by a fit to (5.4), and the results for Λ are given by 12.1 GeV, 4.8 GeV, 2.6 GeV and 1.7 GeV for $g_A = 1.4, 1.6, 1.8$ and 2.0, respectively. From the values of k_0^2/Λ^2 given above, we then obtain the corresponding values for k_0 , $k_0 = 104$ MeV, 119 MeV, 134 MeV and 151 MeV. We see that the curves in Fig. 7 coincide remarkably well, which assures us that the numerical method we use for the integration of the ERGEs does not accumulate numerical errors. The approximate independence of the curves on g_A or Λ even for momenta larger than the respective starting scale is due to the one-loop improvement of the starting action $\hat{\Gamma}_\Lambda$ (or of the functions $f_{1,\Lambda}$ and $f_{2,\Lambda}$ according to (4.4)). In Fig. 8 we show the same curves for $F_{k_0}(p^2)$ for the different values of g_A or Λ , which have been obtained without the one-loop improvement.

Figs. 7 and 8 lead to a series of important observations: First, as we see most clearly from Fig. 8, the




Fig. 2. Results for the k -dependent function $F_k(p^2)$ (5.1), which is related to the potential $V_k(p^2)$ through (5.2), for different values of k^2 , $k^2 = \Lambda^2$, $10^{-1}\Lambda^2$, $10^{-2}\Lambda^2$, $10^{-3}\Lambda^2$, $k^2 = k_0^2 = 7.43 \cdot 10^{-5}\Lambda^2$, as *full lines*, and for $k^2 = \hat{k}^2 = 3.68 \cdot 10^{-5}\Lambda^2$ as a *dashed line*. The bare coupling g_Λ is $g_\Lambda = 1.4$, and p^2 is given in units of Λ^2 . The curves with decreasing values at $p^2 = 0$ correspond to decreasing values of k^2




Fig. 3. As in Fig. 2, with a maximal value of $p^2 = 10^{-1}\Lambda^2$




Fig. 4. As in Fig. 2, with a maximal value of $p^2 = 10^{-2}\Lambda^2$ in order to resolve the small p^2 region




Fig. 5. Fit of our result for $F_{k_0}(p^2)$ (full line) to the parametrization of Richardson [24] (dashed line), with an optimized relation between Λ and $\Lambda_R \cong 400$ MeV such that $\Lambda = 12.1$ GeV




Fig. 6. As in Fig. 5, with a maximal value of $p^2 = 4$ GeV²




Fig. 7. Results for $F_{k_0}(p^2)$ for different values of the bare coupling $g_\Lambda = 1.4$ (full line), $g_\Lambda = 1.6$ (long-dashed line), $g_\Lambda = 1.8$ (short-dashed line) and $g_\Lambda = 2.0$ (dotted line). In each case we determined the physical value of the starting scale Λ by a fit to the parametrization of Richardson, (5.4)




Fig. 8. As in Fig. 7, but without the one-loop improvement of the propagator functions $f_{i,k}$ at the starting scale Λ

curves are essentially unchanged compared to their form at the respective starting scale Λ for momenta $p^2 \gg \Lambda^2$, as they should be. In particular, the curve for $\Lambda = 1.7$ GeV ($g_\Lambda = 2.0$) in Fig. 7 is given in the perturbative region essentially by the one-loop form (4.4), so comparison with, e.g., the curves for $\Lambda = 12.1$ GeV ($g_\Lambda = 1.4$) in both figures shows, that the integration of the ERGEs correctly reproduces the one-loop form in the perturbative domain. More importantly, we find universal behaviour in the non-perturbative domain: For momenta $p^2 \ll \Lambda^2$, i.e. below the respective starting scales, all curves merge in one, as is most impressively demonstrated by Fig. 8, and the form of the resulting curve does not depend on the boundary condition we use. This independence of physics at $p^2 \ll \Lambda^2$ with respect to variations of the starting scale Λ and to the inclusion of irrelevant operators at Λ (as is the case for the one-loop improved boundary conditions) is what we call universality here, in accordance with [23].

Generally, the running coupling constant in the perturbative domain can be parametrized with the help of a RG invariant parameter Λ_{QCD} . The precise value of Λ_{QCD} depends on the definition of the running coupling, and it is RG invariant, in the UV domain, only if the coupling runs according to the two-loop beta function. The scale k_0 can be considered as an example for a dynamically generated scale. Then, the ratio k_0/Λ_{QCD} and hence k_0 itself (in MeV) should be independent of the value of g_Λ for Λ large enough. However, within our parametrization of $\hat{\Gamma}_k$ only a subset of all possible two-loop diagrams is included within a perturbative expansion of the ERGEs. Thus a running coupling g_k (defined through, e.g., the ghost-gluon vertex at vanishing external momenta, but nonvanishing IR cutoff k) and hence g_Λ do not yet run according to two-loop perturbation theory. This explains why k_0 , as given in MeV above, is not yet independent of g_Λ .

Nevertheless, assuming k_0 to be of the order of Λ_{QCD} , already one-loop perturbation theory implies the following relation for large Λ :

$$\frac{1}{(b_0 \ln(\Lambda^2/k_0^2))^{1/2}} = g_\Lambda \left(1 + \mathcal{O}\left(\frac{g_\Lambda^2}{4\pi}\right) \right) \quad (5.5)$$

with $b_0 = 11/(16\pi^2)$. From our results we obtain for the quantity $(b_0 \ln(\Lambda^2/k_0^2))^{-1/2}$ the values 1.23, 1.40, 1.56 and 1.72 for $g_\Lambda = 1.4, 1.6, 1.8$ and 2.0, respectively. Hence (5.5) is satisfied to the required accuracy.

For practical applications of our results on the heavy quark potential $V^{Q\bar{Q}}(p^2)$ it may be useful to see its final explicit form (in momentum space). Taking the QCD colour factors into account, we have with our convention (3.14) for $V_k(p^2)$

$$V^{Q\bar{Q}}(p^2) = \frac{4}{3} V_{k=k_0 \sim 0}(p^2) \quad . \quad (5.6)$$

In Fig. 9 we plot our result for $V^{Q\bar{Q}}(p^2)$ for $0 < p^2 < (5 \text{ GeV})^2$.

Finally, one may ask whether the behaviour of $F_{k_0}(p^2)$ for $p^2 \rightarrow 0$ is due to the form of $f_{1,k_0}(p^2)$ or both $f_{1,k_0}(p^2)$ and $f_{2,k_0}(p^2)$. To this end we plot in Fig. 10 both functions $f_{1,k_0}(p^2)$ and $f_{2,k_0}(p^2)$ (with p^2 rescaled as before) for $g_\Lambda = 1.4$. We see that both functions become small for $p^2 \rightarrow 0$, but it would be grossly misleading (and unphysical) to identify the gluon propagator function $(p^2 f_1(p^2))^{-1}$ with the heavy quark potential and to neglect the dependence of $V(p^2)$ of (3.14) on $f_2(p^2)$, as it is done if the ghost part of the action is neglected, and the STIs are not properly taken into account.

6 Discussion and outlook

In this paper we have performed the integration of the ERGEs for non-abelian gauge theories, with the bare action as the only input. We concentrated on pure SU(3) Yang-Mills theory (or quenched QCD), and the momentum dependence of the gluon and ghost propagators; these quantities determine the heavy quark potential.

The neglect of certain contributions on the r.h. sides of the ERGEs for the 2-point functions turned the system of ERGEs into a closed system of differential equations, which we integrated numerically. A priori the validity of the approximation is difficult to judge in the strong




Fig. 9. The explicit form of our result for the heavy quark potential $V^{Q\bar{Q}}(p^2)$ in momentum space




Fig. 10. Results for the propagator functions f_{1,k_0} (full line) and f_{2,k_0} (dashed line) for $g_\Lambda = 1.4$

coupling regime, where it amounts to the neglect of contributions of the same order in the coupling constant as the ones maintained. Here the use of modified STIs plays an important role: Due to their validity parts of the k -dependent effective action can be determined in two different ways, namely either directly through these identities or by integrating the ERGEs. Without approximations both methods give the same results, but in the presence of approximations a possible difference between the corresponding results can be used to estimate the error induced by the approximations. In our case it was the gluon mass term, which was obtained in two different ways. We used a 10%-deviation in $\partial_k f_{1,k}$ or $\partial_k f_{2,k}$, induced by the different methods to obtain $\partial_k m_k^2$, as an indication for the breakdown of our approximation near the corresponding scale k_0 . Accordingly, our result for $V(p^2) \equiv V_{k_0}(p^2)$ is only consistent for $p^2 \gtrsim k_0^2$, where we estimated $k_0 \sim 100$ MeV. Our main findings are the strong indication of a $1/p^4$ -behaviour of $V(p^2)$ in the trustworthy regime of p^2 , and the fact that the full form of $V(p^2)$ happens to be close to the one proposed by Richardson [24].

In the second section of this paper we discussed the formal relation between ERGEs and SDEs, with the re-

sult that effective actions, which solve SDEs in the presence of an IR cutoff k , can be considered as quasi-fixed points of the ERGEs. Although this formal relation is no longer exact in the presence of truncations of effective actions, it is thus not astonishing, that the behaviour of the gluon propagator, which has been derived here using the ERGEs, has also been shown to be a solution to SDEs in the Landau gauge [17]. We believe, however, that the present method possesses a number of advantages compared to the formalism of the SDEs: Firstly, the regularization of both ultraviolet and infrared singularities in the context of the SDEs generates notorious technical difficulties, which require, to some extent, ad hoc prescriptions, the influence of which on the final result is difficult to control. The present method is, in contrast, free of problems related to UV or IR singularities by construction.

Secondly, a priori both methods require a suitable truncation of the effective action. Within the present method, however, it is technically simpler to include more and more terms; for the present investigation, e.g., we already included the contributions of the 4-gluon vertex and the ghost fields, which were neglected in the SDE approach [17]. We have stressed that the detailed form of the dressed

one-gluon exchange diagram between heavy quarks, which involves the ghost propagator function at the quark-gluon vertex due to the STIs, differs considerably from the form of the inverse gluon propagator alone (in the Landau gauge), which is not a physical quantity by itself. Moreover, it is technically feasible in our formalism to include further terms in the effective action without too much effort.

Finally, within the SDE approach one is only able to check the self-consistency of a particular ansatz; it is practically impossible to prove rigorously, that the chosen ansatz is the only possible one. The integration of the ERGEs, on the other hand, always gives a unique answer, namely the one which is smoothly connected to the corresponding bare action, and to perturbation theory in the UV in the case of asymptotic freedom. Actually, our discussion in Sect. 2 sheds some light on the cases where different solutions of the SDEs exist: on the one hand solutions of the SDEs can be considered as fixed points of the ERGEs for $k \rightarrow 0$, on the other hand it is generally not clear, whether one has obtained an IR stable or an UV stable fixed point. Since only the IR stable solutions can also be obtained by the integration of the ERGEs towards $k \rightarrow 0$, one is lead to the conclusion, that only these are actually physically relevant, which is certainly difficult to check within the SDE approach alone.

Many possibilities exist in order to extend the present approach in the future: firstly, still more terms can be included in the effective action, and collective fields, e.g. for $F_{\mu\nu}^a, F_{\mu\nu}^a$, can be introduced along the lines of [7, 8] in order to reach renormalization scales k below k_0 , and to get a still better understanding of pure Yang-Mills theory free of systematic uncertainties for $p^2 < k_0^2$. Secondly, the quark sector of QCD can be introduced as well. First efforts in this direction have already been made in [8]; there, however, the effects of the gluons have been guessed and parametrized in a phenomenological manner. Now the integration of the ERGEs for the full quark gluon system is within reach (possibly along the lines proposed in [25]), which will allow to study many phenomenologically interesting systems like heavy $q\bar{q}$ -bound states, $q\bar{q}$ -condensates and the meson sector on the basis of just the bare QCD Lagrangian.

Appendix

In this appendix we list some of the functions which are needed in the actual computations. We begin with the 3- and 4-gluon vertices employed in our ansatz (3.10) for $\hat{\Gamma}_k$. They are chosen in such a way as to fulfill the standard STIs in a "minimal" fashion. Explicitly they read

$$\begin{aligned} f_{3,\mu\nu\rho}(p, q, r) &= -\frac{1}{6f_{2,k}(p^2)f_{2,k}(q^2)f_{2,k}(r^2)} \\ &\times \left\{ (q_\mu f_{g,k}(q^2) - r_\mu f_{g,k}(r^2)) \delta_{\nu\rho} \right. \\ &\left. + (r_\nu f_{g,k}(r^2) - p_\nu f_{g,k}(p^2)) \delta_{\rho\mu} \right\} \end{aligned}$$

$$\begin{aligned} &+ (p_\rho f_{g,k}(p^2) - q_\rho f_{g,k}(q^2)) \delta_{\mu\nu} \\ &+ (q_\mu - r_\mu) \frac{f_{g,k}(q^2) - f_{g,k}(r^2)}{q^2 - r^2} (r_\nu q_\rho - q \cdot r \delta_{\nu\rho}) \\ &+ (r_\nu - p_\nu) \frac{f_{g,k}(r^2) - f_{g,k}(p^2)}{r^2 - p^2} (p_\rho r_\mu - r \cdot p \delta_{\rho\mu}) \\ &+ (p_\rho - q_\rho) \frac{f_{g,k}(p^2) - f_{g,k}(q^2)}{p^2 - q^2} \\ &\times (q_\mu p_\nu - p \cdot q \delta_{\mu\nu}) \left. \right\} \end{aligned} \quad (\text{A.1})$$

and

$$\begin{aligned} f_{4,\mu\nu\rho\sigma}(p, q, r, s) &= \frac{1}{4f_{2,k}(p^2)f_{2,k}(q^2)f_{2,k}(r^2)f_{2,k}(s^2)} \\ &\times \left\{ f_{g,k}((p+q)^2) \delta_{\mu\rho} \delta_{\nu\sigma} - 2 \frac{f_{g,k}(p^2) - f_{g,k}(s^2)}{p^2 - s^2} \right. \\ &\times (p_\sigma s_\mu - p \cdot s \delta_{\sigma\mu}) \delta_{\nu\rho} - 2 \frac{f_{g,k}((p+q)^2) - f_{g,k}(p^2)}{(p+q)^2 - p^2} \\ &\times (2p_\nu + q_\nu) (p_\rho \delta_{\sigma\mu} - p_\sigma \delta_{\mu\rho}) + \frac{2}{(r+s)^2 - s^2} \\ &\times \left(\frac{f_{g,k}(p^2) - f_{g,k}((r+s)^2)}{p^2 - (r+s)^2} - \frac{f_{g,k}(p^2) - f_{g,k}(s^2)}{p^2 - s^2} \right) \\ &\left. \times (2p_\nu + q_\nu) (2s_\rho + r_\rho) (p_\sigma s_\mu - p \cdot s \delta_{\sigma\mu}) \right\}, \end{aligned} \quad (\text{A.2})$$

where

$$f_{g,k}(p^2) = f_{1,k}(p^2) f_{2,k}^2(p^2) \quad . \quad (\text{A.3})$$

Observe that we have not symmetrized the expression (A.2) for the four-gluon vertex with respect to momenta and indices, in order to keep it to a reasonable length.

Next we turn to the modified STI for the gluon mass term, (3.15). In diagrammatic form it is given by a number of one-loop diagrams. The integrations over the angular variables of the loop momenta can be performed, and one is left with integrals over $t \equiv q^2$. After a careful consideration of the limit $\alpha \rightarrow 0$, corresponding to the Landau gauge, the following expression for the function ST results:

$$\begin{aligned} \text{ST}(k^2, m_k^2, f_{1,k}, f_{2,k}) &= \frac{3}{4f_{2,k}^2(0)} \int_0^\infty dt \frac{t}{k^2} \left\{ -e^{-tf_{2,k}(t)/k^2} \right. \\ &\times \left[3f_{1,k}(t) \frac{1 - e^{-(tf_{1,k}(t)+m_k^2)/k^2}}{tf_{1,k}(t) + m_k^2} \right. \\ &\left. + \left(3 - 4 \frac{t}{k^2} \frac{\partial}{\partial t} (tf_{2,k}(t)) \right) f_{2,k}(t) \frac{1 - e^{-tf_{2,k}(t)/k^2}}{tf_{2,k}(t)} \right] \\ &\left. + 3e^{-(tf_{1,k}(t)+m_k^2)/k^2} \left[\left(4 - 4 \frac{t}{k^2} \frac{\partial}{\partial t} (tf_{1,k}(t)) \right) \right. \right. \end{aligned}$$

$$\begin{aligned}
& \times f_{1,k}(t) \frac{1 - e^{-(tf_{1,k}(t)+m_k^2)/k^2}}{tf_{1,k}(t) + m_k^2} \\
& + f_{2,k}(t) \frac{1 - e^{-tf_{2,k}(t)/k^2}}{tf_{2,k}(t)} \Big] \\
& - 6 \frac{e^{-(tf_{1,k}(t)+m_k^2)/k^2}}{f_{2,k}^2(t)} t \frac{\partial}{\partial t} \left(f_{1,k}(t) f_{2,k}^2(t) \right) \\
& \times \left[\left(1 + \frac{t}{k^2} \frac{\partial}{\partial t} (tf_{1,k}(t)) - \frac{2}{f_{2,k}(t)} \frac{\partial}{\partial t} (tf_{2,k}(t)) \right) \right] \\
& \times \frac{1 - e^{-(tf_{1,k}(t)+m_k^2)/k^2}}{tf_{1,k}(t) + m_k^2} - t \frac{\partial}{\partial t} (tf_{1,k}(t)) \\
& \times \frac{1 - \left(1 + (tf_{1,k}(t) + m_k^2)/k^2 \right) e^{-(tf_{1,k}(t)+m_k^2)/k^2}}{\left(tf_{1,k}(t) + m_k^2 \right)^2} \Big] \Big\} . \tag{A.4}
\end{aligned}$$

Finally we give the one-loop expressions $\delta f_i^{1\text{-loop}}$ for the gluon and ghost 2-point functions as needed in (4.4) for the improved boundary conditions. They are calculated with the IR cutoff functions R_k from (3.16) (with $f_{1,k}$ and $f_{2,k}$ set equal to one and $m_k^2 = 0$) and obey the renormalization conditions (4.3). The results are, again in the Landau gauge,

$$\begin{aligned}
& \delta f_1^{1\text{-loop}}(p^2) \\
& = \frac{3g_A^2}{(4\pi)^2} \left\{ \frac{13}{6} \left(\ln(x/2) + C \right) - \frac{131}{144} \right. \\
& \quad - \frac{x^4 - 8x^3 + 9x^2 - 20x - 6}{12x^3} e^{-x} \\
& \quad + \frac{x^4 - 16x^3 - 16x^2 - 56x + 4}{12x^3} e^{-x/2} - \frac{x^2 - 7x}{12} \text{Ei}(-x) \\
& \quad \left. + \frac{x^2 - 14x - 52}{24} \text{Ei}(-x/2) \right\} \Big|_{x = \frac{p^2}{\Lambda^2}} \tag{A.5}
\end{aligned}$$

and

$$\begin{aligned}
& \delta f_2^{1\text{-loop}}(p^2) \\
& = \frac{3g_A^2}{(4\pi)^2} \left\{ \frac{3}{4} \left(\ln(x/2) + C \right) - \frac{3}{8} \right. \\
& \quad + \frac{3x + 2}{4x^2} + \frac{x^2 - x + 2}{4x^2} e^{-x} - \frac{x^2 + x + 2}{2x^2} e^{-x/2} \\
& \quad \left. + \frac{x}{4} \text{Ei}(-x) - \frac{x + 3}{4} \text{Ei}(-x/2) \right\} \Big|_{x = \frac{p^2}{\Lambda^2}} , \tag{A.6}
\end{aligned}$$

where C denotes Euler's constant, $C \cong 0.577216$. The exponential integral function Ei is defined by

$$\text{Ei}(-x) = \int_{-\infty}^{-x} dt \frac{e^t}{t} \tag{A.7}$$

for $x > 0$.

References

1. C.D. Roberts and A.G. Williams, Progress in Particle and Nuclear Physics **33** (1994) 477
2. K.G. Wilson and I. Kogut, Phys. Rep. **12** (1974) 75; F. Wegner, in Phase Transitions and Critical Phenomena, Vol. 6, eds. C. Domb and M. Green (Academic Press, NY 1975)
3. J. Polchinski, Nucl. Phys. **B231** (1984) 269; for recent references see J. Comellas, Y. Kubyshev and E. Moreno, FTUAM preprint 96/2, hep-th/9601112
4. F. Wegner and A. Houghton, Phys. Rev. **A8** (1973) 401; J.F. Nicoll, T.S. Chang and H.E. Stanley, Phys. Rev. Lett. **33** (1974) 540; A. Hasenfratz and P. Hasenfratz, Nucl. Phys. **B270** (1986) 687
5. S. Weinberg, in Proceedings of the 1976 International School of Subnuclear Physics, Erice (ed. A. Zichichi)
6. U. Ellwanger, Z. Phys. **C62** (1994) 503
7. U. Ellwanger, Z. Phys. **C58** (1993) 679
8. U. Ellwanger and C. Wetterich, Nucl. Phys. **B423** (1994) 137; D. Jungnickel and C. Wetterich, Phys. Rev. **D53** (1996) 5142; D. Jungnickel, Heidelberg preprint HD-THEP-95-40, hep-ph/9509293
9. B. Warr, Ann. Phys. **183** (1988) 1 and ibid. 59
10. G. Keller and C. Kopper, Phys. Lett. **B273** (1991) 323; G. Keller, Helv. Phys. Acta **66** (1993) 453
11. C. Becchi, in Elementary Particles, Field Theory and Statistical Mechanics, eds. M. Bonini, G. Marchesini and E. Onofri, Parma University 1993; M. Bonini, M. D'Attanasio and G. Marchesini, Nucl. Phys. **B418** (1994) 81, Nucl. Phys. **B421** (1994) 429, Nucl. Phys. **B437** (1995) 163, Phys. Lett. **B346** (1995) 87; M. D'Attanasio and T. Morris, Phys. Lett. **B378** (1996) 213
12. M. Reuter and C. Wetterich, Nucl. Phys. **B417** (1994) 181, ibid. **B427** (1994) 291, Phys. Lett. **B334** (1994) 412, Heidelberg preprint HD-THEP-94-39, hep-th/9411227
13. U. Ellwanger, Phys. Lett. **B335** (1994) 364
14. U. Ellwanger, M. Hirsch and A. Weber, Z. Phys. **C69** (1996) 687
15. M. Baker, J.S. Ball and F. Zachariasen, Nucl. Phys. **B186** (1981) 531, ibid. **B186** (1981) 560; A.I. Alekseev, Yad. Fiz. **33** (1981) 516; W.J. Schoenmaker, Nucl. Phys. **B194** (1982) 535; F. Paccanoni, Nuovo Cim. **A74** (1983) 267
16. J.R. Cudell and D.A. Ross, Nucl. Phys. **B194** (1991) 247
17. S. Mandelstam, Phys. Rev. **D20** (1979) 3223; U. Bar-Gadda, Nucl. Phys. **B163** (1980) 312; N. Brown and M.K. Pennington, Phys. Lett. **B202** (1988) 257, Phys. Rev. **D39** (1989) 2723
18. P.V. Landshoff and O. Nachtmann, Z. Phys. **C39** (1989) 405
19. K. Büttner and M.R. Pennington, Phys. Rev. **D52** (1995) 5220
20. C. Itzykson and J.-B. Zuber, Quantum Field Theory (McGraw-Hill, NY 1980)
21. A. Weber, PhD Thesis, Univ. Heidelberg 1995
22. S.K. Kim and M. Baker, Nucl. Phys. **B164** (1980) 152; J.S. Ball and J.-W. Chiu, Phys. Rev. **D22** (1980) 2550
23. R.D. Ball and R.S. Thorne, Ann. Phys. **236** (1994) 117
24. J.L. Richardson, Phys. Lett. **B82** (1979) 272
25. C. Wetterich, Heidelberg preprint HD-THEP-95-47