

Higher-derivative supergravity in $U(1)$ superspace

Régis Le Dû

Centre de Physique Théorique^a, C.N.R.S. Luminy, case 907, F-13288 Marseille, France
 (allocataire M.E.S.R. e-mail: ledu@cpt.univ-mrs.fr)

Received: 9 June 1997 / Revised version: 12 September 1997 / Published online: 26 February 1998

Abstract. The complete structure of curvature squared terms is analyzed in the context of chirally extended supergravity, with special emphasis on the gravitationally induced Fayet–Iliopoulos D–term. Coupling of (chiral) matter is discussed in relation with a possible extension to $U(1)$ supergravity of the equivalence mechanism between $\mathcal{R} + \alpha\mathcal{R}^2$ and General Relativity coupled to a scalar.

Introduction

Higher derivative supergravity theories [1–5] have been proposed as messengers for supersymmetry breaking through gravitational effects [6]. More recently, the equivalence of $\mathcal{R} + \mathcal{R}^2$ theories to gravity coupled to a scalar [7], in its supersymmetric version [8], has been advocated to provide a supersymmetry breaking device as well [9–11]. All these scenarios are based on traditional supergravity. On the other hand, in the presence of a chiral abelian gauge structure, known as $U(1)$ supergravity [12, 13] a supergravity induced D–term appears naturally [14, 15].

In the present paper, we describe explicitly the complete structure of curvature squared terms in $U(1)$ supergravity, and discuss possibilities to extend the scheme of [9] to this case.

In the first chapter we review shortly General Relativity [16] and Whitt’s mechanism [7]. After an outline of $U(1)$ superspace and the construction of the pure $U(1)$ supergravity action in Chapter 2 we turn to the complete description of curvature squared actions for $U(1)$ supergravity. Our description is based on methods of superspace geometry as reviewed in [17]. The paper closes with a discussion of matter coupling to $U(1)$ supergravity with curvature squared terms.

General Relativity with curvature squared terms

It is known that pure General Relativity is a nonrenormalizable theory [18, 19]. Adding quadratic terms in the curvature tensor allows to construct renormalizable actions [20]. The most general action which contains fourth order derivatives can be written as

$$\mathcal{S} = \gamma \int d^4x \sqrt{-g} (\mathcal{R} + \alpha' \mathcal{R}^2 + \beta' \mathcal{R}^{mn} \mathcal{R}_{mn})$$

$$+ c \mathcal{R}^{mnpq} \mathcal{R}_{mnpq} , \quad (1)$$

where γ is related to the gravitational constant, resp. the Planck mass,

$$\gamma = \frac{-1}{16\pi G_N} = -\frac{1}{2} M_P^2, \quad (2)$$

whereas $\gamma\alpha'$, $\gamma\beta'$ and γc are dimensionless. Taking into account the Gauss–Bonnet combination

$$\mathcal{S}_{GB} = \int d^4x \sqrt{-g} (\mathcal{R}^2 - 4\mathcal{R}^{mn} \mathcal{R}_{mn} + \mathcal{R}^{mnpq} \mathcal{R}_{mnpq}) , \quad (3)$$

which, as a topological invariant, does not depend on the metric and, as a consequence, does not contribute to the equations of motion, this action may be written as

$$\begin{aligned} \mathcal{S}' &= \mathcal{S} - c\gamma \mathcal{S}_{GB} \\ &= \gamma \int d^4x \sqrt{-g} (\mathcal{R} + \alpha \mathcal{R}^2 + \beta \mathcal{R}^{mn} \mathcal{R}_{mn}) , \end{aligned} \quad (4)$$

with constants $\alpha = \alpha' - c$, $\beta = \beta' + 4c$. This theory describes [16, 21] the graviton together with a massive spin two “poltergeist” and a massive physical scalar field. For $\beta = 0$ the poltergeist decouples [8] and the action

$$\mathcal{S}_{\mathcal{R}+\mathcal{R}^2} = \gamma \int d^4x \sqrt{-g} (\mathcal{R} + \alpha \mathcal{R}^2) , \quad (5)$$

which, however, is no longer renormalizable [20], describes a graviton coupled to a massive scalar field [7]. Following [7], one starts from the action

$$\mathcal{S}_{\mathcal{R}+\phi} = \gamma \int d^4x \sqrt{-g} (\mathcal{R} + 2\alpha\phi\mathcal{R} - \alpha\phi^2) . \quad (6)$$

On the one hand, varying with respect to ϕ reproduces the action (5). On the other hand, performing a Weyl rescaling

$$g_{mn} = (1 + 2\alpha\phi)^{-1} g'_{mn} , \quad (7)$$

^a Unité Propre de Recherche 7061

yields

$$\mathcal{S}_{\mathcal{R}+\phi} = \gamma \int d^4x \sqrt{-g'} (\mathcal{R}' + 6\alpha^2(1 + 2\alpha\phi)^{-2} \times \partial'^m \phi \partial'_m \phi - \alpha(1 + 2\alpha\phi)^{-2} \phi^2) , \quad (8)$$

where the presence of the massive scalar field is manifest. It is in this sense that curvature squared gravity is said to be equivalent to General Relativity coupled to a scalar field [7].

$U(1)$ superspace and supergravity

Supergravity is described in terms of the vielbein field $e_m^a(x)$ and the Rarita–Schwinger field $\psi_m^\alpha(x)$ together with a set of auxiliary fields¹ M, \bar{M} and b_a , i.e a complex scalar and a vector. The latter are necessary to close the algebra of local supersymmetry transformations. In conventional supergravity which is given as the supersymmetric generalization of the curvature scalar they describe non propagating degrees of freedom.

On the other hand, in supersymmetric versions of theories with curvature squared terms those fields acquire derivatives and become propagating fields as well.

In the present paper we will extend this scenario to the case of $U(1)$ supergravity, describing an enlarged multiplet with additional components A_m, λ^α and \mathbf{D} . This theory has an additional gauged chiral $U(1)$ symmetry with A_m as gauge potential, λ^α the gaugino superpartner and \mathbf{D} the auxiliary field. One of its interesting features is that it allows the construction of a particular supersymmetric Fayet–Iliopoulos [22] term in the context of supergravity [15].

The superspace formulation of $U(1)$ supergravity [12, 13] is a generalization of that of conventional supergravity [23]: in addition to the Lorentz group in the superspace structure group it contains a chiral $U(1)$. In order to be more explicit we review shortly the salient features of $U(1)$ superspace geometry, following closely [24, 17]. The basic objects are the vielbein E^A , the Lorentz connection ϕ_B^A and the $U(1)$ connection A . They are all one-forms in superspace:

$$E^A = dz^M E_M^A , \quad (9)$$

$$\phi_B^A = dz^M \phi_{MB}^A , \quad (10)$$

$$A = dz^M A_M . \quad (11)$$

Correspondingly, one defines the torsion T^A , the Lorentz curvature R_B^A , and the $U(1)$ fieldstrength F :

$$T^A = dE^A + E^B \phi_B^A + \omega(E^A) E^A A , \quad (12)$$

$$R_B^A = d\phi_B^A + \phi_B^C \phi_C^A , \quad (13)$$

$$F = dA , \quad (14)$$

¹ This set of auxiliary fields corresponds to the so called “old–minimal” formulation. Other choices are possible, as for instance “new–minimal” and “non–minimal” formulations, but they will not be considered here

which are two-forms in superspace. The chiral weights of the vielbein are defined to be

$$\omega(E^a) = 0 , \quad \omega(E^\alpha) = 1 , \quad \omega(E_{\dot{\alpha}}) = -1 . \quad (15)$$

The basic covariant superfields which completely describe torsion, curvature and $U(1)$ fieldstrength are

$$R , \quad R^\dagger , \quad G_a , \quad W_{\alpha\beta\gamma} , \quad \bar{W}^{\dot{\alpha}\dot{\beta}\dot{\gamma}} , \quad \text{and} \quad X^\alpha , \quad \bar{X}_{\dot{\alpha}} . \quad (16)$$

Component fields are defined as lowest components of superfields in the usual way

$$E_m^a | = e_m^a(x) , \quad E_m^\alpha | = \frac{1}{2} \psi_m^\alpha(x) , \quad A_m | = A_m(x) , \quad (17)$$

for the gauge fields. In particular the metric tensor is defined as $g_{mn} = e_m^a e_n^b \eta_{ab}$, with $\eta_{ab} = \text{diag}(-, +, +, +)$. Moreover one has the usual definitions

$$R | = -\frac{M}{6} , \quad R^\dagger | = -\frac{\bar{M}}{6} , \quad G_a | = -\frac{b_a}{3} , \quad (18)$$

in the gravity sector and

$$X_\alpha | = -i\lambda_\alpha , \quad \bar{X}^{\dot{\alpha}} | = i\bar{\lambda}^{\dot{\alpha}} , \quad \mathcal{D}^\alpha X_\alpha | = -2\mathbf{D} , \quad (19)$$

in the $U(1)$ sector. We also define a gauge potential² \tilde{A}_m which is related to A_m by

$$\tilde{A}_m \equiv A_m + \frac{i}{2} b_m , \quad (20)$$

and which will be used as the basic component field from now on. Correspondingly, we define the $U(1)$ covariant derivative $\tilde{\mathcal{D}}_m X$

$$\tilde{\mathcal{D}}_m X = \partial_m X + \omega(X) \tilde{A}_m X = \mathcal{D}_m X + \frac{i}{2} \omega(X) b_m X . \quad (21)$$

The supergravity action is given in compact form as the volume element of $U(1)$ superspace, i.e³

$$\mathcal{S}_1 = -3 \int E . \quad (22)$$

The corresponding component field expression is most conveniently extracted from the generic lagrangian [24]:

$$e^{-1} \mathcal{L}_1 = -\frac{1}{4} \mathcal{D}^2 r | - r | (\bar{M} + \bar{\psi}_m \bar{\sigma}^{mn} \bar{\psi}_n) + \frac{i}{2} (\bar{\psi}_{m\dot{\alpha}} \bar{\sigma}^{m\dot{\alpha}\alpha}) \mathcal{D}_\alpha r | + hc , \quad (23)$$

with the choice

$$r = -3R . \quad (24)$$

² This definition takes into account the constraint: $F^{\dot{\beta}}_\alpha = -3G^{\dot{\beta}}_\alpha$ which is used in [24]

³ Action and lagrangian are related by the relation: $\mathcal{S} = \int d^4x \mathcal{L}$

Using the explicit component field form of $\mathcal{D}_\alpha R|$, $\mathcal{D}^2 R|$, one obtains⁴

$$e^{-1}\mathcal{L}_1 = -\frac{M_P^2}{2}\mathcal{R} + \frac{1}{2}\epsilon^{mnpq}\left(\bar{\psi}_m\bar{\sigma}_n\tilde{\mathcal{D}}_p\psi_q - \psi_m\sigma_n\tilde{\mathcal{D}}_p\bar{\psi}_q\right) - \frac{M_P^2}{3}(M\bar{M} - b^a b_a) + \frac{1}{2}(\bar{\psi}_m\bar{\sigma}^m\lambda - \psi_m\sigma^m\bar{\lambda})M_P + \mathbf{D}M_P^2. \quad (25)$$

Clearly, this lagrangian exhibits the usual Einstein term together with a kinetic term for the gravitino and the auxiliary field terms for M , \bar{M} and b_a as in usual supergravity. One of the new features due to the chiral $U(1)$ is the appearance of \tilde{A}_m in the covariant derivative of the gravitino field

$$\tilde{\mathcal{D}}_m\psi_n^\alpha = \partial_m\psi_n^\alpha + \psi_n^\beta\omega_{m\beta}^\alpha + \psi_n^\alpha\tilde{A}_m. \quad (26)$$

Moreover there is a term linear in \mathbf{D} , which clearly shows that this theory in itself cannot be complete. It is completed in adding a kinetic term for the $U(1)$ gauge multiplet. In this case the term linear in \mathbf{D} can play the role of a Fayet–Iliopoulos term [15]. The superfield action is defined as

$$\mathcal{S}_{X^2} = \frac{1}{8}\int\frac{E}{R}X^\alpha X_\alpha + \frac{1}{8}\int\frac{E}{R^\dagger}\bar{X}_{\dot{\alpha}}\bar{X}^{\dot{\alpha}}. \quad (27)$$

Taking $r = \frac{1}{4}X^\alpha X_\alpha$ in the generic construction (23), one derives the component field expression

$$e^{-1}\mathcal{L}_{X^2} = \frac{1}{2}\mathbf{D}^2 + \tilde{F}^{mn}\tilde{F}_{mn} - \frac{i}{2}\lambda\sigma^m\tilde{\mathcal{D}}_m\bar{\lambda} - \frac{i}{2}\bar{\lambda}\bar{\sigma}^m\tilde{\mathcal{D}}_m\lambda + \text{other fermionic terms}. \quad (28)$$

These two lagrangians are separately invariant under the following supersymmetry transformations (as derived from superspace geometry in the usual way)

$$\delta_\xi e_m^a = i(\xi\sigma^a\bar{\psi}_m + \bar{\xi}\bar{\sigma}^a\psi_m)M_P^{-1}, \quad (29)$$

$$\delta_\xi\psi_m^\alpha = \left(2\tilde{\mathcal{D}}_m\xi^\alpha - i\xi^\alpha b_m - \frac{i}{3}(\xi\sigma^a\bar{\sigma}_m)^\alpha b_a + \frac{i}{3}(\bar{\xi}\bar{\sigma}_m)^\alpha M\right)M_P, \quad (30)$$

$$\delta_\xi\bar{\psi}_{m\dot{\alpha}} = \left(2\tilde{\mathcal{D}}_m\bar{\xi}_{\dot{\alpha}} + i\bar{\xi}_{\dot{\alpha}}b_m + \frac{i}{3}(\bar{\xi}\bar{\sigma}^a\sigma_m)_{\dot{\alpha}} b_a + \frac{i}{3}(\xi\sigma_m)_{\dot{\alpha}}\bar{M}\right)M_P, \quad (31)$$

$$\delta_\xi M = -2i(\xi\lambda)\left(+4(\xi\sigma^{mn}\tilde{\mathcal{D}}_m\psi_n)\right)$$

⁴ In this part, one introduces the Planck mass in order to see how it appears in the lagrangian and also in the supersymmetry transformations. In particular, one can notice that a cosmological constant appears with M_P^4 . In the following, we will take $M_P \equiv 1$

$$+i(\psi_m\xi)b^m + i(\bar{\psi}_m\bar{\sigma}^m\xi)M)M_P^{-1}, \quad (32)$$

$$\delta_\xi\bar{M} = 2i(\bar{\xi}\lambda)\left(+4(\bar{\xi}\bar{\sigma}^{mn}\tilde{\mathcal{D}}_m\bar{\psi}_n) - i(\bar{\psi}_m\bar{\xi})b^m - i(\bar{\xi}\bar{\sigma}^m\psi_m)\bar{M}\right)M_P^{-1}, \quad (33)$$

$$\delta_\xi b_a = \left(\frac{1}{2}(\xi\sigma_a\bar{\sigma}^{mn}\tilde{\mathcal{D}}_m\bar{\psi}_n) + \frac{1}{2}(\xi\sigma_a\bar{\sigma}^{mn}\bar{\psi}_n)b_m\right)M_P^{-1} \left(-\frac{3}{2}(\xi\sigma^{mn}\sigma_a\tilde{\mathcal{D}}_m\bar{\psi}_n) - \frac{3}{2}(\xi\sigma^{mn}\bar{\psi}_n)b_m\right)M_P^{-1} \left(-\frac{i}{2}e_a^m(\xi\sigma^d\bar{\psi}_m)b_d + \frac{i}{2}e_a^m(\xi\psi_m)\bar{M}\right)M_P^{-1} + i(\xi\sigma_a\bar{\lambda}) + h.c., \quad (34)$$

$$\delta_\xi\mathbf{D} = (\bar{\xi}\bar{\sigma}^m\tilde{\mathcal{D}}_m\lambda) - (\xi\sigma^m\tilde{\mathcal{D}}_m\bar{\lambda}) - \frac{i}{2}(\xi\sigma^m\bar{\lambda} + \bar{\xi}\bar{\sigma}^m\lambda)b_m + \frac{1}{2}(\psi_m\sigma^{kl}\sigma^m\bar{\xi} - \bar{\psi}_m\bar{\sigma}^{kl}\bar{\sigma}^m\xi) \times (2i\tilde{F}_{kl}M_P^{-1} + i\psi_k\sigma_l\bar{\lambda}M_P^{-2} + i\bar{\psi}_k\bar{\sigma}_l\lambda M_P^{-2}) + \frac{i}{2}(\bar{\psi}_m\bar{\sigma}^m\xi + \psi_m\sigma^m\bar{\xi})\mathbf{D}M_P^{-1}, \quad (35)$$

$$\delta_\xi\tilde{A}_m = \frac{1}{2}(\bar{\lambda}\sigma_m\xi + \lambda\bar{\sigma}_m\bar{\xi}), \quad (36)$$

$$\delta_\xi\lambda^\alpha = (\xi\sigma^{nm})^\alpha(2i\tilde{F}_{nm} + \psi_n\sigma_m\bar{\lambda}M_P^{-1} + \bar{\psi}_n\bar{\sigma}_m\lambda M_P^{-1}) + i\xi^\alpha\mathbf{D}, \quad (37)$$

$$\delta_\xi\bar{\lambda}_{\dot{\alpha}} = (\bar{\xi}\bar{\sigma}^{nm})_{\dot{\alpha}}(2i\tilde{F}_{nm} + \psi_n\sigma_m\bar{\lambda}M_P^{-1} + \bar{\psi}_n\bar{\sigma}_m\lambda M_P^{-1}) - i\bar{\xi}_{\dot{\alpha}}\mathbf{D}. \quad (38)$$

Here the $U(1)$ fieldstrength

$$\tilde{F}_{kl} = \partial_k\tilde{A}_l - \partial_l\tilde{A}_k, \quad (39)$$

and the covariant derivatives

$$\tilde{\mathcal{D}}_m\lambda^\alpha = \partial_m\lambda^\alpha + \lambda^\beta\omega_{m\beta}^\alpha + \lambda^\alpha\tilde{A}_m, \quad (40)$$

$$\tilde{\mathcal{D}}_m\bar{\lambda}_{\dot{\alpha}} = \partial_m\bar{\lambda}_{\dot{\alpha}} + \bar{\lambda}_{\dot{\beta}}\omega_m^{\dot{\beta}\dot{\alpha}} - \bar{\lambda}_{\dot{\alpha}}\tilde{A}_m. \quad (41)$$

as well as

$$\tilde{\mathcal{D}}_m\xi^\alpha = \partial_m\xi^\alpha + \xi^\beta\omega_{m\beta}^\alpha + \xi^\alpha\tilde{A}_m, \quad (42)$$

$$\tilde{\mathcal{D}}_m\bar{\xi}_{\dot{\alpha}} = \partial_m\bar{\xi}_{\dot{\alpha}} + \bar{\xi}_{\dot{\beta}}\omega_m^{\dot{\beta}\dot{\alpha}} - \bar{\xi}_{\dot{\alpha}}\tilde{A}_m, \quad (43)$$

occur.

Observe that the sum of \mathcal{L}_1 and \mathcal{L}_{X^2} , which might be referred to as pure $U(1)$ supergravity, gives rise to a cosmological constant after diagonalization in the field \mathbf{D} , as discussed in [15,14,25]. In our case this action provides the starting point for the discussion of curvature squared terms and diagonalization should only be performed afterwards.

Curvature squared terms and $U(1)$ supergravity

As is well known [1] curvature squared terms in traditional supergravity are identified in the highest superfield

components of the combinations $W^{\alpha\beta\gamma}W_{\alpha\beta\gamma}$, G^aG_a and RR^\dagger of basic supergravity superfields. In the case of U(1) supergravity a number of modifications arise due to the presence of the $U(1)$ sector in the geometry, as explained in detail in [17]. In order to fix our notations we shall consider here the three superspace actions⁵

$$\mathcal{S}_{W^2} = \int \frac{E}{2R} \left(W^{\alpha\beta\gamma} W_{\alpha\beta\gamma} \right) + h.c. , \quad (44)$$

$$\mathcal{S}_{G^2+2RR^\dagger} = \int E (G^a G_a + 2RR^\dagger) , \quad (45)$$

$$\mathcal{S}_{RR^\dagger} = -3 \int E (36RR^\dagger) . \quad (46)$$

Complete component field expressions can be evaluated in using the generic component field action (23) with the identifications, respectively⁶,

$$r_{W^2} = W^{\alpha\beta\gamma} W_{\alpha\beta\gamma} , \quad (47)$$

$$r_{G^2+2RR^\dagger} = -\frac{1}{8} (\bar{\mathcal{D}}^2 - 8R) (G^a G_a + 2RR^\dagger) , \quad (48)$$

$$r_{RR^\dagger} = \frac{3}{8} (\bar{\mathcal{D}}^2 - 8R) (36RR^\dagger) , \quad (49)$$

for the generic chiral superfield r .

In what follows we shall only discuss the purely bosonic contributions of these actions. Following [17] one obtains

$$e^{-1} \mathcal{L}_{W^2} = \frac{1}{8} \mathcal{W}^{dc,ba} \mathcal{W}_{dc,ba} + \frac{1}{3} (F^{mn} F_{mn}) , \quad (50)$$

$$e^{-1} \mathcal{L}_{G^2+2RR^\dagger} = -\frac{1}{8} \tilde{\mathcal{R}}^{ba} \tilde{\mathcal{R}}_{ba} + \frac{1}{96} \mathcal{R}^2 - \frac{1}{6} \mathbf{D}^2 - \frac{1}{6} (F^{mn} F_{mn} + 2\tilde{F}^{mn} \tilde{F}_{mn}) , \quad (51)$$

$$e^{-1} \mathcal{L}_{RR^\dagger} = -\frac{3}{4} (\mathcal{R} - 2\mathbf{D})^2 + \left(b^m b_m + \frac{1}{2} M \bar{M} \right) \times \mathcal{R} - 2(b^m b_m + 2M \bar{M}) \mathbf{D} + 3\tilde{\mathcal{D}}^m M \tilde{\mathcal{D}}_m \bar{M} - 3(e_a{}^m \tilde{\mathcal{D}}_m b^a)^2 + ib^m (\bar{M} \tilde{\mathcal{D}}_m M - M \tilde{\mathcal{D}}_m \bar{M}) - \frac{1}{3} ((M \bar{M})^2 + M \bar{M} b^a b_a + (b^a b_a)^2) , \quad (52)$$

with the conventions

$$F_{mn} = \tilde{F}_{mn} + \frac{i}{2} B_{mn} , \quad (53)$$

$$B_{mn} = \partial_m b_n - \partial_n b_m , \quad (54)$$

⁵ The combination $G^a G_a + 2RR^\dagger$ which appears in the second action is particularly convenient for discussion of the supersymmetric Gauss–Bonnet invariant

⁶ Component formulation of a lagrangian $\int EX$, where X is a real and $U(1)$ invariant superfield, is obtained by using the generic lagrangian (23), with $r = -\frac{1}{8} (\bar{\mathcal{D}}^2 - 8R) X$, $(\bar{\mathcal{D}}^2 - 8R)$ is called the chiral projector

and where the Weyl tensor $\mathcal{W}_{dc,ba}$, the Ricci tensor $\tilde{\mathcal{R}}_{ca}$ and the curvature scalar \mathcal{R} are identified as usual in the decomposition of the Riemann tensor $\mathcal{R}_{dc,ba}$:

$$\mathcal{R}_{dc,ba} = \mathcal{W}_{dc,ba} + \frac{1}{2} \left(\eta_{db} \tilde{\mathcal{R}}_{ca} - \eta_{da} \tilde{\mathcal{R}}_{cb} - \eta_{cb} \tilde{\mathcal{R}}_{da} + \eta_{ca} \tilde{\mathcal{R}}_{db} \right) + \frac{1}{12} (\eta_{db} \eta_{ca} - \eta_{da} \eta_{cb}) \mathcal{R} . \quad (55)$$

With the first two lagrangians (50) and (51), one can obtain the supersymmetric version of the pure Gauss–Bonnet invariant (3) plus terms involving \mathbf{D}^2 and $\tilde{F}^{mn} \tilde{F}_{mn}$. These new contributions are due to the additional $U(1)$ sector, they can be cancelled by adding \mathcal{L}_{X^2} . As a result, the pure super–Gauss–Bonnet combination is:

$$\mathcal{L}_{GB} = 8\mathcal{L}_{W^2} + 16\mathcal{L}_{G^2+2RR^\dagger} + \frac{16}{3} \mathcal{L}_{X^2} . \quad (56)$$

In terms of component fields this reproduces exactly the combination of equation (3), i.e:

$$e^{-1} \mathcal{L}_{GB} = \mathcal{W}^{dc,ba} \mathcal{W}_{dc,ba} - 2\tilde{\mathcal{R}}^{ba} \tilde{\mathcal{R}}_{ba} + \frac{1}{6} \mathcal{R}^2 . \quad (57)$$

In order to discuss the most general form of $U(1)$ supergravity with curvature squared terms we shall consider the combination⁷

$$\mathcal{L}_{tot} = a_1 \mathcal{L}_1 + a_2 \mathcal{L}_{X^2} + a_3 \mathcal{L}_{RR^\dagger} + a_G \mathcal{L}_{G^2+2RR^\dagger} + a_W \mathcal{L}_{W^2} \quad (58)$$

with \mathcal{L}_1 and \mathcal{L}_{X^2} defined in the previous section. Inspection of the individual contributions shows that diagonalization in the auxiliary field \mathbf{D} of the $U(1)$ sector will introduce additional curvature scalar squared terms. More precisely, defining

$$\hat{\mathbf{D}} \equiv \mathbf{D} + \frac{c}{2} (a_1 + 3a_3 \mathcal{R} - 4a_3 M \bar{M} - 2a_3 b^m b_m) , \quad (59)$$

with

$$c = \left(-3a_3 + \frac{a_2}{2} - \frac{a_G}{6} \right)^{-1} , \quad (60)$$

gives rises to the component field lagrangian

$$e^{-1} \mathcal{L}_{tot} = -\frac{a_1}{2} (1 + 3a_3 c) \mathcal{R} - \frac{a_1}{3} (1 - 6a_3 c) M \bar{M} + \frac{a_1}{3} (1 + 3a_3 c) b^a b_a + \frac{a_W}{8} \mathcal{W}^{dc,ba} \mathcal{W}_{dc,ba} - \frac{a_G}{8} \tilde{\mathcal{R}}^{ba} \tilde{\mathcal{R}}_{ba} + \left(-\frac{3a_3}{4} (1 + 3a_3 c) + \frac{a_G}{96} \right) \mathcal{R}^2 + \left(a_2 - \frac{a_G}{3} \right) \tilde{F}^{mn} \tilde{F}_{mn} + \frac{1}{6} (2a_W - a_G) F^{mn} F_{mn} + 3a_3 \tilde{\mathcal{D}}^m M \tilde{\mathcal{D}}_m \bar{M} + c^{-1} \hat{\mathbf{D}}^2 - 3a_3 (e_a{}^m \tilde{\mathcal{D}}_m b^a)^2$$

⁷ a_i are real constants

$$\begin{aligned}
& +ia_3b^m\left(\overline{M}\tilde{D}_mM - M\tilde{D}_m\overline{M}\right) \\
& +\frac{a_3}{2}(1+12a_3c)\mathcal{R}M\overline{M} \\
& +a_3(1+3a_3c)\mathcal{R}b^ab_a - \frac{a_1^2c}{4} \\
& -\frac{a_3}{3}(1+12a_3c)((M\overline{M})^2 + M\overline{M}b^ab_a) \\
& -\frac{a_3}{3}(1+3a_3c)(b^ab_a)^2. \tag{61}
\end{aligned}$$

The Lorentz and $U(1)$ covariant derivatives appearing here are defined as

$$\tilde{D}_mM = \partial_mM + 2\tilde{A}_mM, \tag{62}$$

$$\tilde{D}_mb^a = \partial_mb^a + b^c\omega_{mc}{}^a, \tag{63}$$

according to the chiral weights:

$$\omega(M) = 2, \quad \omega(b^a) = 0. \tag{64}$$

This is the (bosonic part) of the component field lagrangian which is relevant for the discussion of curvature squared terms in $U(1)$ supergravity. As a first observation consider the special case

$$a_G = 2a_W, \tag{65}$$

which adjusts the relative factor between the squares of the Weyl and the Ricci tensors to that occurring in the Gauss–Bonnet combination. In this case the general action is specified to

$$\begin{aligned}
e^{-1}\mathcal{L}_{tot} = & -\frac{a_1}{2}(1+3a_3c)\mathcal{R} \\
& -\frac{a_1}{3}(1-6a_3c)M\overline{M} + \frac{a_1}{3}(1+3a_3c)b^ab_a \\
& +\frac{a_W}{8}\left(\mathcal{W}^{dc,ba}\mathcal{W}_{dc,ba} - 2\tilde{\mathcal{R}}^{ba}\tilde{\mathcal{R}}_{ba} + \frac{1}{6}\mathcal{R}^2\right) \\
& -\frac{3a_3}{4}(1+3a_3c)\mathcal{R}^2 + 2c^{-1}(1+3a_3c)\tilde{F}^{mn}\tilde{F}_{mn} \\
& +3a_3\tilde{D}^mM\tilde{D}_m\overline{M} + c^{-1}\hat{\mathbf{D}}^2 - 3a_3\left(e_a{}^m\tilde{D}_mb^a\right)^2 \\
& +ia_3b^m\left(\overline{M}\tilde{D}_mM - M\tilde{D}_m\overline{M}\right) \\
& -\frac{a_3}{3}(1+12a_3c)((M\overline{M})^2 + M\overline{M}b^ab_a) \\
& -\frac{a_3}{3}(1+3a_3c)(b^ab_a)^2 \\
& +\frac{a_3}{2}(1+12a_3c)\mathcal{R}M\overline{M} \\
& +a_3(1+3a_3c)\mathcal{R}b^ab_a - \frac{a_1^2c}{4}. \tag{66}
\end{aligned}$$

This action is the $U(1)$ supergravity analogue of the case $\beta = 0$ discussed for the non-supersymmetric case (5).

As an aside, note that in order to obtain the Gauss–Bonnet combination of curvature squared terms one has to cancel the additional \mathcal{R}^2 term. This can be done by choosing either $a_3 = 0$ or $(1+3a_3c) = 0$. In the first case, the lagrangian is reduced to

$$e^{-1}\mathcal{L}_{tot} = -\frac{a_1}{2}\mathcal{R} - \frac{a_1}{3}M\overline{M} + \frac{a_1}{3}b^ab_a$$

$$\begin{aligned}
& +2c^{-1}\tilde{F}^{mn}\tilde{F}_{mn} + c^{-1}\hat{\mathbf{D}}^2 - \frac{a_1^2c}{4} \\
& +\frac{a_W}{8}\left(\mathcal{W}^{dc,ba}\mathcal{W}_{dc,ba} - 2\tilde{\mathcal{R}}^{ba}\tilde{\mathcal{R}}_{ba} + \frac{1}{6}\mathcal{R}^2\right), \tag{67}
\end{aligned}$$

and in the second case one obtains

$$\begin{aligned}
e^{-1}\mathcal{L}_{tot} = & -a_1M\overline{M} - \frac{3a_3}{2}M\overline{M}\mathcal{R} + 3a_3\tilde{D}^mM\tilde{D}_m\overline{M} \\
& -3a_3\hat{\mathbf{D}}^2 - 3a_3\left(e_a{}^m\tilde{D}_mb^a\right)^2 \\
& +ia_3b^m\left(\overline{M}\tilde{D}_mM - M\tilde{D}_m\overline{M}\right) \\
& +\frac{a_1^2}{12a_3} + a_3((M\overline{M})^2 + M\overline{M}b^ab_a) \\
& +\frac{a_W}{8}\left(\mathcal{W}^{dc,ba}\mathcal{W}_{dc,ba} - 2\tilde{\mathcal{R}}^{ba}\tilde{\mathcal{R}}_{ba} + \frac{1}{6}\mathcal{R}^2\right). \tag{68}
\end{aligned}$$

Clearly, the first case describes a generalization of a supergravity action with a correctly normalized (for $a_1 = 1$) curvature scalar term.

The interpretation of the second case is more subtle in that a Weyl rescaling in $M\overline{M}$ should be performed [6, 9] to arrive at a correctly normalized Einstein term.

Finally, the coupling of curvature squared term to traditional supergravity can be recovered from equation (61) in simply switching off the $U(1)$ sector, i.e taking $a_2 = 0$, $\tilde{A}_m = 0$ and substituting $\hat{\mathbf{D}} \equiv \frac{c}{2}(a_1 + 3a_3\mathcal{R} - 4a_3M\overline{M} - 2a_3b^mb_m)$, which eliminates the c -dependence in (61).

Coupling to matter

As chirally extended $U(1)$ supergravity provides a natural framework for a gravity coupled Fayet–Iliopoulos term [14, 15], it is interesting to investigate couplings to chiral matter in this context. This discussion serves at the same time as a prerequisite for the generalization of the Whitt mechanism, as alluded to in the first section, to the case of $U(1)$ supergravity.

To begin with, we consider a single chiral superfield Φ of $U(1)$ weight $\omega(\Phi) = \omega$, and, correspondingly $\bar{\Phi}$ of $U(1)$ weight $\omega(\bar{\Phi}) = -\omega$. Evaluating the supersymmetric action

$$\begin{aligned}
\mathcal{S} = & a_1\mathcal{S}_1 + a_2\mathcal{S}_{X^2} + a_4\int Ef(\Phi, \bar{\Phi}) \\
& + a_5\left(\int \frac{E}{2R}\Phi^x + \int \frac{E}{2R^\dagger}\bar{\Phi}^x\right), \tag{69}
\end{aligned}$$

where x is given in term of the chiral weight: $x = \frac{\omega(R)}{\omega(\Phi)}$ for $\omega(\Phi) \neq 0$. In terms of component fields, one finds, for the purely bosonic contribution,

$$\begin{aligned}
e^{-1}\mathcal{L} = & (a_1 + a_4f(A, \bar{A}))\left(-\frac{1}{2}\mathcal{R} - \frac{1}{3}M\overline{M} + \frac{1}{3}b^mb_m\right) \\
& + \left(a_1 + a_4f(A, \bar{A}) - \frac{3}{4}a_4\omega(f_AA + f_{\bar{A}}\bar{A})\right)\mathbf{D}
\end{aligned}$$

$$\begin{aligned}
& + \frac{a_2}{2} \mathbf{D}^2 + a_2 \tilde{F}^{mn} \tilde{F}_{mn} \\
& + i a_4 b^m \left(f_A \tilde{\mathcal{D}}_m A - f_{\bar{A}} \tilde{\mathcal{D}}_m \bar{A} \right) \\
& + 3 a_4 f_{A\bar{A}} \left(\tilde{\mathcal{D}}^m A \tilde{\mathcal{D}}_m \bar{A} - F \bar{F} \right) \\
& + a_4 \left(f_A M F + f_{\bar{A}} \bar{M} \bar{F} \right) \\
& + a_5 x \left(F A^{(x-1)} + \bar{F} \bar{A}^{(x-1)} \right), \tag{70}
\end{aligned}$$

with the definitions

$$|\Phi\rangle = A, \quad \mathcal{D}^2 |\Phi\rangle = -4F. \tag{71}$$

Clearly, this provides a generalization of the Fayet–Iliopoulos term to the case of matter–coupled chirally extended supergravity, and possible applications to symmetry breaking mechanisms deserve further study.

On the other hand, this action is the starting point for the generalization of the Whitt’s mechanism as well. In order to establish the relation between curvature squared $U(1)$ supergravity with its matter coupled counterpart, linear in the curvature scalar, we shall start from (66) with the particular choice $a_G = a_W = 0$, i.e

$$\mathcal{L} = a_1 \mathcal{L}_1 + a_2 \mathcal{L}_{X^2} + a_3 \mathcal{L}_{RR^\dagger}, \tag{72}$$

which is a supersymmetric version of (5). As to the supersymmetric analogue of (6) we consider

$$\begin{aligned}
\mathcal{S} = & a_1 \mathcal{S}_1 + a_2 \mathcal{S}_{X^2} + a_3 \left(-3 \int E (\Phi \bar{\Phi} + \Lambda + \bar{\Lambda}) \right. \\
& \left. + \int \frac{E}{2R} \Lambda \Phi + \int \frac{E}{2R^\dagger} \bar{\Lambda} \bar{\Phi} \right), \tag{73}
\end{aligned}$$

which has the same appearance as the corresponding action in traditional supergravity [8]. However, in the present context a number of new features appear. In particular, for reasons of consistency with the $U(1)$ gauge structure, the $U(1)$ weights of the chiral superfields Φ and Λ are determined to be:

$$\omega(\Phi) = 2, \quad \omega(\Lambda) = 0. \tag{74}$$

Evaluation of the purely bosonic part of this action in terms of component fields gives

$$\begin{aligned}
e^{-1} \mathcal{L} = & (a_1 + a_3 (A \bar{A} + B + \bar{B})) \\
& \times \left(-\frac{1}{2} \mathcal{R} - \frac{1}{3} M \bar{M} + \frac{1}{3} b^m b_m \right) \\
& + \left(a_1 + a_3 \left(1 - \frac{3}{2} \omega(\Phi) \right) A \bar{A} + a_3 (B + \bar{B}) \right) \mathbf{D} \\
& + \frac{a_2}{2} \mathbf{D}^2 + a_2 \tilde{F}^{mn} \tilde{F}_{mn} \\
& + i a_3 b^m \left(\bar{A} \tilde{\mathcal{D}}_m A - A \tilde{\mathcal{D}}_m \bar{A} + \tilde{\mathcal{D}}_m B - \tilde{\mathcal{D}}_m \bar{B} \right) \\
& + 3 a_3 \tilde{\mathcal{D}}^m A \tilde{\mathcal{D}}_m \bar{A} \\
& + a_3 \left(-3 F \bar{F} + \bar{A} M F + A \bar{M} \bar{F} + F B + \bar{F} \bar{B} \right) \\
& + a_3 \left(G(M + A) + \bar{G}(\bar{M} + \bar{A}) \right. \\
& \left. - A \bar{B} \bar{M} - \bar{A} B M \right), \tag{75}
\end{aligned}$$

with the definitions:

$$|\Lambda\rangle = B, \quad \mathcal{D}^2 |\Lambda\rangle = -4G. \tag{76}$$

Taking into account the equations of motion⁸

$$A = -M, \tag{77}$$

$$\begin{aligned}
F = & A \bar{M} + \frac{1}{2} \mathcal{R} - \frac{1}{3} b^m b_m \\
& + \frac{1}{3} M \bar{M} + i e_a{}^m \tilde{\mathcal{D}}_m b^a - \mathbf{D}, \tag{78}
\end{aligned}$$

reproduces exactly the lagrangian (66) of curvature squared $U(1)$ supergravity.

On the other hand, performing appropriately a Weyl rescaling in the supersymmetric context [23, 9], this action will describe supergravity, with a properly normalized curvature scalar, coupled to two chiral matter multiplets in the presence of a Fayet–Iliopoulos term.

In conclusion, we expect that this mechanism might open new possibilities for scenarios of gravity induced supersymmetry breaking in the presence of curvature squared terms [9].

Acknowledgements. I would like to thank R.Grimm for suggestions and helpful discussions about this work. Thanks also to G.Girardi for conversations on this subject.

References

1. S. Ferrara, B. Zumino. Structure of linearized supergravity and conformal supergravity. Nucl. Phys. **B134** (1978), 301
2. M. Kaku, P.K. Townsend, P. van Nieuwenhuizen. Properties of conformal supergravity. Phys. Rev. **D17** (1978), 3179
3. S. Cecotti, S. Ferrara, L. Girardello, A. Pasquinucci, M. Porrati. Matter coupling in higher–derivative supergravity. Phys. Rev. **D33** (1986), 2504
4. S. Theisen. Fourth order supergravity. Nucl. Phys. **B263** (1986), 687
5. M. de Roo, A. Wiedemann, E. Zijstra. The construction of R^2 actions in $d = 4, n = 1$ supergravity. Class. Quantum Grav. **7** (1990), 1181
6. S. Cecotti, S. Ferrara, L. Girardello, A. Pasquinucci, M. Porrati. Matter coupled supergravity with Gauss–Bonnet invariants: component lagrangian and supersymmetry breaking. Int. J. Mod. Phys. **A3** (1988), 1675
7. B. Whitt. Fourth–order gravity as general relativity plus matter. Phys. Lett. **145B** (1984), 176
8. S. Cecotti. Higher derivative supergravity is equivalent to standard supergravity coupled to matter. Phys. Lett. **190B** (1987), 86
9. A. Hindawi, B.A. Ovrut, D. Waldram. Soft supersymmetry breaking induced by higher–derivative supergravitation in the electroweak standard model. Phys. Lett. **381B** (1996), 154

⁸ In superfield language this corresponds to varying (73) with respect to (the prepotential of) the chiral superfield Λ

10. A. Hindawi, B.A. Ovrut, D. Waldram. Higher derivative gravitation and a new mechanism for supersymmetry breaking in four dimensions. *Prog. Theor. Phys. Suppl.* **123** (1996), 397
11. A. Hindawi, B.A. Ovrut, D. Waldram. Four-dimensional higher derivative supergravity and spontaneous supersymmetry breaking. *Nucl. Phys.* **B476** (1996), 175
12. P. Howe. Supergravity in superspace. *Nucl. Phys.* **B199** (1982), 309
13. M. Müller. Supergravity in $U(1)$ superspace with a two-form gauge potential. *Nucl. Phys.* **B264** (1986), 292
14. D.Z. Freedman. Supergravity with axial-gauge invariance. *Phys. Rev.* **D15** (1977), 1173
15. B. De Wit, P. van Nieuwenhuizen. The auxiliary field structure in chirally extended supergravity. *Nucl. Phys.* **B139** (1978), 216
16. K.S. Stelle. Classical gravity with higher derivatives. *Gen. Rel. Grav.* **9,4** (1978), 353
17. G. Girardi, R. Grimm. The superspace geometry of gravitational Chern-Simons forms and their couplings to linear multiplets: a review. In preparation
18. G. 't Hooft, M. Veltman. One loop divergencies in the theory of gravitation. *Ann. Inst. Henri Poincaré* **20** (1974), 69
19. S. Deser, P. van Nieuwenhuizen. One loop divergences of quantized Einstein-Maxwell fields. *Phys. Rev.* **D10** (1974), 401
20. K.S. Stelle. Renormalization of higher-derivative quantum gravity. *Phys. Rev.* **D16** (1977), 953
21. N.H. Barth, S.M. Christensen. Quantizing fourth-order gravity theories: The functional integral. *Phys. Rev.* **D28** (1983), 1876
22. P. Fayet, J. Iliopoulos. Spontaneously broken supergauge symmetries and Goldstone spinors. *Phys. Lett.* **51B** (1974), 461
23. J. Wess, J. Bagger. *Supersymmetry and Supergravity*. Princeton Series in Physics. Princeton University Press, Princeton, 1983. 2nd edition 1992
24. P. Binétruy, G. Girardi, R. Grimm. *Supergravity and matter: a geometric formulation*. LAPP-TH-275/90, August 1990
25. B. Zumino. Supersymmetry and the vacuum. *Nucl. Phys.* **B89** (1975), 535