

Extracting CKM phases and B_s – \bar{B}_s mixing parameters from angular distributions of non-leptonic B decays

A.S. Dighe¹, I. Dunietz², R. Fleischer³

¹ The Abdus Salam International Centre for Theoretical Physics, I-34100 Trieste, Italy

² Theoretical Physics Division, Fermi National Accelerator Laboratory, Batavia, IL 60510, USA

³ Theory Division, CERN, CH-1211 Geneva 23, Switzerland

Received: 8 April 1998 / Published online: 21 August 1998

Abstract. Suggestions for efficiently determining the lifetimes and mass difference of the light and heavy B_s mesons (B_s^L, B_s^H) from $B_s \rightarrow J/\psi\phi, D_s^{*+}D_s^{*-}$ decays are given. Using appropriate weighting functions for the angular distributions of the decay products (*moment analysis*), one can extract $(\Gamma_H, \Gamma_L, \Delta m)_{B_s}$. Such a moment analysis allows the determination of the relative magnitudes and phases of the CP-odd and CP-even decay amplitudes. Efficient determinations of CP-violating effects occurring in $B_s \rightarrow J/\psi\phi, D_s^{*+}D_s^{*-}$ are discussed in the light of a possible width difference $(\Delta\Gamma)_{B_s}$, and the utility of this method for $B \rightarrow J/\psi K^*, D_s^{*+}\bar{D}^*$ decays is noted. Since our approach is very general, it can in principle be applied to all kinds of angular distributions and allows the determination of all relevant observables, including fundamental CKM (Cabibbo–Kobayashi–Maskawa) parameters, as well as tests of various aspects of the factorization hypothesis. Explicit angular distributions and weighting functions are given, and the general method that can be used for any angular distribution is indicated.

1 Introduction

Strategies for obtaining experimental insights into CP violation and non-factorizable contributions to weak decays are of particular interest for present particle physics. The observables of angular distributions can be obtained in an efficient way by using an *angular moment analysis* [1]–[3]. In this approach, the observed experimental data are weighted by judiciously chosen functions, which project out any desired observable. This strategy is an alternative to the usual *likelihood fit* method [4]. It is demonstrated that the moment analysis extracts all observables of measured angular distributions, such as the ones occurring in weak decays of pseudoscalars [$P \rightarrow V\ell\nu, X_J\ell\nu, VV$, etc.]. This method is of general validity. In our present paper, we apply the formalism to angular distributions [5, 6] of B_s and B meson decays into two vector-meson final states that are caused by $\bar{b} \rightarrow \bar{s}c\bar{c}$ quark-level transitions. By making use of the general formalism outlined in this paper, it is straightforward to derive weighting functions for other exclusive mesonic or baryonic transitions, governed for instance by $b \rightarrow \bar{c}ud, cl\bar{\nu}, ul\bar{\nu}, c \rightarrow \bar{s}du, sl^+\nu, dl^+\nu$.

The mixing between neutral B_s mesons is expected to give rise to CP-even (B_s^L) and CP-odd (B_s^H) mass eigenstates, which may have a perceptible width difference $\Delta\Gamma \equiv \Gamma_H - \Gamma_L$ [7]. Using appropriate weighting functions for the angular distributions of the decay products in the transitions $B_s \rightarrow J/\psi\phi$ and/or $B_s \rightarrow D_s^{*+}D_s^{*-}$, one can extract $(\Gamma_H, \Gamma_L, \Delta m)_{B_s}$.

A characteristic feature of the angular distributions considered in this paper is the fact that they contain terms describing interference effects between CP-even and CP-odd final-state configurations. Because of the lifetime difference, these contributions give rise to a term in the time evolution of the *untagged* rate, which is proportional to [8]:

$$(e^{-\Gamma_H t} - e^{-\Gamma_L t}) \sin \phi_{\text{CKM}}, \quad (1)$$

where ϕ_{CKM} is a weak phase that is introduced through the CKM matrix [9]. In the B_s decays considered in the present paper, ϕ_{CKM} is related to the Wolfenstein parameter η [10]. It is a remarkable feature that time-evolved *untagged* data samples of angular distributions of B_s decays may exhibit CP-violating effects, if $\Delta\Gamma$ is sizeable [8, 11]. This feature may be important, because it provides an alternative to previous investigations, which have shown how to extract $\sin \phi_{\text{CKM}}$ from tagged, time-dependent analyses [12, 13]. This extraction, however, may not be feasible in the near future because it requires tagging and superb vertex detectors, which must resolve the rapid $\Delta m t$ oscillations. In contrast, any dependence on $\Delta m t$ cancels in untagged data samples, which therefore allow feasibility studies with current vertex technology [14].

Concerning tests of the *factorization hypothesis* [15]–[20], we divide the $\bar{b} \rightarrow \bar{s}c\bar{c}$ modes into the following two categories:

- colour-suppressed decays: $B_q \rightarrow J/\psi V$ with $(q, V) \in \{(s, \phi); (d, K^{*0}); (u, K^{*+})\}$ [21, 22].

– colour-allowed decays: $B_q \rightarrow D_s^{*+} \bar{D}_q^*$ with $q \in \{s, d, u\}$ [23,24].

Whereas the validity of the factorization assumption is very doubtful in the colour-suppressed case, it should work much better for the colour-allowed channels because of colour transparency [18]. The latter have furthermore rather tight restrictions from the Heavy Quark Effective Theory (HQET) [25] for the form factors describing the “factorized” hadronic matrix elements of the relevant four-quark operators [23,24,26].

Our paper is organized as follows: in Sect.2 we calculate the transition matrix elements and observables of the angular distributions by using an appropriate low-energy effective Hamiltonian. There we also give estimates for these observables, allowing a comparison with experimental data. The efficient experimental determination of these observables is the subject of Sect.3, where we shall discuss the angular moment analysis. Sections 4 and 5 are devoted to the angular correlations in the colour-suppressed decays $B_s \rightarrow J/\psi \phi$, $B \rightarrow J/\psi K^*$ and the colour-allowed decays $B_s \rightarrow D_s^{*+} D_s^{*-}$, $B \rightarrow D_s^{*+} \bar{D}^*$, respectively. There we give the time evolutions of the angular distributions, appropriate weighting functions, and discuss CP-violating effects. Finally in Sect.6 the main results are summarized.

2 Transition matrix elements and observables

Before we present an efficient method for extracting the observables of the angular distributions from experimental data – the *angular moment analysis* – let us discuss in this section how these observables are calculated and what orders of magnitude we expect for them.

2.1 General aspects

In order to calculate the decay amplitudes of the $\bar{b} \rightarrow \bar{s} c \bar{c}$ transitions considered in this paper, we use an appropriate low-energy effective Hamiltonian, which has the following structure:

$$\mathcal{H}_{\text{eff}} = \frac{G_F}{\sqrt{2}} \left[\sum_{j=u,c} \lambda_j^{(s)} \left\{ Q_1^j C_1(\mu) + Q_2^j C_2(\mu) + \sum_{k=3}^{10} Q_k C_k(\mu) \right\} \right]. \quad (2)$$

Here the quantities $\lambda_j^{(s)} \equiv V_{js} V_{jb}^*$ denote CKM factors,

$$\begin{aligned} Q_1^c &= (\bar{c}_\alpha s_\beta)_{V-A} (\bar{b}_\beta c_\alpha)_{V-A} \\ Q_2^c &= (\bar{c}_\alpha s_\alpha)_{V-A} (\bar{b}_\beta c_\beta)_{V-A} \\ Q_1^u &= (\bar{u}_\alpha s_\beta)_{V-A} (\bar{b}_\beta u_\alpha)_{V-A} \\ Q_2^u &= (\bar{u}_\alpha s_\alpha)_{V-A} (\bar{b}_\beta u_\beta)_{V-A} \end{aligned} \quad (3)$$

are “current–current” operators,

$$\begin{aligned} Q_3 &= (\bar{b}_\alpha s_\alpha)_{V-A} \sum_{q=u,d,s,c,b} (\bar{q}_\beta q_\beta)_{V-A} \\ Q_4 &= (\bar{b}_\alpha s_\beta)_{V-A} \sum_{q=u,d,s,c,b} (\bar{q}_\beta q_\alpha)_{V-A} \\ Q_5 &= (\bar{b}_\alpha s_\alpha)_{V-A} \sum_{q=u,d,s,c,b} (\bar{q}_\beta q_\beta)_{V+A} \\ Q_6 &= (\bar{b}_\alpha s_\beta)_{V-A} \sum_{q=u,d,s,c,b} (\bar{q}_\beta q_\alpha)_{V+A} \end{aligned} \quad (4)$$

describe QCD penguins, while the operators

$$\begin{aligned} Q_7 &= \frac{3}{2} (\bar{b}_\alpha s_\alpha)_{V-A} \sum_{q=u,d,s,c,b} e_q (\bar{q}_\beta q_\beta)_{V+A} \\ Q_8 &= \frac{3}{2} (\bar{b}_\alpha s_\beta)_{V-A} \sum_{q=u,d,s,c,b} e_q (\bar{q}_\beta q_\alpha)_{V+A} \\ Q_9 &= \frac{3}{2} (\bar{b}_\alpha s_\alpha)_{V-A} \sum_{q=u,d,s,c,b} e_q (\bar{q}_\beta q_\beta)_{V-A} \\ Q_{10} &= \frac{3}{2} (\bar{b}_\alpha s_\beta)_{V-A} \sum_{q=u,d,s,c,b} e_q (\bar{q}_\beta q_\alpha)_{V-A} \end{aligned} \quad (5)$$

are “electroweak” penguin operators. Here $V \pm A$ corresponds to $\gamma_\mu (\hat{1} \pm \gamma_5)$ quark currents, Greek indices are associated with the $SU(3)_C$ quark-colour, and the quantities e_q arising in the expressions for the electroweak penguin operators label the electrical quark charges. Nowadays, the Wilson coefficient functions $C_k(\mu)$ of the low-energy effective Hamiltonian \mathcal{H}_{eff} , where $\mu = \mathcal{O}(m_b)$ denotes the usual renormalization scale, are known beyond the leading logarithmic approximation [27].

Since $\lambda_u^{(s)}$ is suppressed with respect to $\lambda_c^{(s)}$ by a CKM factor $\lambda^2 R_b$, where $\lambda = 0.22$ is the Wolfenstein parameter [10] and

$$R_b \equiv \frac{1}{\lambda} \frac{|V_{ub}|}{|V_{cb}|} \quad (6)$$

is constrained by present experimental data to lie within the range $R_b = 0.36 \pm 0.08$ [28,29], and since furthermore the current–current operators Q_1^u, Q_2^u may contribute only through penguin-like matrix elements to $\bar{b} \rightarrow \bar{s} c \bar{c}$ modes, the corresponding transition amplitudes are dominated to an *excellent approximation* by the contribution proportional to $\lambda_c^{(s)}$ (for a detailed discussion, see [30]). In the penguin operators, we neglect the parts of flavour structure different from $(\bar{c}c)(\bar{b}s)$. Then the number of relevant operators reduces from ten to four and the structure of the decay amplitude simplifies considerably.

In order to implement the factorization hypothesis by factorizing the hadronic matrix elements of the four-quark operators Q_k into hadronic matrix elements of quark currents, we have to perform suitable Fierz transformations of the operator basis specified in (3)–(5). Beyond the leading logarithmic approximation one has to be very careful in performing such Fierz transformations, as the Wilson

coefficients depend both on the form of the chosen operator basis and on the applied renormalization scheme [27]. Since we do not use any specific Wilson coefficients to obtain numerical estimates in this paper, we may perform such Fierz transformations and will use a tilde ($\tilde{}$) to indicate Fierz-transformed operators. For a discussion of the renormalization-scheme dependences arising beyond the leading logarithmic approximation and their consistent cancellation in the physical transition amplitudes through certain one-loop matrix elements at $\mu = \mathcal{O}(m_b)$, the reader is referred to [31].

Let us, in the following two subsections, investigate the structure of the hadronic matrix elements of the low-energy effective Hamiltonian (2) for the exclusive colour-suppressed and colour-allowed decays $B_s \rightarrow J/\psi \phi$, $B \rightarrow J/\psi K^*$ and $B_s \rightarrow D_s^{*+} D_s^{*-}$, $B \rightarrow D_s^{*+} \bar{D}^*$, respectively.

2.2 Colour-suppressed decays

If we perform a Fierz transformation of the current-current operators specified in (3), the decay amplitude for $B_q \rightarrow J/\psi V$ ($(q, V) \in \{(s, \phi); (d, K^{*0}); (u, K^{*+})\}$) can be written in the following form:

$$\begin{aligned} & \langle J/\psi(\lambda)V(\lambda) | \mathcal{H}_{\text{eff}} | B_q \rangle \\ &= \frac{G_F}{\sqrt{2}} V_{cs} V_{cb}^* \left[\mathcal{C}_1^{\text{eff}}(\mu) \langle J/\psi(\lambda)V(\lambda) | \tilde{Q}_1^c(\mu) | B_q \rangle \right. \\ & \quad + \mathcal{C}_{1,\text{oct}}^{\text{eff}}(\mu) \langle J/\psi(\lambda)V(\lambda) | \tilde{Q}_{1,\text{oct}}^c(\mu) | B_q \rangle \\ & \quad + \mathcal{C}_5^{\text{eff}}(\mu) \langle J/\psi(\lambda)V(\lambda) | Q_5^c(\mu) | B_q \rangle \\ & \quad \left. + \mathcal{C}_{5,\text{oct}}^{\text{eff}}(\mu) \langle J/\psi(\lambda)V(\lambda) | Q_{5,\text{oct}}^c(\mu) | B_q \rangle \right], \quad (7) \end{aligned}$$

where λ denotes the helicities of the final-state vector mesons and the ‘‘effective’’ Wilson coefficient functions are given by

$$\begin{aligned} \mathcal{C}_1^{\text{eff}}(\mu) &\equiv C_1(\mu) + \frac{1}{3}C_2(\mu) + C_3(\mu) \\ & \quad + \frac{1}{3}C_4(\mu) + C_9(\mu) + \frac{1}{3}C_{10}(\mu) \quad (8) \end{aligned}$$

$$\mathcal{C}_{1,\text{oct}}^{\text{eff}}(\mu) \equiv 2[C_2(\mu) + C_4(\mu) + C_{10}(\mu)] \quad (9)$$

$$\mathcal{C}_5^{\text{eff}}(\mu) \equiv C_5(\mu) + \frac{1}{3}C_6(\mu) + C_7(\mu) + \frac{1}{3}C_8(\mu) \quad (10)$$

$$\mathcal{C}_{5,\text{oct}}^{\text{eff}}(\mu) \equiv 2[C_6(\mu) + C_8(\mu)]. \quad (11)$$

The μ -dependence of these Wilson coefficients is cancelled by that of the hadronic matrix elements appearing in (7). In deriving the transition matrix element in (7), we have used the relations

$$\tilde{Q}_2^c = \frac{1}{3}\tilde{Q}_1^c + 2\tilde{Q}_{1,\text{oct}}^c \quad (12)$$

$$Q_6^c = \frac{1}{3}Q_5^c + 2Q_{5,\text{oct}}^c \quad (13)$$

with

$$\tilde{Q}_1^c = (\bar{c}_\alpha c_\alpha)_{V-A} (\bar{b}_\beta s_\beta)_{V-A} \quad (14)$$

$$\tilde{Q}_{1,\text{oct}}^c = (\bar{c}_\alpha T_{\alpha\beta}^a c_\beta)_{V-A} (\bar{b}_\gamma T_{\gamma\delta}^a s_\delta)_{V-A} \quad (15)$$

and

$$Q_5^c = (\bar{c}_\alpha c_\alpha)_{V+A} (\bar{b}_\beta s_\beta)_{V-A} \quad (16)$$

$$Q_{5,\text{oct}}^c = (\bar{c}_\alpha T_{\alpha\beta}^a c_\beta)_{V+A} (\bar{b}_\gamma T_{\gamma\delta}^a s_\delta)_{V-A}. \quad (17)$$

Here the 3×3 matrices T^a are the $SU(3)_C$ generators, normalized to $\text{tr}(T^a T^b) = \delta^{ab}/2$. As we will see below, the form of the Fierz-transformed operators given above is better suited to analyse the $B_q \rightarrow J/\psi V$ decays since the J/ψ is related to the $(\bar{c}c)$ pieces. The penguin contributions to $\mathcal{C}_1^{\text{eff}}(\mu)$ and $\mathcal{C}_{1,\text{oct}}^{\text{eff}}(\mu)$ are at most $\mathcal{O}(10\%)$ and $\mathcal{O}(1\%)$, respectively, as can be estimated from the values of their Wilson coefficients [27].

If one assumes that J/ψ emerges from the vector parts of the $(\bar{c}c)_{V\pm A}$ quark currents appearing in the operators in (14)–(17), the matrix elements of $\tilde{Q}_{1,(\text{oct})}^c$ and $\tilde{Q}_{5,(\text{oct})}^c$ will be equal and the decay amplitude (7) can be simplified considerably. Moreover, within the framework of naive factorization, we obtain (analogous for Q_5^c and $Q_{5,\text{oct}}^c$):

$$\begin{aligned} & \langle J/\psi(\lambda)V(\lambda) | \tilde{Q}_1^c | B_q \rangle_f \\ &= \langle J/\psi(\lambda) | (\bar{c}c)_{V-A} | 0 \rangle \langle V(\lambda) | (\bar{b}s)_{V-A} | B_s \rangle \quad (18) \end{aligned}$$

$$\begin{aligned} & \langle J/\psi(\lambda)V(\lambda) | \tilde{Q}_{1,\text{oct}}^c | B_q \rangle_f \\ &= \langle J/\psi(\lambda) | (\bar{c}T^a c)_{V-A} | 0 \rangle \langle V(\lambda) | (\bar{b}T^a s)_{V-A} | B_s \rangle, \quad (19) \end{aligned}$$

where summation over colour-indices is understood implicitly. Consequently, since J/ψ is a colour-singlet state, the factorized hadronic matrix elements of the colour-octet operators given in (19) vanish.

2.3 Colour-allowed decays

In the case of the colour-allowed decays $B_q \rightarrow D_s^{*+} \bar{D}_q^*$ ($q \in \{u, d, s\}$), the transition amplitude can be written in a way that is completely analogous to (7):

$$\begin{aligned} & \langle D_s^{*+}(\lambda) \bar{D}_q^*(\lambda) | \mathcal{H}_{\text{eff}} | B_q \rangle \\ &= \frac{G_F}{\sqrt{2}} V_{cs} V_{cb}^* \left[\mathcal{C}_2^{\text{eff}}(\mu) \langle D_s^{*+}(\lambda) \bar{D}_q^*(\lambda) | Q_2^c(\mu) | B_q \rangle \right. \\ & \quad + \mathcal{C}_{2,\text{oct}}^{\text{eff}}(\mu) \langle D_s^{*+}(\lambda) \bar{D}_q^*(\lambda) | Q_{2,\text{oct}}^c(\mu) | B_q \rangle \\ & \quad + \mathcal{C}_6^{\text{eff}}(\mu) \langle D_s^{*+}(\lambda) \bar{D}_q^*(\lambda) | \tilde{Q}_6^c(\mu) | B_q \rangle \\ & \quad \left. + \mathcal{C}_{6,\text{oct}}^{\text{eff}}(\mu) \langle D_s^{*+}(\lambda) \bar{D}_q^*(\lambda) | \tilde{Q}_{6,\text{oct}}^c(\mu) | B_q \rangle \right]. \quad (20) \end{aligned}$$

The corresponding effective Wilson coefficient functions are, however, very different:

$$\begin{aligned} \mathcal{C}_2^{\text{eff}}(\mu) &\equiv \frac{1}{3}C_1(\mu) + C_2(\mu) \\ & \quad + \frac{1}{3}C_3(\mu) + C_4(\mu) + \frac{1}{3}C_9(\mu) + C_{10}(\mu) \quad (21) \end{aligned}$$

$$\mathcal{C}_{2,\text{oct}}^{\text{eff}}(\mu) \equiv 2[C_1(\mu) + C_3(\mu) + C_9(\mu)] \quad (22)$$

$$\mathcal{C}_6^{\text{eff}}(\mu) \equiv \frac{1}{3}C_5(\mu) + C_6(\mu) + \frac{1}{3}C_7(\mu) + C_8(\mu) \quad (23)$$

$$\mathcal{C}_{6,\text{oct}}^{\text{eff}}(\mu) \equiv 2[C_5(\mu) + C_7(\mu)]. \quad (24)$$

In deriving (20), we have used the relations

$$Q_1^c = \frac{1}{3}Q_2^c + 2Q_{2,\text{oct}}^c \quad (25)$$

$$\tilde{Q}_5^c = \frac{1}{3}\tilde{Q}_6^c + 2\tilde{Q}_{6,\text{oct}}^c \quad (26)$$

with

$$Q_2^c = (\bar{c}_\alpha s_\alpha)_{V-A} (\bar{b}_\beta c_\beta)_{V-A} \quad (27)$$

$$Q_{2,\text{oct}}^c = (\bar{c}_\alpha T_{\alpha\beta}^a s_\beta)_{V-A} (\bar{b}_\gamma T_{\gamma\delta}^a c_\delta)_{V-A} \quad (28)$$

and

$$\tilde{Q}_6^c = -2(\bar{c}_\alpha L s_\alpha) (\bar{b}_\beta R c_\beta) \quad (29)$$

$$\tilde{Q}_{6,\text{oct}}^c = -2(\bar{c}_\alpha L T_{\alpha\beta}^a s_\beta) (\bar{b}_\gamma R T_{\gamma\delta}^a c_\delta). \quad (30)$$

Here L and R correspond to the Dirac structures $\hat{1} - \gamma_5$ and $\hat{1} + \gamma_5$, respectively. The D_s^{*+} meson emerges from the $(\bar{c}s)$ pieces of these operators. Since it is a *vector* meson, we have

$$\langle D_s^{*+} | \bar{c}_\alpha L s_\alpha | 0 \rangle = 0, \quad (31)$$

and hence the *factorized* matrix element of \tilde{Q}_6^c vanishes. As in Sect. 2.2, the hadronic matrix elements of the colour-octet operators vanish within the factorization approximation because of their colour-structure.

2.4 Observables of the angular distributions

The hadronic matrix element of a generic four-quark operator \mathcal{Q} between the state vectors $\langle V_1(\lambda)V_2(\lambda) |$ and $|B_q\rangle$ has the following general Lorentz-decomposition [5, 6]:

$$\begin{aligned} \langle V_1(\lambda)V_2(\lambda) | \mathcal{Q} | B_q \rangle &= \epsilon_{V_1,\mu}(\lambda)^* \epsilon_{V_2,\nu}(\lambda)^* \left[a g^{\mu\nu} + \frac{b}{m_{V_1} m_{V_2}} p_{V_2}^\mu p_{V_1}^\nu \right. \\ &\quad \left. + i \frac{c}{m_{V_1} m_{V_2}} \varepsilon^{\mu\nu\alpha\beta} p_{V_1,\alpha} p_{V_2,\beta} \right], \quad (32) \end{aligned}$$

where the symbols $\epsilon(\lambda)$ denote the polarization vectors of the final-state vector mesons V_1 and V_2 . A similar parametrization can be employed to express the transition matrix elements [(7) and (20)], yielding

$$\begin{aligned} a &= \frac{G_F}{\sqrt{2}} V_{cs} V_{cb}^* \left[\mathcal{C}_i^{\text{eff}}(\mu) A_i^f + \mathcal{C}_{i+4}^{\text{eff}}(\mu) A_{i+4}^f \right. \\ &\quad \left. + \mathcal{C}_i^{\text{eff}}(\mu) A_i^{\text{nf}} + \mathcal{C}_{i,\text{oct}}^{\text{eff}}(\mu) A_{i,\text{oct}}^{\text{nf}}(\mu) \right. \\ &\quad \left. + \mathcal{C}_{i+4}^{\text{eff}}(\mu) A_{i+4}^{\text{nf}} + \mathcal{C}_{i+4,\text{oct}}^{\text{eff}}(\mu) A_{i+4,\text{oct}}^{\text{nf}}(\mu) \right] \quad (33) \end{aligned}$$

$$\begin{aligned} b &= \frac{G_F}{\sqrt{2}} V_{cs} V_{cb}^* \left[\mathcal{C}_i^{\text{eff}}(\mu) B_i^f + \mathcal{C}_{i+4}^{\text{eff}}(\mu) B_{i+4}^f \right. \\ &\quad \left. + \mathcal{C}_i^{\text{eff}}(\mu) B_i^{\text{nf}} + \mathcal{C}_{i,\text{oct}}^{\text{eff}}(\mu) B_{i,\text{oct}}^{\text{nf}}(\mu) \right. \\ &\quad \left. + \mathcal{C}_{i+4}^{\text{eff}}(\mu) B_{i+4}^{\text{nf}} + \mathcal{C}_{i+4,\text{oct}}^{\text{eff}}(\mu) B_{i+4,\text{oct}}^{\text{nf}}(\mu) \right] \quad (34) \end{aligned}$$

$$\begin{aligned} c &= \frac{G_F}{\sqrt{2}} V_{cs} V_{cb}^* \left[\mathcal{C}_i^{\text{eff}}(\mu) C_i^f + \mathcal{C}_{i+4}^{\text{eff}}(\mu) C_{i+4}^f \right. \\ &\quad \left. + \mathcal{C}_i^{\text{eff}}(\mu) C_i^{\text{nf}} + \mathcal{C}_{i,\text{oct}}^{\text{eff}}(\mu) C_{i,\text{oct}}^{\text{nf}}(\mu) \right. \\ &\quad \left. + \mathcal{C}_{i+4}^{\text{eff}}(\mu) C_{i+4}^{\text{nf}} + \mathcal{C}_{i+4,\text{oct}}^{\text{eff}}(\mu) C_{i+4,\text{oct}}^{\text{nf}}(\mu) \right], \quad (35) \end{aligned}$$

where the index i distinguishes between colour-suppressed ($i = 1$) and colour-allowed ($i = 2$) decays and ‘‘f’’ and ‘‘nf’’ correspond to ‘‘factorized’’ and ‘‘non-factorized’’ matrix elements, respectively. Note that the factorized amplitudes do not depend on the renormalization scale μ . Since the Wilson coefficients depend on this scale, this already signals the need for non-factorizable contributions to cancel the μ -dependence in (33)–(35) (see e.g. [32] for a further discussion of that point).

In the following sections we will analyse the decays $B_q \rightarrow V_1 V_2$ in terms of *linear polarization states*. The corresponding decay amplitudes take the form [23, 33]

$$\begin{aligned} A(B_q(t) \rightarrow V_1 V_2) &= \frac{A_0(t)}{x} \epsilon_{V_1}^{*L} \epsilon_{V_2}^{*L} - A_{\parallel}(t) \epsilon_{V_1}^{*T} \cdot \epsilon_{V_2}^{*T} / \sqrt{2} \\ &\quad - i A_{\perp}(t) \epsilon_{V_1}^* \times \epsilon_{V_2}^* \cdot \hat{\mathbf{p}}_{V_2} / \sqrt{2}, \quad (36) \end{aligned}$$

where $x \equiv p_{V_1} \cdot p_{V_2} / (m_{V_1} m_{V_2})$ and $\hat{\mathbf{p}}_{V_2}$ is the unit vector along the direction of motion of V_2 in the rest frame of V_1 . Here the time dependences originate from B_q - \bar{B}_q mixing. In our notation, an unmixed B_q meson is present at $t = 0$.

The linear polarization amplitudes at $t = 0$ defined by (36) can be expressed in terms of a , b and c as follows [33]:

$$\begin{aligned} A_0(0) &= -xa - (x^2 - 1)b \\ A_{\parallel}(0) &= \sqrt{2}a \\ A_{\perp}(0) &= \sqrt{2(x^2 - 1)}c. \quad (37) \end{aligned}$$

At time $t = 0$, the angular distributions for $B_q \rightarrow V_1 V_2$ depend on the observables $|A_0(0)|$, $|A_{\parallel}(0)|$, $|A_{\perp}(0)|$ and on the two phases $\delta_1 \equiv \text{Arg}[A_{\parallel}(0)^* A_{\perp}(0)]$ and $\delta_2 \equiv \text{Arg}[A_0(0)^* A_{\perp}(0)]$, which are CP-conserving strong phases that are 0 (mod π) in the absence of final-state interactions (probably not a justifiable assumption for the colour-suppressed modes). Quantitative estimates for these observables will be given in the following subsection.

2.5 Factorization tests and estimates of observables

While the non-factorizable contributions to a , b and c cannot be calculated at present, the evaluation of the factorizable contributions is straightforward. Without yet going into the details of which form factors to employ, the naive factorization assumption yields many testable consequences. For example, time-reversal invariance forces the form factors parametrizing quark currents to be all relatively real. Consequently, naive factorization predicts the same strong phase (mod π) for the three amplitudes $A_0(0)$, $A_{\parallel}(0)$, $A_{\perp}(0)$. It therefore predicts vanishing values of the two observables [17, 19, 20]

$$\text{Im}[A_0^*(0)A_{\perp}(0)] = 0 \quad (38)$$

$$\text{Im}[A_{\parallel}^*(0)A_{\perp}(0)] = 0, \quad (39)$$

and the equality

$$\text{Re}[A_0^*(0)A_{\parallel}(0)] = \pm |A_0(0)A_{\parallel}(0)|. \quad (40)$$

The breakdown of the naive factorization assumption is unequivocally proved if any of the three Eqs. (38)–(40) is not satisfied. Detailed comparisons of polarization amplitudes in non-leptonic and semi-leptonic decays test additional implications of the naive factorization assumption. The phenomenology of detailed studies of the full non-trivial angular distributions is thus much richer than the single factorization test available for a pseudoscalar decaying into two pseudoscalars [18]. While the above equations represent general tests of the factorization assumption, it is also useful to examine the predictions for the observables of the angular distributions for various form factor ansätze.

2.5.1 The colour-suppressed decays $B_q \rightarrow J/\psi V$

The factorized amplitudes for $B_q \rightarrow J/\psi V$ with $(q, V) \in \{(s, \phi); (d, K^{*0}); (u, K^{*+})\}$ are given by [16, 21, 22]:

$$\begin{aligned} A_1^f &= -f_{J/\psi} m_{J/\psi} (m_{B_q} + m_V) A_1^{B_q V}(m_{J/\psi}^2) = A_5^f \\ B_1^f &= 2 \frac{f_{J/\psi} m_{J/\psi}^2 m_V}{m_{B_q} + m_V} A_2^{B_q V}(m_{J/\psi}^2) = B_5^f \\ C_1^f &= 2 \frac{f_{J/\psi} m_{J/\psi}^2 m_V}{m_{B_q} + m_V} V^{B_q V}(m_{J/\psi}^2) = C_5^f \end{aligned} \quad (41)$$

where we have used the notation of Bauer, Stech and Wirbel for the form factors $A_i^{B_q V}(q^2)$ and $V^{B_q V}(q^2)$ of quark currents [16]. The parameter $f_{J/\psi}$ denotes the J/ψ decay constant, which can be determined from the $J/\psi \rightarrow e^+e^-$ rate, yielding $f_{J/\psi} = 395$ MeV.

At present, several methods for obtaining the form factors $A_1(m_{J/\psi}^2)$, $A_2(m_{J/\psi}^2)$ and $V(m_{J/\psi}^2)$ for the $B \rightarrow K^*$ case are on the market. Using $SU(3)$ flavour symmetry of strong interactions, the $B \rightarrow K^*$ form factors can be related to the $B_s \rightarrow \phi$ case. In Table 1 we have collected the form factors proposed by several authors [16, 34, 35], and have moreover given the corresponding predictions for the ratios of observables of the angular distributions. These ratios should suffer less from unknown $SU(3)$ -breaking corrections than the observables themselves. Note that these ratios are independent of the Wilson coefficients within the factorization approach.

The quantity

$$\frac{\Gamma_0(0)}{\Gamma_0(0) + \Gamma_T(0)} \equiv \frac{|A_0(0)|^2}{|A_0(0)|^2 + |A_{\parallel}(0)|^2 + |A_{\perp}(0)|^2} \quad (42)$$

describes the ratio of the longitudinal to the total rate at $t = 0$. Although CDF [36] claims to have measured this quantity, from their untagged data sample, to be $0.56 \pm 0.21(\text{stat.})_{-0.04}^{+0.02}(\text{syst.})$, their claim is valid only if the CP-odd component of $B_s \rightarrow J/\psi \phi$ is negligible, or if the lifetime difference $\Delta\Gamma$ can be ignored.

The 2nd–4th columns of Table 1 are calculated within the framework of naive factorization, i.e. we have inserted (41) into (33)–(35) and have omitted the “nf” terms in order to calculate the amplitudes in (37).

Table 1. Predictions for form factors and $B_s \rightarrow J/\psi \phi$ ($B \rightarrow J/\psi K^*$) observables

Observable	BSW [16]	Soares [34]	Cheng [35]
$A_1^{BK^*}(m_{J/\psi}^2)$	0.46	0.42	0.41
$A_2^{BK^*}(m_{J/\psi}^2)$	0.46	0.43	0.36
$V^{BK^*}(m_{J/\psi}^2)$	0.55	1.08	0.72
$ A_{\parallel}(0) / A_0(0) $	0.81 (0.77)	0.82 (0.78)	0.75 (0.70)
$ A_{\perp}(0) / A_0(0) $	0.41 (0.40)	0.89 (0.88)	0.55 (0.54)
$\Gamma_0(0)/(\Gamma_0(0) + \Gamma_T(0))$	0.55 (0.57)	0.40 (0.42)	0.54 (0.56)
δ_1	π	π	π
δ_2	0	0	0

The form factors given by Soares [34] are obtained from $\bar{D} \rightarrow K^{(*)} l \bar{\nu}_l$ data by using heavy-quark symmetry relations [37] and assuming the monopole momentum-transfer dependence of the BSW model [16]. Some more form-factor models and their predictions are discussed in [38]. Note that the small difference between the $B_s \rightarrow J/\psi \phi$ and $B \rightarrow J/\psi K^*$ results in Table 1 is related to phase-space effects and not to any $SU(3)$ -breaking effects in the corresponding hadronic matrix elements.

Looking at Table 1, we observe that the “factorized” predictions for $|A_{\parallel}(0)|/|A_0(0)|$ are rather stable (≈ 0.8), while $|A_{\perp}(0)|/|A_0(0)|$ depends strongly on the method used for obtaining the form factors. A common feature of all results is $\delta_1 = \pi$ and $\delta_2 = 0$. Therefore a measurement of non-trivial phases δ_1 and δ_2 would imply the presence of strong final-state interactions and non-factorizable contributions.

Whereas the use of the factorization assumption is very questionable in the case of the channels $B_s \rightarrow J/\psi \phi$ and $B \rightarrow J/\psi K^*$, flavour $SU(3)$ symmetry is probably a good working assumption. Thus all the hadronization dynamics of the $B_s \rightarrow J/\psi \phi$ decay, such as the phases δ_1 and δ_2 and magnitudes of the amplitudes

$$A_0(0), A_{\parallel}(0), A_{\perp}(0), \quad (43)$$

can be obtained from the $B \rightarrow J/\psi K^*$ modes.¹ This approach may be helpful to extract the CKM phase ϕ_{CKM} (see (1)), as we will see below.

¹ Although those $SU(3)$ relations are mostly trivial, one subtlety due to quantum-coherence must be emphasized. Because of the $SU(3)$ relations in the unmixed amplitudes

$A_f(B_s \rightarrow J/\psi \phi) = A_f(B \rightarrow J/\psi K^*)$, where $f = 0, \parallel, \perp$, the magnitudes of the amplitudes for B_s^L or B_s^H decays into CP-even or CP-odd $J/\psi \phi$ final-state configurations, respectively, are a factor of $\sqrt{2}$ larger than their corresponding $B \rightarrow J/\psi K^*$ ones. [Here the K^* is seen in a flavour-specific mode. If K^* is neutral and is observed as $\pi^0 K_S$, quantum coherence in $B^0 - \bar{B}^0$ must also be taken into account.] If the CP-even processes dominate, then

$\Gamma(B_s^L \rightarrow J/\psi \phi) \approx 2\Gamma(B \rightarrow J/\psi K^*)$.

Studies of B_s versus B production fractions can thus be undertaken, since the lifetimes will be precisely known

Table 2. Predictions for form factors and $B_s \rightarrow D_s^{*+} D_s^{*-}$ ($B \rightarrow D_s^{*+} \bar{D}^*$) observables

Observable	BSW	HQET _{strict}	HQET _{sym.-break.}
$A_1^{BD^*}(m_{D_s^*}^2)$	0.72	0.70 (0.68)	0.70 (0.68)
$A_2^{BD^*}(m_{D_s^*}^2)$	0.76	0.76 (0.75)	0.54 (0.53)
$V^{BD^*}(m_{D_s^*}^2)$	0.79	0.76 (0.75)	0.89 (0.88)
$ A_{\parallel}(0) / A_0(0) $	0.90 (0.90)	0.91 (0.91)	0.81 (0.80)
$ A_{\perp}(0) / A_0(0) $	0.32 (0.32)	0.32 (0.33)	0.33 (0.34)
$\Gamma_0(0)/(\Gamma_0(0) + \Gamma_T(0))$	0.52 (0.52)	0.52 (0.52)	0.57 (0.57)
δ_1	π	π	π
δ_2	0	0	0

The factorization assumption should work much better for the transitions $B_s \rightarrow D_s^{*+} D_s^{*-}$ and $B \rightarrow D_s^{*+} \bar{D}^*$. Therefore the results presented in the following subsection should be more reliable than those summarized in Table 1.

2.5.2 The colour-allowed decays $B_q \rightarrow D_s^{*+} \bar{D}_q^*$

Using again the same notation as [16], we get the following “factorized” results for the modes $B_q \rightarrow D_s^{*+} \bar{D}_q^*$ ($q \in \{u, d, s\}$) [23, 24, 26]:

$$\begin{aligned} A_2^f &= -f_{D_s^*} m_{D_s^*} (m_{B_q} + m_{D_q^*}) A_1^{B_q D_q^*}(m_{D_s^*}^2), & A_6^f &= 0 \\ B_2^f &= 2 \frac{f_{D_s^*} m_{D_s^*}^2 m_{D_q^*}}{m_{B_q} + m_{D_q^*}} A_2^{B_q D_q^*}(m_{D_s^*}^2), & B_6^f &= 0 \\ C_2^f &= 2 \frac{f_{D_s^*} m_{D_s^*}^2 m_{D_q^*}}{m_{B_q} + m_{D_q^*}} V^{B_q D_q^*}(m_{D_s^*}^2), & C_6^f &= 0. \end{aligned} \quad (44)$$

The parameter $f_{D_s^*}$ is the D_s^* decay constant. The spin symmetry of HQET implies $f_{D_s^*} \approx f_{D_s}$. A recent compilation of measurements of f_{D_s} from $D_s \rightarrow \mu \bar{\nu}$ gives $(241 \pm 21 \pm 30)$ MeV [39].

In the case of $B_q \rightarrow \bar{D}_q^*$ transitions we have rather tight restrictions from HQET (for reviews, see for example [25]) for the corresponding form factors. The following ratios turn out to be useful to implement these HQET constraints [40]:

$$R_1(w) = \left[1 - \frac{q^2}{(m_{B_q} + m_{D_q^*})^2} \right] \frac{V^{BD^*}(q^2)}{A_1^{BD^*}(q^2)} \quad (45)$$

$$R_2(w) = \left[1 - \frac{q^2}{(m_{B_q} + m_{D_q^*})^2} \right] \frac{A_2^{BD^*}(q^2)}{A_1^{BD^*}(q^2)}, \quad (46)$$

where $R_1(w)$ and $R_2(w)$ are defined in such a way that we have

$$R_1(w) = R_2(w) = 1 \quad (47)$$

for all values of w in the *strict* heavy-quark limit. The kinematical variable w is defined by

$$w = \frac{m_{B_q}^2 + m_{D_q^*}^2 - q^2}{2m_{B_q} m_{D_q^*}}. \quad (48)$$

The value of the momentum transfer q^2 relevant for (44) is $q^2 = m_{D_s^*}^2$. The form factor $A_1^{BD^*}(q^2)$ is usually written as

$$A_1^{BD^*}(q^2) = \frac{m_{B_q} + m_{D_q^*}}{2\sqrt{m_{B_q} m_{D_q^*}}} \left[1 - \frac{q^2}{(m_{B_q} + m_{D_q^*})^2} \right] h_{A_1}(w), \quad (49)$$

where $h_{A_1}(w)$ corresponds to the Isgur–Wise function in the strict heavy-quark limit and can be written as

$$h_{A_1}(w) = \mathcal{F}(1) [1 - \rho_{A_1}^2 (w - 1) + \mathcal{O}((w - 1)^2)]. \quad (50)$$

The current status of the normalization $\mathcal{F}(1)$ and of the “slope parameter” $\rho_{A_1}^2$ has been summarized recently by Neubert in [40]. The form factor $A_1^{BD^*}(q^2)$ is protected by Luke’s theorem [41] against $1/m_Q$ corrections at zero recoil. The other form factors $A_2^{BD^*}(q^2)$ and $V^{BD^*}(q^2)$ are not protected by this theorem. From calculations based on HQET one expects a rather weak dependence of $R_1(w)$ and $R_2(w)$ on w and therefore uses

$$R_1(w) = R_1[1 + \mathcal{O}(w - 1)] \quad (51)$$

$$R_2(w) = R_2[1 + \mathcal{O}(w - 1)]. \quad (52)$$

In our analysis we will neglect the w -dependence completely.

Following these lines we have calculated the results for the form factors and ratios of observables, which should receive smaller $SU(3)$ -breaking corrections than the observables themselves, summarized in Table 2. For completeness we have also given the results obtained by applying the BSW model [16] in the 2nd column. In order to calculate the 3rd and 4th columns, we have used $\mathcal{F}(1) = 0.91$ and $\rho_{A_1}^2 = 0.91$ [40]. The columns denoted by HQET_{strict} and HQET_{sym.-break.} correspond to $R_1 = R_2 = 1$ and

$R_1 = 1.18$, $R_2 = 0.71$, respectively, where we have employed the results by Neubert [40] to take into account HQET symmetry-breaking corrections. Within the factorization approximation we obtain the following simple expressions for $A_{\parallel}(0)/A_0(0)$ and $A_{\perp}(0)/A_0(0)$ in terms of the HQET parameters:

$$\frac{A_{\parallel}(0)}{A_0(0)} = \sqrt{2} \left[\frac{m_{D_s^*}}{m_{B_q}} \left(\frac{x^2 - 1}{w + 1} \right) R_2(w) - x \right]^{-1} \quad (53)$$

$$\frac{A_{\perp}(0)}{A_0(0)} = - \left[\frac{m_{D_s^*}}{m_{B_q}} \left(\frac{\sqrt{x^2 - 1}}{w + 1} \right) R_1(w) \right] \frac{A_{\parallel}(0)}{A_0(0)}, \quad (54)$$

where the kinematical variable

$$x = \frac{m_{B_q}^2 - m_{D_q^*}^2 - m_{D_s^*}^2}{2m_{D_q^*}m_{D_s^*}} \quad (55)$$

has been defined after (36).

If we compare Table 2 with Table 1, we note that the results for the observables depend much less on the way of obtaining the form factors. Also the ‘‘old’’ BSW model is in rather good agreement with the HQET predictions, which is quite remarkable. Therefore the results given in Table 2 are more reliable than those collected in Table 1. In this respect it is also important to note that the non-factorizable contributions appearing in (33)–(35) should play a minor role for the colour-allowed decay class and that $\delta_1 = \pi$ and $\delta_2 = 0$ is expected to hold on rather solid ground. Because of the latter feature, CP-violating effects arising in untagged $B_s \rightarrow D_s^{*+} D_s^{*-}$ data samples should be a promising way to extract the weak phase ϕ_{CKM} (see (1)), as has been outlined in detail in [8].

3 The angular-moment analysis

The main focus of this section is the efficient determination of the observables discussed in Sect. 2.4 and 2.5. This can be accomplished by an *angular-moment analysis* [1]. In this approach, the observed data are weighted by judiciously chosen functions, which project out any desired observable. Whereas [1] determines the moments for a few choice angular distributions, using spherical harmonics, this paper indicates how to determine suitable weighting functions for all kinds of angular distributions, using only orthogonality arguments (without invoking spherical harmonics).

Let us denote the angular distribution of a given decay by

$$f(\Theta, \alpha; t) = \sum_i b^{(i)}(\alpha; t) g^{(i)}(\Theta), \quad (56)$$

where α represents all the parameters that are independent of the kinematics, which is described by certain decay angles. In general, the physical process involves an arbitrary number of such angles denoted generically by Θ . For the examples considered in this article, i runs from 1 to 6, and we have

$$\alpha = \{\Gamma_H, \Gamma_L, \Delta m, |A_0(0)|, |A_{\parallel}(0)|, |A_{\perp}(0)|, \delta_1, \delta_2, \phi_{\text{CKM}}\}.$$

All the quantities of interest are encoded in the time evolution of the observables $b^{(i)}(\alpha; t)$. In the following discussion the α - and t -dependence of the $b^{(i)}$'s is implicit wherever not explicitly stated.

The usual method for extracting $b^{(i)}$'s is to use an *unbinned maximum likelihood fit* [4]. Performing such a fit for a given quantity requires some idea of the values of the other quantities. When one deals with limited statistics, one may want to exploit alternative methods that completely decouple the extraction of one observable from all the others. Luckily such a method exists, the angular-moment analysis.

If we can find a weighting function $w^{(i)}(\Theta)$ for each i such that

$$\int [D\Theta] w^{(i)}(\Theta) g^{(k)}(\Theta) = \delta_{ik} = \begin{cases} 0 & \text{for } i \neq k \\ 1 & \text{for } i = k, \end{cases} \quad (57)$$

then the $b^{(i)}$'s can be obtained directly from

$$b^{(i)} = \int [D\Theta] w^{(i)}(\Theta) f_{\text{expt}}(\Theta). \quad (58)$$

Here $[D\Theta]$ denotes the appropriate measure for integrating over all angles Θ , and $f_{\text{expt}}(\Theta)$ denotes the observed full angular distribution. For a small number of events (N), the form of the function $f_{\text{expt}}(\Theta)$ will not be known, but only the values for Θ will be known for each event. In that case the above equation reduces to

$$b^{(i)} = \frac{1}{N} \sum_{\text{events}} w^{(i)}(\Theta). \quad (59)$$

These $b^{(i)}$'s can then be used directly for studying their $(\alpha; t)$ dependence.

That $w^{(k)}$'s can always be found for any angular distribution follows from the linear independence of the $g^{(i)}$'s (they have to be independent for the angular distribution to be legitimate). The vector space \mathcal{V}_k spanned by all $g^{(j)}$'s for $j \neq k$ is a *proper* subspace of the vector space \mathcal{V} spanned by all $g^{(i)}$'s. Then there exists a one-dimensional vector space \mathcal{W} such that $\mathcal{V} = \mathcal{V}_k \oplus \mathcal{W}$ and $\mathcal{V}_k \perp \mathcal{W}$. Here the scalar product is defined as $v_1 \cdot v_2 \equiv \int [D\Theta] v_1(\Theta) v_2(\Theta)$; $w^{(k)}$ is then the element of \mathcal{W} with proper magnitude.

For a given set of $g^{(i)}$'s, the choice of $w^{(i)}$'s need not be unique. We can always take any vector space $\mathcal{V}' \supset \mathcal{V}$ and the corresponding *projection* space \mathcal{W}' such that $\mathcal{V}' = \mathcal{V}_k \oplus \mathcal{W}'$ and $\mathcal{V}_k \perp \mathcal{W}'$. Then any $w^{(k)} \in \mathcal{W}'$ with $w^{(k)} \cdot g^{(k)} = 1$ will serve our purpose.

We now indicate an explicit procedure for finding a set of weighting functions applicable to any given angular distribution. For a theoretical angular distribution of the form $f = \sum_{i=1}^n b^{(i)} g^{(i)}$ (the dependence on angles and time is implicit),

$$w^{(i)} = \sum_{j=1}^n \lambda_{ij} g^{(j)} \quad (60)$$

is a proper weighting function, where the n^2 unknowns λ_{ij} are solutions of the n^2 simultaneous equations

$$\delta_{ik} = \sum_{j=1}^n \lambda_{ij} \int [D\Theta] g^{(j)} g^{(k)}. \quad (61)$$

The existence of such a solution follows from the vector-space arguments given earlier. The $w^{(i)}$'s need not be restricted to the vector space spanned by the n vectors $g^{(j)}$'s, in which case the unknowns λ_{ij} will be underdetermined and more than one set of $w^{(i)}$'s will serve our purpose.

It is crucial to observe that the weighting functions $w^{(i)}$ depend only on the angular terms and not on the values of the observables $b^{(j)}$. The implication is that no matter how complicated the detailed angular distribution, there always exists an angular weighting, which projects out the desired observables alone. We therefore recommend the use of moments whenever one wishes to extract observables from measured angular distributions, such as in weak decays of baryons [2] or pseudoscalars [$P \rightarrow V \ell \nu$, $X_J \ell \nu$, VV , etc.], or strong and electromagnetic decays [3]. The utility of this approach cannot be overemphasized. For instance, the moment analysis allows the study of the q^2 dependence of each of the observables separately in the process $P \rightarrow V \ell \nu$. This could prove useful for the extraction of form factors and the determination of CKM elements, e.g. V_{cb} and V_{ub} .

We note that there exist many legitimate choices of weighting functions. The *optimal* choice depends on the numerical values of the observables [1] and on the detector configuration.

4 The angular distribution of the colour-suppressed decays $B_s \rightarrow J/\psi \phi$ and $B \rightarrow J/\psi K^*$

In this section we give the angular distribution of the decays $B_s \rightarrow J/\psi \phi$ and $B \rightarrow J/\psi K^*$, their time-dependences and appropriate weighting functions.

4.1 The decay $B_s \rightarrow J/\psi(\rightarrow l^+l^-)\phi(\rightarrow K^+K^-)$

An analysis of this process has been performed in [33] in terms of linear polarization states of the final-state vector mesons. The corresponding decay amplitude has the same form as (36). Since the amplitudes $A_{0,\parallel}$ and A_{\perp} are related to CP-even and CP-odd final-state configurations, respectively, they differ in time evolution as well as angular distribution. The angular distribution can be used to separate these components and their time evolution can be studied individually.

The differential decay rate at time t as a function of a generic variable x will be denoted by

$$\frac{d\Gamma(t, x)}{dx} \equiv \frac{1}{N(t)} \frac{d^2 N(t)}{dx dt}. \quad (62)$$

Consequently the normalized number of decays in the intervals $[t, t + \Delta t]$ and $[x, x + \Delta x]$ is given by

$$\frac{d\Gamma(t, x)}{dx} \Delta x \Delta t = \frac{1}{N(t)} \frac{d^2 N(t)}{dx dt} \Delta x \Delta t. \quad (63)$$

4.2 Tagged decays

In the case of $B_s \rightarrow J/\psi \phi$, the three-angle distribution for the decay of an initially present (i.e. tagged) B_s meson takes the form [33]

$$\begin{aligned} & \frac{d^3 \Gamma[B_s(t) \rightarrow J/\psi(\rightarrow l^+l^-)\phi(\rightarrow K^+K^-)]}{d \cos \theta d \varphi d \cos \psi} \\ & \propto \frac{9}{32\pi} \left[2|A_0(t)|^2 \cos^2 \psi (1 - \sin^2 \theta \cos^2 \varphi) \right. \\ & \quad + \sin^2 \psi \{ |A_{\parallel}(t)|^2 (1 - \sin^2 \theta \sin^2 \varphi) + |A_{\perp}(t)|^2 \sin^2 \theta \\ & \quad - \text{Im}(A_{\parallel}^*(t) A_{\perp}(t)) \sin 2\theta \sin \varphi \} \\ & \quad + \frac{1}{\sqrt{2}} \sin 2\psi \{ \text{Re}(A_0^*(t) A_{\parallel}(t)) \sin^2 \theta \sin 2\varphi \\ & \quad \left. + \text{Im}(A_0^*(t) A_{\perp}(t)) \sin 2\theta \cos \varphi \right] . \quad (64) \end{aligned}$$

Throughout this section we will apply the same conventions as in [33], i.e. ϕ moves in x direction in the J/ψ rest frame, the z axis is perpendicular to the decay plane of $\phi \rightarrow K^+K^-$, and $p_y(K^+) \geq 0$. The coordinates (θ, φ) describe the decay direction of l^+ in the J/ψ rest frame and ψ is the angle made by $\mathbf{p}(K^+)$ with the x axis in the ϕ rest frame. With this convention,

$$\begin{aligned} \mathbf{x} &= \mathbf{p}_{\phi}, \quad \mathbf{y} = \frac{\mathbf{p}_{K^+} - \mathbf{p}_{\phi}(\mathbf{p}_{\phi} \cdot \mathbf{p}_{K^+})}{|\mathbf{p}_{K^+} - \mathbf{p}_{\phi}(\mathbf{p}_{\phi} \cdot \mathbf{p}_{K^+})|}, \\ \mathbf{z} &= \mathbf{x} \times \mathbf{y} \\ \sin \theta \cos \varphi &= \mathbf{p}_{\ell^+} \cdot \mathbf{x}, \quad \sin \theta \sin \varphi = \mathbf{p}_{\ell^+} \cdot \mathbf{y}, \\ \cos \theta &= \mathbf{p}_{\ell^+} \cdot \mathbf{z}. \quad (65) \end{aligned}$$

Here, the bold-face characters represent *unit* 3-vectors and everything is measured in the rest frame of J/ψ . Also

$$\cos \psi = -\mathbf{p}'_{K^+} \cdot \mathbf{p}'_{J/\psi}, \quad (66)$$

where the primed quantities are *unit vectors* measured in the rest frame of ϕ .

The time dependence of the right-hand side of (64) can be read off from Table 3, where $\Delta m \equiv m_H - m_L > 0$ is the mass difference of the mass eigenstates B_s^H and B_s^L of the B_s system and $\bar{\Gamma} \equiv (\Gamma_H + \Gamma_L)/2$ denotes their average decay width. The phases $\delta_1 \equiv \text{Arg}(A_{\parallel}(0)^* A_{\perp}(0))$ and $\delta_2 \equiv \text{Arg}(A_0(0)^* A_{\perp}(0))$ are *CP-conserving* strong phases. In the absence of final-state interactions – probably not a justifiable assumption for $B_s \rightarrow J/\psi \phi$ – they are expected to be 0 (mod π).

On the other hand, the quantity $\delta\phi = \phi_{\text{CKM}}$ (see (1)) is a *CP-violating* weak phase, which is introduced through interference effects between $B_s\text{--}\bar{B}_s$ mixing and decay processes. It can be expressed in terms of elements of the

Table 3. Time evolution of the decay $B_s \rightarrow J/\psi(\rightarrow l^+l^-)\phi(\rightarrow K^+K^-)$ of an initially (i.e. at $t = 0$) pure B_s meson

Observable	Time evolution
$ A_0(t) ^2$	$ A_0(0) ^2 \left[e^{-\Gamma_L t} - e^{-\bar{\Gamma} t} \sin(\Delta m t) \delta\phi \right]$
$ A_{\parallel}(t) ^2$	$ A_{\parallel}(0) ^2 \left[e^{-\Gamma_L t} - e^{-\bar{\Gamma} t} \sin(\Delta m t) \delta\phi \right]$
$ A_{\perp}(t) ^2$	$ A_{\perp}(0) ^2 \left[e^{-\Gamma_H t} + e^{-\bar{\Gamma} t} \sin(\Delta m t) \delta\phi \right]$
$\text{Re}(A_0^*(t)A_{\parallel}(t))$	$ A_0(0) A_{\parallel}(0) \cos(\delta_2 - \delta_1) \left[e^{-\Gamma_L t} - e^{-\bar{\Gamma} t} \sin(\Delta m t) \delta\phi \right]$
$\text{Im}(A_{\parallel}^*(t)A_{\perp}(t))$	$ A_{\parallel}(0) A_{\perp}(0) \left[e^{-\bar{\Gamma} t} \sin(\delta_1 - \Delta m t) + \frac{1}{2} (e^{-\Gamma_H t} - e^{-\Gamma_L t}) \cos(\delta_1) \delta\phi \right]$
$\text{Im}(A_0^*(t)A_{\perp}(t))$	$ A_0(0) A_{\perp}(0) \left[e^{-\bar{\Gamma} t} \sin(\delta_2 - \Delta m t) + \frac{1}{2} (e^{-\Gamma_H t} - e^{-\Gamma_L t}) \cos(\delta_2) \delta\phi \right]$

CKM matrix [9, 12] as

$$\exp(i\delta\phi) = \frac{V_{ts}V_{tb}^*V_{cs}^*V_{cb}}{V_{ts}^*V_{tb}V_{cs}V_{cb}^*} \quad (67)$$

and is very small, as can be seen easily by applying the Wolfenstein expansion of the CKM matrix [10]. At leading order in this expansion $\delta\phi$ vanishes. However, taking into account higher-order terms (for a treatment of such terms, see e.g. [42]) gives a non-vanishing result [13, 43]:

$$\delta\phi = 2\lambda^2\eta = \mathcal{O}(0.03). \quad (68)$$

Consequently $\delta\phi$ measures simply the CKM parameter η . Note that $\lambda = \sin\theta_C = 0.22$ is related to the Cabibbo angle. Useful expressions for $\delta\phi$ can be found in [13], where the following relation has been derived:

$$\delta\phi = 2\lambda^2 R_b \sin\gamma. \quad (69)$$

Here γ is one angle of the ‘‘usual’’ unitarity triangle [44]. Consequently, if the CKM-parameter R_b (defined by (6)) is used as an input, $\delta\phi$ allows a determination of γ . That input allows, however, also the determination of η (or γ) from the mixing-induced CP asymmetry of $B_d \rightarrow J/\psi K_S$ measuring $\sin 2\beta$, where β denotes another angle of the unitarity triangle [44]. If one compares these two results for η (or γ) obtained from B_s and B_d modes, respectively, a test of whether the $B_s-\bar{B}_s$ and $B_d-\bar{B}_d$ mixing phases are described by the Standard Model, or receive additional contributions from physics beyond that model can be performed. Needless to note, a measurement of a value of $\delta\phi$ much larger than the Standard Model expectation of $\mathcal{O}(0.03)$ would anyway be a striking signal for new physics in $B_s-\bar{B}_s$ mixing.

An interesting interpretation of $\delta\phi$ has been given in [45]. There it was shown that $\delta\phi$ is related to one angle in a rather squashed (and therefore ‘‘unpopular’’) unitarity triangle. Note that terms of $\mathcal{O}(\delta\phi^2)$ have been neglected in Table 3.

The angular distribution for an initially present \bar{B}_s meson is given by

$$\frac{d^3\Gamma[\bar{B}_s(t) \rightarrow J/\psi(\rightarrow l^+l^-)\phi(\rightarrow K^+K^-)]}{d\cos\theta d\varphi d\cos\psi}$$

$$\begin{aligned} &\propto \frac{9}{32\pi} \left[2|\bar{A}_0(t)|^2 \cos^2\psi(1 - \sin^2\theta \cos^2\varphi) \right. \\ &\quad + \sin^2\psi \{ |\bar{A}_{\parallel}(t)|^2(1 - \sin^2\theta \sin^2\varphi) + |\bar{A}_{\perp}(t)|^2 \sin^2\theta \\ &\quad - \text{Im}(\bar{A}_{\parallel}^*(t)\bar{A}_{\perp}(t)) \sin 2\theta \sin\varphi \} \\ &\quad + \frac{1}{\sqrt{2}} \sin 2\psi \{ \text{Re}(\bar{A}_0^*(t)\bar{A}_{\parallel}(t)) \sin^2\theta \sin 2\varphi \\ &\quad \left. + \text{Im}(\bar{A}_0^*(t)\bar{A}_{\perp}(t)) \sin 2\theta \cos\varphi \right], \quad (70) \end{aligned}$$

where the angles are again defined by (65) and (66). The time dependence of this rate can be obtained easily with the help of Table 4, where terms of $\mathcal{O}(\delta\phi^2)$ have been neglected, as in Table 3. In calculating Tables 3 and 4 we have used the fact that $B_s \rightarrow J/\psi\phi$ (and $\bar{B}_s \rightarrow J/\psi\phi$) is dominated to *excellent accuracy* by a single weak amplitude, as we have seen in Sect. 2. Therefore we have to deal only with *mixing-induced* CP violation and there is no *direct* CP violation, i.e. $|\bar{A}_0(0)| = |A_0(0)|$, $|\bar{A}_{\parallel}(0)| = |A_{\parallel}(0)|$ and $|\bar{A}_{\perp}(0)| = |A_{\perp}(0)|$.

It is important to note that the mass difference Δm can be extracted from time-dependent analyses of tagged $B_s \rightarrow J/\psi\phi$ data samples [33]. Previous experimental feasibility investigations for the extraction of Δm focused entirely on tagged flavour-specific modes of B_s mesons [46, 47].

4.3 Untagged decays

Combining Tables 3 and 4, we find that the time evolution of the *untagged* data sample for $f = J/\psi(\rightarrow l^+l^-)\phi(\rightarrow K^+K^-)$ is given by

$$\begin{aligned} &\frac{d^3\Gamma[f(t)]}{d\cos\theta d\varphi d\cos\psi} \\ &\propto \frac{9}{16\pi} \left[2|A_0(0)|^2 e^{-\Gamma_L t} \cos^2\psi(1 - \sin^2\theta \cos^2\varphi) \right. \\ &\quad \left. + \sin^2\psi \{ |A_{\parallel}(0)|^2 e^{-\Gamma_L t} (1 - \sin^2\theta \sin^2\varphi) \right. \end{aligned}$$

Table 4. Time evolution of the decay $\bar{B}_s \rightarrow J/\psi(\rightarrow l^+l^-)\phi(\rightarrow K^+K^-)$ of an initially (i.e. at $t = 0$) pure \bar{B}_s meson

Observable	Time evolution
$ \bar{A}_0(t) ^2$	$ A_0(0) ^2 \left[e^{-\Gamma_L t} + e^{-\bar{\Gamma} t} \sin(\Delta m t) \delta\phi \right]$
$ \bar{A}_\parallel(t) ^2$	$ A_\parallel(0) ^2 \left[e^{-\Gamma_L t} + e^{-\bar{\Gamma} t} \sin(\Delta m t) \delta\phi \right]$
$ \bar{A}_\perp(t) ^2$	$ A_\perp(0) ^2 \left[e^{-\Gamma_H t} - e^{-\bar{\Gamma} t} \sin(\Delta m t) \delta\phi \right]$
$\text{Re}(\bar{A}_0^*(t)\bar{A}_\parallel(t))$	$ A_0(0) A_\parallel(0) \cos(\delta_2 - \delta_1) \left[e^{-\Gamma_L t} + e^{-\bar{\Gamma} t} \sin(\Delta m t) \delta\phi \right]$
$\text{Im}(\bar{A}_\parallel^*(t)\bar{A}_\perp(t))$	$- A_\parallel(0) A_\perp(0) \left[e^{-\bar{\Gamma} t} \sin(\delta_1 - \Delta m t) - \frac{1}{2} (e^{-\Gamma_H t} - e^{-\Gamma_L t}) \cos(\delta_1) \delta\phi \right]$
$\text{Im}(\bar{A}_0^*(t)\bar{A}_\perp(t))$	$- A_0(0) A_\perp(0) \left[e^{-\bar{\Gamma} t} \sin(\delta_2 - \Delta m t) - \frac{1}{2} (e^{-\Gamma_H t} - e^{-\Gamma_L t}) \cos(\delta_2) \delta\phi \right]$

$$\begin{aligned}
& + |A_\perp(0)|^2 e^{-\Gamma_H t} \sin^2 \theta \} + \frac{1}{\sqrt{2}} \sin 2\psi \{ |A_0(0)||A_\parallel(0)| \\
& \times \cos(\delta_2 - \delta_1) e^{-\Gamma_L t} \sin^2 \theta \sin 2\varphi \} \\
& + \left\{ \frac{1}{\sqrt{2}} |A_0(0)||A_\perp(0)| \cos \delta_2 \sin 2\psi \sin 2\theta \cos \varphi \right. \\
& \left. - |A_\parallel(0)||A_\perp(0)| \cos \delta_1 \sin^2 \psi \sin 2\theta \sin \varphi \right\} \\
& \times \frac{1}{2} (e^{-\Gamma_H t} - e^{-\Gamma_L t}) \delta\phi \Big]. \quad (71)
\end{aligned}$$

Remarkably the time dependence of the untagged rate does not depend on the mass difference Δm . This feature has been discussed within a more general framework in [14]. Consequently, whereas Γ_L and Γ_H can be determined from the untagged data sample, the extraction of $(\Delta m)_{B_s}$ requires tagging. As has already been pointed out in [8], because of the lifetime difference $(\Delta\Gamma)_{B_s}$, the untagged decay rate (71) develops an interesting contribution for $t > 0$, which is proportional to the CP-violating weak phase $\delta\phi$. It originates from the imaginary parts of the interference terms between $A_\perp(t)$ ($\bar{A}_\perp(t)$) and $A_\parallel^*(t)$ ($\bar{A}_\parallel^*(t)$), $A_0^*(t)$ ($\bar{A}_0^*(t)$). If $\Delta\Gamma \equiv \Gamma_H - \Gamma_L$ is in fact sizeable, we are optimistic that it will be possible to measure this effect.

4.4 A closer look at the one-angle distribution

The full three-angle distributions for tagged and untagged $B_s \rightarrow J/\psi(\rightarrow l^+l^-)\phi(\rightarrow K^+K^-)$ decays discussed in the previous subsections are quite complicated. A much simpler case arises if we integrate out the two decay angles φ and ψ in (64), leading to the following *one-angle* distribution [33]:

$$\begin{aligned}
\frac{d\Gamma(t)}{d\cos\theta} & \propto (|A_0(t)|^2 + |A_\parallel(t)|^2) \frac{3}{8} (1 + \cos^2\theta) \\
& + |A_\perp(t)|^2 \frac{3}{4} \sin^2\theta. \quad (72)
\end{aligned}$$

Let us first briefly illustrate the angular moment analysis outlined in Sect. 3 for this transparent one-angle distribution. In this case, we have

$$g^{(1)}(\theta) = \frac{3}{8} (1 + \cos^2\theta), \quad g^{(2)}(\theta) = \frac{3}{4} \sin^2\theta. \quad (73)$$

Consequently, if we choose

$$w^{(1)}(\theta) = 5 \cos^2\theta - 1 \quad \text{and} \quad w^{(2)}(\theta) = 2 - 5 \cos^2\theta, \quad (74)$$

the orthogonality relation

$$\int_{-1}^{+1} d(\cos\theta) w^{(i)}(\theta) g^{(k)}(\theta) = \delta_{ik} \quad (75)$$

is satisfied, and we obtain immediately

$$|A_0(t_j)|^2 + |A_\parallel(t_j)|^2 \propto \sum_i (5 \cos^2\theta_i - 1) \quad (76)$$

$$|A_\perp(t_j)|^2 \propto \sum_i (2 - 5 \cos^2\theta_i), \quad (77)$$

where the summation is over all the events in the same time bin as t_j .

In the case of the *untagged* one-angle distribution, the $\Delta m t$ oscillations proportional to the CP-violating weak phase $\delta\phi$ cancel, and the terms (76) and (77) evolve like $(|A_0(0)|^2 + |A_\parallel(0)|^2) e^{-\Gamma_L t}$ and $|A_\perp(0)|^2 e^{-\Gamma_H t}$, respectively. A fit (now with only one parameter in each time evolution) gives the decay widths Γ_L and Γ_H of the CP-even and CP-odd B_s mass eigenstates, as well as the CP-even and CP-odd rates $|A_0(0)|^2 + |A_\parallel(0)|^2$ and $|A_\perp(0)|^2$, respectively. For limited statistics, one may want to use time moments [48]

$$f^{(n)} = \int_0^\infty dt t^n f(t). \quad (78)$$

The *weighting-functions method* is thus an alternative to the *two-bin method* suggested in [1,33]. Note that we do not need any *a priori* information about the relative magnitudes of CP-even and CP-odd amplitudes.

In the case of *tagged* measurements, the integrated decay rates

$$\Gamma(t) = \int_{-1}^{+1} d(\cos\theta) \frac{d\Gamma(t)}{d\cos\theta} \quad (79)$$

evolve in time for initially present B_s and \bar{B}_s mesons as

$$\begin{aligned} \Gamma(t) \propto & (|A_0(0)|^2 + |A_{\parallel}(0)|^2) e^{-\Gamma_L t} + |A_{\perp}(0)|^2 e^{-\Gamma_H t} \\ & - (|A_0(0)|^2 + |A_{\parallel}(0)|^2 - |A_{\perp}(0)|^2) \\ & \times e^{-\bar{\Gamma} t} \sin(\Delta m t) \delta\phi \end{aligned} \quad (80)$$

and

$$\begin{aligned} \bar{\Gamma}(t) \propto & (|A_0(0)|^2 + |A_{\parallel}(0)|^2) e^{-\Gamma_L t} + |A_{\perp}(0)|^2 e^{-\Gamma_H t} \\ & + (|A_0(0)|^2 + |A_{\parallel}(0)|^2 - |A_{\perp}(0)|^2) \\ & \times e^{-\bar{\Gamma} t} \sin(\Delta m t) \delta\phi, \end{aligned} \quad (81)$$

respectively, where we have used Tables 3 and 4. Consequently, the time-dependent CP asymmetry arising in the decay $B_s \rightarrow J/\psi \phi$ takes the following form:

$$\begin{aligned} a_{\text{CP}}(B_s(t) \rightarrow J/\psi \phi) & \equiv \frac{\Gamma(t) - \bar{\Gamma}(t)}{\Gamma(t) + \bar{\Gamma}(t)} \\ & = - \frac{|A_0(0)|^2 + |A_{\parallel}(0)|^2 - |A_{\perp}(0)|^2}{(|A_0(0)|^2 + |A_{\parallel}(0)|^2) e^{-\Gamma_L t} + |A_{\perp}(0)|^2 e^{-\Gamma_H t}} \\ & \quad \times e^{-\bar{\Gamma} t} \sin(\Delta m t) \delta\phi. \end{aligned} \quad (82)$$

Using the quantitative estimates collected in Table 1, we obtain

$$\frac{|A_{\perp}(0)|^2}{|A_0(0)|^2 + |A_{\parallel}(0)|^2} = 0.1 \dots 0.5. \quad (83)$$

Although these estimates suffer from large hadronic uncertainties, they indicate that it may not be justified to neglect the CP-odd contributions proportional to $|A_{\perp}(0)|^2$ in the time-dependent CP asymmetry (82).

The coefficient of $\sin(\Delta m t) \delta\phi$ in (82) can be experimentally determined [for instance, from the untagged studies outlined above]. Thus the fundamental weak phase $\delta\phi$ can be cleanly extracted once the $\Delta m t$ oscillations are resolved. Future experiments at the Tevatron and the LHC should be able to achieve this goal. Once the $\Delta m t$ oscillations are traced, one can alternatively perform a tagged, one-angle, time-dependent study to separate the CP-even and CP-odd contributions, from each of which $\delta\phi$ can be directly extracted. The efficient extraction of the various observables depends on the detector configuration, so that other possible variations should be considered. The full angular distributions contain, of course, all the available information, and will be determined eventually.

In order to determine $\delta\phi$ from *untagged* $B_s \rightarrow J/\psi \phi$ decays, where the $\Delta m t$ oscillations cancel, the observables corresponding to the interference terms $\text{Im}(A_{\parallel}^*(t)A_{\perp}(t))$ and $\text{Im}(A_0^*(t)A_{\perp}(t))$ must be studied. Valuable information about CP-conserving strong phases can also be obtained, thereby shedding light on the hadronization dynamics of $B_s \rightarrow J/\psi \phi$ and the issue of ‘‘factorization’’, which predicts trivial strong phases. A set of weighting functions applicable to this case is given in Table 5.

Table 5. A set of weighting functions for extracting the observables $b^{(i)}(t)$ of the decays $B_s \rightarrow J/\psi(\rightarrow l^+l^-)\phi(\rightarrow K^+K^-)$ and $B \rightarrow J/\psi(\rightarrow l^+l^-)K^*(\rightarrow \pi K)$

Observables: $b^{(i)}(t)$	$w^{(i)}(\theta, \varphi, \psi)$
$ A_0(t) ^2$	$\frac{1}{2}[5(\cos^2\theta - \sin^2\theta \cos 2\varphi) - 1]$
$ A_{\parallel}(t) ^2$	$\frac{1}{2}[5(\cos^2\theta + \sin^2\theta \cos 2\varphi) - 1]$
$ A_{\perp}(t) ^2$	$2 - 5\cos^2\theta$
$\text{Re}(A_0^*(t)A_{\parallel}(t))$	$\frac{5}{\sqrt{2}} \sin(2\psi) \sin(2\varphi)$
$\text{Im}(A_{\parallel}^*(t)A_{\perp}(t))$	$-\frac{5}{2} \sin(2\theta) \sin\varphi$
$\text{Im}(A_0^*(t)A_{\perp}(t))$	$\frac{25}{4\sqrt{2}} \sin(2\psi) \sin(2\theta) \cos\varphi$

4.5 The decay $B \rightarrow J/\psi(\rightarrow l^+l^-)K^*(\rightarrow \pi K)$

The angular distribution for $B \rightarrow J/\psi(\rightarrow l^+l^-)K^*(\rightarrow \pi K)$ takes the same form as (64) if we use the decay angles specified in (65) and (66) with ϕ replaced by K^* and K^+ replaced by the strange meson.

Using the same angles for $\bar{B} \rightarrow J/\psi(\rightarrow l^+l^-)\bar{K}^*(\rightarrow \pi\bar{K})$, we obtain the analogous angular distribution to the $\bar{B}_s \rightarrow J/\psi(\rightarrow l^+l^-)\phi(\rightarrow K^+K^-)$ case given in (70). The same weighting functions (see Table 5) can therefore be used to determine the corresponding observables in those decays. The comparison of the observables in these two modes would give us an idea of the extent of $SU(3)$ breaking.

If the K^{*0} is observed to decay to the CP eigenstate $\pi^0 K_S$, the time evolution of the corresponding three-angle distributions [(64) and (70)] is given in Tables 6 and 7, respectively [1]. Tables 6 and 7 assume that the unmixed amplitudes depend on a single, unique weak phase, which is justified within the CKM model (see Sect. 2). In these tables, Γ and $\Delta m > 0$ describe B_d - \bar{B}_d mixing. They are related to each other through the mixing parameter $x_d \equiv (\Delta m/\Gamma)_{B_d}$. In analogy to (67), mixing-induced CP violation in $B_d \rightarrow J/\psi(\rightarrow l^+l^-)K^{*0}(\rightarrow \pi^0 K_S)$ [1, 49] measures a weak phase $\tilde{\beta}$, which is given by

$$\exp(-2i\tilde{\beta}) = \frac{V_{td}V_{tb}^* V_{cs}^*V_{cb}}{V_{td}^*V_{tb} V_{cs}V_{cb}^*}. \quad (84)$$

Within the Wolfenstein expansion [10], $\tilde{\beta}$ is equal, to a very good approximation, to the angle β of the ‘‘standard’’ (non-squashed) unitarity triangle [44]. Therefore we have not distinguished between $\tilde{\beta}$ and β in Tables 6 and 7.

Whereas the rates for tagged $B_d \rightarrow J/\psi K_S$ and $\bar{B}_d \rightarrow J/\psi K_S$ events, which are given by

$$\begin{aligned} \Gamma[B_d(t) \rightarrow J/\psi K_S] \\ \propto |A(0)|^2 e^{-\Gamma t} [1 - \sin(2\beta) \sin(\Delta m t)] \end{aligned} \quad (85)$$

$$\begin{aligned} \Gamma[\bar{B}_d(t) \rightarrow J/\psi K_S] \\ \propto |A(0)|^2 e^{-\Gamma t} [1 + \sin(2\beta) \sin(\Delta m t)] , \end{aligned} \quad (86)$$

allow *only* the determination of $\sin(2\beta)$ and of $(\Delta m, \Gamma)_{B_d}$, an analysis of the tagged three-angle distribution for the

Table 6. Time evolution of the decay $B_d \rightarrow J/\psi(\rightarrow l^+l^-)K^{*0}(\rightarrow \pi^0 K_S)$ of an initially (i.e. at $t = 0$) pure B_d meson

Observable	Time evolution
$ A_0(t) ^2$	$ A_0(0) ^2 e^{-\Gamma t} [1 + \sin(2\beta) \sin(\Delta mt)]$
$ A_{\parallel}(t) ^2$	$ A_{\parallel}(0) ^2 e^{-\Gamma t} [1 + \sin(2\beta) \sin(\Delta mt)]$
$ A_{\perp}(t) ^2$	$ A_{\perp}(0) ^2 e^{-\Gamma t} [1 - \sin(2\beta) \sin(\Delta mt)]$
$\text{Re}(A_0^*(t)A_{\parallel}(t))$	$ A_0(0) A_{\parallel}(0) \cos(\delta_2 - \delta_1) e^{-\Gamma t} [1 + \sin(2\beta) \sin(\Delta mt)]$
$\text{Im}(A_{\parallel}^*(t)A_{\perp}(t))$	$ A_{\parallel}(0) A_{\perp}(0) e^{-\Gamma t} [\sin(\delta_1) \cos(\Delta mt) - \cos(2\beta) \cos(\delta_1) \sin(\Delta mt)]$
$\text{Im}(A_0^*(t)A_{\perp}(t))$	$ A_0(0) A_{\perp}(0) e^{-\Gamma t} [\sin(\delta_2) \cos(\Delta mt) - \cos(2\beta) \cos(\delta_2) \sin(\Delta mt)]$

Table 7. Time evolution of the decay $\bar{B}_d \rightarrow J/\psi(\rightarrow l^+l^-)\bar{K}^{*0}(\rightarrow \pi^0 K_S)$ of an initially (i.e. at $t = 0$) pure \bar{B}_d meson

Observable	Time evolution
$ \bar{A}_0(t) ^2$	$ A_0(0) ^2 e^{-\Gamma t} [1 - \sin(2\beta) \sin(\Delta mt)]$
$ \bar{A}_{\parallel}(t) ^2$	$ A_{\parallel}(0) ^2 e^{-\Gamma t} [1 - \sin(2\beta) \sin(\Delta mt)]$
$ \bar{A}_{\perp}(t) ^2$	$ A_{\perp}(0) ^2 e^{-\Gamma t} [1 + \sin(2\beta) \sin(\Delta mt)]$
$\text{Re}(\bar{A}_0^*(t)\bar{A}_{\parallel}(t))$	$ A_0(0) A_{\parallel}(0) \cos(\delta_2 - \delta_1) e^{-\Gamma t} [1 - \sin(2\beta) \sin(\Delta mt)]$
$\text{Im}(\bar{A}_{\parallel}^*(t)\bar{A}_{\perp}(t))$	$- A_{\parallel}(0) A_{\perp}(0) e^{-\Gamma t} [\sin(\delta_1) \cos(\Delta mt) - \cos(2\beta) \cos(\delta_1) \sin(\Delta mt)]$
$\text{Im}(\bar{A}_0^*(t)\bar{A}_{\perp}(t))$	$- A_0(0) A_{\perp}(0) e^{-\Gamma t} [\sin(\delta_2) \cos(\Delta mt) - \cos(2\beta) \cos(\delta_2) \sin(\Delta mt)]$

decay $B_d \rightarrow J/\psi(\rightarrow l^+l^-)K^{*0}(\rightarrow \pi^0 K_S)$ (and its CP-conjugate) yields valuable additional information from the interference terms, as can be seen by looking at Tables 6 and 7:

- $\text{Re}(A_0^*(t)A_{\parallel}(t))$ provides additional information on $\cos(\delta_2 - \delta_1)$.
- $\text{Im}(A_{\parallel}^*(t)A_{\perp}(t))$, $\text{Im}(A_0^*(t)A_{\perp}(t))$ provide additional information both on $\sin \delta_{1(2)}$ and $\cos \delta_{1(2)}$ and on $\cos(2\beta)$. The latter quantity plays an important role to resolve discrete ambiguities in the determination of the CKM angle β [50].

Predictions for these observables are given in Table 1.

The largest data sample for $B_d \rightarrow J/\psi K^{*0}$ is, however, not for $K^{*0} \rightarrow \pi^0 K_S$, but for $K^{*0} \rightarrow \pi^- K^+$. The complete angular distributions and time dependences for the relevant decay modes are given in Appendix A. For charged B decays, the corresponding time and angular distribution is obtained by going to the isospin-related mode and setting $\Delta m = 0$. Experimental studies of these decays are very important, since they probe $\sin(\delta_{1(2)})$ and non-factorizable terms through the observables corresponding to the left-hand sides of (38)–(40) [17, 19, 20]. The relevant information about δ_1 and δ_2 extracted from these B data samples, “tagged” at the time of decay, can be related to $B_s \rightarrow J/\psi \phi$ by using $SU(3)$ flavour symmetry of strong interactions, and allows a determination of $\delta\phi$ from the time evolution of even the *untagged* rate given by (71) [8]. This approach does not involve the assumption of factorization, just $SU(3)$ symmetry arguments. Unfortunately

the corresponding $SU(3)$ -breaking corrections cannot be treated in a quantitative way at present.

5 The angular distribution of the colour-allowed decays $B_s \rightarrow D_s^{*+} D_s^{*-}$ and $B \rightarrow D_s^{*+} \bar{D}^*$

The decay of $D_s^{*\pm}$ is predominantly electromagnetic, i.e. $D_s^{*\pm} \rightarrow D_s^{\pm} \gamma$, whereas \bar{D}^* decays also strongly to $\bar{D} \pi$. Therefore the angular distributions of the two decay modes discussed in this section are quite different from each other. The first step in the decay chain is, however, still of the form $P \rightarrow V_1 V_2$ (as in $B_s \rightarrow J/\psi \phi$) and consequently the terms $A_0, A_{\parallel}, A_{\perp}$ retain the same meanings as in (36) and the same physical significance as in Sect. 4.

5.1 The decay $B_s \rightarrow D_s^{*+}(\rightarrow D_s^+ \gamma) D_s^{*-}(\rightarrow D_s^- \gamma)$

Applying the same convention as in [51], we define the coordinate system as follows: in the rest frame of the decaying B_s meson, let the directions of motion of D_s^{*+} and D_s^{*-} be z' and z'' , respectively. In the plane transverse to z' (or z''), choose any direction as y' and y'' . The directions of x' and x'' are then specified uniquely via $x' = y' \times z'$ and $x'' = y'' \times z''$. Thus, x' and x'' point in opposite directions. Then (θ', φ') is the direction of D_s^+ in the rest frame of D_s^{*+} in the $(x' - y' - z')$ coordinate system, whereas

Table 8. A set of weighting functions for $B_s \rightarrow D_s^{*+}(\rightarrow D_s^+\gamma)D_s^{*-}(\rightarrow D_s^-\gamma)$. Here $C(\theta', \theta'') = (1 + \cos^2 \theta')(1 + \cos^2 \theta'')$

$b^{(i)}(t)$	$w^{(i)}(\theta', \theta'', \chi)$
$ A_0 ^2$	$(-45/92)C(\theta', \theta'') + (245/92)\sin^2 \theta' \sin^2 \theta''$
$ A_{\parallel} ^2$	$(10/23)C(\theta', \theta'') - (45/46)\sin^2 \theta' \sin^2 \theta'' + (25/4)\sin^2 \theta' \sin^2 \theta'' \cos 2\chi$
$ A_{\perp} ^2$	$(10/23)C(\theta', \theta'') - (45/46)\sin^2 \theta' \sin^2 \theta'' - (25/4)\sin^2 \theta' \sin^2 \theta'' \cos 2\chi$
$\text{Re}(A_0^* A_{\parallel})$	$25\sqrt{2}\sin \theta' \sin \theta'' \cos \theta' \cos \theta'' \cos \chi$
$\text{Im}(A_{\parallel}^* A_{\perp})$	$-(25/4)\sin^2 \theta' \sin^2 \theta'' \sin 2\chi$
$\text{Im}(A_0^* A_{\perp})$	$-25\sqrt{2}\sin \theta' \sin \theta'' \cos \theta' \cos \theta'' \sin \chi$

(θ'', φ'') is the direction of D_s^- in the rest frame of D_s^{*-} in the $(x'' - y'' - z'')$ coordinate system.

Since the choice of directions of y' and y'' was completely arbitrary, only the combination $\chi = \varphi' + \varphi''$ of φ' and φ'' is physical and these two angles will appear in the angular distribution only through χ . In terms of the momenta of particles, the angles θ', θ'' and χ can be defined as:

$$\begin{aligned} \cos \theta' &= \mathbf{p}'_{D_s^+} \cdot \mathbf{p}_{D_s^{*+}}, \quad \cos \theta'' = \mathbf{p}''_{D_s^-} \cdot \mathbf{p}_{D_s^{*-}} \\ \sin \theta' \sin \theta'' \cos \chi &= -\cos \theta' \cos \theta'' - \mathbf{p}'_{D_s^+} \cdot \mathbf{p}''_{D_s^-} \\ \sin \theta' \sin \theta'' \sin \chi &= (\mathbf{p}'_{D_s^+} \times \mathbf{p}''_{D_s^-}) \cdot \mathbf{p}_{D_s^{*+}}. \end{aligned} \quad (87)$$

The bold-faced quantities are unit three-vectors. The unprimed quantities are measured in the rest frame of B_s , single-primed quantities in the rest frame of D_s^{*+} , and double-primed quantities in the rest frame of D_s^{*-} .

In terms of these angles, the angular distribution takes the form

$$\begin{aligned} &\frac{d^3 \Gamma}{d \cos \theta' d \cos \theta'' d \chi} \\ &\propto \frac{9}{64\pi} \left\{ 2|A_0|^2 \sin^2 \theta' \sin^2 \theta'' + \frac{1}{2}|A_{\parallel}|^2 [(1 + \cos^2 \theta') \right. \\ &\quad \times (1 + \cos^2 \theta'') + \sin^2 \theta' \sin^2 \theta'' \cos 2\chi] \\ &\quad + \frac{1}{2}|A_{\perp}|^2 [(1 + \cos^2 \theta')(1 + \cos^2 \theta'') \\ &\quad - \sin^2 \theta' \sin^2 \theta'' \cos 2\chi] \\ &\quad - \text{Im}(A_{\parallel}^* A_{\perp}) \sin^2 \theta' \sin^2 \theta'' \sin 2\chi \\ &\quad + 2\sqrt{2} \text{Re}(A_0^* A_{\parallel}) \sin \theta' \sin \theta'' \cos \theta' \cos \theta'' \cos \chi \\ &\quad \left. - 2\sqrt{2} \text{Im}(A_0^* A_{\perp}) \sin \theta' \sin \theta'' \cos \theta' \cos \theta'' \sin \chi \right\}, \end{aligned} \quad (88)$$

where the time dependence of all observables is implicit. It can be read off from Table 3. The weighting functions are listed in Table 8.

The angular distribution for the CP-conjugate process $\bar{B}_s \rightarrow D_s^{*+}(\rightarrow D_s^+\gamma)D_s^{*-}(\rightarrow D_s^-\gamma)$ is given by

$$\begin{aligned} &\frac{d^3 \Gamma}{d \cos \theta' d \cos \theta'' d \chi} \\ &\propto \frac{9}{64\pi} \left\{ 2|\bar{A}_0|^2 \sin^2 \theta' \sin^2 \theta'' + \frac{1}{2}|\bar{A}_{\parallel}|^2 [(1 + \cos^2 \theta') \right. \end{aligned} \quad (89)$$

$$\begin{aligned} &\times (1 + \cos^2 \theta'') + \sin^2 \theta' \sin^2 \theta'' \cos 2\chi] \\ &\quad + \frac{1}{2}|\bar{A}_{\perp}|^2 [(1 + \cos^2 \theta')(1 + \cos^2 \theta'') \\ &\quad - \sin^2 \theta' \sin^2 \theta'' \cos 2\chi] \\ &\quad - \text{Im}(\bar{A}_{\parallel}^* \bar{A}_{\perp}) \sin^2 \theta' \sin^2 \theta'' \sin 2\chi \\ &\quad + 2\sqrt{2} \text{Re}(\bar{A}_0^* \bar{A}_{\parallel}) \sin \theta' \sin \theta'' \cos \theta' \cos \theta'' \cos \chi \\ &\quad \left. - 2\sqrt{2} \text{Im}(\bar{A}_0^* \bar{A}_{\perp}) \sin \theta' \sin \theta'' \cos \theta' \cos \theta'' \sin \chi \right\}. \end{aligned}$$

The time evolution of the various quantities is the same as in Table 4.

As in the case of $B_s \rightarrow J/\psi\phi$, the decay $B_s \rightarrow D_s^{*+}D_s^{*-}$ is dominated by a single weak amplitude. Therefore the analysis of the tagged and untagged decays outlined in Sect. 4.2 and 4.3 remains valid by replacing $(J/\psi, \phi, l^+, l^-, K^+, K^-) \rightarrow (D_s^{*+}, D_s^{*-}, D_s^+, \gamma, D_s^-, \gamma)$. Since this process is colour-allowed, factorization is expected to hold more strongly.

5.2 The decay $B \rightarrow D_s^{*+}(\rightarrow D_s^+\gamma)\bar{D}^*(\rightarrow \bar{D}\pi)$

Whereas the decay of the D_s^{*+} meson is of the form $V \rightarrow P\gamma$, which has the same angular dependence as $V \rightarrow l^+l^-$ for massless leptons, the \bar{D}^* decay belongs to the category $V \rightarrow P_1P_2$. The net angular distribution should therefore have the same form as that for $P \rightarrow V(\rightarrow l^+l^-)V(\rightarrow P_1P_2)$. The angular distribution is thus given by (64), where the definitions of angles are the same as in (65) and (66) with ϕ replaced by \bar{D}^* , l^+ replaced by D_s^+ , and K^+ replaced by the charmed meson arising from the \bar{D}^* decay. The angular distribution for the CP-conjugate decay $\bar{B} \rightarrow D_s^{*-}(\rightarrow D_s^-\gamma)D^*(\rightarrow D\pi)$ is given as in (70). The weighting functions collected in Table 5 can be used to extract the corresponding observables from experimental data.

At this point, a few comments concerning the time evolution of these angular distributions are in order. Let us first consider decays of neutral B_d mesons. Since here the final states are flavour-specific, no interference effects between $B_d-\bar{B}_d$ mixing and decay processes arise in this case. Consequently, the time evolution of the corresponding observables is only governed by the ‘‘mixing’’ of the

initial particle, which is either a pure B_d or \overline{B}_d . For $B_d \rightarrow D_s^{*+}(\rightarrow D_s^+\gamma)D^{*-}(\rightarrow \overline{D}\pi)$ and $\overline{B}_d \rightarrow D_s^{*-}(\rightarrow D_s^-\gamma)D^{*+}(\rightarrow D\pi)$, the time evolution of a generic observable \mathcal{Q} of the angular distribution takes hence the form $\mathcal{Q}(t) = \mathcal{Q}(0) e^{-\Gamma t} \cos^2(\Delta mt/2)$, while we have $\mathcal{Q}(t) = \mathcal{Q}(0) e^{-\Gamma t} \sin^2(\Delta mt/2)$ in the case of $B_d \rightarrow D_s^{*-}(\rightarrow D_s^-\gamma)D^{*+}(\rightarrow D\pi)$ and $\overline{B}_d \rightarrow D_s^{*+}(\rightarrow D_s^+\gamma)D^{*-}(\rightarrow \overline{D}\pi)$. The time evolutions of the untagged flavour-specific decays and the related B^\pm decays can be obtained straightforwardly from these expressions by setting $\Delta m = 0$.

6 Summary

The kinematics of B and B_s meson decays into two vector-particles, which both continue to decay through CP-conserving interactions into two lighter particles, involve three independent decay angles. The time evolution of the coefficients of the corresponding angular distributions contains valuable information about the lifetime and mass differences between the B_s mass eigenstates B_s^H and B_s^L , the relative magnitudes and phases of CP-odd and CP-even decay amplitudes, and CP-violating effects, including the Wolfenstein parameter η and the CKM angle β . The ratios of these coefficients are estimated by using various form-factor models. Determinations of these time-dependent coefficients will be useful in testing these models and furthermore in determining the extent to which factorization or the $SU(3)$ flavour symmetry of strong interactions hold in these decays.

The observables of the angular distributions can be determined from experimental data by an angular-moment analysis in which the data are weighted by judiciously chosen weighting functions in order to arrive *directly* at the observables. At times, this permits the extraction of the fundamental CKM parameters. A method applicable to all kinds of angular distributions is indicated, where the weighting functions can be determined without any *a priori* knowledge of the values of the coefficients. This method is almost as good as the likelihood-fit method for a small number of parameters and is expected to give some reliable results even with low statistics where a likelihood fit to a large number of parameters is inefficient.

The B_s meson decays $B_s \rightarrow J/\psi\phi$, $D_s^{*+}D_s^{*-}$ are considered in the light of a possible width difference $(\Delta\Gamma)_{B_s}$. The observables of their angular distributions can be related to those of the decays $B \rightarrow J/\psi K^*$, $D_s^{*+}\overline{D}^*$ by using the $SU(3)$ flavour symmetry, where B stands for B_d or B^+ . The full angular distributions for all these transitions are given explicitly, and the corresponding weighting functions are specified. The time-dependent observables in all these decays provide information about the corresponding values of $\Delta\Gamma$ and Δm . In addition, the decays of B_s mesons inform us about the Wolfenstein parameter η , while $B_d \rightarrow J/\psi K^*(\rightarrow \pi^0 K_S)$ probes the CKM angle β . Some of the quantities related to the B_s case can even be extracted from *untagged* data samples, where one does not distinguish between initially present B_s or \overline{B}_s mesons. The comparison between coefficients of angular distributions of B_s and B mesons may give us an idea about

$SU(3)$ -breaking effects, while the comparison of colour-suppressed ($B \rightarrow J/\psi V$) and colour-allowed ($B \rightarrow D_s^{*\pm} V$) modes should help in testing the expectation that factorization holds to a greater extent in the latter case.

Acknowledgements. We are grateful to H.J. Lipkin and J.L. Rosner for previous collaborations. We thank D. Atwood, J. Incandela, R. Kutschke, J. Lewis, M. Neubert, K. Ohl, S. Pappas, M. Schmidt, S. Sen, M. Shochet, W. Wester, H. Yamamoto for useful discussions. This work was supported in part by the Department of Energy, Contract Nos. DE-AC02-76CHO3000 and DE FG02 90ER40560.

Appendix

A Angular distributions and time dependences for flavour-specific $B_d \rightarrow J/\psi(\rightarrow l^+l^-) K^*(\rightarrow K^\pm\pi^\mp)$ modes

The angles are defined as in (65), where the ϕ meson is replaced by K^* , and the K^+ meson by the strange meson in the final state. In order to parametrize the corresponding angular distributions, we use the following combinations of trigonometric functions:

$$\begin{aligned} f_1 &= 2 \cos^2 \psi (1 - \sin^2 \theta \cos^2 \varphi) \\ f_2 &= \sin^2 \psi (1 - \sin^2 \theta \sin^2 \varphi) \\ f_3 &= \sin^2 \psi \sin^2 \theta \\ f_4 &= \sin^2 \psi \sin 2\theta \sin \varphi \\ f_5 &= (1/\sqrt{2}) \sin 2\psi \sin^2 \theta \sin 2\varphi \\ f_6 &= (1/\sqrt{2}) \sin 2\psi \sin 2\theta \cos \varphi. \end{aligned} \quad (90)$$

Taking into account $|\overline{A}_f| = |A_f|$ and using the notation $A_f \equiv A_f(0)$, where $f \in \{0, \parallel, \perp\}$, we obtain

$$\frac{d^3 \Gamma[B_d(t) \rightarrow J/\psi(\rightarrow l^+l^-)K^*(\rightarrow K^+\pi^-)]}{d \cos \theta d\varphi d \cos \psi} \quad (91)$$

$$\begin{aligned} &= \frac{9}{32\pi} \cos^2 \left(\frac{\Delta mt}{2} \right) e^{-\Gamma t} \left\{ f_1 |A_0|^2 + f_2 |A_{\parallel}|^2 + f_3 |A_{\perp}|^2 \right. \\ &\quad \left. - f_4 \operatorname{Im}(A_{\parallel}^* A_{\perp}) + f_5 \operatorname{Re}(A_0^* A_{\parallel}) + f_6 \operatorname{Im}(A_0^* A_{\perp}) \right\} \end{aligned}$$

$$\frac{d^3 \Gamma[\overline{B}_d(t) \rightarrow J/\psi(\rightarrow l^+l^-)\overline{K}^*(\rightarrow K^-\pi^+)]}{d \cos \theta d\varphi d \cos \psi} \quad (92)$$

$$\begin{aligned} &= \frac{9}{32\pi} \cos^2 \left(\frac{\Delta mt}{2} \right) e^{-\Gamma t} \left\{ f_1 |A_0|^2 + f_2 |A_{\parallel}|^2 + f_3 |A_{\perp}|^2 \right. \\ &\quad \left. + f_4 \operatorname{Im}(A_{\parallel}^* A_{\perp}) + f_5 \operatorname{Re}(A_0^* A_{\parallel}) - f_6 \operatorname{Im}(A_0^* A_{\perp}) \right\} \end{aligned}$$

$$\frac{d^3 \Gamma[B_d(t) \rightarrow J/\psi(\rightarrow l^+l^-)\overline{K}^*(\rightarrow K^-\pi^+)]}{d \cos \theta d\varphi d \cos \psi} \quad (93)$$

$$\begin{aligned} &= \frac{9}{32\pi} \sin^2 \left(\frac{\Delta mt}{2} \right) e^{-\Gamma t} \left\{ f_1 |A_0|^2 + f_2 |A_{\parallel}|^2 + f_3 |A_{\perp}|^2 \right. \\ &\quad \left. + f_4 \operatorname{Im}(A_{\parallel}^* A_{\perp}) + f_5 \operatorname{Re}(A_0^* A_{\parallel}) - f_6 \operatorname{Im}(A_0^* A_{\perp}) \right\} \end{aligned}$$

$$\frac{d^3\Gamma[\bar{B}_d(t) \rightarrow J/\psi(\rightarrow l^+l^-)K^*(\rightarrow K^+\pi^-)]}{d\cos\theta d\varphi d\cos\psi} \quad (94)$$

$$= \frac{9}{32\pi} \sin^2\left(\frac{\Delta mt}{2}\right) e^{-\Gamma t} \left\{ f_1|A_0|^2 + f_2|A_{\parallel}|^2 + f_3|A_{\perp}|^2 - f_4 \operatorname{Im}(A_{\parallel}^* A_{\perp}) + f_5 \operatorname{Re}(A_0^* A_{\parallel}) + f_6 \operatorname{Im}(A_0^* A_{\perp}) \right\}.$$

References

1. I. Dunietz, H.R. Quinn, A. Snyder, W. Toki, H.J. Lipkin, Phys. Rev. **D43** (1991) 2193
2. P. Bialas et al., Z. Phys. **C57** (1993) 115; J. Hrivnac, R. Lednicky, M. Smizanska, J. Phys. **G21** (1995) 629
3. Q. Shen et al., Phys. Rev. **D52** (1995) 2825
4. Review of Particle Physics, Phys. Rev. **D54** (1996) 159
5. G. Valencia, Phys. Rev. **D39** (1989) 3339
6. G. Kramer, W.F. Palmer, Phys. Rev. **D45** (1992) 193, Phys. Lett. **B279** (1992) 181, Phys. Rev. **D46** (1992) 2969 and 3197, G. Kramer, W.F. Palmer, H. Simma, Nucl. Phys. **B428** (1994) 77; G. Kramer, T. Mannel, W.F. Palmer, Z. Phys. **C55** (1992) 497
7. J.S. Hagelin, Nucl. Phys. **B193** (1981) 123; E. Franco, M. Lusignoli, A. Pugliese, *ibid.* **B194** (1982) 403; L.L. Chau, W.-Y. Keung, M.D. Tran, Phys. Rev. **D27** (1983) 2145; L.L. Chau, Phys. Rep. **95** (1983) 1; A.J. Buras, W. Slominski, H. Steger, Nucl. Phys. **B245** (1984) 369; M.B. Voloshin, N.G. Uraltsev, V.A. Khoze, M.A. Shifman, Yad. Fiz. **46** (1987) 181 [Sov. J. Nucl. Phys. **46** (1987) 112]; A. Datta, E.A. Paschos, U. Türke, Phys. Lett. **B196** (1987) 382; A. Datta, E.A. Paschos, Y.L. Wu, Nucl. Phys. **B311** (1988) 35; M. Lusignoli, Z. Phys. **C41** (1989) 645; R. Aleksan, A. Le Yaouanc, L. Oliver, O. Pène, Y.-C. Raynal, Phys. Lett. **B316** (1993) 567; I. Bigi, B. Blok, M. Shifman, N. Uraltsev, A. Vainshtein, in *B Decays*, ed. S. Stone, 2nd edition (World Scientific, Singapore, 1994), p. 132 and references therein; M. Beneke, G. Buchalla, I. Dunietz, Phys. Rev. **D54** (1996) 4419
8. R. Fleischer, I. Dunietz, Phys. Rev. **D55** (1997) 259
9. N. Cabibbo, Phys. Rev. Lett. **10** (1963) 531; M. Kobayashi, T. Maskawa, Prog. Theor. Phys. **49** (1973) 652
10. L. Wolfenstein, Phys. Rev. Lett. **51** (1983) 1945
11. R. Fleischer, I. Dunietz, Phys. Lett. **B387** (1996) 361
12. D. Du, I. Dunietz, Dan-di Wu, Phys. Rev. **D34** (1986) 3414; I. Dunietz, J.L. Rosner, Phys. Rev. **D34** (1986) 1404; Ya.I. Azimov, N.G. Uraltsev, V.A. Khoze, JETP Lett. **43** (1986) 409; I. Dunietz, Ann. Phys. **184** (1988) 350
13. I. Dunietz, in Proceedings of the Workshop on *B* Physics at Hadron Accelerators, Snowmass, Colorado, eds. P. McBride, C. Shekhar Mishra, Fermilab-CONF-93/267 (Batavia, 1993), p. 83
14. I. Dunietz, Phys. Rev. **D52** (1995) 3048
15. D. Fakirov, B. Stech, Nucl. Phys. **B133** (1978) 315
16. M. Bauer, B. Stech, M. Wirbel, Z. Phys. **C29** (1985) 637, Z. Phys. **C34** (1987) 103
17. J. Körner, G. Goldstein, Phys. Lett. **89B** (1979) 105
18. J.D. Bjorken, Nucl. Phys. **B** (Proc. Suppl.) **11** (1989) 325
19. H. Yamamoto, in same Proceedings as in [13], p. 29
20. T.E. Browder, K. Honscheid, D. Pedrini, Annu. Rev. Nucl. Part. Sci. **46** (1996) 395
21. R. Aleksan, A. Le Yaouanc, L. Oliver, O. Pène, Y.-C. Raynal, Phys. Lett. **B316** (1993) 567
22. J. Bijnens, F. Hoogeveen, Phys. Lett. **B283**, 434 (1992)
23. J.L. Rosner, Phys. Rev. **D42** (1990) 3732
24. T. Mannel, W. Roberts, Z. Ryzak, Phys. Rev. **D44** (1991) 18; Phys. Lett. **B259** (1991) 359
25. H. Georgi, in Proceedings of TASI-91, Boulder, Colorado, 1991, eds. R.K. Ellis, C.T. Hill, J.D. Lykken (World Scientific, Singapore, 1992), p. 589; B. Grinstein, in Proceedings of the Workshop on High Energy Phenomenology, Mexico City, 1991, ed. M.A. Pérez, R. Huerta (World Scientific, Singapore, 1992), p. 161; N. Isgur, M.B. Wise, in *Heavy Flavours*, eds. A.J. Buras, M. Lindner (World Scientific, Singapore, 1992), p. 234; T. Mannel, in Proceedings of the Workshop on QCD, Aachen, Germany, 1992, eds. P.M. Zerwas, H.A. Kastrup (World Scientific, Singapore, 1993), p. 634; M. Neubert, Phys. Rep. **245**, 259 (1994); T. Mannel, J. Phys. **G21** (1995) 1007; M. Shifman, in Proceedings of TASI-95, Boulder, Colorado, 1995, p. 409, ed. D. E. Soper (World Scientific, Singapore, 1996)
26. M. Neubert, B. Stech, preprint CERN-TH/97-99, hep-ph/9705292, to appear in *Heavy Flavours II*, eds. A.J. Buras, M. Lindner (World Scientific, Singapore, 1998)
27. For a review, see G. Buchalla, A.J. Buras, M.E. Lautenbacher, Rev. Mod. Phys. **68** (1996) 1125
28. J. Bartelt et al., Phys. Rev. Lett. **71** (1993) 4111
29. A.J. Buras, R. Fleischer, University of Karlsruhe preprint TTP97-15, hep-ph/9704376, to appear in *Heavy Flavours II*, eds. A.J. Buras, M. Lindner (World Scientific, Singapore, 1998)
30. R. Fleischer, Int. J. Mod. Phys. **A12** (1997) 2459
31. R. Fleischer, Z. Phys. **C58** (1993) 483 and **C62** (1994) 81; G. Kramer, W.F. Palmer, H. Simma, Z. Phys. **C66** (1995) 429 (see also the article by these authors listed in [6])
32. A.J. Buras, Nucl. Phys. **B434** (1995) 606
33. A.S. Dighe, I. Dunietz, H.J. Lipkin, J.L. Rosner, Phys. Lett. **B369** (1996) 144
34. J.M. Soares, Phys. Rev. **D53** (1996) 241
35. H.-Y. Cheng, Z. Phys. **C69** (1996) 647
36. CDF Collaboration, F. Abe et al., Phys. Rev. Lett. **75** (1995) 3068
37. N. Isgur, M.B. Wise, Phys. Rev. **D42** (1990) 2388
38. H.-Y. Cheng, B. Tseng, Phys. Rev. **D51** (1995) 6259
39. J.D. Richman, in Proceedings of the XXVIII International Conference on High Energy Physics – ICHEP96, Warsaw, July 1996, eds. Z. Ajduk, A.K. Wroblewski (World Scientific, Singapore, 1997), p. 143
40. M. Neubert, preprint CERN-TH/97-24, hep-ph/9702375, to appear in *Heavy Flavours II*, eds. A.J. Buras, M. Lindner (World Scientific, Singapore, 1998)
41. M.E. Luke, Phys. Lett. **B252** (1990) 447
42. A.J. Buras, M.E. Lautenbacher, G. Ostermaier, Phys. Rev. **D50** (1994) 3433
43. I. Dunietz, Ph.D. Thesis, Ann. Phys. **184** (1988) 350, and references therein
44. L.L. Chau, W.-Y. Keung, Phys. Rev. Lett. **53** (1984) 1802.; J.D. Bjorken, private communication, 1987; C. Jarlskog, R. Stora, Phys. Lett. **208B** (1988) 268
45. R. Aleksan, B. Kayser, D. London, Phys. Rev. Lett. **73** (1994) 18
46. See the following papers in same Proceedings as in [13]: D.J. Ritchie et al., p. 357; J.E. Skarha, A.B. Wicklund, p. 361; T.H. Burnett, p. 367; T.J. Lawry et al., p. 371; X. Lou, p. 373; C. Baltay et al., p. 377; K. Johns, p. 383

47. See also V. Andreev et al. (The LEP B Oscillations Working Group), "Combined Results on B^0 Oscillations: Update for the Summer 1997 Conferences", LEPBOSC 97/2, August 18, 1997
48. A.S. Dighe, S. Sen, in preparation
49. B. Kayser, M. Kuroda, R.D. Peccei, A.I. Sanda, Phys. Lett. **B237** (1990) 508
50. A.S. Dighe, I. Dunietz, R. Fleischer, Phys. Lett. B **433** (1998) 147
51. R. Kutschke, An angular distribution cookbook (unpublished)