

On particle multiplicities in three-jet events

P. Edén¹, G. Gustafson¹, V. Khoze²

¹ Department of Theoretical Physics, Lund University, Helgonav. 5, 223 62 Lund, Sweden

² INFN-Laboratori Nazionali di Frascati, P.O. Box 13, 00044, Frascati (Roma), Italy

Received: 8 June 1999 / Published online: 14 October 1999

Abstract. A thorough verification of the distinct differences in the properties of quark and gluon jets is considered as one of the most instructive tests of the basic ideas of QCD. In the real life experiments such a comparison appears to be quite a delicate task and various subtle issues require further theoretical efforts. In this paper we discuss in detail the possibility to extract the theoretically adequate information from the particle multiplicity patterns in three-jet events in e^+e^- annihilation.

1 Introduction

As well known, the larger colour charge of gluons ($C_A = N_c = 3$) compared to quarks ($C_F = (N_c^2 - 1)/2N_c = 4/3$) leads to various distinctive differences between the two types of jets, for recent articles see e.g. [1] and the review [2]. Thus, a detailed comparison of the properties of quark and gluon jets provides one of the most instructive tests of the basic ideas of QCD. An experimental verification of these differences has been a subject of quite intensive investigations, especially in the last years, e.g. [3]. However, obtaining of the theoretically adequate information about the properties of the gluon jet appears to be not an easy task. Recall that the analytical QCD results address the comparison between the energetic gluon and quark jets emerging from the point-like colourless sources, and that (unlike the $q\bar{q}$ case) the pure high energy gg events at present are not available experimentally¹.

So far, most studies of the structure of gluon jets have been performed in three-jet events of e^+e^- annihilation. As a rule, these rely on a jet finding procedure both for selection of the $q\bar{q}g$ events and for a separation between the jets in an event. Without special care, such an analysis is inherently ambiguous and may suffer from the lack of the direct correspondence to the underlying theory. Recently some more sophisticated approaches have been exploited (see e.g. [3, 5-8]) which allow better theoretical significance. There are still a number of issues which are frequently overlooked in the present gluon jet analyses and some further theoretical efforts are required. First of all, this concerns particle multiplicity distributions in the jets. Clarification of these issues is the main aim of this paper. More detailed description of the theoretical framework can be found in [1].

¹ In principle, it is possible to create a pure source of the colour singlet gg events at a future linear e^+e^- collider through the process $\gamma\gamma \rightarrow gg$ [4]

In particular the following problems are addressed.

1. Different approaches to the three-jet studies employ different definitions of the $q\bar{q}g$ kinematics. In particular, this concerns such a key variable as a transverse momentum scale of the gluon, p_\perp . Our first issue here is to discuss an exact definition of this quantity, which governs radiation from the gluon.
2. The definition of the three-jet topology with the gluon registered at a given p_\perp imposes an obvious requirement that there are no other subjects in the event with the transverse momentum exceeding p_\perp . We have to investigate quantitatively the impact of this requirement on the jet sample.
3. To calculate predictions from perturbative QCD, using the assumption of local parton hadron duality (LPHD) [9], a cutoff is needed for the infrared singularities. As discussed in detail in [1] such a cutoff depends on the soft hadronization process and can *not* be uniquely specified from perturbative QCD alone. Thus, the result is necessarily model dependent.

In what follows we discuss these three issues successively in Sects. 2, 3 and 4, and in Sect. 5 we study their effect on analyses of 3-jet events in e^+e^- annihilation.

2 Definition of p_\perp

In the simplest case of soft radiation, p_\perp can be easily defined, as the quark and antiquark specify a unique direction. For large p_\perp gluons, however, the q and \bar{q} get recoils such that there is no obvious direction against which the transverse momentum should be measured. To have well defined expressions such a direction has to be specified. In the Lund dipole formalism [10-16] p_\perp has been defined according to (subscript Lu for Lund)

$$p_{\perp\text{Lu}}^2 \equiv \frac{s_{qg}s_{g\bar{q}}}{s}, \quad (1)$$

where s_{qg} denotes the squared mass of the quark-gluon system etc. In this particular frame the gluon rapidity is given by the expression

$$y = \frac{1}{2} \ln\left(\frac{s_{\text{qg}}}{s_{\text{g}\bar{\text{q}}}}\right). \quad (2)$$

The kinematically allowed region is given by

$$p_{\perp\text{Lu}} < \frac{\sqrt{s}}{2}; \quad |y| < \ln\left(\frac{\sqrt{s}}{p_{\perp\text{Lu}}}\right) \equiv \frac{1}{2}(L - \kappa_{\text{Lu}});$$

$$L \equiv \ln\left(\frac{s}{\Lambda^2}\right), \quad \kappa_{\text{Lu}} \equiv \ln\left(\frac{p_{\perp\text{Lu}}^2}{\Lambda^2}\right). \quad (3)$$

These variables have the advantage that the phase space element usually expressed in the scaled energy variables x_q and $x_{\bar{q}}$ is exactly given by the simple relation

$$sdx_q dx_{\bar{q}} = dp_{\perp\text{Lu}}^2 dy. \quad (4)$$

As discussed in Sect. 5, $p_{\perp\text{Lu}}$ may also work well as a scale parameter in the QCD cascade.

An alternative definition has also been used in the literature, e.g. by the Leningrad group [17,18]

$$p_{\perp\text{Le}}^2 \equiv \frac{s_{\text{qg}} s_{\text{g}\bar{\text{q}}}}{s_{\text{q}\bar{\text{q}}}}. \quad (5)$$

This definition corresponds to the gluon transverse momentum in the $q\bar{q}$ cms (with respect to the $q\bar{q}$ direction). It is notable that in this frame the gluon rapidity is also exactly given by the expression in (2). The two p_{\perp} -definitions agree for soft gluons, but deviate for harder gluons. While $p_{\perp\text{Lu}}$ is always bounded by $\sqrt{s}/2$, $p_{\perp\text{Le}}$ has no kinematic upper limit in the massless case.

3 Bias from restrictions on subjet transverse momenta

The effect of a cutoff in p_{\perp} has been discussed previously [11,19]. Here we give a brief review of the results, in order to end the section with an investigation of the numerical importance of subleading terms. These are essential for a correct analysis of three-jet events, which will be discussed in Sect. 5.

To see the qualitative features of the bias we first study $e^+e^- \rightarrow q\bar{q}$ events within the Leading Log approximation (LLA). The quark and antiquark emit gluons according to the well-known radiation pattern

$$dn_g \approx C_F \frac{\alpha_s}{\pi} \frac{dx_q dx_{\bar{q}}}{(1-x_q)(1-x_{\bar{q}})} = C_F \frac{\alpha_s(p_{\perp}^2)}{\pi} \frac{dp_{\perp}^2}{p_{\perp}^2} dy$$

$$\equiv C_F \frac{\alpha_s(\kappa)}{\pi} d\kappa dy; \quad \kappa \equiv \ln(p_{\perp}^2/\Lambda^2). \quad (6)$$

We have here used (4), and in the following we define p_{\perp} and y according to (1) and (2), unless otherwise stated.

Due to colour coherence the hadronic multiplicity $N_g(\kappa)$ in a gluon jet depends on the p_{\perp} of the gluon and

not on its energy (see, e.g., [17,18]). Summing up the contributions from all gluons in a cascade we arrive at the average multiplicity $N_{q\bar{q}}(L = \ln(s/\Lambda^2))$ in the original $q\bar{q}$ system [13, 15–18] ([15–18] include also nonleading terms.)

$$N_{q\bar{q}}(L) \approx \int_{\kappa_0}^L d\kappa \int_{-\frac{1}{2}(L-\kappa)}^{\frac{1}{2}(L-\kappa)} dy C_F \frac{\alpha_s(\kappa)}{\pi} N_g(\kappa)$$

$$= \int_{\kappa_0}^L d\kappa (L - \kappa) C_F \frac{\alpha_s(\kappa)}{\pi} N_g(\kappa). \quad (7)$$

(We have here introduced a lower cutoff κ_0 for the integral over transverse momentum. This point will be discussed in Sect. 4.) Taking the derivative with respect to L we find

$$N'_{q\bar{q}}(L) \approx \int_{\kappa_0}^L d\kappa C_F \frac{\alpha_s(\kappa)}{\pi} N_g(\kappa). \quad (8)$$

Consider now a sample of events selected in such a way that there are *no* subjets with $p_{\perp} > p_{\perp\text{cut}}$. (Within a k_{\perp} -based cluster scheme with a resolution parameter $p_{\perp\text{cut}}$, this means that there are only two primary q and \bar{q} jets.) To obtain the multiplicity $N_{q\bar{q}}(L, \kappa_{\text{cut}})$ in this biased sample, we must restrict the κ integral in (7) to the region $\kappa < \kappa_{\text{cut}}$. We then find [11]

$$N_{q\bar{q}}(L, \kappa_{\text{cut}}) \approx N_{q\bar{q}}(\kappa_{\text{cut}}) + (L - \kappa_{\text{cut}}) N'_{q\bar{q}}(\kappa_{\text{cut}}). \quad (9)$$

The first term corresponds to two cones around the q and \bar{q} jet directions. Here the p_{\perp} of the emissions is limited by the kinematical constraint in (3) rather than by κ_{cut} . It also corresponds exactly to an unbiased $q\bar{q}$ system with cms energy $p_{\perp\text{cut}}$. The second term describes a central rapidity plateau of width $(L - \kappa_{\text{cut}})$, in which the limit for gluon emission is given by the constraint κ_{cut} . This expression for a two-jet event can be generalized for a biased multi-jet configuration, and a similar discussion applies also to the multiplicity variance, cf. [11]. (Similar equations for biased two-jet and three-jet events were later discussed also in [19].)

The average particle multiplicity in the selected two-jet sample is smaller than in an unbiased sample. The modification due to the bias is similar to the suppression from a Sudakov form factor. It is formally $\mathcal{O}(\alpha_s)$, but it also contains a factor $\ln^2(s/p_{\perp}^2)$. Thus, it is small for large p_{\perp} values but it becomes significant for smaller p_{\perp} . This clearly demonstrates that the multiplicity in this restricted case depends on *two* scales, \sqrt{s} and $p_{\perp\text{cut}}$. The p_{\perp} of an emitted gluon is related to the virtual mass of the radiating parent quark. Therefore, the two scales $\sqrt{s}/2$ and $p_{\perp\text{cut}}$ represent the energy and virtuality of the quark and antiquark initiating the jets.

Though the LLA result in (9) describes the qualitative features of the bias, subleading corrections are needed for a quantitative analysis. Within the Modified Leading Log approximation (MLLA) [20], subleading terms are included, which affect the prediction for the unbiased multiplicities and, thus, implicitly also the biased multiplicity in (9). Furthermore, it is in [1] shown that the expression in (9) for the biased multiplicity is explicitly changed




Fig. 1. The effect from the bias due to a constraint $p_{\perp \text{cut}}$ on emitted subjects, at 90 GeV energy. The figure shows the ratio of biased over unbiased multiplicities as a function of $p_{\perp \text{cut}}$. The results for LLA and MLLA relations ((9) and (10), respectively) differ significantly from each other. The result of the MLLA relation in (10), using the p_{\perp} definition in (1), is in good agreement with ARIADNE MC and Durham cluster algorithm results

when MLLA corrections are considered. An unbiased system should be restored when $p_{\perp \text{cut}}$ approaches the kinematical limit $\sqrt{s}/2$, but the r.h.s. of (9) equals the unbiased quantity $N_{q\bar{q}}(L)$ only when $p_{\perp \text{cut}} = \sqrt{s}$. The relation consistent with the MLLA is [1]

$$N_{q\bar{q}}(L, \kappa_{\text{cut}}) \approx N_{q\bar{q}}(\kappa_{\text{cut}} + c_q) + (L - \kappa_{\text{cut}} - c_q) \times N'_{q\bar{q}}(\kappa_{\text{cut}} + c_q); \quad c_q = \frac{3}{2}. \quad (10)$$

The bias is illustrated in Fig.1. The dotted line shows results from the ARIADNE MC [14], when the Durham cluster algorithm [21] is used to define a biased sample of events classified as two-jet events with a y_{cut} equal to $p_{\perp \text{cut}}^2/s$. The MC results agree well with the prediction of (10), where for $p_{\perp \text{cut}}$ we have used the p_{\perp} definition in (1) (solid line). The predicted effect is below 5% for $p_{\perp \text{cut}} > 20$ GeV, but increases rapidly for smaller $p_{\perp \text{cut}}$.

Figure 1 presents also the result using the LLA relation in (9) (dashed line). To elucidate the effect of the differences between (9) and (10), we have used the same expression for the unbiased quantities $N_{q\bar{q}}$ and $N'_{q\bar{q}}$. (These are obtained by a simple fit to ARIADNE MC results, which are in good agreement with the MLLA.) As seen, the subleading terms are important; the LLA relation significantly overestimates the effect. To our knowledge experimental data for this bias have not been presented. Such data should be obtainable in a rather straightforward analysis, which thus readily could test the accuracy of the MC result or the MLLA relation.

4 Infrared cutoffs

Gluon radiation diverges for collinear and soft emissions. Therefore, to estimate the hadronic multiplicity from the

assumption of LPHD [9], a cutoff is needed. Naturally, the cutoff must be Lorentz invariant. For collinear emissions a single Feynman diagram dominates, and there are two possibilities, the virtual mass, μ , of the emitting parent parton or the transverse momentum, p_{\perp} , of the emitted gluon measured relative to the parent parton direction. These quantities are connected by the relation

$$p_{\perp}^2 = \mu^2 z(1-z), \quad (11)$$

where z equals the light cone momentum fraction taken by the emitted gluon. The transverse momentum is directly related to the formation time, and, therefore, we regard this as the most natural choice for a cutoff. (For a further discussion see [1].)

For soft emissions no obvious cutoff is available, however. As several Feynman diagrams contribute and interfere, there is no unique parent parton. Consequently μ^2 or p_{\perp}^2 cannot be uniquely specified and, therefore, cannot be directly used. (Obviously a cut in energy is not possible, as this is not Lorentz invariant.)

For soft emissions from a single $q\bar{q}$ colour dipole a cutoff in p_{\perp} is still the natural choice if measured in the cms, where the q and \bar{q} move back to back. For emissions from a more complicated state the situation simplifies greatly in the large- N_c limit, as many interference terms disappear. In this limit the emission corresponds to a set of *independent* colour dipoles [22, 12]. The natural choice for the cutoff is then p_{\perp} in the cms of the emitting dipole (measured with respect to the dipole direction). We note that this implies that the soft gluons connect the hard partons in exactly the same way as the string in the string fragmentation model [23], which illustrates the connection between perturbative QCD and the string model [17].

For the physical case with 3 colours, extra interference terms appear with relative magnitude $1/N_c^2$ [17, 24]. Here nonplanar Feynman diagrams contribute, and it is impossible to uniquely specify a parent parton or a relevant p_{\perp} . Thus, a more fundamental understanding of confinement is needed to specify the cutoff, which cannot be determined from perturbative QCD alone [1]. In hadronization models the $1/N_c^2$ interference terms correspond to the problem of ‘‘colour reconnection’’, and different models have been proposed [25]. None of these can be motivated from first principles, and only experimental data can differentiate among the various models.

In spite of the formal uncertainties, the success of current Monte Carlo programs [26, 14] indicate that the colour suppressed interference terms do not have a very large effect. This is also supported by recent searches by OPAL of the reconnection effects in hadronic Z events [6]. In most parton cascade formalisms, a cascade cutoff motivated in the large- N_c limit is used also for finite N_c . The colour interference effects are accounted for by reducing the colour factor from $N_c/2$ to C_F in regions collinear with quarks and antiquarks, and, due to colour coherence, also in some parts of the central rapidity region. We note, however, that some subtle interference phenomena, as a matter of principle, cannot be absorbed into a probabilistic scheme, see [24] for details. These are still awaiting a thorough experimental test.

5 Formalism for three-jet events

After these general discussions we are now ready to consider three-jet $q\bar{q}g$ systems. To simplify the discussion we first study the large- N_c limit. The emission of softer gluons from a $q\bar{q}$ system corresponds then to two dipoles which emit gluons independently. If a gluon jet is resolved with transverse momentum p_\perp , this imposes a constraint on the emission of subjets from the two dipoles. Thus, the contribution from each dipole is determined by an expression like (10). For relatively soft primary gluons the constraint should be given by $p_{\perp\text{cut}} = p_{\perp g}$. For hard gluons $p_{\perp\text{Lu}}$ is of the same order as its parent quark virtuality, and in [27] it is shown that $\mathcal{O}(\alpha_s^2)$ matrix elements are well described if $p_{\perp\text{Lu}}$ is used as an ordering parameter for the perturbative cascade. This is also indicated by the successful applications of the ARIADNE MC. We will, therefore, assume that the constraint on further emissions is well described by the identification $p_{\perp\text{cut}} = p_{\perp\text{Lu}}$. The multiplicity in a $q\bar{q}$ dipole with an upper limit on p_\perp can, just as for the $q\bar{q}$ case discussed in Sect. 3, be described as two forward jet regions and a central plateau.

We note that if the three-jet events were selected using a cluster algorithm with a *fixed* resolution scale, then the constraint on subjet transverse momenta, $p_{\perp\text{cut}}$, would be smaller than the p_\perp of the gluon jet (as the gluon jet was resolved). In this case most jet definitions give three jets which are all biased [11,19]. We will, however, here focus on three-jet configurations obtained by iterative clustering until exactly three jets remain, *without* a specified resolution scale, where hence the constraint on subjet p_\perp is described by $p_{\perp\text{cut}} = p_{\perp\text{Lu}}$. As we will see, this implies that the bias on the gluon jet is negligible, which makes this selection procedure suitable for an investigation of unbiased gluon jets.

For finite N_c the different dipoles in a multi-parton configuration can not be completely independent of each other. However, encouraged by the success of MC programs, let us assume that the main effect of finite N_c is that the colour factor, which determines softer gluon emission, is reduced from $N_c/2$ to C_F in the domains where the emission is dominated by radiation from the quark or the antiquark leg. Let us assume that a rapidity range Y_q in the $q\bar{q}$ dipole is similar to a corresponding range in a $q\bar{q}$ dipole, while the remaining range $L_{q\bar{q}} - Y_q$ is similar to a range in one half of a $g\bar{g}$ system. The corresponding ranges in the $g\bar{q}$ dipole are $Y_{\bar{q}}$ and $L_{g\bar{q}} - Y_{\bar{q}}$. This implies that the total multiplicity in the $q\bar{q}g$ event corresponds to the expression

$$N_{q\bar{q}g} = N_{q\bar{q}}(Y_q + Y_{\bar{q}}, \kappa_{\text{Lu}}) + \frac{1}{2} N_{g\bar{g}}(L_{q\bar{q}} + L_{g\bar{q}} - Y_q - Y_{\bar{q}}, \kappa_{\text{Lu}}). \quad (12)$$

For the constraint $p_{\perp\text{cut}}$ we have here written κ_{Lu} , which is appropriate for the selection procedure discussed above.

As discussed in Sect. 4, the size of Y_q and $Y_{\bar{q}}$ cannot be uniquely determined within perturbative QCD. Possibly the most natural choice is to assume that the quantity $Y_q + Y_{\bar{q}}$ corresponds to the energy in the $q\bar{q}$ subsystem




Fig. 2. $N_{q\bar{q}g}$ as a function of $p_{\perp\text{Lu}}$ for $\sqrt{s_{q\bar{q}}} = 60$ GeV. The different predictions from (14a,b) and (15) illustrates the importance of the bias at moderate p_\perp

[18], which implies

$$Y_q + Y_{\bar{q}} \approx \ln(s_{q\bar{q}}/A^2) \equiv L_{q\bar{q}}. \quad (13a)$$

The relation in (13a) can be regarded as an educated guess, but a finite shift cannot be excluded. In [15] it is assumed that

$$Y_q + Y_{\bar{q}} \approx \ln(s/A^2) = L, \quad (13b)$$

which agrees with (13a) to leading order. For relatively soft gluons we have $s_{q\bar{q}} \approx s$, and in this case (13a) and (13b) are approximately equivalent. The assumption in (13a) implies that the energy scale for the gluon term is given by $L_{qg} + L_{g\bar{q}} - L_{q\bar{q}} = \kappa_{\text{Le}}$. Similarly we get from (13b) the corresponding gluonic energy scale κ_{Lu} .

The effect of the p_\perp constraint is rather different in the two terms in (12). For the gluon term the energy scale is in general only slightly larger than the bias scale κ_{Lu} . This implies that in most cases the bias can be disregarded in this term. Inserting the different assumptions in (13a) and (13b) into (12) then gives

$$N_{q\bar{q}g} \approx N_{q\bar{q}}(L_{q\bar{q}}, \kappa_{\text{Lu}}) + \frac{1}{2} N_{g\bar{g}}(\kappa_{\text{Le}}), \quad (14a)$$

$$N_{q\bar{q}g} \approx N_{q\bar{q}}(L, \kappa_{\text{Lu}}) + \frac{1}{2} N_{g\bar{g}}(\kappa_{\text{Lu}}). \quad (14b)$$

We note that the consistency between (14a) and (14b) follows from the fact that the total rapidity range in the two dipoles, $L_{qg} + L_{g\bar{q}}$, can be expressed in two different ways by the equalities $L_{qg} + L_{g\bar{q}} = L_{q\bar{q}} + \kappa_{\text{Le}} = L + \kappa_{\text{Lu}}$. In particular, we see from these equalities that the argument in $N_{g\bar{g}}$ has to be $p_{\perp\text{Le}}^2$ in (14a) and $p_{\perp\text{Lu}}^2$ in (14b), and not e.g. $(2p_\perp)^2$.

The leading effect of a finite shift in $Y_q + Y_{\bar{q}}$ is colour-suppressed, and therefore not expected to be large. However, subleading corrections introduce a difference between the results of (14a) and (14b). This is seen in Fig.2, where the difference is approximately 1 particle for $\sqrt{s_{q\bar{q}}} = 60$ GeV. In the calculations of $N_{q\bar{q}g}$ in Fig.2, we have used the expressions in [1] for the multiplicities $N_{q\bar{q}}$ and $N_{g\bar{g}}$. These include MLLA corrections and recoil effects, which




Fig. 3. The prediction for N_{gg} , obtained by subtracting from $N_{q\bar{q}g}$ the quark contribution $N_{q\bar{q}}$, changes significantly if the bias in the $q\bar{q}$ -term is neglected. The figure shows the effect for $\sqrt{s_{q\bar{q}}} = 60\text{GeV}$, with $N_{q\bar{q}g}$ given by (14a)

implies that $N_{gg} < 2N_{q\bar{q}}$ for accessible energies. Consequently, the result for $N_{q\bar{q}g}$ grows with the assumed value of $Y_q + Y_{\bar{q}}$.

While the bias is not serious for the gluon term in (12), it is more important for the $q\bar{q}$ term. Focusing on events with comparatively large values of p_{\perp} , where the bias is less essential, and using the assumption in (13a), we arrive at the result of [18]:

$$N_{q\bar{q}g}(s, p_{\perp Le}^2) = \left[N_{q\bar{q}}(s_{q\bar{q}}) + \frac{1}{2}N_{gg}(p_{\perp Le}^2) \right] \times (1 + \mathcal{O}(\alpha_s)). \quad (15)$$

The bias is formally of order α_s , and is here taken into account by the factor $(1 + \mathcal{O}(\alpha_s))$. The result of this expression, neglecting the $\mathcal{O}(\alpha_s)$ term, is also shown in Fig.2. The effect of the bias corresponds to less than one charged particle for $p_{\perp cut}$ larger than $\sim 10\text{GeV}$, but becomes much more important for smaller $p_{\perp cut}$ -values.

An alternative way to express this result is the effect on extracting N_{gg} from data for $N_{q\bar{q}g}$, as illustrated in Fig.3. N_{gg} can be extracted by subtracting the biased quark multiplicity $N_{q\bar{q}}(L_{q\bar{q}}, \kappa_{Lu})$ from $N_{q\bar{q}g}$, here assumed to be described by (14a). Neglecting the bias in the subtracted $N_{q\bar{q}}$ term gives a significantly different result. The relative effect of the bias is in this case larger, and it exceeds 20% for $p_{\perp} < 15\text{GeV}$. Furthermore, to get a reliable result for N_{gg} , the relevance of subleading terms in the biased quark multiplicity needs to be well understood. For the solid line in Fig.3, the MLLA relation in (10) is used to subtract the $q\bar{q}$ contribution from the total multiplicity. Instead using the LLA relation in (9) would give a prediction for N_{gg} which is about three charged particles higher for most values of $p_{\perp cut}$.

Although the effect of the bias is very important for small p_{\perp} , we also see from Figs.2 and 3 that it can be neglected for large p_{\perp} -values, where, thus, the results in [18] and (15) can be safely used. This implies e.g. that the bias is negligible in gluon systems defined as the hemisphere opposite to two quasi-collinear quark jets, thoroughly investigated by OPAL [5,6].

It would be very interesting to compare the results in Figs.2 and 3 to experiments. Experimental data on $N_{q\bar{q}g}$ can be directly compared to the Monte Carlo or MLLA results in Fig.2, Data on the difference $N_{q\bar{q}g} - N_{q\bar{q}}$ can be compared either to the predictions in Fig.3 or to experimental results for N_{gg} obtained through one of the methods described in [1]. We have compared the results in Fig.2 with MC simulations, where the p_{\perp} scale is determined by the Durham cluster algorithm. The MC results (not shown) indicate that an analysis based on jet reconstruction is accurate enough to illustrate the effects of the bias, but perhaps not to distinguish between the assumptions in (14a) and (14b). We also note that the effects described here may have a phenomenological impact on the recent analysis of $N_{q\bar{q}g}$ [8], which employs the two-scale dependence.

6 Conclusion

A series of subtle effects influence an analysis of the difference between quark and gluon jets in a real life experiment. In this letter we discuss and clarify effects associated with

- the definition of p_{\perp} ,
- the bias from restrictions on subjet p_{\perp} ,
- the problem that infrared cutoffs cannot be uniquely defined from perturbative QCD.

We also demonstrate the impact of these effects on the analysis of three-jet events in e^+e^- annihilation.

Acknowledgements. We thank K. Hamacher, W. Ochs, R. Orava and T. Sjöstrand for useful discussions. This work was supported in part by the EU Fourth Framework Programme ‘Training and Mobility of Researchers’, Network ‘Quantum Chromodynamics and the Deep Structure of Elementary Particles’, contract FMRX-CT98-0194 (DG 12-MIHT).

References

1. P. Edén, G. Gustafson, *JHEP* **09** (1998) 015
2. V.A. Khoze, W. Ochs, *Int. J. Mod. Phys.* **A12** (2949) 1997
3. For recent experimental reviews, see e.g. K. Hamacher, hep-ex/9809018; J.W. Gary, hep-ex/9903020, and references therein.
4. V.A. Khoze, Plenary talk at Workshop on Physics with Linear Colliders, Saariselka, Finland, Sep 9-14, 1991. Published in Saariselka Workshop 1991:547.
5. J.W. Gary, *Phys. Rev.* **D49** (1994) 4503
6. OPAL collaboration, CERN-EP/99-028, to appear in Eur. Phys. J. C
7. ALEPH collaboration, *Z. Phys.* **C76** (1997) 191
8. DELPHI collaboration, CERN-EP/99-03, to appear in Phys. Lett. B
9. Ya.I. Asimov, Yu.L. Dokshitzer, V.A. Khoze, S.I. Troyan, *Z. Phys.* **C27** (1985) 65; Ya.I. Asimov, Yu.L. Dokshitzer, V.A. Khoze, S.I. Troyan, *Z. Phys.* **C31** (1986) 213

10. G. Gustafson, U. Pettersson, *Nucl. Phys.* **B306** (1988) 746
11. B. Andersson, G. Gustafson, A. Nilsson, C. Sjögren, *Z. Phys.* **C49** (1991) 79
12. G. Gustafson, *Phys. Lett.* **B175** (1986) 453
13. B. Andersson, P. Dahlquist, G. Gustafson, *Phys. Lett.* **B214** (1988) 604
14. L. Lönnblad, *Comp. Phys. Comm.* **71** (1992) 15
15. G. Gustafson, *Nucl. Phys.* **B392** (1993) 251
16. G. Gustafson, M. Olsson, *Nucl. Phys.* **B406** (1993) 293
17. Ya.I. Asimov, Yu.L. Dokshitzer, V.A. Khoze, S.I. Troyan, *Phys. Lett.* **B165** (1985) 147; Ya.I. Asimov, Yu.L. Dokshitzer, V.A. Khoze, S.I. Troyan, *Sov. J. Nucl. Phys.* **43** (1986) 95
18. Yu.L. Dokshitzer, V.A. Khoze, S.I. Troyan, *Sov. J. Nucl. Phys.* **47** (1988) 881
19. S. Catani, Yu.L. Dokshitzer, F. Fiorani, B.R. Webber, *Nucl. Phys.* **B383** (1992) 419
20. Yu.L. Dokshitzer, V.A. Khoze, A.H. Mueller and S.I. Troyan, Basics of Perturbative QCD, (*Editions Frontières, Gif-sur-Yvette, 1991*)
21. S. Catani et al., *Phys. Lett.* **B269** (1991) 432
22. G. Gustafson, *Z. Phys.* **C15** (1982) 155; Yu.L. Dokshitzer, V.A. Khoze, S.I. Troyan, *Sov. J. Nucl. Phys.* **46** (1987) 712
23. B. Andersson, G. Gustafson, G. Ingelman, T. Sjöstrand, *Phys. Rep.* **97** (1983) 31
24. Yu.L. Dokshitzer, V.A. Khoze, S.I. Troyan, *Sov. J. Nucl. Phys.* **50** (1989) 505
25. For a review, see A. Ballestrero et al., *J. Phys.* **G24** (1998) 365
26. M. Bengtsson, T. Sjöstrand, *Comp. Phys. Comm.* **39** (1986) 347; T. Sjöstrand, *Comp. Phys. Comm.* **82** (1994) 74; G. Marchesini et al., *Comp. Phys. Comm.* **67** (1992) 465
27. B. Andersson, G. Gustafson, C. Sjögren, *Nucl. Phys.* **B380** (1992) 391