

Hadronization corrections to helicity components of the fragmentation function

T. Sjöstrand¹, O. Smirnova^{2,a}, Ch. Zacharitou Jarlskog²

¹ Department of Theoretical Physics, Lund University, Sölvegatan 14A, 22362 Lund, Sweden

² Department of Elementary Particle Physics, Lund University, P.O. Box 118, 22100 Lund, Sweden

Received: 11 April 2001 / Revised version: 4 May 2001 /

Published online: 29 June 2001 – © Springer-Verlag / Società Italiana di Fisica 2001

Abstract. In the hadronic decays of Z^0 , gluon emission leads to the appearance of the longitudinal component of the fragmentation function, F_L . Measurement of F_L and the transverse component, F_T , could thus provide insight into the gluon fragmentation function. However, hadronization corrections at low x can be significant. Here we present a method of accounting for such corrections, using the Jetset event generator as illustration.

1 Introduction

Studies of fragmentation functions have always been important, since these distributions cannot be predicted theoretically, but can only be measured experimentally, and consecutively be described by phenomenological models. Hadronic decays of γ^*/Z^0 provide a particularly convenient set of events for analysis and interpretation. Helicity components of the fragmentation function, measured in such events, can be used in various QCD studies, e.g., the extraction of the gluon fragmentation function, and the evaluation of α_s . However, existing theoretical calculations being restricted to the perturbative region, hadronization corrections must be taken into account. In what follows, methods for applying such corrections, using the Jetset event generator [1] as an example, will be discussed.

Consider the angular distribution in the process $e^+e^- \rightarrow \gamma^*/Z^0 \rightarrow q\bar{q}$ in its rest frame. Assuming that the final quark and antiquark are not charge-tagged, i.e. that the forward–backward asymmetry is not accessed, the cross section can be written as [2]

$$\frac{d\sigma}{d(\cos\theta)} = \frac{3}{8}(1 + \cos^2\theta)\sigma_T + \frac{3}{4}\sin^2\theta\sigma_L. \quad (1)$$

Here σ_T (σ_L) is the cross section associated with a transverse (longitudinal) gauge boson polarization state with respect to the $q\bar{q}$ axis, and θ is the polar angle of a particle with respect to the incoming lepton axis. To lowest order, only mass effects contribute to a non-vanishing σ_L , but only for the vector part of the cross section, and there only with a coefficient $\sigma_L/\sigma_T = 2m_q^2/E_{\text{cm}}^2$. Even for the b quarks this gives a negligible σ_L contribution at the energies around the Z^0 peak. Therefore, σ_L effectively starts

in $\mathcal{O}(\alpha_s)$ of perturbation theory, associated with the emission of gluons.

Since partons are not directly observable, one may define a hadron-level analogue of (1) [3],

$$\frac{d^2\sigma^h}{dx d(\cos\theta)} = \frac{3}{8}(1 + \cos^2\theta)\frac{d\sigma_T}{dx} + \frac{3}{4}\sin^2\theta\frac{d\sigma_L}{dx}. \quad (2)$$

Here x would preferably be associated with the energy fraction taken by a hadron, $x_E = 2E/E_{\text{cm}}$, so that $\sum x_E = 2$ in each event. Experimentally it is more convenient to use the momentum fraction x_p . The transverse and longitudinal fragmentation functions are defined by normalization to the total cross section $\sigma_{\text{tot}} = \sigma_T + \sigma_L$ [4],

$$F_T(x) = \frac{1}{\sigma_{\text{tot}}}\frac{d\sigma_T}{dx}, \quad F_L(x) = \frac{1}{\sigma_{\text{tot}}}\frac{d\sigma_L}{dx}. \quad (3)$$

The former is dominated by the fragmentation of quark jets, whereas the latter receives a major contribution from gluon fragmentation. Therefore an experimental determination of $F_L(x)$ is a first step towards an extraction of the gluon fragmentation function, alternative to what is offered by more direct methods in 3-jet events [5]. Several experimental $F_L(x)$ studies have also been presented [6].

A complication is that hadrons are not moving in the direction of their imagined mother parton. Already in lowest order of perturbation theory, for $q\bar{q}$ 2-jet events, non-perturbative hadronization gives an effective p_\perp smearing that induces a nonvanishing $F_L(x)$ even where none is expected. Furthermore, the association of a hadron to a single mother parton is not in agreement with our current best understanding of the hadronization process, where it is rather the color field between a color-connected pair of partons (a string piece [7], or a cluster [8]) that mediates the hadron production. Therefore the structure of smearing effects may become rather nontrivial. Obviously, the

^a On leave from JINR, 141980 Dubna, Russia




Fig. 1. $F_T(x_p)$ and $F_L(x_p)$ for corrected (smeared) hadron angles (dashed lines) and for uncorrected (non-smeared) hadron angles (solid lines)

effects are especially important at small x , which is also the region where one would hope to have some sensitivity to the gluon fragmentation function. The string picture also casts in doubt the concept of a gluon fragmentation function defined from inclusive quantities, since the string fragmentation of a parton depends on the angles to other color-connected partons.

The issue of hadronization corrections to fragmentation functions was addressed in [4,9]. The emphasis was on the σ_L that can be extracted from $\int_0^1 F_L(x)xdx$ rather than on $F_L(x)$ itself, however. Therefore we here address the question how hadronization affects $F_L(x)$ (and $F_T(x)$). One main conclusion is that a simple smearing approach is not sufficient to describe hadronization effects. Thus it appears impossible to define a completely model independent, hadronization-smearing-corrected $F_L(x)$, that could be used to extract a gluon fragmentation function. We further suggest a correction procedure, based on a cluster search strategy, that should give a less model dependent $F_L(x)$, but at the price of introducing the cluster resolution scale y as a new parameter in the problem.

2 The simple smearing

The string model description of $q\bar{q}$ events introduces a Gaussian transverse momentum smearing of primary hadrons, $\propto \exp(-p_\perp^2/2\sigma^2)d^2p_\perp$, where $\sigma \approx 0.36$ GeV [1, 10]. Many primary hadrons are unstable and decay further; this distorts the original Gaussian spectrum and reduces the average p_\perp . Since decay products have smaller $p_{||}$, the momentum parallel to the jet axis, some correlation is also introduced between p_\perp and $p_{||}$. Therefore no simple parameterization is proposed, but instead a Monte Carlo simulation with Jetset 7.4 [1] is used to histogram the amount of angular smearing for different x_p bins.

It is now assumed that this smearing should be applied both to quark and gluon jets, so that a hadron will not move exactly in the direction of its mother parton. There are obvious shortcomings to equating different kinds of jets, like that gluon jets have a lower energy and do not contain decays of charm and bottom hadrons, but those particular issues only introduce moderate corrections. More severe objections can be raised to the associ-

ation of hadrons to individual partons, as we will discuss further in the next section, but forget for the moment.

Had particles not been smeared in p_\perp , but parallel with their parton of origin, then the shape of the angular distribution

$$F(x_p, \cos \theta) = \frac{3}{8}(1 + \cos^2 \theta)F_T(x_p) + \frac{3}{4}\sin^2 \theta F_L(x_p) \quad (4)$$

in a bin of x_p could be used to extract $F_T(x_p)$ and $F_L(x_p)$ in that bin. The above mentioned smearing will now modify this. The two angular shapes, $(3/8)(1 + \cos^2 \theta)$ and $(3/4)\sin^2 \theta$, both normalized to unity, are therefore convoluted with the x_p -dependent smearing distributions, characterized by a distribution in the smearing angle θ^{sm} and an isotropic azimuthal distribution φ^{sm} . That is, a parton at an angle θ^p will produce a hadron at an angle θ^h , where

$$\cos \theta^h = \cos \theta^p \cos \theta^{\text{sm}} - \sin \theta^p \sin \theta^{\text{sm}} \cos \varphi^{\text{sm}}. \quad (5)$$

Data can now be fitted both to the “non-smeared” angular distribution form, (4), and to the convoluted (“smeared”) analogue. The resulting distributions for F_T and F_L , obtained using the Jetset generated events, are shown in Fig.1. The effect on F_T and F_L is visible for hadron momenta below 10% of the beam energy. The low-momentum region is affected the most, giving F_L values reduced up to one order of magnitude.

3 Objections to the simple smearing

The above smearing procedure is correct to lowest order in α_s , i.e. it describes how 2-jet events can induce a non-vanishing $F_L(x_p)$. We know, however, that hadronization of 3-jet events cannot be described in terms of a simple incoherent sum of three q , \bar{q} and g jets. One example is the string/drag effect [11,12], i.e. that particle production is suppressed in the angular region between the q and \bar{q} and enhanced in the other two regions, well confirmed experimentally [13]. High-momentum hadrons still essentially follow the separate parton directions, but low-momentum ones are significantly affected. These are the ones where the angular smearing effects are large to begin with. It is well-known that the string effect leads to more 2-jetlike events, e.g. in terms of thrust T , than implied




Fig. 2. Event-by-event hadronization corrections $\Delta F_L(\bar{T}) = F_L^h(\bar{T}) - F_L^p(\bar{T})$ for different simulations: the solid line corresponds to Jetset 7.4 PS, the dashed one to Jetset 7.4 ME (both using the string fragmentation), and the dotted to Jetset 7.4 ME with independent fragmentation scheme

by symmetric smearing [14]. The reason is to be found in the enhanced production of particles between two color-connected partons that are close in angle, leading to them seemingly being even closer, e.g. that the opening angle between the reconstructed jets typically is smaller than that between the original partons.

There is a nontrivial topology dependence on string effects, especially when multiple gluon emission is considered. The issue is therefore best studied in an event generator, by comparing angular distributions on the parton ($i = p$ below) and on the hadron ($i = h$) level. As a simple measure of the jettiness of events we use $\bar{T} = 1 - T$, defined on the parton level of each event. The integrated x_E spectrum is decomposed as

$$\begin{aligned} & \left(\frac{d\sigma_{\text{tot}}}{d\bar{T}} \right)^{-1} \int_0^1 \frac{d^3\sigma^i}{d\bar{T}dx_E d(\cos\theta)} x_E dx_E \\ & = \frac{3}{8} (1 + \cos^2\theta) F_T^i(\bar{T}) + \frac{3}{4} \sin^2\theta F_L^i(\bar{T}), \quad (6) \end{aligned}$$

where the x_E -weighting ensures a common normalization $F_T^i(\bar{T}) + F_L^i(\bar{T}) = 2$ at parton and hadron level (values of F_T^i and F_L^i most conveniently are obtained by weighting each particle with an appropriate angular factor [4]). Then $\Delta F_L(\bar{T}) = F_L^h(\bar{T}) - F_L^p(\bar{T})$ is a simple measure of the hadronization impact on F_L . This quantity is shown in Fig. 2, for one realistic simulation and two toy ones, for $u\bar{u}$ events at 91.2 GeV. In the realistic case, a parton shower is used to generate multiparton configurations, followed by string fragmentation. The shower develops down to a cut-off scale $Q_0 \approx 1$ GeV, so that also events in the first bin, $\bar{T} < 0.01$, can contain some gluons. The other two histograms are based on $\mathcal{O}(\alpha_s)$ matrix elements, where only 2- and 3-parton configurations are generated, with a cut $\bar{T} > 0.01$ on the latter to avoid the singularities of the 3-parton matrix element. Thus the first bin here represents pure 2-parton events. While one simulation is again based on string fragmentation, the other assumes isotropic smearing around the jet axes, basically the independent fragmentation scheme of Hoyer et al. [15,14].

(The same fragmentation parameters, tuned to the shower model, have been used in all three cases. A retuning of parameters for the $\mathcal{O}(\alpha_s)$ simulations would have given a larger nonperturbative p_\perp width σ to cover for the lack of perturbative gluons, and so would have implied even larger fragmentation smearing.)

The isotropic smearing is, as expected, giving a rather constant hadronization correction $\Delta F_L(\bar{T})$. There is some jump up in going from two to three jets that are smeared, followed by a slow but steady drop with \bar{T} , since the longitudinal component itself is increasing in importance with \bar{T} and therefore gives an increasing hadronization smearing of the longitudinal component onto the transverse one rather than only the other way around. By contrast, the string fragmentation provides a much steeper drop of ΔF_L with \bar{T} , kicking in immediately when going from two to three partons, and enhanced in the shower simulation relative to the simpler $\mathcal{O}(\alpha_s)$ one. At large \bar{T} the overall hadronization correction can even turn negative. Averaging over the \bar{T} spectrum (with mean value $\langle \bar{T} \rangle \approx 0.05$), we conclude that the typical hadronization smearing contribution is only about a third of the naively expected one, as obtained from 2-parton results. (Qualitatively this agrees with and probably explains a similar observation in [4] of smaller-than-expected hadronization corrections when using Herwig [16].) That is, if hadronization corrections are viewed as a power series in α_s , the $\mathcal{O}(\alpha_s)$ term is of opposite sign and almost as large as the $\mathcal{O}(1)$ one.

It should be remembered, however, that this is integrated over all x_E , and that we have no similar way of addressing results in specific x bins, since the parton and hadron x spectra are quite different. Thus it is likely that the $F_L(x_p)$ derived in the previous section is an underestimation, just like an $F_L(x_p)$ found without any smearing corrections is likely to be an overestimation, but it appears impossible to find the “correct” $F_L(x_p)$ without making detailed assumptions about the hadronization process.

4 Clustering

Given the problems with the above smearing recipe, we introduce a new strategy, based on the clustering approach. In a nutshell, we propose to rotate all hadrons to the direction of the cluster they belong to, as an approximate way of removing hadronization smearing effects. Only thereafter is $F_L(x)$ extracted from this modified $\cos\theta$ distribution. The strategy is explained further in the following.

In clustering algorithms, nearby hadrons are combined to form clusters/jets, in a way that should reflect the underlying partonic state, to some approximation. The combination process is controlled by (at least) one separation parameter, call it y_{min} , such that the final state contains no pair of clusters closer to each other than that. Clustering algorithms can be applied also to a partonic state, and here y_{min} provides a regularization of soft and collinear divergences in the perturbative cross sections. It is then meaningful to calculate the distribution of partons at a factorization scale $\mu^2 = y_{\text{min}} E_{\text{cm}}^2$, and define scale dependent fragmentation functions parameterizing the sub-

sequent soft-perturbative and nonperturbative hadronization. The latter should obey standard QCD evolution equations, starting from some unknown nonperturbative form at a low reference scale.

Over the years many cluster algorithms have been proposed [17], each with its strengths and weaknesses. In this article we adopt the Durham one [18], which is a standard for many perturbative calculations. The distance measure between two clusters i and j is

$$y_{ij} = \frac{2 \min(E_i^2, E_j^2)(1 - \cos \theta_{ij})}{E_{\text{vis}}^2}, \quad (7)$$

so that $y_{ij}^{1/2}$ roughly corresponds to the relative transverse momentum, scaled to the total visible energy E_{vis} ($= E_{\text{cm}}$ for an ideal detector).

If we begin by considering a simple $q\bar{q}$ event, it should reconstruct back to two clusters, unless y_{min} has been chosen very small. Since the momentum of a cluster is given by the vector sum of its constituent hadrons, it would resum opposite and compensating p_{\perp} kicks imparted to hadrons in the fragmentation process. The cluster direction should therefore be a better measure of the $q\bar{q}$ axis than that provided by the individual hadron momenta. It is the angular distribution of this axis that relates back to the polarization character of the $\gamma^*/Z^0 \rightarrow q\bar{q}$ decay, and that we want to be reflected in our extracted $F_L(x)$ and $F_T(x)$. Therefore it would be an improvement to rotate all hadrons in a cluster to sit along the cluster direction. That is, the θ of a hadron is redefined while its x value is unchanged.

At this level there is no contradiction with the smearing approach studied earlier. Then we smeared the simple partonic angular shapes to arrive at realistic hadronic ones to compare with data, now we un-smear the hadronic angles to approach the simple partonic distributions. There is one advantage, however: the clustering approach is not sensitive to the width of the p_{\perp} distribution, i.e. the σ parameter, unlike the smearing procedure. Of course, the p_{\perp} width still affects the typical error between the $q\bar{q}$ and cluster axes.

When considering multijet production, the y_{min} choice does become relevant, with $\mu^2 = y_{\text{min}} E_{\text{cm}}^2$ acting as a factorization scale, as noted above. For a large y_{min} all activity is clustered into the two quark jets, and neither the gluon structure nor $F_L(x)$ would be probed. For $y_{\text{min}} \rightarrow 0$ each hadron or parton is a cluster unto itself, and we are back at the starting point. So obviously some intermediate scale is to be preferred. Given that the typical hadronization p_{\perp} width is ~ 0.4 GeV, with a tail to larger values, one would conclude that clustering up to $p_{\perp} \sim 1$ GeV would be a sensible minimum to eliminate the bulk of the hadronization p_{\perp} smearing. At the Z^0 peak this translates into $y_{\text{min}} \gtrsim 0.0001$. In the upper end, we want to stay with a picture of multiple gluon emission as the norm, i.e. retain $F_L(x)$ as an inclusive quantity, in order not to overlap with traditional studies of gluon jets in exclusive 3-jet events. Since the average number of clusters per event is three for $y_{\text{min}} \approx 0.0025$, we conclude that $0.0001 \lesssim y_{\text{min}} \lesssim 0.0025$ is




Fig. 3. Event-by-event hadronization corrections $\Delta\sigma_L/\sigma_{\text{tot}} = (\sigma_L^h - \sigma_L^p)/\sigma_{\text{tot}}$ for different y_{min} scales (Jetset 7.4 PS and string fragmentation)

a reasonable range over which to study a scale dependent $F_L(x, \mu)$.

In Fig. 3, the distribution of event-by-event x_E -weighted and -integrated hadronization corrections $\Delta\sigma_L/\sigma_{\text{tot}} = (\sigma_L^h - \sigma_L^p)/\sigma_{\text{tot}}$ is shown for some different y_{min} scales, for events generated with parton showers and string fragmentation (superscripts h and p stand for hadron and parton level, respectively). We note the significant width of these distributions, showing that event-by-event fluctuations in the hadronization process are important and can be of either sign. Even if small by comparison, the mean $\langle \Delta\sigma_L/\sigma_{\text{tot}} \rangle$ does show a systematic bias, positive for small y_{min} and negative for large y_{min} . That is, at small y_{min} the hadronization smearing wins over the string effects, while it is the other way around for large y_{min} – but remember that this is only true when averaging over many events. Nevertheless, one possible criterion for a good choice of y_{min} would be where the two effects cancel, which then gives $y_{\text{min}} \approx 0.0002$, i.e. $\mu \approx 1.3$ GeV. While a sensible reference value, one should not take this particular value too seriously, since it is for one specific model, and for one specific set of model parameters. Somewhat different parameter values, like for the parton shower cut-off $Q_0 \approx 1$ GeV, defining the parton level of the events studied, could lead to slightly different “preferred” μ values.

It is important to note that we here have been considering the x -integrated quantity. This is of relevance if one e.g. would like to extract an α_s from an σ_L measurement, and so this deserves our interest. For the purpose of determining the differential x distribution, $F_L(x)$, however, one would have reason to fear that any bias could have an x dependence that would not be caught. In the string model, a string piece connecting two partons is boosted by an increasing velocity vector as the relative opening angle between the partons is decreased, and so the string effects spread upwards to larger x values. A warning signal is then that ΔF_L does depend quite significantly on \bar{T} , Fig. 4, i.e. clustering does not reduce the \bar{T} dependence noted in Fig. 2, but mainly shifts the overall level. Since \bar{T} probes




Fig. 4. Hadronization corrections $\Delta F_L(\bar{\tau}) = F_L^h(\bar{\tau}) - F_L^p(\bar{\tau})$ for different y_{\min} scales (Jetset 7.4 PS and string fragmentation)




Fig. 5. Ratio $\Delta F_L/F_L = (F_L - F_L^{\text{cluster}})/F_L$ for various y_{\min} values: the dashed line corresponds to $y_{\min} = 0.0001$, the solid one to $y_{\min} = 0.0002$, the dotted one to $y_{\min} = 0.0005$ and the dash-dotted one to $y_{\min} = 0.0025$

the topology of events, we also do expect this topology to reflect itself in an x dependence of hadronization corrections. As in the previous studies, this dependence is then likely to show up mainly in the lower end of the x range. At larger x , hadrons are rather well aligned with the jet axes, so, even with x -weighting, the few particles out there give a small contribution to the $\langle \Delta F_L \rangle$. In Fig. 5, the relative difference between the inclusive $F_L(x_p)$ and $F_L^{\text{cluster}}(x_p)$, obtained by replacing hadron angles with cluster angles for different y_{\min} values, is shown. It is clearly seen, indeed, that the hadronization corrections are only important at low x_p , unless y_{\min} is chosen too high. For the transverse component of the fragmentation function, $F_T(x_p)$, corrections have the same absolute amplitude but the opposite sign.

On the up side, the string effect has its perturbative equivalent in the color dipole [12]. That is, asymmetries also exist in the production of soft gluons around the direction of the harder partons of an event. Such soft parton emission, below the cut-off scale $Q_0 \approx 1$ GeV we have used, would thus largely fill in the same regions as the non-

perturbative hadron production, and with the same topology dependence. If one takes local parton-hadron duality [19] seriously, this equivalence should come very close. Our proposed strategy, to reset the θ angle of particles to that of the cluster they belong to, would be applicable also to such perturbatively calculated parton topologies.

5 Summary

The coherence phenomenon [20] kills the concept of gluon fragmentation functions that can be defined independently of the environment they are found in. The “hump-backed” shape of inclusive x spectra [19, 21] is an excellent illustration: by coherence the multiplication of partons/hadrons at small x is much less than if the hard partons could radiate/hadronize independently. The immediate consequence is that the expected “softer gluon than quark jets” picture is difficult to test. This impacts both on studies of gluon jets directly in identified 3-jet events and indirectly via $F_L(x)$. In this article we have illustrated some of these issues for the latter observable.

It appears safe to conclude that a straightforward extraction of $F_L(x)$ from hadron angular distributions exaggerates the rate of particles at small x that should be attributed to gluon jets, since even the hadronization of pure $q\bar{q}$ events induces a “false” $F_L(x)$ by p_{\perp} smearing. We have also here shown that a symmetric smearing around jet axes introduces a bias in the other direction, since it misses important string/drag effects that tend to make 3-jet events more 2-jetlike. In summary, there is no model independent extraction of a unique $F_L(x)$, especially not at small x values.

We therefore propose to introduce a scale dependent quantity $F_L(x, \mu^2)$. Particles are clustered, e.g. with the Durham algorithm, and thereafter assigned the θ angle of the cluster they belong to, while retaining their x value. Thus μ^2 sets an “un-smearing” scale, below which p_{\perp} fluctuations are killed. We find that $\mu \approx 1.3$ GeV gives opposite and compensating p_{\perp} smearing and string effects in Jetset simulations. While the exact number certainly is model dependent, the order is a sensible one, given that the average hadronization p_{\perp} is of the order of 0.4 GeV. If this then sets a reasonably lower limit, an upper one is related to the desire to stay away from the region of exclusive 2- or 3-jet events. Over an intermediate range, one could imagine several experimental determinations providing the scale dependence. We also remind the reader that, so far, our studies have only been intended for LEP1 energies. Coverage of a wider energy range, e.g. at LEP2, introduces s as a further scale of the process and allows more differential tests.

We have not considered here how $F_L(x)$ should be used to extract a gluon fragmentation function or, more likely, check one extracted e.g. from exclusive 3-jet data. In conventional quark and gluon fragmentation function parameterizations [22], transverse momentum degrees of freedom are ignored in leading order, and only enter by next-to-leading order perturbative corrections, which cannot fully account for nonperturbative jet broadening [4]. On the

other hand, we have seen how, for $F_L(x, \mu^2)$, string effects largely cancel the nonperturbative jet broadening for factorization scales in the range around 1 GeV. Since many parameterizations actually make their ansatz for quark and gluon fragmentation functions at a starting scale in that order, it would appear that data analyzed according to the suggestions in this article could provide an interesting input for these theoretical studies. If nothing else, it would allow a better understanding to be gained of systematic errors coming from the choice of methodology.

References

1. T. Sjöstrand, *Computer Phys. Commun.* **82**, 74 (1994)
2. A.C. Hirschfeld, G. Kramer, *Nucl. Phys. B* **74**, 211 (1974)
3. G. Altarelli et al., *Nucl. Phys. B* **160**, 301 (1979); *Phys. Rep.* **81**, 1 (1982)
4. P. Nason, B.R. Webber, *Nucl. Phys. B* **421**, 473 (1994), erratum *B* **480**, 755 (1996)
5. OPAL Coll., G. Abbiendi et al., *Eur. Phys. J. C* **11**, 217 (1999); DELPHI Coll., P. Abreu et al., *Eur. Phys. J. C* **13**, 573 (2000); ALEPH Coll., R. Barate et al., *Eur. Phys. J. C* **17**, 1 (2000)
6. TASSO Coll., R. Brandelik et al., *Phys. Lett. B* **114**, 65 (1982); OPAL Coll., R. Akers et al., *Z. Phys. C* **68**, 203 (1995); ALEPH Coll., D. Buskulic et al., *Phys. Lett. B* **357**, 487 (1995); DELPHI Coll., P. Abreu et al., *Eur. Phys. J. C* **6**, 19 (1999)
7. B. Andersson, G. Gustafson, G. Ingelman, T. Sjöstrand, *Phys. Rep.* **97**, 31 (1983); B. Andersson, *The Lund model* (Cambridge University Press, 1998)
8. B.R. Webber, *Nucl. Phys. B* **238**, 492 (1984)
9. Yu.L. Dokshitzer, B.R. Webber, *Phys. Lett. B* **352**, 451 (1995); M. Dasgupta, B.R. Webber, *Nucl. Phys. B* **484**, 247 (1997); M. Beneke, V.M. Braun, L. Magnea, *Nucl. Phys. B* **497**, 297 (1997)
10. I.G. Knowles et al., in *Physics at LEP2*, edited by G. Altarelli, T. Sjöstrand, F. Zwirner, CERN 96-01, Vol. 2, p. 103
11. B. Andersson, G. Gustafson, T. Sjöstrand, *Phys. Lett. B* **94**, 211 (1980)
12. Ya.I. Azimov, Yu.L. Dokshitzer, V.A. Khoze, S.I. Troyan, *Phys. Lett. B* **165**, 147 (1985)
13. JADE Coll., W. Bartel et al., *Z. Phys. C* **21**, 37 (1983); OPAL Coll., R. Akers et al., *Z. Phys. C* **68**, 531 (1995); DELPHI Coll., P. Abreu et al., *Z. Phys. C* **70**, 179 (1996); ALEPH Coll., R. Barate et al., *Phys. Rep.* **294**, 1 (1998)
14. T. Sjöstrand, *Z. Phys. C* **26**, 93 (1984)
15. P. Hoyer, P. Osland, H.G. Sander, T.F. Walsh, P.M. Zerwas, *Nucl. Phys. B* **161**, 349 (1979)
16. G. Marchesini, B.R. Webber, G. Abbiendi, I.G. Knowles, M.H. Seymour, L. Stanco, *Computer Phys. Commun.* **67**, 465 (1992)
17. For a review see e.g. S. Moretti, L. Lönnblad, T. Sjöstrand, *J. High Energy Phys.* **08**, 001 (1998)
18. S. Catani, Yu.L. Dokshitzer, M. Olsson, G. Turnock, B.R. Webber, *Phys. Lett. B* **269**, 432 (1991)
19. Ya.I. Azimov, Yu.L. Dokshitzer, V.A. Khoze, S.I. Troyan, *Z. Phys. C* **27**, 65 (1985), *Z. Phys. C* **31**, 213 (1986)
20. A.H. Mueller, *Phys. Lett. B* **104**, 161 (1981); B.I. Ermolaev, V.S. Fadin, *JETP Lett.* **33**, 269 (1981)
21. Yu.L. Dokshitzer, V.S. Fadin, V.A. Khoze, *Phys. Lett. B* **115**, 242 (1982); *Z. Phys. C* **15**, 325 (1982); *Z. Phys. C* **18**, 37 (1983); A. Bassetto, M. Ciafaloni, G. Marchesini, A.H. Mueller, *Nucl. Phys. B* **207**, 189 (1982)
22. J. Binnewies, B.A. Kniehl, G. Kramer, *Z. Phys. C* **65**, 471 (1995); *Phys. Rev. D* **52**, 4947 (1995); P.J. Rijken, W.L. van Neerven, *Phys. Lett. B* **386**, 422 (1996); *Phys. Lett. B* **392**, 207 (1997); *Nucl. Phys. B* **487**, 233 (1997); B.A. Kniehl, G. Kramer, B. Pötter, *Nucl. Phys. B* **582**, 514 (2000); S. Kretzer, *Phys. Rev. D* **62**, 054001 (2000)