

The radiative return at small angles: virtual corrections

J.H. Kühn^{1,2,a}, G. Rodrigo^{1,b}

¹ Theory Division, CERN, 1211 Geneva 23, Switzerland

² Institut für Theoretische Teilchenphysik, Universität Karlsruhe, 76128 Karlsruhe, Germany

Received: 24 April 2002 /

Published online: 12 July 2002 – © Springer-Verlag / Società Italiana di Fisica 2002

Abstract. Virtual corrections for electron–positron annihilation into one real and one off-shell photon of invariant mass Q^2 are evaluated. Special attention is paid to those configurations where the real photon is collinear with the beam direction. This calculation is an important ingredient of a Monte Carlo program, which simulates events with tagged photons from initial-state radiation, including NLO corrections.

1 Introduction

The total cross section for electron–positron annihilation into hadrons is one of the fundamental observables in particle physics. Its high energy behaviour provides one of the first and still most convincing arguments for the point-like nature of quarks. Its normalization was evidence for the existence of quarks of three different colours, and the recent, precise measurements even allow for an excellent determination of the strong coupling at very high [1] and intermediate energies (e.g. [2] and references therein) through the influence of QCD corrections.

Weighted integrals over the cross section with properly chosen kernels are, furthermore, a decisive input for electroweak precision tests. This applies, for example, to the electromagnetic coupling at higher energies or to the anomalous magnetic moment of the muon.

Of particular importance for these two applications is the low energy region, say from threshold up to centre-of-mass (cms) energies of approximately 3 GeV and 10 GeV, respectively. Recent measurements based on energy scans between 2 and 5 GeV have improved the accuracy in part of this range. However, similar, or even further improvements below 2 GeV would be highly welcome. The region between 1.4 GeV and 2 GeV, in particular, is poorly studied and no collider will cover this region in the near future. Improvements or even an independent cross-check of the precise measurements of the pion form factor in the low energy region by the CMD2 and DM2 collaborations would be extremely useful, since this dominates in the analysis of the muon anomalous magnetic moment.

Experiments at present electron–positron colliders operate mostly at fixed energies, albeit with enormous lumi-

nosity, with BaBar and BELLE at 10.6 GeV and KLOE at 1.02 GeV as most prominent examples.

This peculiar feature allows the use of the radiative return, i.e. the reaction

$$e^+(p_1) + e^-(p_2) \rightarrow \gamma(k) + \gamma^*(Q) (\rightarrow \text{hadrons}), \quad (1)$$

to explore a wide range of Q^2 in a single experiment [3–9].

Nominally masses of the hadronic system between $2m_\pi$ and the cms energy of the experiment are accessible. In practice it is useful to consider only events with a hard photon — tagged or untagged — to clearly identify the reaction, which lowers the energy significantly.

The study of events with photons emitted under both large and small angles, and thus at a significantly enhanced rate, is particularly attractive for the $\pi^+\pi^-$ final state with its clear signature, an investigation performed at present at DAΦNE [10–12]. Events with a tagged photon, emitted under a large angle with respect to the beam, have a clear signature and are thus particularly suited to the analysis of hadronic final states of higher multiplicity [13].

The inclusion of radiative corrections is essential for the precise extraction of the cross section, which is necessarily based on a Monte Carlo simulation. A first program, called EVA, was constructed some time ago [3]. It includes initial-state radiation (ISR), final-state radiation (FSR), their interference, and the dominant radiative corrections from additional collinear radiation through structure function techniques [14].

The complete NLO corrections have recently been implemented in a program called PHOKHARA [15]. Both programs, however, were designed to simulate reactions with tagged photons, i.e. at least one photon was required to be emitted under large angles.

An important ingredient in the extension of the NLO Monte Carlo program PHOKHARA to small photon angles is the evaluation of the virtual corrections to the reaction (1) in the limit $m_e^2/s \ll 1$, which are equally valid

^a e-mail: jk@particle.uni-karlsruhe.de

^b Supported in part by E.U. TMR grant HPMF-CT-2000-00989; e-mail: german.rodrigo@cern.ch

for large and small angles. Compact results for the one-loop two-, three- and four-point functions that enter this calculation can be found in the literature [16,17] for arbitrary values of m_e^2/s . However, the combination of these analytical expressions with the relevant coefficients is numerically unstable in the limit of small mass and angles. A compact, numerically stable result, valid for an arbitrarily small photon angle, is therefore required. As a consequence of the highly singular kinematic coefficients, terms proportional to m_e^2 and even m_e^4 must be kept in the expansion, which after angular integration will contribute to the total cross section even in the limit $m_e^2/s \rightarrow 0$.

The present paper extends the analysis of [18] where the corrections from virtual and soft photons were presented for the case of large angles. In Sect. 2 we recall the basic definitions and describe the systematic procedure used in the expansion of the results for small m_e^2/s and small angles simultaneously. The analytic results for real and imaginary parts of the leptonic tensor, expressed in an angular momentum basis, are presented in Sect. 3 and compared with results for related quantities that can be found in the literature. After summation over the polarizations of the virtual photons, our result agrees with the one of Berends, Burgers and van Neerven [20,21]. The result of Kuraev, Merenkov and Fadin [22] for the real part of the tensor, which was obtained for virtual Compton scattering $\gamma^* + e^- \rightarrow \gamma + e^-$ is related to our case by proper analytic continuation; and indeed after analytic continuation we find agreement for the real part¹. Section 4 contains our summary and the conclusions. The mass-dependent terms proportional to m_e^2 and m_e^4 , expressed in the Cartesian basis, are given in Appendix A. The scalar loop integrals needed for the calculation are listed in Appendix B.

2 The leptonic tensor for the radiative return

Consider the e^+e^- annihilation process

$$e^+(p_1) + e^-(p_2) \rightarrow \gamma^*(Q) + \gamma(k_1), \quad (2)$$

where the virtual photon decays into a hadronic final state, $\gamma^*(Q) \rightarrow \text{hadrons}$, and the real one is emitted from the initial state. The differential rate can be cast into the product of a leptonic and a hadronic tensor and the corresponding factorized phase space

$$d\sigma = \frac{1}{2s} L_{\mu\nu} H^{\mu\nu} d\Phi_2(p_1, p_2; Q, k_1) d\Phi_n(Q; q_1, \dots, q_n) \frac{dQ^2}{2\pi}, \quad (3)$$

where $d\Phi_n(Q; q_1, \dots, q_n)$ denotes the hadronic n -body phase space, including all the statistical factors coming from the hadronic final state.

For an arbitrary hadronic final state, the matrix element for the diagrams in Fig. 1 can be written

$$M_0 = M_0^\mu J_\mu,$$




Fig. 1. Initial-state radiation in the annihilation process $e^+e^- \rightarrow \gamma + \text{hadrons}$ at the Born level

where J_μ is the hadronic current and M_0^μ is the leptonic current in lowest order. Summing over the polarizations of the real photon, averaging over the polarizations of the initial e^+e^- state, and using current conservation, $Q_\mu J^\mu = 0$, the leptonic tensor

$$L_0^{\mu\nu} = \overline{M_0^\mu M_0^{\nu\dagger}}$$

can be written in the following form:

$$L_0^{\mu\nu} = \frac{(4\pi\alpha/s)^2}{q^4} \left[\left(\frac{2m^2q^2(1-q^2)^2}{y_1^2y_2^2} - \frac{2q^2 + y_1^2 + y_2^2}{y_1y_2} \right) g^{\mu\nu} + \left(\frac{8m^2}{y_2^2} - \frac{4q^2}{y_1y_2} \right) \frac{p_1^\mu p_1^\nu}{s} + \left(\frac{8m^2}{y_1^2} - \frac{4q^2}{y_1y_2} \right) \frac{p_2^\mu p_2^\nu}{s} - \left(\frac{8m^2}{y_1y_2} \right) \frac{p_1^\mu p_2^\nu + p_1^\nu p_2^\mu}{s} \right], \quad (4)$$

with

$$y_i = \frac{2k_1 \cdot p_i}{s}, \quad m^2 = \frac{m_e^2}{s}, \quad q^2 = \frac{Q^2}{s}. \quad (5)$$

It is symmetric under the exchange of the electron and the positron momenta. Expressing the bilinear products y_i by the photon emission angle in the cms frame

$$y_{1,2} = \frac{1-q^2}{2} (1 \mp \beta \cos \theta), \quad \beta = \sqrt{1-4m^2},$$

and rewriting the two-body phase space

$$d\Phi_2(p_1, p_2; Q, k_1) = \frac{1-q^2}{32\pi^2} d\Omega, \quad (6)$$

it is evident that expression (4) contains several singularities: soft singularities for $q^2 \rightarrow 1$ and collinear singularities for $\cos \theta \rightarrow \pm 1$. The former are avoided by requiring a minimal photon energy. The latter are regulated by the electron mass.

The physics of the hadronic system, whose description is model-dependent, enters only through the hadronic tensor

$$H^{\mu\nu} = J^\mu J^{\nu\dagger}, \quad (7)$$

where the hadronic current has to be parametrized through form factors. For two charged pions in the final state, the current

$$J_{2\pi}^\mu = ieF_{2\pi}(Q^2) (q_{\pi^+} - q_{\pi^-})^\mu, \quad (8)$$

where q_{π^+} and q_{π^-} are the momenta of the π^+ and π^- respectively, is determined by only one function, the pion

¹ We disagree, however, with (30) of the translated version [23] which contains a missprint

form factor $F_{2\pi}$ [24]. The hadronic current for four pions exhibits a more complicated structure and has been discussed in [5, 25, 26].

At NLO, the leptonic tensor receives contributions both from one-loop corrections arising from the insertion of virtual photon lines in the tree diagrams of Fig. 1 and from the emission of an extra real photon from the initial state. In this paper, we consider only the emission of soft photons. The implementation of these results in the program PHOKHARA and the discussion of their physical consequences will be considered in a separate work [27].

At NLO, the leptonic tensor has the following general form²:

$$L^{\mu\nu} = \frac{(4\pi\alpha/s)^2}{q^4} \left[a_{00} g^{\mu\nu} + a_{11} \frac{p_1^\mu p_1^\nu}{s} + a_{22} \frac{p_2^\mu p_2^\nu}{s} + a_{12} \frac{p_1^\mu p_2^\nu + p_2^\mu p_1^\nu}{s} + i\pi a_{-1} \frac{p_1^\mu p_2^\nu - p_2^\mu p_1^\nu}{s} \right]. \quad (9)$$

Terms proportional to Q^μ are absent as a consequence of current conservation. The scalar coefficients a_{ij} and a_{-1} allow the following expansion

$$a_{ij} = a_{ij}^{(0)} + \frac{\alpha}{\pi} a_{ij}^{(1)}, \quad a_{-1} = \frac{\alpha}{\pi} a_{-1}^{(1)}. \quad (10)$$

The LO coefficients $a_{ij}^{(0)}$ can be directly read from (4)

$$\begin{aligned} a_{00}^{(0)} &= \frac{2m^2 q^2 (1-q^2)^2}{y_1^2 y_2^2} - \frac{2q^2 + y_1^2 + y_2^2}{y_1 y_2}, \\ a_{11}^{(0)} &= \frac{8m^2}{y_2^2} - \frac{4q^2}{y_1 y_2}, \quad a_{22}^{(0)} = a_{11}^{(0)}(y_1 \leftrightarrow y_2), \\ a_{12}^{(0)} &= -\frac{8m^2}{y_1 y_2}. \end{aligned} \quad (11)$$

The imaginary antisymmetric piece proportional to a_{-1} appears for the first time at NLO. The leptonic tensor therefore remains fully symmetric only at LO.

As an alternative one can replace the Cartesian basis ((9)) by a basis derived from the three circular polarization vectors of the virtual photon ε_L and ε_\pm , defined through

$$\begin{aligned} \varepsilon_L^\mu &= \frac{2(q^\mu q^\nu - g^{\mu\nu} q^2)}{\sqrt{s} q^2 (1-q^2)} k_{1\nu}, \\ \varepsilon_1^\mu &= \frac{2(q^\mu q^\nu - g^{\mu\nu} q^2)}{\sqrt{s} q^2 (1-q^2) \beta \sin \theta} \\ &\quad \times ((y_1 - q^2 y_2) p_{1\nu} - (y_2 - q^2 y_1) p_{2\nu}), \\ \varepsilon_2^\mu &= \frac{2(q^\mu q^\nu - g^{\mu\nu} q^2)}{\sqrt{s^3} q^2 (1-q^2) \beta \sin \theta} \varepsilon_{\nu\eta\rho\sigma} k_1^\eta p_1^\rho p_2^\sigma, \\ \varepsilon_\pm^\mu &= \frac{1}{\sqrt{2}} (\varepsilon_1^\mu \pm i\varepsilon_2^\mu), \end{aligned} \quad (12)$$

where $\varepsilon_{\nu\eta\rho\sigma}$ is the four-dimensional totally antisymmetric Levi-Civita tensor, $\varepsilon_{0123} = -1$. The leptonic tensor is thus

² The present convention differs from the one in [18] by a factor $1/(y_1 y_2)$

given by

$$L^{\mu\nu} = \frac{(4\pi\alpha/s)^2}{q^4} \sum a_{ij} \varepsilon_i^{*\mu} \varepsilon_j^\nu, \quad i, j = L, \pm. \quad (13)$$

Only four of the scalar coefficients are independent

$$\begin{aligned} a_{L-} &= a_{L+}, & a_{-L} &= a_{+L} = a_{L+}^*, \\ a_{--} &= a_{++}, & a_{+-} &= a_{-+}. \end{aligned}$$

The trace of the leptonic tensor

$$L^{\mu\nu} (q_\mu q_\nu - g_{\mu\nu} q^2) = \frac{(4\pi\alpha/s)^2}{q^2} (a_{LL} + 2a_{++}) \quad (14)$$

is related to the cross section after angular averaging of the hadronic tensor.

The relations between the components in the Cartesian and the circular basis read as follows:

$$\begin{aligned} a_{LL} &= -a_{00} + \frac{1}{4q^2(1-q^2)^2} [(y_2 - q^2 y_1)^2 a_{11} \\ &\quad + (y_1 - q^2 y_2)^2 a_{22} + 2(y_1 - q^2 y_2)(y_2 - q^2 y_1) a_{12}], \\ a_{L+} &= \frac{\beta \sin \theta}{4\sqrt{2}q^2(1-q^2)} [(y_2 - q^2 y_1) a_{11} - (y_1 - q^2 y_2) a_{22} \\ &\quad + (1+q^2)(y_1 - y_2) a_{12} - i\pi a_{-1}], \\ a_{+-} &= \frac{\beta^2 \sin^2 \theta}{8} (a_{11} + a_{22} - 2a_{12}), \\ a_{++} &= a_{+-} - a_{00}. \end{aligned} \quad (15)$$

Conversely

$$\begin{aligned} a_{00} &= a_{+-} - a_{++}, \\ a_{11} &= \frac{4}{(1-q^2)^2} \left[q^2 (a_{LL} + a_{+-} - a_{++}) \right. \\ &\quad + \frac{2\sqrt{2}q^2(y_1 - q^2 y_2)}{(1-q^2)\beta \sin \theta} \text{Re}(a_{L+}) \\ &\quad \left. + \frac{2(y_1 - q^2 y_2)^2}{(1-q^2)^2 \beta^2 \sin^2 \theta} a_{+-} \right], \\ a_{22} &= a_{11}(y_1 \leftrightarrow y_2), \quad a_{12} = \frac{a_{11} + a_{22}}{2} - \frac{4a_{+-}}{\beta^2 \sin^2 \theta}, \\ a_{-1} &= -\frac{4\sqrt{2}q^2}{\pi(1-q^2)\beta \sin \theta} \text{Im}(a_{L+}). \end{aligned} \quad (16)$$

The scalar coefficients in the circular basis are given at LO by

$$\begin{aligned} a_{LL}^{(0)} &= \frac{2q^2 \beta^2 \sin^2 \theta}{y_1 y_2}, \\ a_{L+}^{(0)} &= \frac{\sqrt{q^2}(y_1 - y_2)\beta \sin \theta}{\sqrt{2}(1-q^2)y_1 y_2} \left(1 + q^2 - \frac{2m^2(1-q^2)^2}{y_1 y_2} \right), \\ a_{+-}^{(0)} &= \frac{\beta^2 \sin^2 \theta}{y_1 y_2} \left(-q^2 + \frac{m^2(1-q^2)^2}{y_1 y_2} \right), \\ a_{++}^{(0)} &= a_{+-}^{(0)} + \frac{2q^2 + y_1^2 + y_2^2}{y_1 y_2} - \frac{2m^2 q^2 (1-q^2)^2}{y_1^2 y_2^2}. \end{aligned} \quad (17)$$




Fig. 2. One-loop corrections to initial-state radiation in e^+e^- annihilation with the emission of a virtual photon

The one-loop matrix elements (Fig. 2) contribute to the leptonic tensor through their interference with the lowest order diagrams (Fig. 1). They contain ultraviolet (UV) and infrared (IR) divergences, which are regularized using dimensional regularization in $D = 4 - 2\epsilon$ dimensions. The UV divergences are renormalized in the on-shell scheme. The IR divergences are cancelled by adding the contribution of an extra soft photon emitted from the initial state and integrated in the phase space up to an energy cutoff $E_\gamma < w\sqrt{s}$ far below \sqrt{s} . The result, which is finite, depends, however, on this soft photon cutoff. Only the contribution from hard photons with energy $E_\gamma > w\sqrt{s}$ would cancel this dependence.

The algebraic manipulations have been carried out with the help of the *FeynCalc* Mathematica package [28]. Using standard techniques [29], it automatically reduces the evaluation of the one-loop contribution to the calculation of a few scalar one-loop integrals and performs the Dirac algebra.

Since we consider the small angular region, mass terms proportional to y_i^{-2} and even y_i^{-3} arise. Terms proportional to m^2 and even m^4 must be kept, if they are multiplied by y_i^{-2} and y_i^{-3} respectively. In the expansion of the one-loop integrals, functions that depend on the ratio m^2/y_i cannot be expanded, in contrast to functions of m^2 or y_i separately. To arrive at a systematic approach we therefore make the replacements $m^2 \rightarrow \lambda m^2$, $y_i \rightarrow \lambda y_i$, perform the expansion for small λ up to the appropriate order, and set $\lambda = 1$ at the end.

3 The NLO leptonic tensor

Combining the one-loop and the soft contribution we now arrive at the leptonic tensor in NLO. It will be convenient

to split the coefficients $a_{ij}^{(1)}$ into a part that contributes at large angles and a part proportional to m^2 and m^4 , denoted by $a_{ij}^{(1,0)}$ and $a_{ij}^{(1,m)}$ respectively:

$$a_{ij}^{(1)} = a_{ij}^{(0)} \left[-\log(4w^2)[1 + \log(m^2)] - \frac{3}{2} \log\left(\frac{m^2}{q^2}\right) - 2 + \frac{\pi^2}{3} \right] + a_{ij}^{(1,0)} + a_{ij}^{(1,m)}. \quad (18)$$

The factor proportional to the LO coefficients $a_{ij}^{(0)}$ contains the usual soft and collinear logarithms. The expressions are particularly compact in the circular basis. For completeness we also repeat the results for $a_{ij}^{(1,0)}$, which can be found in [18]³, albeit in the Cartesian basis:

$$a_{LL}^{(1,0)} = \frac{2q^2}{(1-q^2)^2} \left\{ -\frac{y_1}{1-y_1} - 2\log(q^2) + 2L(y_1) + \left[1 - \frac{1-q^2}{1-y_2} - \frac{q^2}{(1-y_2)^2} \right] \times \log\left(\frac{y_1}{q^2}\right) + [y_1 \leftrightarrow y_2] \right\}, \quad (19)$$

$$a_{L+}^{(1,0)} = \frac{\beta \sin \theta}{4\sqrt{2q^2(1-q^2)}} \left\{ \frac{y_1(1+q^2)^2}{(1-y_1)(1-y_2)} - \frac{(1-q^2)^2}{y_2} - \frac{2q^2(2+q^2)}{y_2} \log(q^2) + \left[2(1-q^2) \left(\frac{1}{y_1} - \frac{q^2}{y_2} \right) - \frac{2(1+q^2)}{1-y_2} - \frac{q^2(1+q^2)^2}{(1-y_2)^2} \right] \log\left(\frac{y_1}{q^2}\right) - 2q^2 \left[\frac{1+2q^2}{y_1} + \frac{y_1}{y_2^2} \right] L(y_1) - \frac{i\pi q^2}{y_1 y_2} \left[\frac{2\log(1-y_1)}{y_1} + \frac{1-q^2}{1-y_1} + \frac{q^2}{(1-y_1)^2} \right] - [y_1 \leftrightarrow y_2] \right\}, \quad (20)$$

$$a_{++}^{(1,0)} = \frac{1}{2} \left\{ \frac{1-y_2}{y_1} + \frac{1}{(1-q^2)^2} \left(\frac{1+q^4}{1-y_1} - 2 + 2\log(q^2) \right) + \frac{1}{1-q^2} \left[\frac{1+q^4}{1-q^2} \left(\frac{q^2}{(1-y_2)^2} - 1 \right) + \frac{3-q^4}{1-y_2} \right] \times \log\left(\frac{y_1}{q^2}\right) + \frac{2}{1-q^2} \left[\frac{(1+q^4)y_2}{(1-q^2)y_1} + \frac{2}{y_2} \right] \times L(y_1) + [y_1 \leftrightarrow y_2] \right\}, \quad (21)$$

$$a_{+-}^{(1,0)} = \frac{q^2}{(1-q^2)^2} \left\{ \frac{1+q^4}{2q^2(1-y_1)} - \frac{1}{q^2} - \frac{y_2}{y_1} \log(q^2) - \left[1 + \frac{(1-q^2)^2}{y_2} - \frac{q^2(1-q^2)}{1-y_2} - \frac{1+q^4}{2(1-y_2)^2} \right] \times \log\left(\frac{y_1}{q^2}\right) - \left[2 + \frac{(1-q^2)^2}{y_2^2} \right] L(y_1) + [y_1 \leftrightarrow y_2] \right\}, \quad (22)$$

³ The result for the imaginary part of a_{L+} differs from a_{-1} in the original version of [18]

with

$$L(y_i) = Li_2\left(\frac{-y_i}{q^2}\right) - Li_2\left(1 - \frac{1}{q^2}\right) + \log(q^2 + y_i) \log\left(\frac{y_i}{q^2}\right),$$

where Li_2 is the Spence or dilogarithm function. The result in the Cartesian basis has been given in [18]. The terms proportional to powers of m^2 are given by

$$a_{LL}^{(1,m)} = 0, \quad (23)$$

$$a_{L+}^{(1,m)} = \frac{\sqrt{q^2} \beta \sin \theta}{4\sqrt{2}} \left\{ \frac{4m^2}{y_1^2} \left[\log(q^2) \log\left(\frac{y_1^4}{m^4 q^2}\right) + 4Li_2(1 - q^2) + \frac{3}{2} \left(Li_2\left(1 - \frac{y_1}{m^2}\right) - \frac{\pi^2}{6} \right) \right] - n\left(y_1, \frac{2(1 - 3q^2)}{q^2}\right) + \frac{2m^2 N(y_1)}{y_1(m^2(1 - q^2) - y_1)} - [y_1 \leftrightarrow y_2] \right\}, \quad (24)$$

$$a_{++}^{(1,m)} = -\frac{m^2 q^2}{y_1^2} \left[\log(q^2) \log\left(\frac{y_1^4}{m^4 q^2}\right) + 4Li_2(1 - q^2) + Li_2\left(1 - \frac{y_1}{m^2}\right) - \frac{\pi^2}{6} \right] - \frac{m^2(1 - q^2)}{y_1^2} \times \left[1 - \log\left(\frac{y_1}{m^2}\right) + \frac{m^2}{y_1} \left(Li_2\left(1 - \frac{y_1}{m^2}\right) - \frac{\pi^2}{6} \right) \right] + \frac{q^2}{2} n\left(y_1, \frac{1 - 3q^2}{q^2}\right) + [y_1 \leftrightarrow y_2], \quad (25)$$

and

$$a_{+-}^{(1,m)} = \frac{\beta^2 \sin^2 \theta}{8} \left\{ \frac{4m^2}{y_1^2} \left[\log(q^2) \log\left(\frac{y_1^4}{m^4 q^2}\right) + 4Li_2(1 - q^2) + 2 \left(Li_2\left(1 - \frac{y_1}{m^2}\right) - \frac{\pi^2}{6} \right) \right] - \frac{1 - q^2}{q^2} n\left(y_1, \frac{3 - 7q^2}{1 - q^2}\right) + \frac{2m^2(1 - q^2)}{y_1(m^2(1 - q^2) - y_1)} \times \left[\frac{1}{q^2} \log\left(\frac{y_1}{m^2}\right) + \frac{\log(q^2)}{1 - q^2} \right] + \left(\frac{3 - q^2}{1 - q^2} + \frac{m^2}{m^2(1 - q^2) - y_1} \right) N(y_1) \right\} + [y_1 \leftrightarrow y_2]. \quad (26)$$

The coefficient a_{L+} is antisymmetric with respect to the exchange $[y_1 \leftrightarrow y_2]$, while all the others are symmetric. Only $a_{L+}^{(1,0)}$ contributes to the imaginary part. Notice that the mass-suppressed terms are all real. The functions $n(y_i, z)$ and $N(y_i)$ are defined through

$$n(y_i, z) = \frac{m^2}{y_i(m^2 - y_i)} \left[1 + z \log\left(\frac{y_i}{m^2}\right) \right] + \frac{m^2}{(m^2 - y_i)^2} \log\left(\frac{y_i}{m^2}\right), \quad (27)$$

$$N(y_i) = \log(q^2) \log\left(\frac{y_i}{m^2}\right) + Li_2(1 - q^2) + Li_2\left(1 - \frac{y_i}{m^2}\right) - \frac{\pi^2}{6}. \quad (28)$$

The apparent singularity of n inside the limits of phase space is compensated by the zero in the numerator. For the numerical evaluation in the region y_i close to m^2 we use

$$n(y_i, z)|_{y_i \rightarrow m^2} = \frac{1}{y_i} \left[1 + z \log\left(\frac{y_i}{m^2}\right) \right] - \frac{1}{m^2} \sum_{n=0} \left(\frac{1}{n+2} + \frac{z}{n+1} \right) \left(1 - \frac{y_i}{m^2} \right)^n. \quad (29)$$

For the conversion from the circular to the Cartesian basis, and to ensure finite results in the limit $\sin \theta \rightarrow 0$, it is important that a_{L+} and a_{+-} vanish at $\sim \sin \theta$ and $\sin^2 \theta$ respectively. This corresponds to the requirement that the factors in curly brackets do not diverge for small $\sin \theta$, i.e. in the limit $(m^2(1 - q^2) - y_i) \rightarrow 0$. This is guaranteed by the behaviour of $N(y_i)$ for $y_i \rightarrow m^2(1 - q^2)$:

$$\frac{m^2 N(y_i)}{m^2(1 - q^2) - y_i} \Big|_{y_i \rightarrow m^2(1 - q^2)} = -\frac{\log(1 - q^2)}{q^2} - \frac{\log(q^2)}{1 - q^2}, \quad (30)$$

The results for $a_{ij}^{(1,m)}$ in the Cartesian basis are listed in Appendix A.

We note that the imaginary part of $L_{\mu\nu}$, which is present in the coefficients a_{L+} or a_{-1} only, is of interest for those cases where the hadronic current receives contributions from different amplitudes with non-trivial relative phases. This is possible, e.g. for final states with three or more mesons or for $p\bar{p}$ production.

4 Tests of the result

After summation over the polarizations of the virtual photon the differential rate is given by the properly contracted leptonic tensor:

$$\begin{aligned} & \frac{q^2}{(4\pi\alpha/s)^2} L^{\mu\nu}(q_\mu q_\nu - g_{\mu\nu} q^2) = a_{LL} + 2a_{++} \\ & = -3 a_{00} + \left(\frac{1}{4q^2} (1 - y_1)^2 - m^2 \right) a_{11} \\ & + \left(\frac{1}{4q^2} (1 - y_2)^2 - m^2 \right) a_{22} \\ & + \left(\frac{1}{2q^2} (1 - y_1)(1 - y_2) - (1 - 2m^2) \right) a_{12} \\ & = L_0 \left\{ 1 + \frac{\alpha}{\pi} \left[-\log(4w^2) [1 + \log(m^2)] \right. \right. \\ & \left. \left. - \frac{3}{2} \log\left(\frac{m^2}{q^2}\right) - 2 + \frac{\pi^2}{3} \right] \right\} + \frac{\alpha}{\pi} \left\{ 4 \left(\frac{1 + (1 - y_2)^2}{2y_1 y_2} L(y_1) \right. \right. \\ & \left. \left. + \frac{1 - 2q^2}{2(1 - q^2)^2} \log(q^2) + \left[1 - \frac{y_1 - 2y_2}{2(1 - y_2)} - \frac{y_1 y_2}{4(1 - y_2)^2} \right] \right) \right\} \end{aligned}$$

$$\begin{aligned} & \times \log\left(\frac{y_1}{q^2}\right) - \frac{1}{2(1-q^2)} + \frac{1}{4(1-y_1)} + \frac{1-y_2}{4y_1} \\ & + [y_1 \leftrightarrow y_2] - 2a_{++}^{(1,m)} \Big\}, \end{aligned} \quad (31)$$

where

$$\begin{aligned} L_0 &= \frac{q^2}{(4\pi\alpha/s)^2} L_0^{\mu\nu} (q_\mu q_\nu - q^2 g_{\mu\nu}) \\ &= -\frac{2}{y_1 y_2} \left[2q^2 + y_1^2 + y_2^2 - \frac{2m^2 q^2 (1-q^2)^2}{y_1 y_2} \right], \end{aligned} \quad (32)$$

$L_0^{\mu\nu}$ being the leptonic tensor at LO. In L_0 only the relevant terms in the limit $m^2 \rightarrow 0$ have been kept. Large angle terms and mass corrections are in agreement with Berends et al. [20,21]. After proper analytic continuation and using the substitutions

$$t \rightarrow -y_1 + i\eta, \quad s \rightarrow -y_2 + i\eta, \quad u \rightarrow s + i\eta, \quad (33)$$

the leptonic tensor in (9) is closely related⁴ to the tensor $T_{\mu\nu}$, which describes ‘‘virtual Compton scattering’’ and has been calculated by Kuraev et al. [22]. After performing this analytical continuation, the results are in agreement (However, the result for T_{12} printed in [23] contains a typo and would give rise to a discrepancy.).

5 Conclusions

Compact analytical formulae have been obtained for the virtual corrections to the process $e^+e^- \rightarrow \gamma\gamma^*$, which are valid for photon emission under both large and small angles. After proper analytic continuation the results are in agreement with those obtained in [22] for the reaction $\gamma^* + e^- \rightarrow \gamma + e^-$. In polarization averaged form they are in agreement with those for $e^+e^- \rightarrow \gamma Z$ from [20,21].

Acknowledgements. We would like to thank G. Cataldi, A. Denig, S. Di Falco, W. Kluge, S. Müller, G. Venanzoni and B. Valeriani for reminding us constantly of the importance of this work for the experimental analysis and H. Czyż and N.P. Merenkov for very useful discussions. Work supported in part by BMBF under grant number 05HT9VKB0 and E.U. EURODAΦNE network TMR project FMRX-CT98-0169.

Appendix

A The leptonic tensor in the Cartesian basis

For the convenience of the reader we list the mass-suppressed terms $a_{ij}^{(1,m)}$ also in the Cartesian basis. The large-angle contributions have been given in [18]. The component proportional to $g_{\mu\nu}$ reads

$$a_{00}^{(1,m)} = -a_{++}^{(1,m)}, \quad (34)$$

⁴ We thank N.P. Merenkov for drawing our attention to this reference

see (25). The coefficient a_{+-} does not contribute at this order to a_{00} . The remaining components are given by

$$\begin{aligned} a_{11}^{(1,m)} &= \frac{q^2}{1-q^2} \left\{ \frac{4m^2}{y_1^2} \left[1 - \log\left(\frac{y_1}{m^2}\right) \right. \right. \\ &+ \left. \frac{m^2}{y_1} \left(Li_2\left(1 - \frac{y_1}{m^2}\right) - \frac{\pi^2}{6}\right) \right] - n(y_1, 1) \\ &+ \frac{2m^2 q^2}{y_1(m^2(1-q^2) - y_1)} \left[\frac{1}{q^2} \log\left(\frac{y_1}{m^2}\right) + \frac{\log(q^2)}{1-q^2} \right. \\ &+ \left. \left(1 + \frac{m^2}{m^2(1-q^2) - y_1} \right) N(y_1) \right] \Big\} + \\ &+ \frac{1}{1-q^2} \left\{ \frac{4m^2(1-q^2)}{y_2^2} \left[\log(q^2) \log\left(\frac{y_2^4}{m^4 q^2}\right) \right. \right. \\ &+ \left. 4Li_2(1-q^2) + 2 \left(Li_2\left(1 - \frac{y_2}{m^2}\right) - \frac{\pi^2}{6}\right) \right] \\ &+ \frac{4m^2 q^2}{y_2^2} \left[1 - \log\left(\frac{y_2}{m^2}\right) + \left(1 + \frac{m^2}{y_2} \right) \right. \\ &\times \left. \left(Li_2\left(1 - \frac{y_2}{m^2}\right) - \frac{\pi^2}{6}\right) \right] \\ &- \frac{1-2q^4}{q^2} n\left(y_2, \frac{3-8q^2+6q^4}{1-2q^4}\right) \\ &+ \frac{2m^2}{y_2(m^2(1-q^2) - y_2)} \left[\frac{1}{q^2} \log\left(\frac{y_2}{m^2}\right) + \frac{\log(q^2)}{1-q^2} \right. \\ &+ \left. \left(3 + \frac{m^2}{m^2(1-q^2) - y_2} \right) N(y_2) \right] \Big\}, \end{aligned} \quad (35)$$

$$a_{22}^{(1,m)} = a_{11}^{(1,m)}(y_1 \leftrightarrow y_2), \quad (36)$$

$$\begin{aligned} a_{12}^{(1,m)} &= \frac{q^2}{1-q^2} \left\{ \frac{4m^2}{y_1^2} \left[1 - \log\left(\frac{y_1}{m^2}\right) \right. \right. \\ &+ \left. \left(\frac{1}{2} + \frac{m^2}{y_1} \right) \left(Li_2\left(1 - \frac{y_1}{m^2}\right) - \frac{\pi^2}{6}\right) \right] \\ &- \frac{1-q^2}{q^2} n\left(y_1, \frac{1}{1-q^2}\right) + \frac{2m^2}{y_1(m^2(1-q^2) - y_1)} \\ &\times \left[\frac{1}{q^2} \log\left(\frac{y_1}{m^2}\right) + \frac{\log(q^2)}{1-q^2} \right. \\ &+ \left. \left(2 + \frac{m^2}{m^2(1-q^2) - y_1} \right) N(y_1) \right] \Big\} \\ &+ [y_1 \leftrightarrow y_2]. \end{aligned} \quad (37)$$

and

$$a_{-1}^{(1,m)} = 0. \quad (38)$$

B Scalar one-loop integrals

A few two-, three-, and four-point scalar one-loop integrals enter our calculation. Expression for the two-point scalar integrals are simple and well known. The notation from [18], where the corresponding results valid for large photon angles ($m^2 \ll 1, q^2, y_i$) can be found, is used in the

following. The general three-point scalar one-loop integral is defined by

$$C_0(p_a^2, (p_a - p_b)^2, p_b^2, m_1^2, m_2^2, m_3^2) = -i16\pi^2\mu^{4-D} \quad (39)$$

$$\times \int \frac{d^D k}{(2\pi)^D} \frac{1}{[k^2 - m_1^2][(k - p_a)^2 - m_2^2][(k - p_b)^2 - m_3^2]}.$$

Four different three-point scalar one-loop integrals are needed

$$\begin{aligned} C01 &= C_0((p_i - k_1)^2, 0, m_e^2, 0, m_e^2, m_e^2), \\ C02 &= C_0(m_e^2, s, m_e^2, 0, m_e^2, m_e^2), \\ C03 &= C_0((p_i - k_1)^2, Q^2, m_e^2, 0, m_e^2, m_e^2), \\ C04 &= C_0(Q^2, s, 0, m_e^2, m_e^2, m_e^2), \end{aligned} \quad (40)$$

$i = 1, 2$, and one scalar box

$$D0 = -i16\pi^2\mu^{4-D} \int \frac{d^D k}{(2\pi)^D} \left\{ \left(1\right) / \left(k^2[(k + p_i)^2 - m_e^2]\right) \right. \\ \left. \times \left[(k + p_i - k_1)^2 - m_e^2\right] \left[(k - p_j)^2 - m_e^2\right] \right\}, \quad (41)$$

with $j \neq i$.

The following simple expressions are used, from where the limits ($m^2 \ll 1, q^2, y_i$) or ($m^2 \ll 1, q^2$ but $m^2 \sim y_i$) are obtained:

$$\begin{aligned} C01 &= \frac{s^{-1}}{y_i} \left[Li_2\left(1 - \frac{y_i}{m^2}\right) - \frac{\pi^2}{6} \right], \\ C02 &= \frac{s^{-1}}{\beta} \left[\left(\Delta - 2\log(\beta) - \frac{\log(c)}{2} \right) \log(c) \right. \\ &\quad \left. - 2Li_2(c) - \frac{2\pi^2}{3} + i\pi \left(\Delta - 2\log(\beta) \right) \right], \\ C04 &= \frac{s^{-1}}{1 - q^2} \left[\frac{\log(c)}{2} - \frac{\log(c_q)}{2} + i\pi \log\left(\frac{c}{c_q}\right) \right], \\ D0 &= \frac{s^{-2}}{\beta y_i} \left[- \left(\Delta + 2\log\left(\frac{m}{y_i}\right) \right) \log(c_q) + \log^2(c) \right. \\ &\quad \left. + 2Li_2\left(1 - \frac{c_q}{c}\right) + 2Li_2(1 - c c_q) - Li_2(1 - c_q^2) - \pi^2 \right. \\ &\quad \left. - i\pi \left(\Delta + 2\log\left(\frac{m}{c y_i}\right) \right) \right], \end{aligned} \quad (42)$$

with

$$\Delta = \frac{(4\pi)^\varepsilon}{\varepsilon \Gamma(1 - \varepsilon)} \left(\frac{\mu^2}{s} \right)^\varepsilon, \quad (43)$$

and

$$\begin{aligned} \beta &= \sqrt{1 - 4m^2}, & c &= \frac{1 - \beta}{1 + \beta}, \\ \beta_q &= \sqrt{1 - \frac{4m^2}{q^2}}, & c_q &= \frac{1 - \beta_q}{1 + \beta_q}. \end{aligned} \quad (44)$$

Our expression for the C03 function is rather cumbersome:

$$C03 = \frac{s^{-1}}{q^2(z_1 - z_2)} \left[\log\left(\frac{y_i}{q^2}\right) \log\left(\frac{(1 - z_1)z_2}{(1 - z_2)z_1}\right) \right.$$

$$\begin{aligned} &+ \left\{ Li_2\left(\frac{1}{z_1}\right) - \log((z_3 - z_1)(z_4 - z_1)) \log\left(1 - \frac{1}{z_1}\right) \right. \\ &+ Li_2\left(\frac{1 - z_1}{z_3 - z_1}\right) - Li_2\left(\frac{-z_1}{z_3 - z_1}\right) + Li_2\left(\frac{1 - z_1}{z_4 - z_1}\right) \\ &\left. - Li_2\left(\frac{-z_1}{z_4 - z_1}\right) - [z_1 \leftrightarrow z_2] \right\}, \end{aligned} \quad (45)$$

where

$$\begin{aligned} z_{1,2} &= \frac{q^2 + y_i}{2q^2} \left(1 \pm \sqrt{1 - \frac{4(m^2 - i\eta)q^2}{(q^2 + y_i)^2}} \right), \\ z_{3,4} &= \frac{1}{2} \left(1 \pm \sqrt{1 - \frac{4(m^2 - i\eta)}{q^2}} \right), \end{aligned} \quad (46)$$

being

$$Im(C03) = \frac{\pi}{q^2(z_1 - z_2)} \log\left(\frac{(z_3 - z_1)(z_4 - z_2)}{(z_4 - z_1)(z_3 - z_2)}\right), \quad (47)$$

its imaginary part. As for the other scalar one-loop functions, this expression is expanded up to terms of order m^2 and m^4 .

References

1. [LEP Collaborations], hep-ex/0103048
2. J. H. Kühn, M. Steinhauser, Nucl. Phys. B **619**, 588 (2001) [hep-ph/0109084]
3. S. Binner, J. H. Kühn, K. Melnikov, Phys. Lett. B **459**, 279 (1999) [hep-ph/9902399]
4. K. Melnikov, F. Nguyen, B. Valeriani, G. Venanzoni, Phys. Lett. B **477**, 114 (2000) [hep-ph/0001064]
5. H. Czyż, J. H. Kühn, Eur. Phys. J. C **18**, 497 (2001) [hep-ph/0008262]
6. S. Spagnolo, Eur. Phys. J. C **6**, 637 (1999)
7. V. A. Khoze, M. I. Konchatnij, N. P. Merenkov, G. Pancheri, L. Trentadue, O. N. Shekhovtsova, Eur. Phys. J. C **18**, 481 (2001) [hep-ph/0003313]
8. A. Höfer, J. Gluza, F. Jegerlehner, Eur. Phys. J. C **24**, 51 (2002) [hep-ph/0107154]
9. V. A. Khoze, M. I. Konchatnij, N. P. Merenkov, G. Pancheri, L. Trentadue, O. N. Shekhovtsova, hep-ph/0202021
10. A. Aloisio et al. [KLOE Collaboration], hep-ex/0107023
11. A. Denig et al. [KLOE Collaboration], eConf **C010430**, T07 (2001) [hep-ex/0106100]
12. M. Adinolfi et al. [KLOE Collaboration], hep-ex/0006036
13. E. P. Solodov [BABAR collaboration], eConf **C010430**, T03 (2001) [hep-ex/0107027]
14. M. Caffo, H. Czyż, E. Remiddi, Nuovo Cim. A **110**, 515 (1997) [hep-ph/9704443]; Phys. Lett. B **327**, 369 (1994)
15. G. Rodrigo, H. Czyż, J. H. Kühn, M. Szopa, Eur. Phys. J. C **24**, 71 (2002) [hep-ph/0112184]
16. G. 't Hooft, M. J. Veltman, Nucl. Phys. B **153**, 365 (1979)
17. W. Beenakker, A. Denner, Nucl. Phys. B **338**, 349 (1990)
18. G. Rodrigo, A. Gehrmann-De Ridder, M. Guilleaume, J. H. Kühn, Eur. Phys. J. C **22**, 81 (2001) [hep-ph/0106132]

19. G. Rodrigo, *Acta Phys. Polon. B* **32**, 3833 (2001) [hep-ph/0111151]
20. F. A. Berends, G. J. Burgers, W. L. van Neerven, *Phys. Lett. B* **177**, 191 (1986)
21. F. A. Berends, W. L. van Neerven, G. J. Burgers, *Nucl. Phys. B* **297**, 429 (1988); Erratum *Nucl. Phys. B* **304**, 921 (1988)
22. E. A. Kuraev, N. P. Merenkov, V. S. Fadin, *Yad. Fiz.* **45**, 782 (1987)
23. E. A. Kuraev, N. P. Merenkov, V. S. Fadin, *Sov. J. Nucl. Phys.* **45**, 486 (1987)
24. J. H. Kühn, A. Santamaria, *Z. Phys. C* **48**, 445 (1990)
25. R. Decker, M. Finkemeier, P. Heiliger, H. H. Jonsson, *Z. Phys. C* **70**, 247 (1996) [hep-ph/9410260]
26. G. Ecker, R. Unterdorfer, hep-ph/0203075
27. G. Rodrigo, H. Czyż, J. H. Kühn, in preparation
28. R. Mertig, M. Böhm, A. Denner, *Comput. Phys. Commun.* **64**, 345 (1991)
29. G. Passarino, M. Veltman, *Nucl. Phys. B* **160**, 151 (1979)