

Theoretical aspects of Higgs physics at the LHC

Mauro Moretti^a

Department of Physics, Ferrara University and INFN Ferrara

Received: 31 October 2003 / Accepted: 14 November 2003 /
Published Online: 3 December 2003 – © Springer-Verlag / Società Italiana di Fisica 2003

Abstract. The strategies recently developed to study Higgs boson properties at the LHC are reviewed. It is shown how to obtain model-independent determinations of couplings to fermions and gauge bosons by exploiting different production and decay channels. We consider the case of Weak Boson Fusion Higgs production with $H \rightarrow b\bar{b}$

PACS. 14.80.Bn Standard-Model Higgs boson – 13.85.Hd Inelastic Scattering: many particle final state

1 Introduction

The LHC will allow the discovery of the Higgs boson and the study of its mass, width and couplings to fermions and gauge bosons. While the decay channels $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ^{(*)} \rightarrow 4l$ will allow a direct mass measurement at the 0.1% level over a wide range of masses [1], the total width can only be determined with about 10% accuracy by direct measurement of the decay $H \rightarrow ZZ^{(*)} \rightarrow 4l$ for $m_H > 200$ GeV, (the Higgs width for lower Higgs masses being too small with respect to the detector resolution). An indirect measurement of the total width can be performed also in the low mass region by exploiting the available production and decay mechanisms at the LHC. Several studies have been performed to improve on the strategy originally proposed in [2] for the determination of the Higgs boson properties. We will briefly review the progress recently made in this field. The main focus will be on the mass window 115–200 GeV, the one preferred by electroweak precision data and by supersymmetry.

2 Theoretical calculations

In order to disentangle a signal from backgrounds, a good understanding of uncertainties in theoretical predictions is necessary. The QCD corrections, at least at next-to-leading order (NLO), are known for all production channels, the most recent calculations being the NNLO calculation¹ for the gluon fusion process in the limit of heavy top-quark mass [3, 4] and the NLO corrections for the process $pp/p\bar{p} \rightarrow t\bar{t}H + X$ [5]. At present, the uncertainties arising from QCD uncertainties (combining the residual scale dependence with the error from parton distribution

functions) can be estimated to be $\pm 20\%$ for gluon fusion, $\pm 5\%$ for WBF, $\pm 10\%$ for associated production.

Concerning the backgrounds, several processes with low final state parton multiplicity (corresponding to important irreducible backgrounds) are available at NLO, namely $q\bar{q} \rightarrow \gamma\gamma$ [6], $gg \rightarrow \gamma\gamma$ [7], $pp(\bar{p}) \rightarrow Wb\bar{b}$, $pp(\bar{p}) \rightarrow Zb\bar{b}$ [8], $pp(\bar{p}) \rightarrow Wjj$, $pp(\bar{p}) \rightarrow Zjj$ [9], $pp(\bar{p}) \rightarrow VV$ [10] and QCD $H + jj$ production via gluon fusion [11]. Some of these calculations are already implemented in NLO Monte Carlo programs. In the case of multiparton final states, the methods developed for NLO calculations cannot be applied, because of the complexity of the calculations for processes with many external legs. Recently some effort has been devoted to the realization of LO Monte Carlo event generators based on exact matrix element calculations, interfaced to the shower evolution Monte Carlo programs producing the real final state hadrons [12, 13, 14, 15, 16].

3 Higgs couplings to fermions and gauge bosons

In principle, the Higgs coupling to a given fermion family f , could be obtained from the following relation:

$$R(H \rightarrow f\bar{f}) = \int Ldt \cdot \sigma(pp \rightarrow H) \cdot \frac{\Gamma_f}{\Gamma},$$

where $R(H \rightarrow f\bar{f})$ is the Higgs production rate in a given final state, $\int Ldt$ is the integrated luminosity, $\sigma(pp \rightarrow H)$ is the Higgs production cross section, while Γ and Γ_f are the total and partial Higgs widths respectively. Aiming at model-independent coupling determinations, one needs to consider ratios of couplings, which are experimentally accessible through the measurements of ratios of rates for different final states, because the total Higgs cross-section and width cancel in the ratios (as well as the luminosity and all the QCD uncertainties related to the initial state).

^a I acknowledge the financial support of the European Union under the contract HPRN-CT-2000-00149

¹ The K factor for the NLO contribution is 2 making desirable the knowledge of NNLO contribution




Fig. 1. Relative accuracy (%) on the individual rates Γ_i expected at the LHC (from [28]). See the text for a detailed description of the panels

Up to now detailed studies on signal and backgrounds for several channels have been performed, namely $gg \rightarrow H, (H \rightarrow \gamma\gamma, ZZ, WW)$ [1, 17, 18, 19], $qq \rightarrow qqH, (H \rightarrow \gamma\gamma, \tau\tau, WW)$ [20, 21, 22, 23, 24], $pp \rightarrow t\bar{t}H, (H \rightarrow b\bar{b}, WW, \tau\tau)$ [25, 26, 27, 28] and $pp \rightarrow WH, H \rightarrow b\bar{b}$ [29]. Each process depends on two² Higgs couplings, one from the Higgs boson production and one from the Higgs boson decay.

Therefore, every production and decay channel provides a measurement of the ratio $Z_j^{(i)} = \Gamma_i \Gamma_j / \Gamma$, where $i = g, W, t$ indicates the particles involved in the production process while the index $j = b, \tau, W, Z, g, \gamma$ refers to the decay process. In case of $m_H < 140$ GeV, the above mentioned channels allow to express the individual rates $\Gamma_t, \Gamma_b, \Gamma_\tau, \Gamma_W, \Gamma_g$ and Γ_γ as functions of the observables $Z_j^{(i)}$ and of the total Higgs width Γ [28]. With the assumption that the total width is saturated by the known channels $\Gamma = \Gamma_b + \Gamma_\tau + \Gamma_W + \Gamma_Z + \Gamma_g + \Gamma_\gamma$ (otherwise new processes would be observed independently of any precision study), an expression for Γ can be obtained in terms of the measured quantities $Z_j^{(i)}$ [28]. Figure 1 [28] summarizes the relative accuracy on the individual rates Γ_i expected in the model-independent scenario as well as in a scenario with Γ_b/Γ_τ fixed to its SM value, assuming a total integrated luminosity of 200 fb^{-1} . The upper plots show the accuracies obtained without including any theoretical error, while the lower plots show the same accuracies including a systematic theoretical error of 20% for the $gg \rightarrow H$ channel, of 5% for the $qq \rightarrow qqH$, and of 10% for the $pp \rightarrow t\bar{t}H$. As can be seen, the total Higgs width can be indirectly determined in the low mass region with a precision of the order of 30% in a model-

² This is not true in the case of the weak boson fusion channel where one needs to assume that the ratio of HWW and HZZ is the same as in the SM

independent way while the Higgs couplings can be determined with accuracies between 7% and 25%. In the case of $140 < m_H < 200$ GeV, the gluon fusion, weak boson fusion and $t\bar{t}H$ associated production processes, with the Higgs boson decaying only to gauge bosons, allow an indirect determination of Γ_W and Γ with a precision of the order of 10% [2, 30]. In this Higgs mass range, however, there is no handle to study the Higgs Yukawa couplings to b quarks and τ leptons. The assumption $\Gamma_Z/\Gamma_W = z_{SM}$ can be tested at the 20–30% level, for $m_H > 130$ GeV, by measuring the ratio $Z_Z^{(g)}/Z_W^{(g)}$ [30], and it can even be tested with the same level of accuracy for lower Higgs boson masses by comparing the two ratios $Z_b^{(WH)}/Z_b^{(t)}$ and $Z_\tau^{(W)}/Z_\tau^{(t)}$ [28]. For $m_H > 140$ GeV, with luminosities of the order of 300 fb^{-1} , the ratio Γ_t/Γ_g can be tested in a model-independent measuring $Z_W^{(t)}/Z_W^{(g)}$ [27].

4 $H \rightarrow b\bar{b}$ via weak boson fusion

To improve the analysis of the $Hb\bar{b}$ Yukawa coupling, one can consider the decay of an Higgs, produced via Weak Boson Fusion, into $b\bar{b}$ pairs [31].

Signal and background event estimates are based on a leading order partonic calculation of the matrix elements (ME) obtained with the event generator ALPGEN [12]. The background sources considered include:

1. QCD production of $b\bar{b}jj$ final states, where j indicates a jet originating from a light quark (u, d, s, c) or a gluon;
2. QCD production of $jjjj$ final states;
3. associated production of $Z^*/\gamma^* \rightarrow b\bar{b}$ and light jets

along with multiple interaction events ($pp \oplus pp, pp \oplus pp \oplus pp \dots$) giving rise to final states of the kind $b\bar{b}jj$ and $jjjj$. In order to satisfy the requirements of optimization of the signal significance (S/\sqrt{B}) and compatibility with trigger and data acquisition constraints, different selection criteria have been considered. The sensitivity can be as large as 5 for Higgs masses close to the exclusion limit given by LEP searches but the ratio S/B is only a fraction of a percent. This implies that the background will have to be known with accuracies at the permille level. The background should therefore be determined entirely from data. The large rate of $b\bar{b}jj$ from single and multiple interactions and the smoothness of their mass distribution will allow to estimate their size with enough statistical accuracy, without significant systematic uncertainties.

The situation is different in the case of the backgrounds from the tails of the Z decays. The Z mass peak is sufficiently close to m_H to possibly distort the m_{bb} spectrum and spoil the ability to accurately reconstruct the noise level from data. These background rates are at most comparable to the signal at low m_H . A 10% determination of these final states, which should be easily achievable using the $(Z \rightarrow \ell^+\ell^-)jj$ control sample and folding in the detector energy resolution for jets, should therefore be sufficient to fix these background with the required accuracy.

Table 1. The statistical significance of the determination of the branching ratio Γ_b/Γ and of the b -quark Yukawa coupling in the configurations **a** and **b** (see [31] for a description of the event selections), for an integrated luminosity of 600 fb^{-1} . The p_T^j cut on jets is $p_T^j > 60 \text{ GeV}$. The case of $p_T^j > 80 \text{ GeV}$ [31] doesn't affect sizeably the results. The probability to mistag a light-jet as b -jet is assumed to be 0.01

	m_H (GeV)	115	120	140
(a)	$\delta\Gamma_b/\Gamma$	0.33	0.35	0.71
	$\delta y_{Hbb}/y_{Hbb}$	0.58	0.51	0.56
(b)	$\delta\Gamma_b/\Gamma$	0.20	0.19	0.37
	$\delta y_{Hbb}/y_{Hbb}$	0.36	0.30	0.29

Concerning the multiple interactions, in the simplest case of two overlapping events ($pp \oplus pp$), there are four possible (including mistagging effects) combinations of events leading to a $b\bar{b}jj$ background. The largest contribution arises from $(jj\bar{b}) \oplus (j\bar{b}j)$ events, where the $b\bar{b}$ mass spectrum has a broad peak in the middle of the signal region. The absolute rate of these events can be determined if the distribution of the beam-line z vertex separation between the two overlapping events can be determined with a resolution of of 5-10 mm. The number of these events is significantly lowered using the higher threshold of 80 GeV for the forward jets.

Table 1 summarizes the accuracy reachable in the $\mathcal{B}(H \rightarrow b\bar{b})$ and in the $Hb\bar{b}$ Yukawa coupling for the case of two different event selections (described in detail in [31]), assuming that the coupling HWW is the one predicted by the Standard Model or determined in other reactions studied in the literature.

An integrated luminosity of 600 fb^{-1} is considered. The $H \rightarrow b\bar{b}$ decay in the WBF channel could be used together with other processes already examined in the literature for a model independent determination of the ratio of Yukawa couplings $y_{Hbb}/y_{H\tau\tau}$ [32].

As a conclusion of the analysis presented in [31], the $H \rightarrow b\bar{b}$ channel produced in association with two jets is suggested as an additional channel to be exploited for the measurements of the Higgs couplings to fermions.

5 Summary

During the last few years there has been a dramatic improvement in both theoretical and experimental studies of several Higgs boson production and decay channels at the LHC. A strategy has been designed to study, in a model-independent way, the Higgs couplings to fermions and bosons, which allows also, with little theoretical assumption, an indirect determination of the total Higgs width. The main results of a very recent analysis of the $H \rightarrow b\bar{b}$ channel in Weak Boson Fusion production have been reviewed, pointing out its importance for the determination of the $Hb\bar{b}$ Yukawa coupling.

Acknowledgements. I wish to acknowledge M.L. Mangano, F. Piccinini, R. Pittau and A. Polosa for fruitful collabora-

tion. I wish to thank the organizers for the kind invitation and for the pleasant atmosphere during the Workshop.

References

1. ATLAS Collaboration: Tech. Rep. CERN/LHCC/99-15, CERN, 1999
2. R. Kinnunen et al.: Phys. Rev. D **62** 013009 (2000)
3. M. Grazzini: hep-ph/0209302; S. Catani, D. de Florian, and M. Grazzini: JHEP 0105 025 (2001); and JHEP 0201 015 (2002); G. Bozzi, S. Catani, D. de Florian, and M. Grazzini: hep-ph/0302104; S. Catani, D. de Florian, M. Grazzini, and P.Nason: hep-ph/0306211
4. W.B. Kilgore: these proceedings; R.V. Harlander and W.B. Kilgore: Phys. Rev. D **64**, 025 (2001); Phys. Rev. Lett. **88** 201801 (2002); C. Anastasiou and K. Melnikov: Nucl. Phys. **B** 646 220 (2002); V. Ravindran, J. Smith, and W.L. Van Neerven: hep-ph/0302135
5. W. Beenakker et al.: Phys. Rev. Lett. **87**, 201805 (2001); S. Dawson and L. Reina: Phys. Rev. Lett. **87** 201804 (2001); S. Dawson, L. Reina, and D. Wackerth: Phys. Rev. D **65**, 053017 (2002); S. Dawson, C. Jackson, L.H. Orr, L. Reina, and D. Wackerth: hep-ph/0305087
6. T. Binoth et al.: Eur. Phys. J. C **16**, 311 (2000); hep-ph/0203064
7. Z. Bern, L. Dixon, and C. Schmidt: hep-ph/0206194
8. R.K. Ellis and S. Veseli: Phys. Rev. D **60**, 011501 (1999); J. Campbell: hep-ph/0105226
9. J. Campbell and R.K. Ellis: hep-ph/0202176
10. S. Frixione and B.R. Webber: hep-ph/0204244
11. V. Del Duca et al.: Nucl. Phys. B **616**, 367 (2001)
12. M.L. Mangano et al.: hep-ph/0206293
13. A. Pukhov et al.: hep-ph/9908288
14. T. Ishikawa et al., MINAMI-TATEYA Group Coll.: KEK-92-19;
15. T. Stelzer and W.F. Long: Comput. Phys. Commun. **81**, 357 (1994); F. Maltoni and T. Stelzer: hep-ph/0208156
16. B.P. Kersevan and E. Richter-Was: hep-ph/0201302
17. CMS Collaboration: Tech. Rep. CERN/LHCC/94-38, CERN, 1994
18. M. Dittmar and H.K. Dreiner: Phys. Rev. D **55**, 167 (1997); hep-ph/9703401
19. D. Denegri et al.: hep-ph/0112045
20. D. Rainwater and D. Zeppenfeld: JHEP 12 005 (1997)
21. D. Rainwater, D. Zeppenfeld, and K. Hagiwara: Phys. Rev. D **59**, 014037 (1999)
22. T. Plehn, D. Rainwater, and D. Zeppenfeld: Phys. Rev. D **61**, (2000)
23. D. Rainwater and D. Zeppenfeld: Phys. Rev. D **60**, 113004 (1999)
24. N. Kauer et al.: Phys. Lett. B **503**, 113 (2001)
25. E. Richter-Was and M. Sapinski: Acta Phys. Polon. B **30**, 1001 (1999)
26. V. Drollinger, T. Müller, and D. Denegri: hep-ph/0111312
27. F. Maltoni, D. Rainwater, and S. Willenbrock: hep-ph/0202205
28. A. Belyaev and L. Reina: hep-ph/0205270
29. V. Drollinger, T. Müller, and D. Denegri: hep-ph/0201249
30. D. Zeppenfeld: hep-ph/0203123
31. M.L. Mangano et al.: hep-ph/0210261
32. D. Rainwater, D. Zeppenfeld, and K. Hagiwara: Phys. Rev. D **59**, 014037 (1999)