

Complete description of polarization effects in emission of a photon by an electron in the field of a strong laser wave

D.Yu. Ivanov^{1,a}, G.L. Kotkin^{2,b}, V.G. Serbo^{2,c}

¹ Sobolev Institute of Mathematics, Novosibirsk, 6300090, Russia

² Novosibirsk State University, Novosibirsk, 630090 Russia

Received: 13 February 2004 /

Published online: 23 June 2004 – © Springer-Verlag / Società Italiana di Fisica 2004

Abstract. We consider the emission of a photon by an electron in the field of a strong laser wave. Polarization effects in this process are important for a number of physical problems. The probability of this process for circularly or linearly polarized laser photons and for arbitrary polarization of all other particles is calculated. We obtain the complete set of functions which describe such a probability in a compact invariant form. Besides this, we discuss in some detail the polarization effects in the kinematics relevant to the problem of $e \rightarrow \gamma$ conversion at $\gamma\gamma$ and γe colliders.

1 Introduction

Compton scattering,

$$e(p) + \gamma(k) \rightarrow e(p') + \gamma(k'), \quad (1)$$

was one of the first processes calculated in quantum electrodynamics at the end of 1920's. The analysis of polarization effects in this reaction is now included in text books (see, for example, [1], §87). Nevertheless, the complete description of cross sections with polarization of both initial and final particles has been considered in detail only recently (see [2–4] and literature therein). One interesting application is the collision of an ultra-relativistic electron with a beam of polarized laser photons. In this case the Compton effect is the basic process for obtaining high-energy photons for contemporary experiments in nuclear physics (photo-nuclear reactions with polarized photons) and for future $\gamma\gamma$ and γe colliders [5]. The importance of the particle polarization is clearly seen from the fact that in comparison with the unpolarized case the number of final photons with maximum energy is nearly doubled when the helicities of the initial electron and photon are opposite [3].

With the growth of the laser field intensity, an electron starts to interact coherently with n laser photons,

$$e(q) + n \gamma_L(k) \rightarrow e(q') + \gamma(k'); \quad (2)$$

thus the Compton scattering becomes non-linear. Such a process with absorption of $n = 1, 2, 3, 4$ linearly polarized laser photons was observed in a recent experiment at

SLAC [6]. The polarization properties of the process (2) are important for a number of problems, for example, for laser beam cooling [7] and, especially, for future $\gamma\gamma$ and γe colliders (see [8,9] and the literature therein). In the latter case the non-linear Compton scattering must be taken into account in simulations of the processes in the conversion region. For comprehensive simulation, including processes of multiple electron scattering, one has to know not only the differential cross section of the non-linear Compton scattering with a given number of the absorbed laser photons n , but energy, angles and polarization of final photons and electrons as well. The method of calculation for such cross sections was developed by Nikishov and Ritus [10]. It is based on the exact solution of the Dirac equation in the field of the external electromagnetic plane wave. Some particular polarization properties of this process were considered in [10–15] and have already been included in the existing simulation codes [14,16]. Another approach was used in [17], where the total cross section and the spectrum for the non-linear Compton scattering, summed over numbers of the absorbed laser photons n and over spin states of the final particles, has been obtained.

In the present paper we give a complete description of the non-linear Compton scattering for the case of circularly or linearly polarized laser photons and arbitrary polarization of all other particles¹. We follow the method of Nikishov and Ritus in the form presented in [1], §101. In the next section we describe in detail the kinematics of the process (2). The effective differential cross section is obtained in Sect. 3 in the compact invariant form, including the polarization of all particles. In Sect. 4 we consider

^a e-mail: d-ivanov@math.nsc.ru

^b e-mail: kotkin@math.nsc.ru

^c e-mail: serbo@math.nsc.ru

¹ Preliminary results of this work have been submitted to the hep-ph archive [18]

process (2) in the reference frame relevant for $\gamma\gamma$ and γe colliders. The limiting cases are discussed in Sect. 5. In Sect. 6 we present some numerical results obtained for the range of parameters close to those in the existing projects of the $\gamma\gamma$ and γe colliders. In the last section we summarize our results and compare them with those known in the literature. In Appendix A we show that our results coincide in the limit of weak laser field with those for the linear Compton scattering.

2 Kinematics

2.1 Parameter of non-linearity

Let us consider the interaction of an electron with a monochromatic plane wave described by the 4-potential $A_\mu(x)$. The corresponding electric and magnetic fields are \mathbf{E} and \mathbf{B} , the frequency is ω , and let F be the root-mean-squared field strength,

$$F^2 = \langle \mathbf{B}^2 \rangle = \langle \mathbf{E}^2 \rangle.$$

The invariant parameter describing the intensity of the laser field (the parameter of non-linearity) is defined via the mean value of the squared 4-potential:

$$\xi = \frac{e}{mc^2} \sqrt{-\langle A_\mu(x) A^\mu(x) \rangle}, \quad (3)$$

where e and m are the electron charge and the mass, and c is the velocity of light. We use this definition of ξ both for the circularly and linearly polarized laser photons².

The origin of this parameter can be explained as follows. The electron oscillates in the transverse direction under the influence of the force $\sim eF$ and for the time $\sim 1/\omega$ acquires the transverse momentum $\sim p_\perp = eF/\omega$; thus for the longitudinal motion the effective electron mass is $m_* = \sqrt{m^2 + (p_\perp/c)^2}$. The ratio of the momentum p_\perp to mc is the natural dimensionless parameter,

$$\xi = \frac{eF}{mc\omega}. \quad (4)$$

This parameter can be expressed also via the density n_L of photons in the laser wave:

$$\xi^2 = \left(\frac{eF}{m\omega c} \right)^2 = \frac{4\pi\alpha\hbar^2}{m^2 c \omega} n_L, \quad (5)$$

where \hbar is the Plank constant and $\alpha = e^2/(\hbar c) \approx 1/137$.

2.2 Invariant variables

From the classical point of view, the oscillated electron emits harmonics with frequencies $n\omega$, where $n = 1, 2, \dots$

² Note that our definition of this parameter for the case of the linear polarization differs from the one used in [10] by a factor $1/\sqrt{2}$

Their intensities at small ξ^2 are proportional to $(\mathbf{E}^2)^n \propto \xi^{2n}$; the polarization properties of these harmonics depend on the polarizations of the laser wave and the initial electron³. From the quantum point of view, this radiation can be described as the non-linear Compton scattering with absorption of n laser photons. When describing such a scattering, one has to take into account that in a laser wave the 4-momenta p and p' of the free initial and final electrons are replaced by the 4-quasi-momenta q and q' (similar to the description of a particle motion in a periodic potential field in non-relativistic quantum mechanics),

$$q = p + \xi^2 \frac{m^2 c^2}{2pk} k, \quad q' = p' + \xi^2 \frac{m^2 c^2}{2p'k} k, \quad (6)$$

$$q^2 = (q')^2 = (1 + \xi^2) m^2 c^2 \equiv m_*^2 c^2.$$

In particular, the energy of the free incident electron E is replaced by the quasi-energy

$$cq_0 = E + \xi^2 \frac{m^2 c^2}{2pk} \hbar\omega. \quad (7)$$

As a result, we deal with the reaction (2) for which the conservation law reads

$$q + nk = q' + k'. \quad (8)$$

From this it follows that all the kinematic relations which occur for the linear Compton scattering will apply to the process considered here if the electron momenta p and p' are replaced by the quasi-momenta q and q' and the incident photon momentum k by the 4-vector nk . Since $qk = pk$, it is convenient to use the same invariant variables as for the linear Compton scattering (compare [3]):

$$x = \frac{2pk}{m^2 c^2}, \quad y = \frac{kk'}{pk}. \quad (9)$$

Moreover, many kinematic relations can be obtained from that for the linear Compton scattering by the replacement $\omega \rightarrow n\omega$, $m \rightarrow m_*$. In particular, we introduce the auxiliary combinations

$$s_n = 2\sqrt{r_n(1-r_n)}, \quad c_n = 1 - 2r_n, \quad (10)$$

where

$$r_n = \frac{y(1+\xi^2)}{(1-y)nx}. \quad (11)$$

It is useful to note that these invariants have a simple notation, namely

$$s_n = \sin \tilde{\theta}, \quad c_n = \cos \tilde{\theta}, \quad r_n = \sin^2(\tilde{\theta}/2), \quad (12)$$

³ The polarization of the first harmonic is related to the tensor of the second rank $\langle E_i E_j^* \rangle$; in this case one needs only three Stokes parameters. The polarization properties of higher harmonics are connected with the tensors of higher rank $\langle E_i E_j \dots E_k^* \rangle$, and in this case one needs more parameters for their description. It is one of the reasons why the non-linear Compton effect has been considered so far only for 100% polarized laser beam, mainly for circular or linear polarization.

where $\tilde{\theta}$ is the photon scattering angle in the frame of reference where the initial electron is at rest on average ($\mathbf{q} = 0$, $q_0 = m_*c$). Therefore,

$$0 \leq s_n, r_n \leq 1; \quad -1 \leq c_n \leq 1. \quad (13)$$

The maximum value of the variable y for the reaction (2) is

$$y \leq y_n = \frac{nx}{nx + 1 + \xi^2}. \quad (14)$$

The value of y_n is close to 1 for large n , but for a given n it decreases with the growth of the non-linearity parameter ξ^2 . With this notation one can rewrite s_n in the form

$$s_n = \frac{2}{y_n(1-y)} \sqrt{y(y_n - y)(1 - y_n)}, \quad (15)$$

from which it follows that

$$s_n \rightarrow 0 \quad \text{at} \quad y \rightarrow y_n \quad \text{or} \quad y \rightarrow 0. \quad (16)$$

The usual notion of the cross section is not applicable for the reaction (2) and usually its description is given in terms of the probability of the process per second $\dot{W}^{(n)}$. However, for the procedure of simulation in the conversion region as well as for the simple comparison with the linear case, it is useful to introduce the ‘‘effective cross section’’ given by the definition

$$d\sigma^{(n)} = \frac{d\dot{W}^{(n)}}{j_n}, \quad (17)$$

where

$$j_n = \frac{(qnk)c^2}{q_0 n \hbar \omega} n_L = \frac{m^2 c^4 x}{2q_0 \hbar \omega} n_L$$

is the flux density of colliding particles. Contrary to the usual cross section, this effective cross section does depend on the laser beam intensity, i.e. on the parameter ξ^2 . The total effective cross section is defined as the sum over harmonics, corresponding to the reaction (2) with a given number n of absorbed laser photons⁴:

$$d\sigma = \sum_n d\sigma^{(n)}. \quad (18)$$

2.3 Invariant polarization parameters

The invariant description of the polarization properties of both the initial and the final photons can be performed in the standard way (see [1], §87). We define a pair of unit 4-vectors

$$e^{(1)} = \frac{N}{\sqrt{-N^2}}, \quad e^{(2)} = \frac{P}{\sqrt{-P^2}}, \quad (19)$$

⁴ In this formula and below the sum is over those n which satisfy the condition $y < y_n$, i.e. this sum runs from some minimal value n_{\min} up to $n = \infty$, where n_{\min} is determined by the equation $y_{n_{\min}-1} < y < y_{n_{\min}}$.

where⁵

$$\begin{aligned} N^\mu &= \varepsilon^{\mu\alpha\beta\gamma} P_\alpha (k' - nk)_\beta K_\gamma, \\ P_\alpha &= (q + q')_\alpha - \frac{(q + q')K}{K^2} K_\alpha, \quad K_\alpha = nk_\alpha + k'_\alpha, \\ \sqrt{-N^2} &= m^3 xy \frac{n s_n}{r_n} \sqrt{1 + \xi^2}, \quad \sqrt{-P^2} = m \frac{s_n}{r_n} \sqrt{1 + \xi^2}. \end{aligned}$$

The 4-vectors $e^{(1)}$ and $e^{(2)}$ are orthogonal to each other and to the 4-vectors k and k' ,

$$e^{(i)} e^{(j)} = -\delta_{ij}, \quad e^{(i)} k = e^{(i)} k' = 0; \quad i, j = 1, 2.$$

Therefore, they are fixed with respect to the scattering plane of the process.

Let ξ_j be the Stokes parameters for the initial photon which are defined with respect to the 4-vectors $e^{(1)}$ and $e^{(2)}$. As for the polarization of the final photon, it is necessary to distinguish the polarization $\xi_j^{(f)}$ of the final photon as resulting from the scattering process itself from the detected polarization ξ'_j which enters the effective cross section and which essentially represents the properties of the detector as selecting one or other polarization of the final photon (for details see [1], §65). Both these Stokes parameters, $\xi_j^{(f)}$ and ξ'_j , are also defined with respect to the 4-vectors $e^{(1)}$ and $e^{(2)}$.

Let ζ be the polarization vector of the initial electrons. As with the final photon, it is necessary to distinguish the polarization $\zeta^{(f)}$ of the final electron as such from the polarization ζ' that is selected by the detector. The vectors ζ and ζ' enter the effective cross section. They also determine the electron-spin 4-vectors

$$\begin{aligned} a &= \left(\frac{\zeta \mathbf{p}}{m}, \zeta + \mathbf{p} \frac{\zeta \mathbf{p}}{m(E + m)} \right), \\ a' &= \left(\frac{\zeta' \mathbf{p}'}{m}, \zeta' + \mathbf{p}' \frac{\zeta' \mathbf{p}'}{m(E' + m)} \right) \end{aligned} \quad (20)$$

and the mean helicity of the initial and final electrons

$$\lambda_e = \frac{\zeta \mathbf{p}}{2|\mathbf{p}|}, \quad \lambda'_e = \frac{\zeta' \mathbf{p}'}{2|\mathbf{p}'|}. \quad (21)$$

Now we have to define invariants which describe the polarization properties of the initial and the final electrons. For the electrons it is a more complicated task than it is for the photons. For the linear Compton scattering, the relatively simple description was obtained in [2] using invariants which have a simple meaning in the center-of-mass system. However, this frame of reference is not convenient for the description of the non-linear Compton scattering, since it has actually to vary with the change of the number of the absorbed laser photons n .

Since $ap = a'p' = 0$, we can decompose the electron polarization 4-vectors over three convenient unit 4-vectors e_j and e'_j , $j = 1, 2, 3$, the projections on which determine the

⁵ Below we use the system of units in which $c = 1$, $\hbar = 1$.

polarization properties of the electrons. Our choice is based on the experience obtained in [4, 15]. We exploit two ideas. First of all, the 4-vector $e^{(1)}$ is orthogonal to the 4-vectors k , k' , p and p' ; therefore, the invariants $\zeta_1 = -ae^{(1)}$ and $\zeta'_1 = -a'e^{(1)}$ are the transverse polarizations of the initial and the final electrons perpendicular to the scattering plane. Further, it is not difficult to check that the invariant $ak/(2mx)$ is the mean helicity of the initial electron in the frame of reference, in which the electron momentum \mathbf{p} is anti-parallel to the initial photon momentum \mathbf{k} . Analogously, the invariant $a'k/[2mx(1-y)]$ is the mean helicity of the final electron in the frame of reference, in which the momentum of the final electron \mathbf{p}' is anti-parallel to the initial photon momentum \mathbf{k} . It is important that this interpretation is valid for any number of absorbed laser photons. Furthermore, we will show that for the practically important case, relevant for the $e \rightarrow \gamma$ conversion, these frames of reference almost coincide.

As a result, we define the two sets of unit 4-vectors

$$\begin{aligned} e_1 &= e^{(1)}, & e_2 &= -e^{(2)} - \frac{\sqrt{-P^2}}{m^2x}k, \\ e_3 &= \frac{1}{m} \left(p - \frac{2}{x}k \right); \\ e'_1 &= e^{(1)}, & e'_2 &= -e^{(2)} - \frac{\sqrt{-P^2}}{m^2x(1-y)}k, \\ e'_3 &= \frac{1}{m} \left(p' - \frac{2}{x(1-y)}k \right). \end{aligned} \quad (22)$$

These vectors satisfy the conditions

$$e_i e_j = -\delta_{ij}, \quad e_j p = 0; \quad e'_i e'_j = -\delta_{ij}, \quad e'_j p' = 0. \quad (23)$$

It allows us to represent the 4-vectors a and a' in the following covariant form:

$$a = \sum_{j=1}^3 \zeta_j e_j, \quad a' = \sum_{j=1}^3 \zeta'_j e'_j, \quad (24)$$

where

$$\zeta_j = -ae_j, \quad \zeta'_j = -a'e'_j. \quad (25)$$

The invariants ζ_j and ζ'_j describe completely the polarization properties of the initial electron and the detected polarization properties of the final electron, respectively. To clarify the meaning of the invariants ζ_j , it is useful to note that

$$\zeta_j = \boldsymbol{\zeta} \mathbf{n}_j, \quad (26)$$

where the corresponding 3-vectors are

$$\mathbf{n}_j = \mathbf{e}_j - \frac{\mathbf{p}}{E+m} e_{j0} \quad (27)$$

with e_{j0} being a time component of the 4-vector e_j defined in (22). Using the properties (23) of the 4-vectors e_j , one can check that

$$\mathbf{n}_i \mathbf{n}_j = \delta_{ij}. \quad (28)$$

As a result, the polarization vector $\boldsymbol{\zeta}$ has the form

$$\boldsymbol{\zeta} = \sum_{j=1}^3 \zeta_j \mathbf{n}_j. \quad (29)$$

Analogously, for the final electron the invariants ζ'_j can be represented as

$$\zeta'_j = \boldsymbol{\zeta}' \mathbf{n}'_j, \quad \mathbf{n}'_j = \mathbf{e}'_j - \frac{\mathbf{p}'}{E'+m} e'_{j0}, \quad \mathbf{n}'_i \mathbf{n}'_j = \delta_{ij}, \quad (30)$$

such that

$$\boldsymbol{\zeta}' = \sum_{j=1}^3 \zeta'_j \mathbf{n}'_j. \quad (31)$$

In what follows, we will often consider non-linear Compton scattering in the frame of reference in which an electron performs a head-on collision with laser photons, i.e. in which $\mathbf{p} \parallel (-\mathbf{k})$. We call this the ‘‘collider system’’. In this frame of reference we choose the z -axis along the initial electron momentum \mathbf{p} . Azimuthal angles φ , β and β' of vectors \mathbf{k}' , $\boldsymbol{\zeta}$ and $\boldsymbol{\zeta}'$ are defined with respect to one fixed x -axis.

3 Cross section in the invariant form

3.1 General relations

The effective differential cross section can be represented in the following invariant form:

$$d\sigma(\boldsymbol{\zeta}, \boldsymbol{\xi}, \boldsymbol{\zeta}', \boldsymbol{\xi}') = \frac{r_e^2}{4x} \sum_n F^{(n)} d\Gamma_n, \quad (32)$$

$$d\Gamma_n = \delta(q + nk - q' - k') \frac{d^3k'}{\omega'} \frac{d^3q'}{q'_0},$$

where $r_e = \alpha/m$ is the classical electron radius, and

$$\begin{aligned} F^{(n)} &= F_0^{(n)} + \sum_{j=1}^3 \left(F_j^{(n)} \xi'_j + G_j^{(n)} \zeta'_j \right) \\ &+ \sum_{i,j=1}^3 H_{ij}^{(n)} \zeta'_i \xi'_j. \end{aligned} \quad (33)$$

Here the function $F_0^{(n)}$ describes the total cross section for a given harmonic n , summed over spin states of the final particles:

$$\sigma^{(n)}(\boldsymbol{\zeta}, \boldsymbol{\xi}) = \frac{r_e^2}{x} \int F_0^{(n)} d\Gamma_n. \quad (34)$$

The terms $F_j^{(n)} \xi'_j$ and $G_j^{(n)} \zeta'_j$ in (33) describe the polarization of the final photons and the final electrons, respectively. The last terms, $H_{ij}^{(n)} \zeta'_i \xi'_j$, stand for the correlation of the final particles' polarizations.

By (32) and (33) one can deduce the polarization of the final photon $\xi_j^{(f)}$ and electron $\zeta_j^{(f)}$ resulting from the scattering process itself. According to the usual rules (see [1], §65), we obtain the following expression for the Stokes parameters of the final photon (summed over polarization states of the final electron):

$$\xi_j^{(f)} = \frac{F_j}{F_0}, \quad F_0 = \sum_n F_0^{(n)}, \quad F_j = \sum_n F_j^{(n)}; \quad (35)$$

$$j = 1, 2, 3.$$

The polarization of the final electron (summed over polarization states of the final photon) is given by the invariants

$$\zeta_j^{(f)} = \frac{G_j}{F_0}, \quad G_j = \sum_n G_j^{(n)}; \quad (36)$$

therefore, its polarization vector is

$$\zeta^{(f)} = \sum_{j=1}^3 \frac{G_j}{F_0} \mathbf{n}'_j. \quad (37)$$

In the similar way, the polarization properties for a given harmonic n are described by

$$\xi_j^{(n)(f)} = \frac{F_j^{(n)}}{F_0^{(n)}}, \quad \zeta_j^{(n)(f)} = \frac{G_j^{(n)}}{F_0^{(n)}}. \quad (38)$$

3.2 The results for the circularly polarized laser photons

In this subsection we consider the case of 100% circularly polarized laser beam. The electromagnetic laser field is described by the 4-potential,

$$A_\mu(x) = \frac{m}{e} \xi \left[e_\mu^{(1)} \cos(kx) + P_c e_\mu^{(2)} \sin(kx) \right], \quad (39)$$

where the unit vectors $e_\mu^{(1,2)}$ are given in (19) and P_c is the degree of the circular polarization of the laser wave or the initial photon helicity. Therefore, the Stokes parameters of the laser photon are

$$\xi_1 = \xi_3 = 0, \quad \xi_2 = P_c = \pm 1. \quad (40)$$

We have calculated the coefficients $F_j^{(n)}$, $G_j^{(n)}$ and $H_{ij}^{(n)}$ using the standard technique presented in [1], §101. The necessary traces have been calculated using the package MATHEMATICA. In the considered case of the 100% circularly polarized ($P_c = \pm 1$) laser beam, almost all dependence on the non-linearity parameter ξ^2 accumulates in three functions:

$$f_n \equiv f_n(z_n) = J_{n-1}^2(z_n) + J_{n+1}^2(z_n) - 2J_n^2(z_n),$$

$$g_n \equiv g_n(z_n) = \frac{4n^2 J_n^2(z_n)}{z_n^2}, \quad (41)$$

$$h_n \equiv h_n(z_n) = J_{n-1}^2(z_n) - J_{n+1}^2(z_n),$$

where $J_n(z)$ is the Bessel function. The functions (41) depend on x , y and ξ via the single argument

$$z_n = \frac{\xi}{\sqrt{1+\xi^2}} n s_n. \quad (42)$$

For the small value of this argument one has

$$f_n = g_n = h_n = \frac{(z_n/2)^{2(n-1)}}{[(n-1)!]^2} \quad \text{at } z_n \rightarrow 0; \quad (43)$$

in particular,

$$f_1 = g_1 = h_1 = 1 \quad \text{at } z_1 = 0. \quad (44)$$

It is useful to note that this argument is small for small ξ^2 , as well as for small or for large values of y :

$$z_n \rightarrow 0 \quad \text{either at } \xi^2 \rightarrow 0, \\ \text{or at } y \rightarrow 0, \quad \text{or at } y \rightarrow y_n. \quad (45)$$

The results of our calculations are the following. The function $F_0^{(n)}$, related to the total cross section (34), reads

$$F_0^{(n)} = \left(\frac{1}{1-y} + 1-y \right) f_n - \frac{s_n^2}{1+\xi^2} g_n \\ - \left[\frac{y s_n}{\sqrt{1+\xi^2}} \zeta_2 - \frac{y(2-y)}{1-y} c_n \zeta_3 \right] h_n P_c. \quad (46)$$

The polarization of the final photons $\xi_j^{(f)}$ is given by (35), where

$$F_1^{(n)} = \frac{y}{1-y} \frac{s_n}{\sqrt{1+\xi^2}} h_n P_c \zeta_1, \quad (47)$$

$$F_2^{(n)} = \left(\frac{1}{1-y} + 1-y \right) c_n h_n P_c - \frac{y s_n c_n}{\sqrt{1+\xi^2}} g_n \zeta_2 \\ + y \left(\frac{2-y}{1-y} f_n - \frac{s_n^2}{1+\xi^2} g_n \right) \zeta_3,$$

$$F_3^{(n)} = 2(f_n - g_n) + s_n^2(1+\Delta) g_n \\ - \frac{y}{1-y} \frac{s_n}{\sqrt{1+\xi^2}} h_n P_c \zeta_2;$$

here and below we use the notation

$$\Delta = \frac{\xi^2}{1+\xi^2}. \quad (48)$$

The polarization of the final electrons $\zeta_j^{(f)}$ is given by (36) and (37), with

$$G_1^{(n)} = G_\perp^{(n)} \zeta_1, \quad (49)$$

$$G_2^{(n)} = G_\perp^{(n)} \zeta_2 - \frac{y s_n}{(1-y)\sqrt{1+\xi^2}} (c_n g_n \zeta_3 + h_n P_c),$$

$$G_3^{(n)} = G_\parallel^{(n)} + \frac{y s_n c_n}{\sqrt{1+\xi^2}} g_n \zeta_2,$$

where we introduced the notation

$$G_{\perp}^{(n)} = 2f_n - \frac{s_n^2}{1 + \xi^2} g_n, \quad (50)$$

$$G_{\parallel}^{(n)} = \left[\left(\frac{1}{1-y} + 1-y \right) f_n - \left(1 + \frac{y^2}{1-y} \right) \frac{s_n^2}{1 + \xi^2} g_n \right] \zeta_3 + \frac{y(2-y)c_n}{1-y} h_n P_c. \quad (51)$$

Finally, the correlations of the final particles' polarizations are

$$H_{11}^{(n)} = \frac{y s_n}{\sqrt{1 + \xi^2}} h_n P_c + \frac{y}{1-y} [(2-y)(f_n - g_n) + (1-y + \Delta) s_n^2 g_n] \zeta_2 - \frac{y c_n s_n}{\sqrt{1 + \xi^2}} g_n \zeta_3,$$

$$H_{21}^{(n)} = -\frac{y}{1-y} \times \{ (2-y)(f_n - g_n) + [1 + (1-y)\Delta] s_n^2 g_n \} \zeta_1,$$

$$H_{31}^{(n)} = \frac{y c_n s_n}{(1-y)\sqrt{1 + \xi^2}} g_n \zeta_1, \quad H_{12}^{(n)} = 2c_n h_n P_c \zeta_1,$$

$$H_{22}^{(n)} = -\frac{y c_n s_n}{(1-y)\sqrt{1 + \xi^2}} g_n + 2c_n h_n P_c \zeta_2 - \frac{y s_n}{(1-y)\sqrt{1 + \xi^2}} h_n P_c \zeta_3, \quad (52)$$

$$H_{32}^{(n)} = \frac{y}{1-y} \left[(2-y)f_n - \frac{s_n^2}{1 + \xi^2} g_n \right] + \frac{y s_n}{\sqrt{1 + \xi^2}} h_n P_c \zeta_2 + \frac{2-2y+y^2}{1-y} c_n h_n P_c \zeta_3,$$

$$H_{13}^{(n)} = \left[\frac{2-2y+y^2}{1-y} (f_n - g_n) + \left(1 + \frac{1-y+y^2}{1-y} \Delta \right) s_n^2 g_n \right] \zeta_1,$$

$$H_{23}^{(n)} = -\frac{y s_n}{\sqrt{1 + \xi^2}} h_n P_c + \left[\frac{2-2y+y^2}{1-y} (f_n - g_n) + \left(\frac{1-y+y^2}{1-y} + \Delta \right) s_n^2 g_n \right] \zeta_2 + \frac{y c_n s_n}{\sqrt{1 + \xi^2}} g_n \zeta_3,$$

$$H_{33}^{(n)} = -\frac{y c_n s_n}{(1-y)\sqrt{1 + \xi^2}} g_n \zeta_2 + [2(f_n - g_n) + (1 + \Delta) s_n^2 g_n] \zeta_3.$$

3.3 The results for the linearly polarized laser photons

Here we consider the case of the 100% linearly polarized laser beam. The electromagnetic laser field is described by the 4-potential

$$A^\mu(x) = A^\mu \cos(kx), \quad A^\mu = \frac{\sqrt{2}m}{e} \xi e_L^\mu, \quad e_L e_L = -1, \quad (53)$$

where e_L^μ is the unit 4-vector describing the polarization of the laser photons, which can be expressed in the covariant form via the unit 4-vectors $e^{(1,2)}$ given in (19) as follows:

$$e_L = e^{(1)} \sin \varphi - e^{(2)} \cos \varphi. \quad (54)$$

The invariants

$$\sin \varphi = -e_L e^{(1)}, \quad \cos \varphi = e_L e^{(2)} \quad (55)$$

have a simple meaning in the collider system. For the problem discussed it is convenient to choose the x -axis of this frame of reference along the direction of the laser linear polarization, i.e. along the vector $\mathbf{e}_{L\perp}$. With such a choice, the quantity φ is the azimuthal angle of the final photon in the collider system (analogously, the azimuthal angles β and β' of the electron polarizations ζ and ζ' are also defined with respect to this x -axis).

The Stokes parameters ξ_i are defined with respect to the $x'y'z'$ -axes which are fixed to the scattering plane. The x' -axis is perpendicular to the scattering plane:

$$x' \parallel \mathbf{k} \times \mathbf{k}'; \quad (56)$$

the y' -axis is in that plane

$$y' \parallel \mathbf{k} \times (\mathbf{k} \times \mathbf{k}') = -\omega^2 \mathbf{k}'_{\perp}. \quad (57)$$

The azimuthal angle of the linear polarization of the laser photon in the collider system equals zero with respect to the xyz -axes, and it is $\varphi + (3\pi/2)$ with respect to the $x'y'z'$ -axes. Therefore, in the considered case of the 100% linearly polarized laser beam one has

$$\xi_1 = -\sin 2\varphi, \quad \xi_2 = 0, \quad \xi_3 = -\cos 2\varphi. \quad (58)$$

When calculating the effective cross section (32), we found that almost all dependence on the non-linearity parameter ξ^2 accumulates in three functions:

$$\begin{aligned} \tilde{f}_n &= 4[A_1(n, a, b)]^2 - 4A_0(n, a, b)A_2(n, a, b), \\ \tilde{g}_n &= \frac{4n^2}{z_n^2} [A_0(n, a, b)]^2, \\ \tilde{h}_n &= \frac{4n}{a} A_0(n, a, b) A_1(n, a, b), \end{aligned} \quad (59)$$

where the functions $A_k(n, a, b)$ were introduced in [10] as follows:

$$A_k(n, a, b) \quad (60)$$

$$= \int_{-\pi}^{\pi} \cos^k \psi \exp [i(n\psi - a \sin \psi + b \sin 2\psi)] \frac{d\psi}{2\pi}.$$

The arguments of these functions are

$$a = e \left(\frac{Ap}{kp} - \frac{Ap'}{kp'} \right), \quad b = \frac{1}{8} e^2 A^2 \left(\frac{1}{kp} - \frac{1}{kp'} \right), \quad (61)$$

or, expressed in terms of the non-linearity parameter, they are

$$a = \sqrt{2} \xi m \left(\frac{e_{LP}}{kp} - \frac{e_{LP'}}{kp'} \right), \quad b = \frac{y}{2(1-y)x} \xi^2. \quad (62)$$

In the collider system one has

$$a = -z_n \sqrt{2} \cos \varphi, \quad (63)$$

where z_n is defined in (42). From the definition of the functions (60) one can deduce that

$$A_k(n, -a, b) = (-1)^{n+k} A_k(n, a, b);$$

therefore, the functions \tilde{f}_n , \tilde{g}_n and \tilde{h}_n are even functions of the variable $\cos \varphi$. To find the photon spectrum, one needs also the functions (59) averaged over the azimuthal angle φ :

$$\langle \tilde{f}_n \rangle = \int_0^{2\pi} \tilde{f}_n \frac{d\varphi}{2\pi}, \quad \langle \tilde{g}_n \rangle = \int_0^{2\pi} \tilde{g}_n \frac{d\varphi}{2\pi}. \quad (64)$$

Among the functions A_0 , A_1 and A_2 there are useful relations [10]

$$\begin{aligned} A_1(n, a, b) &= \frac{1}{2} [A_0(n-1, a, b) + A_0(n+1, a, b)], \\ A_2(n, a, b) &= \frac{1}{4} [A_0(n-2, a, b) + 2A_0(n, a, b) \\ &\quad + A_0(n+2, a, b)], \\ (n-2b) A_0(n, a, b) - a A_1(n, a, b) \\ &\quad + 4b A_2(n, a, b) = 0. \end{aligned} \quad (65)$$

To find the behavior of $A_0(n, a, b)$ for small values of its arguments, one can use the expansion into the series of Bessel functions

$$A_0(n, a, b) = \sum_{s=-\infty}^{s=+\infty} J_{n+2s}(a) J_s(b). \quad (66)$$

As a consequence, for small values of $\xi^2 \rightarrow 0$ or $y \rightarrow 0$ we have

$$a \propto \sqrt{y \xi^2}, \quad b \propto y \xi^2, \quad A_k(n, a, b) \propto (y \xi^2)^{|n-k|/2} \quad (67)$$

and

$$\tilde{f}_n, \tilde{g}_n, \tilde{h}_n \propto (y \xi^2)^{n-1}. \quad (68)$$

In particular, at $\xi^2 = 0$ or at $y = 0$

$$\tilde{f}_1 = \langle \tilde{f}_1 \rangle = \langle \tilde{g}_1 \rangle = \tilde{h}_1 = 1, \quad \tilde{g}_1 = 1 + \cos 2\varphi. \quad (69)$$

The results of our calculations are the following. First, we define the auxiliary functions

$$\begin{aligned} X_n &= \tilde{f}_n - (1 + c_n) \left[(1 - \Delta r_n) \tilde{g}_n - \tilde{h}_n \cos 2\varphi \right], \\ Y_n &= (1 + c_n) \tilde{g}_n - 2\tilde{h}_n \cos^2 \varphi, \\ V_n &= \tilde{f}_n \cos 2\varphi \\ &\quad + 2(1 + c_n) \left[(1 - \Delta r_n) \tilde{g}_n - 2\tilde{h}_n \cos^2 \varphi \right] \sin^2 \varphi, \end{aligned} \quad (70)$$

where c_n , r_n and Δ are defined in (10), (11) and (48), respectively.

The term $F_0^{(n)}$, related to the total cross section (34), reads

$$F_0^{(n)} = \left(\frac{1}{1-y} + 1-y \right) \tilde{f}_n - \frac{s_n^2}{1+\xi^2} \tilde{g}_n. \quad (71)$$

The polarization of the final photons $\xi_j^{(f)}$ is given by (35), where

$$\begin{aligned} F_1^{(n)} &= 2X_n \sin 2\varphi, \quad F_3^{(n)} = -2V_n + \frac{s_n^2}{1+\xi^2} \tilde{g}_n, \\ F_2^{(n)} &= \frac{y s_n}{\sqrt{1+\xi^2}} \left(\tilde{h}_n \zeta_1 \sin 2\varphi - Y_n \zeta_2 \right) \\ &\quad + y \left(\frac{2-y}{1-y} \tilde{f}_n - \frac{s_n^2}{1+\xi^2} \tilde{g}_n \right) \zeta_3, \end{aligned} \quad (72)$$

The polarization of the final electrons $\zeta_j^{(f)}$ is given by (36) and (37), with

$$\begin{aligned} G_1^{(n)} &= \tilde{G}_\perp^{(n)} \zeta_1 + \frac{y s_n}{(1-y)\sqrt{1+\xi^2}} \tilde{h}_n \zeta_3 \sin 2\varphi, \\ G_2^{(n)} &= \tilde{G}_\perp^{(n)} \zeta_2 - \frac{y s_n}{(1-y)\sqrt{1+\xi^2}} Y_n \zeta_3, \\ G_3^{(n)} &= \tilde{G}_\parallel^{(n)} - \frac{y s_n}{\sqrt{1+\xi^2}} \left(\tilde{h}_n \zeta_1 \sin 2\varphi - Y_n \zeta_2 \right), \end{aligned} \quad (73)$$

where we use the notation

$$\begin{aligned} \tilde{G}_\perp^{(n)} &= 2\tilde{f}_n - \frac{s_n^2}{1+\xi^2} \tilde{g}_n, \\ \tilde{G}_\parallel^{(n)} &= \left[\left(\frac{1}{1-y} + 1-y \right) \tilde{f}_n \right. \\ &\quad \left. - \left(1 + \frac{y^2}{1-y} \right) \frac{s_n^2}{1+\xi^2} \tilde{g}_n \right] \zeta_3. \end{aligned} \quad (74)$$

At last, the correlations of the final particles' polarizations are

$$\begin{aligned} H_{11}^{(n)} &= \frac{2-2y+y^2}{1-y} X_n \zeta_1 \sin 2\varphi \\ &\quad - y \left(\frac{2-y}{1-y} V_n - \frac{s_n^2}{1+\xi^2} \tilde{g}_n \right) \zeta_2 - \frac{y s_n}{\sqrt{1+\xi^2}} Y_n \zeta_3, \end{aligned}$$

$$\begin{aligned}
H_{21}^{(n)} &= \frac{y}{1-y} \left[(2-y)V_n - \frac{s_n^2}{1+\xi^2} \tilde{g}_n \right] \zeta_1 \\
&+ \frac{2-2y+y^2}{1-y} X_n \zeta_2 \sin 2\varphi \\
&+ \frac{y s_n}{\sqrt{1+\xi^2}} \tilde{h}_n \zeta_3 \sin 2\varphi, \\
H_{31}^{(n)} &= \frac{y s_n}{(1-y)\sqrt{1+\xi^2}} \left(Y_n \zeta_1 - \tilde{h}_n \zeta_2 \sin 2\varphi \right) \\
&+ 2X_n \zeta_3 \sin 2\varphi, \\
H_{12}^{(n)} &= \frac{y s_n}{(1-y)\sqrt{1+\xi^2}} \tilde{h}_n \sin 2\varphi, \\
H_{22}^{(n)} &= -\frac{y s_n}{(1-y)\sqrt{1+\xi^2}} Y_n, \\
H_{32}^{(n)} &= \frac{y}{1-y} \left[(2-y)\tilde{f}_n - \frac{s_n^2}{1+\xi^2} \tilde{g}_n \right], \quad (76) \\
H_{13}^{(n)} &= -\left(\frac{2-2y+y^2}{1-y} V_n - \frac{s_n^2}{1+\xi^2} \tilde{g}_n \right) \zeta_1 \\
&- \frac{y(2-y)}{1-y} X_n \zeta_2 \sin 2\varphi + \frac{y s_n}{\sqrt{1+\xi^2}} \tilde{h}_n \zeta_3 \sin 2\varphi, \\
H_{23}^{(n)} &= \frac{y(2-y)}{1-y} X_n \zeta_1 \sin 2\varphi \\
&- \left(\frac{2-2y+y^2}{1-y} V_n - \frac{1-y+y^2}{1-y} \frac{s_n^2}{1+\xi^2} \tilde{g}_n \right) \zeta_2 \\
&+ \frac{y s_n}{\sqrt{1+\xi^2}} Y_n \zeta_3, \\
H_{33}^{(n)} &= -\frac{y s_n}{(1-y)\sqrt{1+\xi^2}} \left(\tilde{h}_n \zeta_1 \sin 2\varphi + Y_n \zeta_2 \right) \\
&- \left(2V_n - \frac{s_n^2}{1+\xi^2} \tilde{g}_n \right) \zeta_3.
\end{aligned}$$

4 Going to the collider system

4.1 Exact relations

As an example of the application of the above formulae, let us consider the non-linear Compton scattering in the collider system defined in Sect. 2.3. In such a frame of reference the unit vectors \mathbf{n}_j , defined in (27), have a simple form:

$$\begin{aligned}
\mathbf{n}_1 &= \frac{\mathbf{p} \times \mathbf{p}'}{|\mathbf{p} \times \mathbf{p}'|} = \frac{\mathbf{k} \times \mathbf{k}'}{|\mathbf{k} \times \mathbf{k}'|}, \quad \mathbf{n}_2 = \frac{\mathbf{p} \times \mathbf{n}_1}{|\mathbf{p} \times \mathbf{n}_1|} = \frac{\mathbf{k}'_{\perp}}{|\mathbf{k}'_{\perp}|}, \\
\mathbf{n}_3 &= \frac{\mathbf{p}}{|\mathbf{p}|}, \quad (77)
\end{aligned}$$

and the invariants ζ_j are equal to

$$\begin{aligned}
\zeta_1 &= \boldsymbol{\zeta} \mathbf{n}_1 = \zeta_{\perp} \sin(\varphi - \beta), \quad \zeta_2 = \boldsymbol{\zeta} \mathbf{n}_2 = \zeta_{\perp} \cos(\varphi - \beta), \\
\zeta_3 &= \boldsymbol{\zeta} \mathbf{n}_3 = 2\lambda_e. \quad (78)
\end{aligned}$$

Here \mathbf{k}'_{\perp} and $\boldsymbol{\zeta}_{\perp}$ stands for the transverse components of the vectors \mathbf{k}' and $\boldsymbol{\zeta}$ with respect to the vector \mathbf{p} , and $\zeta_{\perp} = |\boldsymbol{\zeta}_{\perp}|$. Therefore, in this frame of reference, ζ_1 is the transverse polarization of the initial electron perpendicular to the scattering plane, ζ_2 is the transverse polarization in that plane and ζ_3 is the doubled mean helicity of the initial electron.

A phase volume element in (32) is equal to

$$d\Gamma = dy d\varphi, \quad (79)$$

and the differential cross section, summed over spin states of the final particles, is

$$\frac{d\sigma^{(n)}(\boldsymbol{\zeta}, \boldsymbol{\xi})}{dy d\varphi} = \frac{r_e^2}{x} F_0^{(n)}. \quad (80)$$

Integrating this expression over φ and then over y , we find

$$\frac{d\sigma^{(n)}(\boldsymbol{\zeta}, \boldsymbol{\xi})}{dy} = \frac{2\pi r_e^2}{x} \langle F_0^{(n)} \rangle, \quad (81)$$

$$\sigma^{(n)}(\boldsymbol{\zeta}, \boldsymbol{\xi}) = \frac{2\pi r_e^2}{x} \int_0^{y_n} \langle F_0^{(n)} \rangle dy,$$

where for the circular polarization

$$\begin{aligned}
\langle F_0^{(n)} \rangle &= \left(\frac{1}{1-y} + 1-y \right) f_n - \frac{s_n^2}{1+\xi^2} g_n \\
&+ \frac{y(2-y)}{1-y} c_n h_n \zeta_3 P_c \quad (82)
\end{aligned}$$

and for the linear polarization

$$\langle F_0^{(n)} \rangle = \left(\frac{1}{1-y} + 1-y \right) \langle \tilde{f}_n \rangle - \frac{s_n^2}{1+\xi^2} \langle \tilde{g}_n \rangle. \quad (83)$$

This means that for the circularly polarized laser photons the differential $d\sigma^{(n)}(\boldsymbol{\zeta}, \boldsymbol{\xi})/dy$ and the total $\sigma^{(n)}(\boldsymbol{\zeta}, \boldsymbol{\xi})$ cross sections for a given harmonic n do not depend on the transverse polarization of the initial electron. For the linearly polarized laser photons, these cross sections do not depend at all on the polarization of the initial electron. These properties are similar to those in the linear Compton scattering.

The polarization of the final photon is given by (35). The polarization vector of the final electron is determined by (37), but, unfortunately, the unit vectors \mathbf{n}'_j in this equation have no simple form similar to (77). Therefore, we have to find the characteristics that are usually used for description of the electron polarization – the mean helicity of the final electron λ'_e and its transverse (to the vector \mathbf{p}') polarization $\boldsymbol{\zeta}'_{\perp}$ in the considered collider system. For this purpose we introduce unit vectors $\boldsymbol{\nu}_j$, one of them is directed along the momentum of the final electron \mathbf{p}' and two others are in the plane transverse to this direction (compare with (77)):

$$\begin{aligned}
\boldsymbol{\nu}_1 &= \mathbf{n}_1, \quad \boldsymbol{\nu}_2 = \frac{\mathbf{p}' \times \mathbf{n}_1}{|\mathbf{p}' \times \mathbf{n}_1|}, \quad \boldsymbol{\nu}_3 = \frac{\mathbf{p}'}{|\mathbf{p}'|}; \\
\boldsymbol{\nu}_i \boldsymbol{\nu}_j &= \delta_{ij}. \quad (84)
\end{aligned}$$

After that one has

$$\begin{aligned}\zeta' \nu_1 &= \zeta'_\perp \sin(\varphi - \beta'), \quad \zeta' \nu_2 = \zeta'_\perp \cos(\varphi - \beta'), \\ \zeta' \nu_3 &= 2\lambda'_e.\end{aligned}\quad (85)$$

Therefore, $\zeta' \nu_1$ is the transverse polarization of the final electron perpendicular to the scattering plane, $\zeta' \nu_2$ is the transverse polarization in that plane and $\zeta' \nu_3$ is the doubled mean helicity of the final electron.

Let us discuss the relation between the projection $\zeta' \nu_j$, defined above, and the invariants ζ'_j , defined in (25) and (30). In the collider system the vectors ν_1 and \mathbf{n}'_1 coincide:

$$\nu_1 = \mathbf{n}'_1; \quad (86)$$

therefore,

$$\zeta' \nu_1 = \zeta'_1. \quad (87)$$

Two other unit vectors \mathbf{n}'_2 and \mathbf{n}'_3 are in the scattering plane and they can be obtained from the vectors ν_2 and ν_3 by the rotation around the axis ν_1 by the angle $(-\Delta\theta)$:

$$\begin{aligned}\zeta' \nu_2 &= \zeta'_2 \cos \Delta\theta + \zeta'_3 \sin \Delta\theta, \\ \zeta' \nu_3 &= \zeta'_3 \cos \Delta\theta - \zeta'_2 \sin \Delta\theta,\end{aligned}\quad (88)$$

where

$$\cos \Delta\theta = \nu_3 \mathbf{n}'_3 \quad \sin \Delta\theta = -\nu_3 \mathbf{n}'_2. \quad (89)$$

As a result, the polarization vector of the final electron (37) is expressed as follows:

$$\begin{aligned}\zeta^{(f)} &= \nu_1 \frac{G_1}{F_0} + \nu_2 \left(\frac{G_2}{F_0} \cos \Delta\theta + \frac{G_3}{F_0} \sin \Delta\theta \right) \\ &+ \nu_3 \left(\frac{G_3}{F_0} \cos \Delta\theta - \frac{G_2}{F_0} \sin \Delta\theta \right).\end{aligned}\quad (90)$$

At the end of this subsection we consider the relation between the used collider system and the centre-of-mass system (CMS) for a given harmonic n , defined by the requirements

$$\mathbf{p}' = -\mathbf{k}', \quad \mathbf{p} = -n(1 - r_n \Delta) \mathbf{k} \quad (91)$$

(in accordance with the conservation law (8) rewritten in the form $p + n(1 - r_n \Delta)k = p' + k'$). In this frame of reference the invariants $\zeta'_j(\text{CMS})$ are defined analogously to the invariants ζ_j :

$$\zeta'_j(\text{CMS}) = -a'_j e'_j(\text{CMS}), \quad (92)$$

where

$$\begin{aligned}e'_1(\text{CMS}) &= e^{(1)}, \quad e'_2(\text{CMS}) = e^{(2)} + \frac{\sqrt{-P^2}}{2p'k'} k' \\ e'_3(\text{CMS}) &= \frac{1}{m} \left(p' - \frac{m^2}{p'k'} k' \right).\end{aligned}\quad (93)$$

It is not difficult to find the relation between these invariants and the invariants ζ'_j , defined in (25):

$$\zeta'_1(\text{CMS}) = \zeta'_1, \quad \zeta'_2(\text{CMS}) = \cos \theta_n^* \zeta'_2 + \sin \theta_n^* \zeta'_3,$$

$$\zeta'_3(\text{CMS}) = \cos \theta_n^* \zeta'_3 - \sin \theta_n^* \zeta'_2, \quad (94)$$

where θ_n^* is the photon scattering angle in the rest system of the final electron ($\mathbf{p}' = 0$):

$$\begin{aligned}\cos \theta_n^* &= 1 - \frac{2r_n}{1 + (1 - r_n) \xi^2}, \\ \sin \theta_n^* &= \frac{s_n}{(1 - r_n \Delta) \sqrt{1 + \xi^2}},\end{aligned}\quad (95)$$

with Δ defined in (48).

4.2 Approximate formulae

All the above formulae are exact. In this subsection we give some approximate formulae useful for application to the important case of high-energy $\gamma\gamma$ and γe colliders. It is expected (see, for example, the TESLA project [19]) that in the conversion region of these colliders, an electron with the energy $E \sim 100$ GeV performs a head-on collisions with laser photons having the energy $\omega \sim 1$ eV per a single photon. In this case the most important kinematic range corresponds to almost back-scattered final photons, i.e. the initial electron is ultra-relativistic and the final photon is emitted at a small angle θ_γ with respect to the z -axis (chosen along the momentum \mathbf{p} of the initial electron):

$$E \gg m, \quad \theta_\gamma \ll 1. \quad (96)$$

In this approximation we have

$$x \approx \frac{4E\omega}{m^2}, \quad y \approx \frac{\omega'}{E} \approx 1 - \frac{E'}{E}; \quad (97)$$

therefore, (80) gives us the distribution of the final photons over the energy and the azimuthal angle. Besides, the photon emission angle for the reaction (2) is (see Fig. 1)




Fig. 1. Photon scattering angles θ_γ for different harmonics n versus the final photon energy ω' at $x = 4.8$ and $\xi^2 = 3$. The dashed curve corresponds to $\xi^2 = 0$

$$\theta_\gamma \approx \frac{m}{E} \sqrt{nx + 1 + \xi^2} \sqrt{\frac{y_n}{y} - 1}, \quad (98)$$

and $\theta_\gamma \rightarrow 0$ at $y \rightarrow y_n$. For a given y , the photon emission angle increases with the increase of n . The electron scattering angle is small,

$$\theta_e \approx \frac{y\theta_\gamma}{1-y} \leq \frac{2n}{\sqrt{1+\xi^2}} \frac{\omega}{m}. \quad (99)$$

Let us consider the azimuthal dependence of the detected Stokes parameters ξ'_j for the final photon. These parameters are defined with respect to the $x'y'z'$ -axes which are fixed to the scattering plane. The x' -axis is the same as for the initial photon and perpendicular to the scattering plane:

$$x' \parallel \mathbf{k} \times \mathbf{k}'; \quad (100)$$

the y' -axis is in that plane

$$y' \parallel \mathbf{k}' \times (\mathbf{k} \times \mathbf{k}'). \quad (101)$$

At small emission angles of the final photon $\theta_\gamma \ll 1$, the final photon moves almost along the direction of the z -axis and $\mathbf{k} \times \mathbf{k}'$ azimuth is approximately equal to $\varphi + (3\pi/2)$. Let ξ'_j be the detected Stokes parameters for the final photons, fixed to the xyz -axes of the collider system. These Stokes parameters are connected with ξ'_j by the following relations:

$$\begin{aligned} \xi'_1 &\approx -\check{\xi}'_1 \cos 2\varphi + \check{\xi}'_3 \sin 2\varphi, & \xi'_2 &= \check{\xi}'_2, \\ \xi'_3 &\approx -\check{\xi}'_3 \cos 2\varphi - \check{\xi}'_1 \sin 2\varphi. \end{aligned} \quad (102)$$

Inserting these relations into the effective cross section, we find the Stokes parameters resulting from the scattering process itself and fixed to the xyz -axes of the collider system:

$$\begin{aligned} \check{\xi}_1^{(f)} &\approx -\frac{F_1}{F_0} \cos 2\varphi - \frac{F_3}{F_0} \sin 2\varphi, & \check{\xi}_2^{(f)} &= \frac{F_2}{F_0}, \\ \check{\xi}_3^{(f)} &\approx -\frac{F_3}{F_0} \cos 2\varphi + \frac{F_1}{F_0} \sin 2\varphi. \end{aligned} \quad (103)$$

Note that (102) and (103) become exact in the limit $y \rightarrow y_n$.

It is not difficult to check that, for the considered case, the angle $\Delta\theta$ between vector $\boldsymbol{\nu}_3$ and \mathbf{n}'_3 is very small:

$$\Delta\theta \approx |\boldsymbol{\nu}_{3\perp} - \mathbf{n}'_{3\perp}| \approx \frac{m\theta_e}{2E'} \leq \frac{n}{\sqrt{1+\xi^2}} \frac{\omega}{E'}. \quad (104)$$

This means that the invariants ζ'_j in (30) almost coincide with the projections $\zeta' \boldsymbol{\nu}_j$, defined in (85),

$$\begin{aligned} \zeta'_1 &\approx \zeta'_\perp \sin(\varphi - \beta'), & \zeta'_2 &\approx \zeta'_\perp \cos(\varphi - \beta'), \\ \zeta'_3 &\approx 2\lambda'_e, \end{aligned} \quad (105)$$

and that the exact equation (90) for the polarization of the final electron can be replaced with a high accuracy by the approximate equation

$$\zeta^{(f)} \approx \sum_{j=1}^3 \frac{G_j}{F_0} \boldsymbol{\nu}_j. \quad (106)$$

Up to now we dealt with the head-on collisions of the laser photon and the initial electron beams, when the collision angle α_0 between the vectors \mathbf{p} and $(-\mathbf{k})$ was equal to zero. The detailed consideration of the case $\alpha_0 \neq 0$ is given in Appendix B; see also [20]. We present here the summary of such a consideration. If $\alpha_0 \neq 0$, the longitudinal component of the vector \mathbf{k} becomes $k_z = -\omega \cos \alpha_0$ and it appeared to be the transverse (to the momentum \mathbf{p}) component \mathbf{k}_\perp . However, this transverse component is small, $|\mathbf{k}_\perp| \lesssim \omega \ll m$, therefore the transverse momenta of the final particles almost compensate each other, $\mathbf{p}'_\perp = \mathbf{k}_\perp - \mathbf{k}'_\perp \approx -\mathbf{k}'_\perp$. The invariants x and y become (compare with (97))

$$x \approx \frac{4E\omega}{m^2} \cos^2 \frac{\alpha_0}{2}, \quad y \approx \frac{\omega'}{E} \approx 1 - \frac{E'}{E}. \quad (107)$$

The polarization parameters of the initial and the final electrons and photons conserve their forms (78), (105), (40), (58), (102) and (103). Therefore, the whole dependence on α_0 enters the effective cross section and the polarizations only via the quantity x (107).

4.3 Averaged polarization of the final particles

The polarization of the final photon and electron, averaged over the azimuthal angle φ in the collider system, is important in many applications. To find it, one can use the same method as for the linear Compton scattering [21,3,4]). Let us consider, for definiteness, the case of the circularly polarized laser photons. Substituting the exact equations (78) for ζ_j and the approximate equations (102) and (105) for ξ'_j, ζ'_j into (33), we obtain the explicit dependence of the effective cross section (32) on φ . Then we take into account that the terms

$$\begin{aligned} \sum_{j=1}^3 F_j^{(n)} \xi'_j &\approx \left(-F_1^{(n)} \cos 2\varphi - F_3^{(n)} \sin 2\varphi \right) \check{\xi}'_1 \\ &+ F_2^{(n)} \check{\xi}'_2 + \left(F_1^{(n)} \sin 2\varphi - F_3^{(n)} \cos 2\varphi \right) \check{\xi}'_3 \end{aligned} \quad (108)$$

after averaging over φ become equal to

$$\sum_{j=1}^3 \langle F_j^{(n)} \xi'_j \rangle \approx \langle F_2^{(n)} \rangle \check{\xi}'_2, \quad (109)$$

where

$$\begin{aligned} \langle F_2^{(n)} \rangle &= \left(\frac{1}{1-y} + 1 - y \right) c_n h_n P_c \\ &+ y \left(\frac{2-y}{1-y} f_n - \frac{g_n^2}{1+\xi^2} g_n \right) \zeta_3. \end{aligned} \quad (110)$$

After a similar averaging of the terms $\sum_j G_j^{(n)} \zeta'_j$, we obtain

$$\frac{d\sigma^{(n)}(\zeta, \xi, \zeta', \xi')}{dy} \quad (111)$$

$$\approx \frac{\pi r_e^2}{2x} \left(\langle F_0^{(n)} \rangle + \langle F_2^{(n)} \rangle \check{\xi}'_2 + G_{\perp}^{(n)} \zeta_{\perp} \zeta'_{\perp} + G_{\parallel}^{(n)} \zeta'_3 \right. \\ \left. + \sum_{i,j=1}^3 \langle H_{ij}^{(n)} \zeta'_i \xi'_j \rangle \right),$$

where $\langle F_0^{(n)} \rangle$, $G_{\perp}^{(n)}$ and $G_{\parallel}^{(n)}$ are given in (82), (50) and (51), respectively. According to the usual rules, we obtain from this equation the averaged polarization of the final particles for the case of the circularly polarized laser photons:

$$\langle \check{\xi}_1^{(n)(f)} \rangle \approx 0, \quad \langle \check{\xi}_2^{(n)(f)} \rangle \approx \frac{\langle F_2^{(n)} \rangle}{\langle F_0^{(n)} \rangle}, \quad \langle \check{\xi}_3^{(n)(f)} \rangle \approx 0, \\ \langle \zeta_{\perp}^{(n)(f)} \rangle \approx \frac{G_{\perp}^{(n)}}{\langle F_0^{(n)} \rangle} \zeta_{\perp}, \quad \langle \zeta_3^{(n)(f)} \rangle \approx \frac{G_{\parallel}^{(n)}}{\langle F_0^{(n)} \rangle}. \quad (112)$$

In a similar way we obtain the averaged polarization of the final particles for the case of the linearly polarized laser photons⁶:

$$\langle \check{\xi}_1^{(n)(f)} \rangle \approx 0, \quad \langle \check{\xi}_2^{(n)(f)} \rangle \approx \frac{\langle F_2^{(n)} \rangle}{\langle F_0^{(n)} \rangle}, \\ \langle \check{\xi}_3^{(n)(f)} \rangle \approx \frac{1}{\langle F_0^{(n)} \rangle} \left[2\langle \tilde{f}_n \rangle - (2 + 2c_n - s_n^2 \Delta) \langle \tilde{g}_n \rangle \right. \\ \left. + (1 + c_n)^2 \langle \tilde{g}_n \cos 2\varphi \rangle \right], \quad (113) \\ \langle \zeta_{\perp}^{(n)(f)} \rangle \approx \frac{\langle \tilde{G}_{\perp}^{(n)} \rangle}{\langle F_0^{(n)} \rangle} \zeta_{\perp}, \quad \langle \zeta_3^{(n)(f)} \rangle \approx \frac{\langle \tilde{G}_{\parallel}^{(n)} \rangle}{\langle F_0^{(n)} \rangle},$$

where $\langle F_0^{(n)} \rangle$, $\tilde{G}_{\perp}^{(n)}$, $\tilde{G}_{\parallel}^{(n)}$ are given in (83), (74) and (75), and

$$\langle F_2^{(n)} \rangle = y \left(\frac{2-y}{1-y} \langle \tilde{f}_n \rangle - \frac{s_n^2}{1+\xi^2} \langle \tilde{g}_n \rangle \right) \zeta_3. \quad (114)$$

Note that the averaged Stokes parameters of the final photon do not depend on ζ_{\perp} and that the averaged polarization vector of the final electron is not equal to zero only if $P_c \neq 0$ or if $\zeta \neq 0$. These properties are similar to those in linear Compton scattering.

5 Limiting cases

In this section we consider several limiting cases in which description of the non-linear Compton scattering is essentially simplified.

(i) The case of $\xi^2 \rightarrow 0$. At small ξ^2 all harmonics with

⁶ To perform such an averaging, we take into account that the functions \tilde{f}_n , \tilde{g}_n and \tilde{h}_n are even functions of the variable $\cos \varphi$. Therefore, $\langle \tilde{g}_n \sin \varphi \rangle = 0$, since the averaged function is odd under the replacement $\varphi \rightarrow -\varphi$, and $\langle \tilde{g}_n \cos \varphi \rangle = \langle \tilde{h}_n \cos \varphi \rangle = 0$, since the averaged function is odd under the replacement $\varphi \rightarrow \pi - \varphi$.

$n > 1$ disappear due to the properties (43), (45), (68), (69) and we have

$$d\sigma^{(n)}(\zeta, \xi, \zeta', \xi') \propto \xi^{2(n-1)} \quad \text{at} \quad \xi^2 \rightarrow 0. \quad (115)$$

We have checked that in this limit our expression for the effective cross section of the first harmonic, $d\sigma^{(1)}$, coincides with the result known for the linear Compton effect; see Appendix A.

(ii) The case of $y \rightarrow 0$ for arbitrary x and ξ^2 . This limit corresponds to forward scattering ($\theta_e = 0$, $\theta_\gamma = \pi$). We expect that in this limit an electron as well as a photon does not change its polarization. Indeed, at small y all harmonics with $n > 1$ disappear due to the properties (43), (45), (68) and (69), and we have

$$d\sigma^{(n)}(\zeta, \xi, \zeta', \xi') \propto y^{n-1} \quad \text{at} \quad y \rightarrow 0, \quad (116)$$

and we found that

$$F_0 = 2, \quad F_j = 2\xi_j, \quad G_j = 2\zeta_j, \quad H_{ij} = 2\zeta_i \xi_j; \quad (117)$$

therefore,

$$\frac{d\sigma(\zeta, \xi, \zeta', \xi')}{dy d\varphi} = \frac{r_e^2}{2x} (1 + \zeta\zeta')(1 + \xi\xi'). \quad (118)$$

It follows from this equation that

$$\zeta^{(f)} = \zeta, \quad \xi^{(f)} = \xi, \quad (119)$$

in accordance with our expectation.

Note that for the case of circularly polarized laser photons the equations, similar to (119), hold also for each harmonic separately: $\zeta^{(n)(f)} = \zeta$, $\xi^{(n)(f)} = \xi$.

(iii) The case of $x \rightarrow 0$ for arbitrary ξ^2 , which corresponds to the classical limit. In this limit an electron does not change its polarization. For a certain harmonic with $n x \ll 1$ one has $y \leq y_n < n x \ll 1$ and

$$G_j^{(n)} = F_0^{(n)} \zeta_j, \quad H_{ij}^{(n)} = \zeta_i F_j^{(n)}; \quad (120)$$

therefore, in this limit the effective cross section both for the circularly and linearly polarized laser photons has the form

$$\frac{d\sigma^{(n)}(\zeta, \xi, \zeta', \xi')}{dy d\varphi} \quad (121) \\ = \frac{r_e^2}{4x} \left(F_0^{(n)} + \sum_{j=1}^3 F_j^{(n)} \xi'_j \right) \left(1 + \sum_{i=1}^3 \zeta_i \zeta'_i \right).$$

As a consequence, the polarization of the final electron for a given harmonic n is

$$\zeta^{(n)(f)} = \sum_{i=1}^3 \zeta_i \mathbf{n}'_i, \quad (122)$$

where the unit vectors \mathbf{n}'_i are defined in (30). To show that $\zeta^{(n)(f)}$ coincides with ζ , one has to demonstrate that the unit vectors \mathbf{n}'_i coincide with the unit vectors \mathbf{n}_i from (27).

This can easily be seen, for example, in the frame of reference in which $\mathbf{p} = 0$ and in which the photon energies are small in the considered limit, $\omega' < n\omega \ll m$, as well as the momentum of the final electron, $|\mathbf{p}'| \ll m$. As a result,

$$\zeta^{(n)(f)} = \zeta, \quad (123)$$

in accordance with our expectation.

(iv) The case of $\theta_\gamma \rightarrow 0$ for the circularly polarized laser photons, $P_c = \pm 1$. At small θ_γ the energy of the final photon is close to its maximum, $y \rightarrow y_n$, and the functions (41) become equal to

$$f_n = g_n = h_n = \gamma_n (y_n - y)^{n-1}, \quad (124)$$

$$\gamma_n = (n^n/n!)^2 \left[\frac{\xi^2}{y_n(1-y_n)(1+\xi^2)} \right]^{n-1}.$$

Therefore, all harmonics with $n > 1$ disappear⁷ and only the photons of the first harmonic can be emitted along the direction of the initial electron beam:

$$\frac{d\sigma^{(n)}(\zeta, \xi, \zeta', \xi')}{dyd\varphi} = \frac{r_e^2}{4x} F^{(n)} \propto f_n \propto (y_n - y)^{n-1}$$

at $y \rightarrow y_n$, (125)

with

$$\frac{F^{(n)}}{f_n} = \frac{2 - 2y_n + y_n^2}{1 - y_n} (1 + \zeta_3 \zeta'_3)(1 - P_c \xi'_2) \quad (126)$$

$$- \frac{y_n(2 - y_n)}{1 - y_n} (\zeta_3 + \zeta'_3)(P_c - \xi'_2)$$

$$+ 2(\zeta_\perp \zeta'_\perp)(1 - P_c \xi'_2).$$

In this limit, the final photons are circularly polarized,

$$\xi_1^{(n)(f)} = 0, \quad \xi_2^{(n)(f)} = -P_c, \quad \xi_3^{(n)(f)} = 0, \quad (127)$$

and the polarization of the final electrons is

$$\zeta_\perp^{(n)(f)} = \frac{2(1 - y_n)}{2 - (2y_n - y_n^2)(1 + \zeta_3 P_c)} \zeta_\perp, \quad (128)$$

$$\zeta_3^{(n)(f)} = \frac{2\zeta_3 - (2y_n - y_n^2)(\zeta_3 + P_c)}{2 - (2y_n - y_n^2)(1 + \zeta_3 P_c)};$$

⁷ The vanishing for the strict backward scattering of all harmonics with $n \geq 3$ is due to the conservation of the z component of the total angular momentum J_z in this limit: the initial value of $J_z = \lambda_e - n\lambda_L$ cannot be equal for $n \geq 3$ to the final value of $J_z = \lambda'_e + \lambda'_\gamma$ (here λ_e (λ'_e) is the helicity of the initial (final) electron and λ_L (λ'_γ) is the helicity of the laser (final) photon). For $n = 2$ this argument does not help, since $J_z = \pm 3/2$ can be realized for the initial and final states. The vanishing of the second harmonics for the strict backward scattering is a specific feature of the process, related to the facts that $J_z = \pm 3/2$ can be realized only if $\lambda'_\gamma = -\lambda_L$ and that the polarization vector $e^{(\lambda'_\gamma)*} = -e^{(\lambda_L)}$ is orthogonal not only to the 4-vectors k, k' but to the 4-vectors p, p' and $e^{(\lambda_L)}$ as well.

in particular, at $\zeta_3 = \pm 1$ the helicity of the electron is conserved in the non-linear Compton scattering, $\zeta_3^{(n)(f)}(y \rightarrow y_n) \rightarrow \zeta_3$. Note that the function $F^{(n)}$ does not depend on the azimuthal angle φ and (127) and (128) are in accordance with the results (112) for the polarizations averaged over φ .

Let us consider in more detail this limit for the first harmonic. The spectrum of the first harmonic,

$$\frac{d\sigma^{(1)}(\zeta, \xi)}{dy} = \frac{2\pi r_e^2}{x(1-y_1)} [2 - (2y_1 - y_1^2)(1 + \zeta_3 P_c)]$$

at $y \rightarrow y_1$, (129)

depends on the non-linearity parameter ξ^2 only via

$$y_1 = \frac{x}{x + 1 + \xi^2}.$$

For the case of ‘‘good polarization’’, $\zeta_3 P_c = -1$, the first harmonic has a peak at $y \rightarrow y_1$ of the value

$$\frac{d\sigma^{(1)}(\zeta, \xi)}{dy} = \frac{4\pi r_e^2}{x} \left(1 + \frac{x}{1 + \xi^2} \right)$$

at $y \rightarrow y_1$ and $\zeta_3 P_c = -1$. (130)

At large $x \gg 1 + \xi^2$ the quantity y_1 tends to 1,

$$1 - y_1 = \frac{1 + \xi^2}{x + 1 + \xi^2} \ll 1, \quad (131)$$

and the function $F^{(1)}$ becomes large:

$$F^{(1)} = \frac{1}{1 - y_1}$$

$$\times [(1 + \zeta_3 \zeta'_3)(1 - P_c \xi'_2) - (\zeta_3 + \zeta'_3)(P_c - \xi'_2)]. \quad (132)$$

Therefore, integrating the effective cross section for the first harmonic, $d\sigma^{(1)} \propto 1/(1-y)$, near its maximum, i.e. in the region

$$\frac{1 + \xi^2}{x} \ll 1 - y \ll 1, \quad (133)$$

we find (with logarithmic accuracy) the total cross section

$$\sigma(\zeta, \xi, \zeta', \xi')$$

$$= \frac{\pi r_e^2}{2x} [(1 + \zeta_3 \zeta'_3)(1 - P_c \xi'_2) - (\zeta_3 + \zeta'_3)(P_c - \xi'_2)]$$

$$\times \ln \frac{x}{1 + \xi^2}. \quad (134)$$

If one sums this expression over the spin states of the final particles, one arrives at the result for the cross section:

$$\sigma(\zeta, \xi) = \frac{2\pi r_e^2}{x} (1 - \zeta_3 P_c) \ln \frac{x}{1 + \xi^2}, \quad (135)$$

which is in accordance with that obtained by a different method in [17].

(v) The case of $\theta_\gamma \rightarrow 0$ for the linearly polarized laser

photons. At $y \rightarrow y_n$, the argument a tends to zero, while the argument b tends to the non-zero constant, $b \rightarrow n\xi^2/[2(1+\xi^2)]$; the function $A_0(n, a, b) \propto \sqrt{y_n - y}$ vanishes for odd n and $A_0(n, a, b) \rightarrow (-1)^{n/2} J_{n/2}(b)$ for even n . As a result, the spectra and the polarizations are different in this limit for odd and even harmonics.

Odd harmonics. The spectrum in this case,

$$\begin{aligned} \frac{d\sigma^{(n)}(\zeta, \xi)}{dy} &= \frac{2\pi r_e^2}{x(1-y_n)} \\ &\times (2 - 2y_n + y_n^2) [J_{(n-1)/2}(b) - J_{(n+1)/2}(b)]^2 \\ &\text{at } y \rightarrow y_n, \end{aligned} \quad (136)$$

depends on the non-linearity parameter ξ^2 via y_n and via the argument b , but the spectrum does not depend on the initial electron polarization. In this limit, the final photons have circular polarization, proportional to ζ_3 , and linear polarization along the direction of the linear polarization of the laser photons,

$$\begin{aligned} \zeta_1^{(n)(f)} &= 0, \quad \zeta_2^{(n)(f)} = \frac{2y_n - y_n^2}{2 - 2y_n + y_n^2} \zeta_3, \\ \zeta_3^{(n)(f)} &= \frac{2(1 - y_n)}{2 - 2y_n + y_n^2}. \end{aligned} \quad (137)$$

The polarization of the final electrons in this case is

$$\begin{aligned} \zeta_{\perp}^{(n)(f)} &= \frac{2(1 - y_n)}{2 - 2y_n + y_n^2} \zeta_{\perp}, \\ \zeta_3^{(n)(f)} &= \zeta_3; \end{aligned} \quad (138)$$

in particular, the final electrons have the same mean helicity as that for the initial one.

Even harmonics. The spectrum reads

$$\begin{aligned} \frac{d\sigma^{(n)}(\zeta, \xi)}{dy} &= \frac{2\pi r_e^2}{x(1-y_n)} \frac{2y_n^2}{\xi^2} [J_{n/2}(b)]^2 \\ &\text{at } y \rightarrow y_n. \end{aligned} \quad (139)$$

In this limit the final photons have only circular polarization equal to ζ_3 , and the final electrons have mean helicity opposite to that for the initial one:

$$\begin{aligned} \zeta_1^{(n)(f)} &= 0, \quad \zeta_2^{(n)(f)} = \zeta_3, \quad \zeta_3^{(n)(f)} = 0, \\ \zeta_{\perp}^{(n)(f)} &= 0, \quad \zeta_3^{(n)(f)} = -\zeta_3. \end{aligned} \quad (140)$$

Note that (137), (138) and (140) are in accordance with the results (113) for the polarizations averaged over φ .

6 Numerical results related to $\gamma\gamma$ and γe colliders

In this section we present some examples which illustrate the dependence of the differential cross sections and the polarizations, obtained in the previous sections, on the final photon energy. We restrict ourselves to properties of the high-energy photon beam which are of most importance for the future $\gamma\gamma$ and γe colliders. The used parameters are close to those in the TESLA project [19]. In particular,

$$x \approx 4E\omega/m^2 = 4.8,$$

and the non-linearity parameter ξ^2 (5) is chosen either the same as in the TESLA project, $\xi^2 = 0.3$, or larger by one order of magnitude, $\xi^2 = 3$, just to illustrate the tendencies. In the figures below we use the notation

$$\sigma_0 = \pi r_e^2 \approx 2.5 \cdot 10^{-25} \text{ cm}^2.$$

The case of circularly polarized laser photons (Figs. 2, 3, 4, 5). The spectra of the few first harmonics are shown in Fig. 2 for the case of ‘‘a good polarization’’, when helicities of the laser photon and the initial electron are opposite, $\zeta_3 P_c = -1$. At a small intensity of the laser wave ($\xi^2 = 0.3$, Fig. 2a) the main contribution is given by photons of the first harmonic and the probability for generation of the




Fig. 2. Energy spectra of final photons for different harmonics n at $\zeta_3 P_c = -1$ and **a** $\xi^2 = 0.3$; **b** $\xi^2 = 3$. The dashed curves correspond to $\xi^2 = 0$




Fig. 3. Mean helicity of the final photons for different harmonics n versus the final photon energy ω' at $P_c = -1$, $\zeta_3 = 1$ and $\xi^2 = 3$. The dashed curve corresponds to $\xi^2 = 0$

higher harmonics is small. However, with the growth of the non-linearity parameter ($\xi^2 = 3$, Fig. 2b), the maximum energy for the first harmonic decreases and the peak of this harmonic at $y = y_1$ (see (130)) decreases as well. As for the higher harmonics, with the rise of ξ^2 (Fig. 2b) we see an increase of the yield of photons with energies higher than the maximum energy of the first harmonic. As a result, the total spectrum becomes considerably wider.

The mean helicities of the final photons for the few first harmonics, $\xi_2^{(n)(f)}$, are shown in Figs. 3, 4a and 5a. In Fig. 3 we compare the two cases $\xi^2 = 3$ and $\xi^2 = 0$. Each harmonic is 100% polarized near its maximum $y = y_n$, therefore, the whole γ beam at $x > 0.5$ has a high degree of circular polarization. However, for linear Compton scattering, the final photon in the region of high energies, $y \geq 0.8$, is almost 100% circularly polarized, while for the non-linear

case, $\xi^2 = 3$, only harmonics with $n \geq 4$ contribute in this energy range. But the cross sections corresponding to these harmonics are small (see Fig. 2b).

There is an interesting feature related to the linear polarization of the final photons; see Figs. 4b and 5b. Such a polarization is described by the Stokes parameters $\xi_1^{(n)(f)}$ and $\xi_3^{(n)(f)}$. These parameters averaged over the azimuthal angle φ (see (112)) vanish. On the other hand, the degree of the linear polarization for a given angle φ may be close to 100%, especially in the case when the initial electron has a transverse polarization as well (see Fig. 5b). It is not excluded that this property of our process may result in a sizable dependence of the luminosities of $\gamma\gamma$ collisions at a photon collider on the mutual orientation (parallel or perpendicular) of the linear polarizations of the colliding photons. Such a dependence is of importance, for example, for the study of a Higgs boson production at a $\gamma\gamma$ collider. As far as we know, this possibility has not been discussed so far even for linear Compton scattering.

The case of the linearly polarized laser photons (Figs. 6, 7 and 8). The spectra of the first few harmonics for this case are shown in Fig. 6. They differ considerably from those for the case of the circularly polarized laser photons shown in Fig. 2. First of all, in the considered case the spectra do not depend on the polarization of the initial electrons. The maximum of the first harmonic at $y = y_1$ now is about two times smaller than that on Fig. 2. Besides, the harmonics with $n > 1$ do not vanish at $y = y_n$ contrary to such harmonics on Fig. 2.

For the first few harmonics the mean helicities of the final photons, $\xi_2^{(n)(f)}$, are shown at $\zeta_3 = 1$ in Fig. 7a for $\xi^2 = 0.3$ and in Fig. 7b for $\xi^2 = 3$. We would like to direct attention to the surprising fact that each harmonic is almost 100% circularly polarized near the high-energy part of the spectrum. The curves on Fig. 7 are given for the azimuthal angle $\varphi = 0$, for other values of φ these curves would look a bit different, but not too much.




Fig. 4. The Stokes parameters of the final photons for different harmonics n versus the final photon energy ω' at $\zeta_3 = 1$, $P_c = -1$, $\xi^2 = 0.3$: **a** the mean helicity $\xi_2^{(n)(f)}$; **b** the linear polarization $\xi_3^{(n)(f)}$ transverse to the scattering plane. The Stokes parameter obeys $\xi_1^{(n)(f)} = 0$ in this case




Fig. 5. The same as in Fig. 4, but for $\zeta_1 = 0$, $\zeta_2 = 0.7$ and $\zeta_3 = 0.7$




Fig. 6. Energy spectra of final photons for different harmonics n in the case of the linearly polarized laser photons and **a** $\xi^2 = 0.3$; **b** $\xi^2 = 3$. The dashed curves correspond to $\xi^2 = 0$




Fig. 7. The mean helicity $\xi_2^{(n)(f)}$ of the final photons for different harmonics n versus the final photon energy ω' at $\zeta_3 = 1$ and **a** $\xi^2 = 0.3$; **b** $\xi^2 = 3$. The laser photons are linearly polarized. The scattering plane is parallel to the direction of the laser photon polarization




Fig. 8. The Stokes parameters $\xi_3^{(n)(f)}$ of the final photons for different harmonics n versus the final photon energy ω' at **a** $\xi^2 = 0.3$; **b** $\xi^2 = 3$. The laser photons are linearly polarized. The scattering plane is perpendicular to the direction of the laser photon polarization. The Stokes parameter obeys $\xi_1^{(n)(f)} = 0$ in this case




Fig. 9. The Stokes parameters $\langle \xi_3^{(n)(f)} \rangle$ of the final photons for different harmonics n , averaged over the azimuthal angle φ , versus the final photon energy ω' at **a** $\xi^2 = 0.3$; **b** $\xi^2 = 3$. The laser photons are linearly polarized. The averaged Stokes parameter $\langle \xi_1^{(n)(f)} \rangle = 0$ in this case

The degree of linear polarization of the final photons is not large in the high-energy part of the spectrum, but it becomes rather high in the middle and the low part of the spectrum; see Fig. 8. Certainly, the direction of this polarization depends on the azimuthal angle φ , and as a result the linear polarization averaged over φ is substantially smaller, see Fig. 9, than the one on Fig. 8. Nevertheless, the non-trivial effects, related to the high degree of linear polarization present at a certain φ , do exist. For example, it was shown [22] that for linear Compton scattering this feature leads to an important effect for the luminosity of $\gamma\gamma$ collisions.

7 Summary and comparison with other papers

Our main results are given by (46)–(52) for circularly and by (71)–(76) for linearly polarized laser beams. They are

expressed in terms of the 16 functions F_0 , F_j , G_j and H_{ij} with $i, j = 1 \div 3$, which completely describe the polarization properties of the non-linear Compton scattering in a compact invariant form. The functions F_0 and F_j enter the total cross section (34) and (81), the differential cross sections (80) and (81) and the Stokes parameters of the final photons (35). The polarization of the final electrons is described by the functions G_j , which enter the polarization vector $\zeta^{(f)}$ given by the exact equations (37) and (90), and the approximate equations (106).

Besides, we considered the kinematics and the approximate formulae relevant for the problem of $e \rightarrow \gamma$ conversion at the $\gamma\gamma$ and γe colliders. In particular, we discuss the polarization of the final photons and electrons averaged over the azimuthal angle φ (see (112) and (113)) and the spectra and the polarization of the final particles in the limit of small and large energies of the final photons (Sect. 5). For circularly polarized laser photons and for large values of

the parameter $x \gg 1 + \xi^2$, we obtain (with logarithmic accuracy) the simple analytical expression for the total cross section (134). In Sect. 6 we present the numerical results and discuss several interesting features in the final photon polarization, which may be useful for the analysis of the luminosity of $\gamma\gamma$ collisions.

Let us compare our results with those obtained earlier. *Circular polarization of laser photons.* In the literature we found the results which can be compared with our functions F_0, F_2 and G_j , namely, in [10] (the function F_0 only at $\zeta_j = 0$), in [12–14] (the functions F_0 and F_2 only at $\zeta_1 = \zeta_2 = 0$) and in [15] (the function F_0 for arbitrary ζ_j). For these cases our results coincide with the above mentioned ones. For polarization of the final electron, our results differ slightly from those in [14, 15]. Namely, the function G_2 at $\zeta_1 = \zeta_2 = 0$ given in [14] and the functions G_j for arbitrary ζ_j obtained in [15] coincide with ours. However, the polarization vector $\zeta^{(f)}$ in the collider system is obtained in [14, 15] only in the approximate form equivalent to our approximate (106).

Linear polarization of laser photons. For this case we found in the literature the expressions which can be compared with our functions F_0, F_1, F_2, F_3 . The results in [10] are in agreement with our function F_0 (71), and the results in [11] are in agreement with our functions F_j (72).

The correlation of the final particles' polarizations are described by the functions H_{ij} given in (52) and (76). We did not find in the literature any results for these functions both for the case of the circular and the linear polarization.

Acknowledgements. We are grateful to I. Ginzburg, M. Galynskii, A. Milshtein, S. Polityko and V. Telnov for useful discussions. This work is partly supported by INTAS (code 00-00679) and RFBR (code 03-02-17734 and 02-02-17884); D.Yu.I. acknowledges the support of Alexander von Humboldt Foundation.

Appendix A: Limit of the weak laser field

At $\xi^2 \rightarrow 0$, the cross section (32) has the form

$$d\sigma(\zeta, \xi, \zeta', \xi') = \frac{r_e^2}{4x} F d\Gamma, \quad (141)$$

$$d\Gamma = \delta(p + k - p' - k') \frac{d^3k'}{\omega'} \frac{d^3p'}{E'},$$

where

$$F = F_0 + \sum_{j=1}^3 (F_j \xi'_j + G_j \zeta'_j) + \sum_{i,j=1}^3 H_{ij} \zeta'_i \xi'_j. \quad (142)$$

To compare it with the cross section for linear Compton scattering, we should take into account that the Stokes parameters of the initial photon have the values (40) for the circular polarization and the values (58) for the linear polarization. Besides, our invariants c_1, s_1, r_1 and the auxiliary functions (70) transform at $\xi^2 \rightarrow 0$ to

$$c_1 \rightarrow c = 1 - 2r, \quad s_1 \rightarrow s = 2\sqrt{r(1-r)},$$

$$r_1 \rightarrow r = \frac{y}{x(1-y)}, \quad (143)$$

$$X_1 \rightarrow -c, \quad Y_1 \rightarrow c(1 - \xi_3), \quad V_1 \rightarrow -\xi_3.$$

Our functions

$$\begin{aligned} F_0 &= \frac{1}{1-y} + 1 - y - s^2(1 - \xi_3) \\ &\quad - y \left(s \zeta_2 - \frac{2-y}{1-y} c \zeta_3 \right) \xi_2, \\ F_1 &= 2c\xi_1 + \frac{y}{1-y} s \zeta_1 \xi_2, \end{aligned} \quad (144)$$

$$\begin{aligned} F_2 &= \left(\frac{1}{1-y} + 1 - y \right) c \xi_2 - ysc \zeta_2 \\ &\quad + y \left(\frac{2-y}{1-y} - s^2 \right) \zeta_3 - ysc \zeta_1 \xi_1 + ys(c\zeta_2 + s\zeta_3) \xi_3, \\ F_3 &= s^2 - \frac{y}{1-y} s \zeta_2 \xi_2 + (1 + c^2) \xi_3 \end{aligned}$$

coincide with those in [3]. Our functions

$$G_1 = (1 + c^2 + s^2 \xi_3) \zeta_1 - \frac{y}{1-y} s \zeta_3 \xi_1, \quad (145)$$

$$\begin{aligned} G_2 &= -\frac{ys}{1-y} \xi_2 + (1 + c^2 + s^2 \xi_3) \zeta_2 \\ &\quad - \frac{ysc}{1-y} (1 - \xi_3) \zeta_3, \end{aligned}$$

$$\begin{aligned} G_3 &= yc \frac{2-y}{1-y} \xi_2 + ysc \zeta_1 \xi_1 + ysc(1 - \xi_3) \zeta_2 \\ &\quad + \left[1 + \left(\frac{1}{1-y} - y \right) (c^2 + s^2 \xi_3) \right] \zeta_3 \end{aligned}$$

coincide with the functions Φ_j given by (31) in [4]. Finally, we check that our functions

$$\begin{aligned} H_{11} &= ysc\xi_2 + \frac{2-2y+y^2}{1-y} c \zeta_1 \xi_1 \\ &\quad + y \left[\frac{2-y}{1-y} \xi_3 + s^2(1 - \xi_3) \right] \zeta_2 - ysc(1 - \xi_3) \zeta_3, \\ H_{21} &= -\frac{y}{1-y} [s^2 + (1 - y + c^2)\xi_3] \zeta_1 \\ &\quad + \frac{2-2y+y^2}{1-y} c \xi_1 \zeta_2 - ysc \xi_1 \zeta_3, \\ H_{31} &= \frac{ycs}{1-y} (1 - \xi_3) \zeta_1 + \frac{ys}{1-y} \xi_1 \zeta_2 + 2c \xi_1 \zeta_3, \\ H_{12} &= 2c \zeta_1 \xi_2 - \frac{ys}{1-y} \xi_1, \\ H_{22} &= -\frac{ycs}{1-y} (1 - \xi_3) + 2c \zeta_2 \xi_2 - \frac{ys}{1-y} \zeta_3 \xi_2, \\ H_{32} &= \frac{y}{1-y} [2 - y - s^2(1 - \xi_3)] + ysc \zeta_2 \xi_2 \\ &\quad + \frac{2-2y+y^2}{1-y} c \zeta_3 \xi_2, \end{aligned} \quad (146)$$

$$\begin{aligned}
H_{13} &= s^2 \zeta_1 + \left(1 + c^2 + \frac{y^2}{1-y}\right) \xi_3 \zeta_1 \\
&\quad - yc \frac{2-y}{1-y} \xi_1 \zeta_2 - ys \xi_1 \zeta_3, \\
H_{23} &= -ys \xi_2 + y \frac{2-y}{1-y} c \xi_1 \zeta_1 + \frac{1-y+y^2}{1-y} s^2 \zeta_2 \\
&\quad + \left(1 + c^2 + \frac{y^2 c^2}{1-y}\right) \xi_3 \zeta_2 + ysc(1-\xi_3) \zeta_3, \\
H_{33} &= \frac{ys}{1-y} \xi_1 \zeta_1 - \frac{y sc}{1-y} (1-\xi_3) \zeta_2 \\
&\quad + s^2 \zeta_3 + (1+c^2) \xi_3 \zeta_3
\end{aligned}$$

coincide with the corresponding functions in [2]. Making such a comparison, one has to take into account that the set of unit 4-vectors used in our paper (see (19) and (22)) and the one used in [2] are different. The relations between our notation and the notation of [2], which are marked below by the super-index G, are the following:

$$\begin{aligned}
\xi_{1,3}^G &= -\xi_{1,3}, \quad \xi_2^G = \xi_2, \quad \xi'_{1,3}{}^G = -\xi'_{1,3}, \quad \xi_2'^G = \xi_2', \\
\zeta_{1,3}^G &= \zeta_{3,1}, \quad \zeta_2^G = -\zeta_2, \\
\zeta_1'^G &= c\zeta_3' - s\zeta_2', \quad \zeta_2'^G = -c\zeta_2' - s\zeta_3', \quad \zeta_3'^G = \zeta_1'.
\end{aligned} \tag{147}$$

Besides, in [2] there is a misprint, namely, the factor 1/4 should be inserted in the right-hand-side of the equation which is analogous to our (141).

Appendix B: The case of non-zero beam collision angle

In this appendix we consider the case when the beam collision angle α_0 between the vectors \mathbf{p} and $(-\mathbf{k})$ is not equal to zero. We omit systematically terms of the order of ω/m , m/E , θ_γ and θ_e . In this approximation the vectors (19) take the form

$$\begin{aligned}
\mathbf{e}_\perp^{(1)} &= \frac{\mathbf{p} \times \mathbf{p}'}{|\mathbf{p} \times \mathbf{p}'|}, \quad \mathbf{e}_\perp^{(2)} = -\frac{\mathbf{k}'_\perp}{|\mathbf{k}'_\perp|}, \\
e_0^{(j)} &= e_z^{(j)} = \frac{\mathbf{e}_\perp^{(j)} \mathbf{k}_\perp}{(1 + \cos \alpha_0) \omega}, \\
e^{(i)} e^{(j)} &= -\mathbf{e}_\perp^{(i)} \mathbf{e}_\perp^{(j)} = -\delta_{ij},
\end{aligned} \tag{148}$$

i.e. the transverse components $\mathbf{e}_\perp^{(j)}$ do not differ from their values at $\alpha_0 = 0$, and zero and the z components of $e^{(j)}$ vanish in the limit $\mathbf{k}_\perp \rightarrow 0$. As a consequence, the vector

$$\mathbf{e}_{L\perp} = \mathbf{e}_\perp^{(1)} \sin \varphi - \mathbf{e}_\perp^{(2)} \cos \varphi \tag{149}$$

also does not differ from its value at $\alpha_0 = 0$. This means that the parameter φ in all the equations of Sect. 3.3 is the same azimuthal angle of the final photon counted from the x -axis chosen along the direction of the vector $\mathbf{e}_{L\perp}$. Therefore,

the Stokes parameters of the laser photons conserve their values (40) and (58).

The Stokes parameters of the final photons are determined with respect to the transverse to the vector \mathbf{k}' components of the polarization 4-vectors $e^{(1)}$ and $(-e^{(2)})$. At small emission angles $\theta_\gamma \ll 1$, the final photon moves almost along the direction of the z -axis and, therefore, the components of $e^{(j)}$, which are transverse to the momentum \mathbf{k}' , coincide within the used accuracy with the vectors $\mathbf{e}_\perp^{(j)}$ from (148). This means that the Stokes parameters of the final photons also conserve their forms (102) and (103).

Finally, using (148) it is not difficult to check that the polarization parameters of the electrons ζ_j and ζ_j' , defined by (25), (26) and (30), conserve their forms (78) and (105). Therefore, the whole dependence on α_0 enters into the effective cross section and the polarizations only via the quantity $x = (4E\omega/m^2) \cos^2(\alpha_0/2)$.

Finally, one needs to discuss the relation between the direction of the laser wave linear polarization and the angle φ which enters (149) and the equations in Sect. 3.3. Let us decompose the vector \mathbf{e}_L along the photon momentum \mathbf{k} and in the plane orthogonal to that momentum:

$$\mathbf{e}_L = \tilde{\mathbf{e}}_L + c\mathbf{k}, \quad \tilde{\mathbf{e}}_L \mathbf{k} = 0. \tag{150}$$

It is important to note that the electric field of the laser wave is directed along the vector $\tilde{\mathbf{e}}_L$. Therefore, this vector, $\tilde{\mathbf{e}}_L$ (not the vector $\mathbf{e}_{L\perp}$!), determines the direction of the linear polarization of the laser photons. To parametrize this direction we introduce the angle γ ,

$$\tilde{\mathbf{e}}_L = \mathbf{a} \sin \gamma + \mathbf{b} \cos \gamma, \tag{151}$$

$$\mathbf{a} = \frac{\mathbf{k} \times \mathbf{p}}{|\mathbf{k} \times \mathbf{p}|}, \quad \mathbf{b} = \frac{\mathbf{k} \times \mathbf{a}}{|\mathbf{k} \times \mathbf{a}|},$$

$$c = -\frac{\sin \alpha_0}{(1 + \cos \alpha_0)\omega} \cos \gamma.$$

Here the two unit vectors \mathbf{a} and \mathbf{b} are orthogonal to each other and to the vector \mathbf{k} . Besides, the vector \mathbf{a} is orthogonal to the colliding plane, defined by the vectors \mathbf{p} and $(-\mathbf{k})$, while the vector \mathbf{b} is in that plane. The colliding plane should be distinguished from the scattering plane, defined by the vectors \mathbf{p} and \mathbf{k}'_\perp . Let the angle of the scattering plane counted from the colliding plane be φ' . Using the above equations it is not difficult to find that

$$\mathbf{e}_{L\perp} = \mathbf{a} \sin \gamma - \frac{\mathbf{k}_\perp}{|\mathbf{k}_\perp|} \cos \gamma. \tag{152}$$

This means that the angle γ which parametrizes the direction of the linear polarization is simultaneously the angle of the vector $\mathbf{e}_{L\perp}$ counted from the colliding plane. Therefore we conclude that the parameter φ is equal to the difference of two angles, the angle between the two planes and the angle γ ,

$$\varphi = \varphi' - \gamma. \tag{153}$$

References

1. V.B. Berestetskii, E.M. Lifshitz, L.P. Pitaevskii, Quantum electrodynamics (Nauka, Moscow 1989)
2. A.G. Grozin, Proc. Joint Inter. Workshop on High Energy Physics and Quantum Field Theory (Zvenigorod), edited by B.B. Levchenko, Moscow State Univ. (Moscow, 1994) p. 60; Using REDUCE in high energy physics (Cambridge University Press (1997))
3. I.F. Ginzburg, G.L. Kotkin, S.L. Panfil, V.G. Serbo, *Yad. Fiz.* **38**, 1021 (1983);
I.F. Ginzburg, G.L. Kotkin, S.L. Panfil, V.G. Serbo, V.I. Telnov, *Nucl. Instr. Meth.* **219**, 5 (1984)
4. G.L. Kotkin, S.I. Polityko, V.G. Serbo, *Nucl. Instr. Meth. A* **405**, 30 (1998)
5. I.F. Ginzburg, G.L. Kotkin, V.G. Serbo, V.I. Telnov, *Nucl. Instr. Meth.* **205**, 47 (1983)
6. C. Bula, K. McDonald et al., *Phys. Rev. Lett.* **76**, 3116 (1996)
7. V.I. Telnov, *Phys. Rev. Lett.* **78**, 4757 (1997), erratum **80**, 2747 (1998); *Nucl. Instr. Meth. A* **455**, 80 (2000)
8. I.F. Ginzburg, G.L. Kotkin, S.I. Polityko, *Sov. Nucl. Phys.* **40**, 1495 (1984)
9. M. Galynskii, E. Kuraev, M. Levchuk, V. Telnov, *Nucl. Instrum. Meth. A* **472**, 267 (2001)
10. A.I. Nikishov, V.I. Ritus, *Zh. Eksp. Teor. Fiz.* **46**, 776 (1964); *Trudy FIAN* 111 (1979) (Proc. Lebedev Institute, in Russian)
11. Ya.T. Grinchishin, M.P. Rekalov, *Yad. Fiz.* **40**, 181 (1984)
12. M.V. Galynskii, S.M. Sikach, *Sov. Phys. JETP* **101**, 441 (1992) (in English)
13. Yu.S. Tsai, *Phys. Rev. D* **48**, 96 (1993)
14. K. Yokoya, *CAIN2e Users Manual*. The latest version is available from <ftp://lcdev.kek.jp/pub/Yokoya/cain235/CainMan235.pdf>
15. E. Bol'shedvorsky, S. Polityko, A. Misaki, *Prog. Theor. Phys.* **104**, 769 (2000)
16. V.I. Telnov, *Nucl. Instr. Meth. A* **355**, (1995)
17. V.N. Baier, V.M. Katkov, A.I. Milshtein, V.M. Strakhovenko, *Zh. Eksp. Teor. Fiz.* (in Russian) **69**, 783 (1975)
18. D.Yu. Ivanov, G.L. Kotkin, V.G. Serbo, hep-ph/0310325; hep-ph/0311222
19. R. Brinkmann et al., *Nucl. Instr. Meth. A* **406**, 13 (1998)
20. Ya.T. Grinchishin, *Yad. Fiz.* **36**, 1450 (1982)
21. I.I. Goldman, V.A. Khoze, *ZhETF* **57**, 918 (1969); *Phys. Lett. B* **29**, 426 (1969)
22. S. Petrosyan, V. Serbo, V. Telnov, Nontrivial effects in linear polarization at photon colliders. ECFA Study Physics and Detectors for a Linear Collider (November 13–16, 2003, Montpellier, France)