

SuperWIMP dark matter scenario in light of WMAP

Fei Wang¹, Jin Min Yang^{1,2}

¹ Institute of Theoretical Physics, Academia Sinica, Beijing 100080, China

² Department of Physics, Tohoku University, Aoba-ku, Sendai 980, Japan

Received: 17 August 2004 / Revised version: 9 September 2004 /

Published online: 3 November 2004 – © Springer-Verlag / Società Italiana di Fisica 2004

Abstract. The heavy gravitino in the minimal supergravity (mSUGRA) models is likely to be the lightest supersymmetric particle (LSP). Produced from the late decays of the metastable weakly interacting massive particles (WIMPs) such as the lightest neutralinos, the stable gravitinos can be plausible candidates for the cold dark matter in the universe. Such gravitino dark matter can naturally evade the current detection experiments due to its superweak couplings. However, this scenario must be subjected to the constraints from the big bang nucleosynthesis (BBN) predictions for light element abundances as well as the Wilkinson microwave anisotropy probe (WMAP) data for the relic density. Assuming the popular case in which the lightest neutralino is the next-to-lightest supersymmetric particle (NLSP), we find that requiring BBN predictions for light element abundances to agree with the WMAP data can impose upper and lower mass bounds on both the gravitino LSP and the neutralino NLSP. A scan over the mSUGRA parameter space, subjected to the BBN constraints, the WMAP data and the $b \rightarrow s\gamma$ bounds, shows that the low $\tan\beta$ ($\lesssim 40$) region as well as the region accessible at the CERN Large Hadron Collider (LHC) will be severely constrained. Such stringent constraints on the parameter space might be instructive for testing this scenario in future collider experiments.

1 Introduction

The nature of the dark matter is one of the mysteries in today's physical science. It has been intensively explored both theoretically and experimentally. Studies showed that the cosmic dark matter is plausibly composed of non-baryonic weakly interacting massive particles (WIMPs) [1]. While the standard model of particle physics cannot provide a candidate for the dark matter WIMP, the popular supersymmetric theory with R -parity conservation can provide a good candidate, i.e., the lightest supersymmetric particle (LSP). So far the widely studied scenario is that the lightest neutralino is assumed to be the LSP. However, despite the overwhelming popularity of this scenario, other possibilities should not be ignored due to the following reasons. On the one hand, the success of such neutralino dark matter scenario may be spoiled by the problems caused by the gravitino in the reheating era [2]. On the other hand, the neutralino dark matter scenario has not yet been confirmed by current experiments [3].

One possible scenario other than the neutralino LSP one is that the gravitino is assumed to be the LSP. Such gravitino LSPs can form warm or cold cosmic dark matter, depending on the gravitino mass.

(i) In some low-energy SUSY breaking models, like the gauge mediated SUSY breaking (GMSB) models, the gravitino can be as light as keV, much lighter than other supersymmetric particles. It can thus form the warm dark matter. Note that the recent WMAP data imposed severe

constraints on the dark matter type. As analyzed in [4], while a very tiny component of dark matter can be the hot neutrinos, the warm dark matter is ruled out due to the detected early re-ionization of the universe at a redshift $z \approx 0.20$. Therefore, the scenario of the warm dark matter gravitino is not favored by recent observations.

(ii) In the popular mSUGRA models, the gravitino mass is unspecified and only known to be of the weak scale. Such a heavy gravitino is possibly the LSP and can form the cold dark matter in the universe. In contrast to the highly constrained scenarios, in which the gravitino is produced as a thermal relic [5] or produced during reheating [6], a new scenario, assuming the gravitino to be produced from the late decays of the thermal relic WIMPs, was recently proposed in [7].

In this article we focus our attention on this new gravitino dark matter scenario. Since the heavy gravitinos couple gravitationally, they are naturally the so-called superweakly interacting massive particles (SuperWIMPs). As a plausible candidate for the cold dark matter in the universe [7], the gravitino SuperWIMP can naturally evade the current dark matter detection experiments due to its superweak couplings. However, this scenario must be subjected to the constraints from the big bang nucleosynthesis (BBN) as well as the WMAP data [8].

(1) The late decays of WIMPs (like neutralinos) into gravitino SuperWIMPs will release electromagnetic (EM) and hadronic energy. Such an energy release will alter the BBN predictions for light element abundances [9, 10]. Requiring

the resulted predictions for light element abundances to agree with the measured values will impose strong constraints on the gravitino dark matter scenario.

(2) WMAP precisely measured many quantities, especially the total matter density and the baryon density,

$$\Omega_m h^2 = 0.135_{-0.009}^{+0.008}, \quad \Omega_b h^2 = 0.0224_{-0.009}^{+0.009}. \quad (1.1)$$

From such results we can deduce the 2σ range for the cold dark matter density

$$\Omega_{\text{CDM}} h^2 = 0.1126_{-0.0181}^{+0.0161}, \quad (1.2)$$

which is dramatically more accurate than previous results and agrees quite well with other approaches. Such precise measurements will impose strong constraints on the parameter space of the gravitino dark matter scenario.

The aim of this article is to examine these constraints on the parameter space of mSUGRA in this new gravitino dark matter scenario. Assuming the popular case that the lightest neutralino is the NLSP, we will examine the constraints on the mSUGRA parameter space from the BBN light element abundances, the WMAP data of relic density as well as the $b \rightarrow s\gamma$ branching ratio data.

2 Constraints on the parameter space

In the new scenario [7], the NLSPs freeze out with a thermal relic density Ω_{NLSP} and then decay to the gravitino at time $10^4 \sim 10^8$ s. Thus the relic density of gravitino dark matter is obtained by

$$\Omega_{\tilde{G}} = \frac{m_{\tilde{G}}}{M_{\text{NLSP}}} \Omega_{\text{NLSP}}. \quad (2.1)$$

The late decays of the NLSPs will release energy which will alter the light element abundances. In fact, the later injection of high energy photons with the stopping energy inversely proportional to the temperature by scattering off the background photon will dissociate the existing light elements. If carefully chosen, such an injecting of EM energy can destruct the light element abundances to the proper values. It is well known that the BBN predictions for light element abundances are quite successful for most light elements. Yet the BBN predictions for ${}^4\text{He}$ and ${}^7\text{Li}$ (especially ${}^7\text{Li}$) seemingly do not agree with the WMAP data [11]. Requiring such a discrepancy to be settled by the energy released from the late-decaying NLSPs and, at the same time, requiring such an energy release not to spoil the successful BBN predictions for other light elements, the constraints on the NLSP lifetime (in seconds) and EM energy release can be obtained [9]:

$$1.5 \times 10^6 \text{ s} \leq \tau \leq 4 \times 10^6 \text{ s}, \quad (2.2)$$

$$0.8 \times 10^{-9} \text{ GeV} \leq \zeta_{\text{EM}} \leq 1.5 \times 10^{-9} \text{ GeV}, \quad (2.3)$$

where

$$\zeta_{\text{EM}} = \epsilon_{\text{EM}} B_{\text{EM}} Y_{\text{NLSP}} \quad (2.4)$$

is the emitted EM energy density, with ϵ_{EM} being the initial EM energy release from the decay, B_{EM} being the branching fraction of the decay into EM components, and $Y_{\text{NLSP}} = n_{\text{NLSP}}/n_\gamma$ being the NLSP number density normalized to the BG photon number density. In the derivation of the above bounds, the ratio $\eta \equiv n_B/n_\gamma$ is fixed to be 6×10^{-10} , and the bounds on the primordial abundances of light elements are taken to be [9]

$$1.3 \times 10^{-5} < \text{D}/\text{H} < 5.3 \times 10^{-5}, \quad (2.5)$$

$$0.227 < Y_p < 0.249, \quad (2.6)$$

$$9.0 \times 10^{-11} < {}^7\text{Li}/\text{H} < 2.8 \times 10^{-10}, \quad (2.7)$$

$${}^6\text{Li}/{}^7\text{Li} \lesssim 0.07, \quad (2.8)$$

$${}^6\text{Li}/\text{H} \lesssim 2 \times 10^{-11}, \quad (2.9)$$

where Y_p denotes ${}^4\text{He}$ abundance.

The dominant decay of neutralino ($\tilde{\chi}$) NLSP into gravitino (\tilde{G}) LSP is through $\tilde{\chi} \rightarrow \gamma\tilde{G}$ with a rate given by

$$\Gamma(\tilde{\chi} \rightarrow \gamma\tilde{G}) = \frac{|N_{11} \cos \theta_W + N_{12} \sin \theta_W|^2}{48\pi M_*^2} \frac{m_{\tilde{\chi}}^5}{m_{\tilde{G}}^2} \times \left[1 - \frac{m_{\tilde{G}}^2}{m_{\tilde{\chi}}^2} \right]^3 \left[1 + 3 \frac{m_{\tilde{G}}^2}{m_{\tilde{\chi}}^2} \right], \quad (2.10)$$

where $m_{\tilde{G}}$ is the gravitino LSP mass, $m_{\tilde{\chi}}$ is the neutralino NLSP mass, $M_* = 1/\sqrt{8\pi G_N} \simeq 2.4 \times 10^{18}$ GeV is the reduced Planck scale, θ_W is the weak mixing angle, and N_{ij} denotes the matrix element projecting the i th neutralino into Bino ($j = 1$), Wino ($j = 2$) and Higgsinos ($j = 3, 4$). For such a decay we have

$$B_{\text{EM}} = 1, \quad (2.11)$$

$$\epsilon_{\text{EM}} = \frac{m_{\tilde{\chi}}^2 - m_{\tilde{G}}^2}{2m_{\tilde{\chi}}}. \quad (2.12)$$

We would like to make some clarifications about our numerical calculations.

(1) In the calculation of the thermal relic density of the NLSPs, we considered the general mixing of neutralinos and used the package Microomega [13], which includes all tree level contributions to the scattering amplitudes¹.

(2) In the scan over the mSUGRA parameter space, we used SuSpect2.0 [16] to obtain the sparticle masses in mSUGRA models, which includes one-loop corrections to sparticle masses and two-loop corrections to Higgs masses.

(3) When deriving the constraints on the mSUGRA parameter space, we consider the BBN constraints in (2.2) and (2.3), the WMAP data of relic dark matter density in (1.2) as well as the $b \rightarrow s\gamma$ bound [17]:

$$2.16 \times 10^{-4} < BF(b \rightarrow s\gamma) < 4.34 \times 10^{-4}. \quad (2.13)$$

¹ In the present version of Microomega, the package FeynHiggs [14] is used to calculate the Higgs masses and Hdecay [15] is used to include important QCD corrections to the Higgs decays.




Fig. 1. The region (shaded) allowed by BBN to account for ${}^7\text{Li}$ and ${}^4\text{He}$ abundance in the scenario of gravitino LSP and neutralino NLSP

Furthermore, we fixed the common trilinear coupling $A_0 = 0$ since our results are not sensitive to it.

(4) In our calculation we also considered the muon anomalous magnetic moment α_μ . Since so far much theoretical uncertainty exists in α_μ predictions², we did not use α_μ to set constraints and, instead, we only gave the range of the mSUGRA contributions corresponding to each case. This will be useful when α_μ uncertainty is further reduced in the future.

The bounds on the masses of the neutralino NLSP and gravitino LSP are shown in Fig.1. One sees that BBN results give rather stringent upper and lower bounds:

$$\begin{aligned} 430 \text{ GeV} &\leq M_{\text{LSP}} \leq 600 \text{ GeV}, \\ 630 \text{ GeV} &\leq M_{\text{NLSP}} \leq 840 \text{ GeV}. \end{aligned} \quad (2.14)$$

We found that the neutralino NLSP is quite bino-like ($N_{11} \gtrsim 0.99$) in the allowed regions in Fig. 1

The allowed regions in the plane of m_0 versus $m_{1/2}$ are shown in Fig. 2 for plus sign of μ and Fig. 3 for minus sign of μ . From Fig. 2 we see that for plus sign of μ much of the parameter space with $\tan\beta \lesssim 40$ is ruled out. For example, for $\tan\beta = 10$, only a narrow strip survives, i.e., $225 \text{ GeV} \lesssim m_0 \lesssim 300 \text{ GeV}$ and $1020 \text{ GeV} \lesssim m_{1/2} \lesssim 1270 \text{ GeV}$ (the corresponding contribution to α_μ is $1.38 \lesssim \Delta\alpha_\mu \times 10^{10} \lesssim 2.05$). When $\tan\beta$ increases to 50, the allowed region gets quite large, i.e., $740 \text{ GeV} \lesssim m_0 \lesssim 1730 \text{ GeV}$ and $1060 \text{ GeV} \lesssim m_{1/2} \lesssim 1720 \text{ GeV}$ (the corresponding contribution to α_μ is $2.18 \lesssim \Delta\alpha_\mu \times 10^{10} \lesssim 6.32$).

² For example, one group [18] gives $11.5 \leq \Delta\alpha_\mu \times 10^{10} \leq 60.7$ or $-16.7 \leq \Delta\alpha_\mu \times 10^{10} \leq 49.1$, depending on the calculation approaches. Another group [19] gives $\delta\alpha_\mu \times 10^{10} = 12.4_{-8.3}^{+8.3}$ by using τ decay data to determine the vacuum polarization.




Fig. 2. The regions (shaded) allowed by the WMAP data, BBN constraints as well as $b \rightarrow s\gamma$ data. From left to right corresponds to $\tan\beta = 10, 20, 30, 40, 50$, respectively. The sign of μ is assumed to be plus




Fig. 3. Same as Fig. 2, but for minus sign of μ

For a minus sign of μ , similar results are obtained. We see from Fig. 3 that for $\tan\beta \lesssim 30$, the constraints are quite stringent. For example, for $\tan\beta = 10$ the constraints are $220 \text{ GeV} \lesssim m_0 \lesssim 305 \text{ GeV}$ and $1020 \text{ GeV} \lesssim m_{1/2} \lesssim 1280 \text{ GeV}$ (the corresponding contribution to α_μ is $-6.07 \lesssim \Delta\alpha_\mu \times 10^{10} \lesssim -3.7$). When $\tan\beta$ increases to 40, the constraints are weakened to $1060 \text{ GeV} \lesssim m_0 \lesssim 1530 \text{ GeV}$, $1290 \text{ GeV} \lesssim m_{1/2} \lesssim 1795 \text{ GeV}$ (the corresponding contribution to α_μ is $-5.45 \lesssim \Delta\alpha_\mu \times 10^{10} \lesssim -1.90$).

It is interesting to note that if $\Delta\alpha_\mu$ is required to be positive (negative), then the allowed regions in the $\mu < 0$ ($\mu > 0$) case will be completely ruled out. Therefore, further scrutiny of α_μ will provide meaningful constraints on the parameter space of mSUGRA.

It should be pointed out that the above stringent constraints in Figs. 2 and 3 were obtained under the require-

ment that the gravitinos from the late decays of the NLSP make up all the dark matter. If other particles like axions are also a component of dark matter, then the constraints will be changed. Moreover, if the relic gravitino dark matter are partly produced during reheating³, the constraints will also be changed. In order to forbid gravitino (with mass range in (2.14)) production during reheating, the reheating temperature T_R must be lower than a certain value. Following the analyses in [6], we evaluated such a limit and found $T_R \lesssim 10^9$ GeV.

Note that in our analysis we only considered the consequence of EM energy release from the late decays of WIMPs and required such an EM energy release to settle the discrepancy between the BBN predictions and the observed values for light element abundances. However, as can be seen in the allowed $M_{\text{NLSP}}/M_{\text{LSP}}$ region of Fig. 1, the decay $\text{NLSP} \rightarrow \tilde{G} + Z$ is also possible and thus will cause hadronic energy release from the subsequent Z decays. Such a hadronic energy release can be quite dangerous, since it can alter the BBN predictions for the light element abundances.

In [21] the authors studied the bounds from the hadronic energy release and found that such hadronic bounds may be stringent⁴ due to the overproduction of D . However, as pointed out in [21], for the best fit region with lifetime τ between 10^6 s and 10^7 s, the D overproduction from hadronic energy is possibly canceled by the overdestruction from EM energy and thus the hadronic bounds become less stringent and subject to a large uncertainty. In such a region the EM energy effects are dominant [21]. Since in our study we focused on the best fit region, we did not consider the hadronic effects.

3 Conclusions

We examined the constraints on the newly proposed dark matter scenario, in which the gravitino is assumed to be the LSP and produced from the late decays of metastable NLSP. Although such gravitino dark matter can naturally evade the current detection experiments due to its super-weak couplings, we found, however, that this scenario is subjected to stringent constraints from the BBN predictions for light element abundances as well as the WMAP data for the relic density. Assuming the popular case that the lightest neutralino is the NLSP in mSUGRA models, we found that the low $\tan\beta$ ($\lesssim 40$) region as well as the region accessible at the LHC are severely constrained.

The popular mSUGRA models will be explored in future colliders like the LHC. For this purpose, it is important to know which part of the parameter space is viable and

thus should be primarily explored. In this regard, the stringent cosmological constraints on the mSUGRA parameter space obtained in this work will be useful. Especially, if the ongoing and planned dark matter detection experiments fail to find any dark matter signal, it will imply that the dark matter interactions are too weak and thus the gravitino dark matter scenario will be favored. Then, to test this scenario at colliders, the stringent cosmological constraints on the parameter space will be quite instructive. This would serve as a good example that the studies in cosmology and astrophysics can shed some light on particle collider physics.

On the other hand, the LHC could explore mSUGRA parameter space up to $m_{1/2} \sim 1400$ GeV (700 GeV) for small (large) values of m_0 , assuming 100 fb^{-1} of integrated luminosity [20]. If the LHC results finally restrain the parameter space to one of the regions obtained in this work, then it implies that the gravitino dark matter scenario is favored. In this sense, the studies in particle physics can provide some insights in the understanding of dark matter in cosmology.

We note that our study in this work is just illustrative instead of exhaustive. We assumed the popular case that the lightest neutralino is the NLSP in mSUGRA models. Actually, other superparticles, like the tau-slepton, are also likely to be the NLSP in mSUGRA models. If the tau-slepton is assumed to be the NLSP, there are some theoretical uncertainties in its decay modes and the corresponding energy release.

Note added. While we are preparing this manuscript, some other preprints [21] appeared, where the constraints on the gravitino dark matter scenario are studied. We found that the studies in [21] are quite exhaustively, where the scenarios of neutralino NLSP, stau NLSP as well as sneutrino NLSP are all considered. Compared with the studies in [21], the characteristic of our study is that we performed a scan over the mSUGRA parameter space and presented the allowed regions in terms of original mSUGRA parameters. In addition, the BBN constraints on the EM energy release are more stringent in our study since we required such an energy release to settle the discrepancy between the BBN prediction and the WMAP data for the ${}^7\text{Li}$ abundance.

Acknowledgements. We thank Shufang Su for useful comments and Yuqi Lee for debugging the linking code between Micromega and SuSpect. We also thank Junjie Cao and Guangping Gao for helpful discussions. This work is supported in part by National Natural Science Foundation of China (NNSFC) and by a visiting program (No. L03517) of Japan Society for the Promotion of Science.

References

1. For a recent review, see, e.g., C. Munoz, hep-ph/0309346
2. S. Weinberg, Phys. Rev. Lett. **48**, 1303 (1982); E. Holtmann, M. Kawasaki, K. Kohri, T. Moroi, Phys. Rev. D **60**, 023506 (1999); M. Kawasaki, K. Kohri, T. Moroi, Phys. Rev. D **63**, 103502 (2001)

³ The gravitinos can also be produced as a thermal relic at very early times. However, in the context of inflation, the universe inflated between that early time and now, which would dilute any gravitino thermal relic density.

⁴ In their study they used the bounds $2.4 \times 10^{-5} < D/H < 3.2 \times 10^{-5}$, which are more aggressive than the bounds of (2.5) used by us to obtain the best fit region.

3. D.S. Akerib et al., astro-ph/0405033
4. A.B. Lahanas, D.V. Nanopoulos, Phys. Lett. B **568**, 55 (2003)
5. H. Pagels, J.R. Primack, Phys. Rev. Lett. **48**, 223 (1982)
6. M. Bolz, A. Brandenburg, W. Buchmuller, Nucl. Phys. B **606**, 518 (2001); T. Moroi, H. Murayama, M. Yamaguchi, Phys. Lett. B **303**, 289 (1993)
7. J.L. Feng, A. Rajaraman, F. Takayama, Phys. Rev. Lett. **91**, 011302 (2003); Phys. Rev. D **68**, 063504 (2003); J.L. Feng, hep-ph/0308201
8. WMAP Collaboration, Astrophys. J. Suppl. **148**, 1 (2003); **148**, 175 (2003); **248**, 195 (2003)
9. R.H. Cyburt, J. Ellis, B.D. Fields, K.A. Olive, Phys. Rev. D **67**, 103521 (2003)
10. M.H. Reno, D. Seckel, Phys. Rev. D **37**, 3441 (1988); K. Kohri, Phys. Rev. D **64**, 043515 (2001); S. Dimopoulos, R. Esmailzadeh, L.J. Hall, G.D. Starkman, Nucl. Phys. B **311**, 699 (1989); M. Kawasaki, K. Kohri, T. Moroi, astro-ph/0402490
11. B. Fields, Primordial Nucleosynthesis in light of WMAP, CFA colloquium, University of Illinois
12. W. Buchmuller, K. Hamaguchi, M. Ratz, T. Yanagida, hep-ph/0403203
13. G. Belanger, F. Boudjema, A. Pukhov, A. Semenov, Comput. Phys. Commun. **149**, 103 (2002)
14. M. Frank, S. Heinemeyer, W. Hollik, G. Weiglein, hep-ph/0212037
15. A. Djouadi, J. Kalinowski, M. Spira, Comput. Phys. Commun. **108**, 56 (1998)
16. A. Djouadi, J.-L. Kneur, G. Moultaka, hep-ph/0211331
17. H. Baer, C. Balazs, J. Ferrandis, X. Tata, Phys. Rev. D **64**, 035004 (2001)
18. K. Hagiwara, A.D. Martin, D. Nomura, T. Teuber, Phys. Lett. B **557**, 69 (2003)
19. M. Davier, S. Eidelman, A. Hocker, Z. Zhang, Eur. Phys. J. C **27**, 497 (2003)
20. H. Baer, A. Belyaev, T. Krupovnickas, A. Mustafayev, hep-ph/0403214
21. J.L. Feng, S. Su, F. Takayama, hep-ph/0404198; hep-ph/0404231