

Is constituent quark scaling a unique sign of parton recombination?

M. Bleicher^{1,a}, X. Zhu^{2,3}

¹ Institut für Theoretische Physik, J.W. Goethe Universität, Max von Laue Strasse 1, 60438 Frankfurt am Main, Germany

² Frankfurt Institute for Advanced Studies (FIAS), Max von Laue Strasse 1, 60438 Frankfurt am Main, Germany

³ Physics Department, Tsinghua University, Beijing 100084, China

Received: 1 August 2006 /

Published online: 2 December 2006 – © Springer-Verlag / Società Italiana di Fisica 2006

Abstract. We study the transverse momentum dependence of the anisotropic flow v_2 for π , K , nucleon, Λ , Ξ and Ω for Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV within two independent string-hadron transport approaches (RQMD and UrQMD). Although both models reach only 60% of the absolute magnitude of the measured v_2 , they both predict the particle-type dependence of v_2 as observed by the RHIC experiments: v_2 exhibits a hadron-mass hierarchy in the low p_T region and a number-of-constituent-quark (NCQ) dependence in the intermediate p_T region. The failure of the hadronic models to reproduce the absolute magnitude of the observed v_2 indicates that transport calculations of heavy-ion collisions at RHIC must incorporate interactions among quarks and gluons in the early, hot and dense phase. The presence of an NCQ scaling in the string-hadron model results suggests that the particle-type dependences observed in heavy-ion collisions at intermediate p_T might be related to the hadronic cross sections in vacuum rather than to the hadronization process itself.

1 Motivation

To create extremely hot and dense matter with partons as its fundamental components – called the quark–gluon plasma (QGP) – is a major goal of current and future high-energy heavy-ion collisions experiments at SPS, RHIC and LHC [2]. However, due to the complex nature of the relativistic nucleus–nucleus reactions, the QGP, if it has been created, escapes direct detection. Therefore, in order to distinguish the existence and later on to investigate the properties of the new state of matter, one must find observables which allow one to deduce the properties of the intermediate (QGP) state from the final state hadrons. To unravel such observables for this new form of matter has been a focal point of theoretical and experimental studies over the last decade [3]. Measurements of the collective motion of hadrons produced in high-energy nuclear collisions have long been suggested as a valuable tool to gain information about the nature of the constituents and the equation of state of the system in the early stage of the reaction [4–13]. Specifically, strange and multi-strange hadron elliptic and radial flow results seem to indicate that the observed collectivity originates from a partonic phase¹ [14]. Furthermore, elliptic and radial flow measurements for heavy-flavor hadrons like J/Ψ and D mesons

will test the hypothesis of early thermalization in these collisions [15–17].

At RHIC, measurements [18, 19] of the elliptic flow v_2 and the nuclear modification factor $R_{AA,CP}$ for identified particles have led to the conclusion that hadrons ought to be formed via the coalescence or recombination of massive quarks [20–28]. A cornerstone of this conclusion is the observed NCQ scaling of the flow of baryons versus mesons. Because this interpretation addresses key issues in high-energy nuclear collisions such as deconfinement and chiral symmetry restoration, it is of utmost importance to conduct a systematic study of other possible explanations for the observed particle-type dependences.

For the present analysis we employ two independent hadron–string transport models, RQMD (v2.4) and UrQMD (v2.2) [29–32], to study the effect of hadronic cross sections, kinematics etc. on the particle-type dependence of v_2 . We present the v_2 values of π , K , p , Λ , Ξ and Ω for Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV. We find that – although the hadronic models give only 60% of the magnitude of v_2 – both models reproduce the gross features of the particle-type dependence, including the mass ordering at $p_T < 1$ GeV/ c and the NCQ dependence at $p_T > 1$ GeV/ c . Within both models, the NCQ dependence is related to the additive-quark model for hadronic cross sections. These findings imply that detailed comparisons between data and theory are necessary to disentangle hadronic and partonic effects at intermediate transverse momenta, 1.5 GeV/ $c < p_T < 5$ GeV/ c . For a transport model discus-

^a e-mail: bleicher@th.physik.uni-frankfurt.de

¹ The term partonic is used to denote any kind of deconfined matter made of quarks and gluons.

sion of the quantitative accuracy and validity of the various methods to extract elliptic flow values from the data, the reader is referred to [9–12].

As one can observe from Fig. 1, the build-up of the elliptic flow in these models is tightly connected to the hadron–hadron scattering rate (top/bottom figures). Initial valence (di-) quark scatterings before 1 fm/c do only yield a negligible amount of v_2 . Figure 2 shows the time evolution of the elliptic flow for mesons and baryons for low p_T (bottom) and high p_T (top). Already at early times, a separation between meson and baryon elliptic flow is visible, leading to the mass hierarchy at low transverse momenta and to the stronger elliptic flow of baryons (compared to mesons) at high p_T .




Fig. 1. UrQMD calculation for Au + Au interactions at $\sqrt{s_{NN}} = 200$ GeV. Top: meson and baryon v_2 versus time at midrapidity. Bottom: collision rates as a function of time for hadron+hadron (dashed line), hadron+quark/di-quark (full line) and quark/di-quark+quark/di-quark (dotted line)




Fig. 2. UrQMD calculation for Au + Au interactions at $\sqrt{s_{NN}} = 200$ GeV. Bottom: meson and baryon v_2 versus time at midrapidity and $p_T \leq 1$ GeV/c. Top: meson and baryon v_2 versus time at midrapidity and $p_T \geq 2$ GeV/c

2 Model results

Within the framework of the hadronic transport approach, a typical heavy-ion collision proceeds schematically in three stages, i.e. the pre-hadronic (strings and constituent (di-) quarks) stage, the hadronic pre-equilibrium stage, the evolution towards hadronic kinetic equilibrium and freeze-out. The pre-hadronic stage involves the initial excitation and fragmentation of color strings and ropes. At the highest RHIC energy, this stage lasts for about 0.5–1.5 fm/c and the effective transverse pressure/EOS is rather soft. During the late hadronic stage, the hadronic system approaches local kinetic equilibrium, followed by an approach to free streaming, where the system escapes equilibrium due to dilution of the hadronic gas: the mean free path

of the hadrons exceeds the finite size of the system [29–33], the free streaming hadrons decay and feed down to the lightest species.

Figure 3 presents the model results on the centrality dependence of the charged hadron v_2 values along with measurements from the STAR collaboration [34, 35]. Both hadronic transport models (UrQMD v2.2 and RQMD v2.4) reach about 60% of the measured v_2 values only, although the centrality dependences are very similar to the data. There is a small variance between the two models, which we consider as an estimate of the systematic errors in such model calculations. Although the v_2 values from the hadronic transport model also depend on the formation time of hadrons from strings [9, 10], the failure of both hadronic transport models to describe quantitatively the magnitude of v_2 is a strong indication that there are interactions amongst pre-hadronic constituents (partons) present in nature (but not in the hadron–string models discussed here), which are responsible for the large v_2 values observed in the experiments [36–38]. When rescattering between the hadrons is turned off (full circles), v_2 vanishes completely, because repulsive vector interactions are not included into the present simulations [4, 5, 9, 10].

Let us now analyze the temporal structure of the elliptic flow’s development. Figure 4 shows the v_2 values as a function of the freeze-out time for pions in minimum biased Au + Au interactions at $\sqrt{s_{NN}} = 200$ GeV. The different symbols denote different transverse momentum intervals decreasing from top to bottom. One clearly observes a strong correlation between freeze-out time and elliptic flow: particles that decouple earlier have a larger v_2 value than those that freeze out later. The lower p_T particles tend to freeze out later, and their v_2 continues to evolve late in the evolution (a similar correlation has been observed earlier in low-energy collisions [39]) The decrease in the v_2 towards later times is related to the reduction of the coordinate-space anisotropy with time – i.e. the system




Fig. 3. Charged hadron v_2 versus the number of participating nucleons in Au + Au interactions at $\sqrt{s_{NN}} = 200$ GeV. Experimental data [34, 35] from the 4-particle cumulant method is shown as stars. RQMD results are depicted by *filled circles* (full calculation) and *open circles* (without rescattering). The UrQMD calculations are shown as *open triangles*




Fig. 4. Elliptic flow (v_2) of pions from RQMD versus their freeze-out time for several p_T windows for minimum bias Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV. The *dashed-line* represents the p_T integrated distribution

has become more spherical than it was at earlier times. The v_2 values of pions with higher p_T saturate sooner and tend to reflect the earlier stage of the collision more strongly. Thus, within the model dynamics, the final v_2 is mostly driven by the early stages of the reaction; the v_2 values at high p_T are closer to the initial/early v_2 values than the v_2 values at lower p_T .

3 Particle-type dependence

How much of the observed NCQ scaling features can be reproduced by the hadronic models? In both dynamical approaches, finite (vacuum) cross sections are used to model the strong interactions in the hadron–string cascade. Unlike the simplistic Cooper–Frye freeze-out treatment, in most hydrodynamic calculations the transition from strongly interacting matter to free streaming is determined here by the interplay of the local particle density and the energy dependent cross section of the individual hadrons. It is well known that a proper treatment of the gradual freeze-out is crucial for the finally observed hadron distributions. It was pointed out that the hydrodynamical results on flow depend strongly on the proper kinetic treatment of the freeze-out process and cannot be approximated by isotherms [40–42].

However, the major shortcoming of the present hadron–string approach is the lack of the early partonic interactions which are important for the early dynamics in ultra-relativistic heavy-ion collisions [3–5, 9, 10, 43]. In order to take care of both partonic and hadronic interactions in high-energy nuclear collisions, a combination of the hydrodynamic model for the early stage dynamics (the “per-

fect” fluid stage) with a hadronic transport model for the later stage plus freeze-out has been proposed [44–49]. Figure 5 shows the collision centrality dependence of the p_T dependent v_2 values for π , K , p , and Λ . Both the hydrodynamic behavior (in the low p_T region) and a hadron-type dependence (in the intermediate- p_T region) are clearly predicted in all centrality bins. This “crossing and subsequent splitting” between meson and baryon elliptic flow as well as the breakdown of the hydrodynamical mass scaling at high transverse momenta was first predicted within the UrQMD model [9, 10] and has later been observed in the experimental data. It is important to note that the more recent explanations of this effect (the suggested “number-of-constituent-quark” scaling) is not a unique feature of the “quark recombination/coalescence” assumption: hadronic interactions alone have quantitatively (at the correct p_T values) predicted this hadron-type dependence.

Let us explore the p_T dependence of the event anisotropy parameters in detail. Figure 6 shows the calculated unscaled and scaled v_2 values of various hadrons versus the unscaled and scaled transverse momenta, p_T , of the various hadrons. On the left hand side (Fig. 6a) one can see that at lower transverse momenta, $p_T \leq 1.5$ GeV/c, the heavier hadrons exhibit smaller v_2 values than the lighter hadrons: hadron transport theory predicts mass ordering. The Ξ and Ω v_2 values from the UrQMD calculations are also included. They are the lowest of all v_2 values for $p_T \leq 2$ GeV/c. Such a mass ordering is exactly what is observed in the experimental data [18] and is in accord with hydrodynamic calculations [50]. Hence, hadronic interactions, which do take place at later stages of the collisions, also do contribute to the observed collective motion.




Fig. 5. RQMD results of π , K , p , and Λ v_2 from Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV. (a) Minimum bias collisions: at about $p_T \sim 1.2$ GeV/c, baryon and meson v_2 are crossing each other; (b) 40%–80%; (c) 20%–40%; (d) 0%–20%




Fig. 6. (a) Hadron v_2 from minimum bias Au + Au collisions at $\sqrt{s_{NN}} = 200$ GeV; (b) scaled hadron v_2 results are shown. The n_q refers to the number of constituent quarks. Symbols represent results from the UrQMD (v2.2) model for various hadron species. At low $p_T/n_q \leq 0.5$ GeV/c, π does not follow the scaling perhaps caused by the resonance decay [17]. In the higher p_T region, K meson seems to fall off the scaling curve due to the comparatively small hadronic cross sections in the model

At higher p_T values, this mass dependence gives way to the $v_2(p_T)$ dependence on the hadron type (i.e. meson or baryon). Here, it is interesting to note that the Ω -baryons seemingly acquire a significant amount of v_2 in the model calculations. In addition, there is also a clear but small difference for kaons and pions in v_2 values at $p_T \geq 1.5$ GeV/c. This particle-type dependence, rather than the otherwise dominating particle mass dependence, is also observed in the data [18]. It is important to note that the ϕ -meson has a mass that is very close to the mass of the baryons p and Λ , and, indeed, recent experimental results on the ϕ 's v_2 values are similar to other mesons [51]. However, in the hadronic transport model, about 2/3 of the ϕ -mesons are formed via $K\bar{K}$ coalescence, which is not necessarily the dominant process in heavy-ion collisions [52]. Therefore, the v_2 values of the ϕ -meson are not shown in Fig. 6. It should also be noted that in the high p_T region, $p_T \geq 2.5$ GeV/c, all v_2 values start to decrease. This indicates that the system is deviating from an ideal hydrodynamic behavior. This trend is best seen in the right, “scaled”, plot in Fig. 6. Such a drop has been observed in the data.

The test of the NCQ scaling hypothesis is shown in Fig. 6b, which depicts the scaled hadron values, v_2/n_q . The scaling factor is NCQ, in accord with the coalescence approach [26–28]. For mesons and baryons, $n_q = 2$ and $n_q = 3$, respectively. The NCQ scaling is clearly observed in both RQMD (not shown here) and UrQMD model calculations except for the pions. This surprising result and its implications for the frequently invoked recombination/coalescence hypothesis will be discussed in the last section.

However, one should note that there are experimentally distinguishable differences (when differential Ω elliptic flow data with good statistics become available) between “real” NCQ scaling from coalescence and the approximate NCQ scaling due to the cross section hi-

erarchy discussed here. In the coalescence approach one expects identical elliptic flow of all baryons irrespective of their type. In contrast, the present AQM scaling picture leads to an observable ordering of the elliptic flow at fixed p_T according to the strangeness content. This can indeed be observed in Fig. 6, where an ordering of the elliptic flow values can be observed: $v_2(N) > v_2(\Lambda) > v_2(\Xi) > v_2(\Omega)$.

4 Discussion and conclusions

The particle and energy density is highest at the center of the created fireball in relativistic nuclear collisions – initially, there is an angular dependent matter density gradient. The repulsive interactions among the constituents will therefore push matter to move outwards. In this way, the collective flow develops in nucleus–nucleus collisions [3–8]. We would like to stress that flow means matter and energy flow. It is independent of the type of particles, either partons or hadrons, or different kinds of hadrons. Hence, by studying the collective motion of the produced hadrons one can, in principle, extract the information of early collision dynamics [6, 7, 57]. In general, one expects that the final elliptic flow,

$$v_2(p_T) \propto \int_t \int_{\Sigma} \sigma(\rho, p_T) \otimes \rho_{\Sigma}(t, x, y, p_T) dA_{\Sigma}(x, y) dt, \quad (1)$$

where Σ denotes the hyper-surface where hadrons are emitted, will depend on σ , i.e. the interaction cross section, which, in principle, depends on the particle type, c.m. angle and relative momenta. The specific particle density depends on the collision time t , location, and momentum. For short mean free paths, the transverse flow is intimately related to the pressure, which in turn depends on the density and temperature of the matter under study [3–10]. Indeed, the frequent rescatterings among the hadrons can lead to hydrodynamic-like mass ordering in the low p_T region.

At the higher transverse momenta, $p_T \geq 1.5 \text{ GeV}/c$, the particles escape quickly from the system to low density, in effect leading to early freeze-out and lack of development of the hydrodynamics, and the details of the interaction cross sections are most important. As the cross sections depend on the particle type, for mesons or baryons to first approximation given by the constituent quark model [26], we do expect roughly a 2 : 3 scaling of the meson-to-baryon elliptic flow from transport calculations.

The hadronic models underpredicted the strength of v_2 at RHIC, because early partonic interactions (except from quark coalescence during the string break-up) are not included in the model. The early stage with highest density and smallest mean free paths is “missing”. This shortcoming of the hadronic models clearly demonstrates the need for the early, dense partonic interactions in heavy-ion collisions at RHIC.

5 Summary

In summary, the hadronic transport models UrQMD and RQMD have been employed to study the elliptic flow of hadrons in Au + Au collisions at the highest RHIC energy. We have analyzed the v_2 values as a function of collision centrality, transverse momenta and collision time for various meson and baryon species, including multi-strange baryons. Both hadron–string transport models fail by 40% to exhaust the absolute value of v_2 , probably due to the lack of (partonic) interactions in the initial, hot and dense stage. However, because of the hadronic re-interactions, the hadron-mass hierarchy is qualitatively well reproduced in the low p_T region. Rescattering is the key that leads to the quasi-ideal hydrodynamic appearance in $v_2(p_T)$. Also in the intermediate p_T region, the hadron-type dependence (NCQ scaling) is predicted by both hadronic transport models. Here, this dependence is due to the hadronic cross sections which do roughly scale with the number of constituent quarks, in accord with the additive-quark model. This finding challenges the recent interpretation of NCQ scaling as a unique deconfinement signature. Thus, further tests of the deconfinement-plus-recombination hypothesis are necessary with high precision v_2 measurements of resonance hadrons like K^* , ρ , Δ , Λ^* , and Ξ^* .

Acknowledgements. This work is based on [1].

The authors would like to thank Y. Lu, H. Stöcker, N. Xu, P. Sorensen, H. Petersen, C.M. Ko, A. Poskanzer, H.G. Ritter, and K. Schweda for enlightening discussions. This work was supported by BMBF and GSI, Germany.

References

1. Y. Lu et al., J. Phys. G **32**, 1121 (2006) [arXiv:nucl-th/0602009]
2. See Proceedings of Quark Matter 2005, Budapest, Hungary, 2005
3. S.A. Bass, M. Gyulassy, H. Stöcker, W. Greiner, J. Phys. G **25**, R1 (1999) [arXiv:hep-ph/9810281]
4. H. Stöcker, M. Gyulassy, J. Boguta, Phys. Lett. B **103**, 269 (1981)
5. H. Stöcker, W. Greiner, Phys. Rep. **137**, 277 (1986)
6. H. Sorge, Phys. Rev. Lett. **78**, 2309 (1997)
7. J.-Y. Ollitrault, Phys. Rev. D **46**, 229 (1992)
8. W. Reisdorf, H.G. Ritter, Ann. Rev. Nucl. Part. Sci. **47**, 663 (1997)
9. M. Bleicher, H. Stöcker, Phys. Lett. B **526**, 309 (2002) [arXiv:hep-ph/0006147]
10. H. Stöcker, Nucl. Phys. A **750**, 121 (2005) [arXiv:nucl-th/0406018]
11. X. Zhu, M. Bleicher, H. Stöcker, Phys. Rev. C **72**, 064911 (2005) [arXiv:nucl-th/0509081]
12. X. Zhu, M. Bleicher, H. Stöcker, arXiv:nucl-th/0601049
13. S. Mrowczynski, E.V. Shuryak, Acta Phys. Pol. B **34**, 4241 (2003) [arXiv:nucl-th/0208052]
14. N. Xu, Nucl. Phys. A **751**, 109 (2005)
15. B. Svetitsky, A. Uziel, Phys. Rev. D **55**, 2616 (1997) [arXiv:hep-ph/9606284]
16. B. Svetitsky, A. Uziel, arXiv:hep-ph/9709228

17. X. Dong et al., Phys. Lett. B **597**, 328 (2004)
18. STAR Collaboration, J. Adams et al., Phys. Rev. Lett. **92**, 052302 (2004)
19. PHENIX Collaboration, S.S. Adler et al., Phys. Rev. Lett. **91**, 182301 (2003)
20. S. Scherer et al., Prog. Part. Nucl. Phys. **42**, 279 (1999)
21. S.A. Bass et al., Prog. Part. Nucl. Phys. **42**, 313 (1999) [arXiv:nucl-th/9810077]
22. M. Hofmann, M. Bleicher, S. Scherer, L. Neise, H. Stöcker, W. Greiner, Phys. Lett. B **478**, 161 (2000) [arXiv:nucl-th/9908030]
23. M. Hofmann, J.M. Eisenberg, S. Scherer, M. Bleicher, L. Neise, H. Stöcker, W. Greiner, arXiv:nucl-th/9908031
24. S. Scherer, M. Hofmann, M. Bleicher, L. Neise, H. Stöcker, W. Greiner, New J. Phys. **3**, 8 (2001) [arXiv:nucl-th/0106036]
25. S. Scherer, H. Stöcker, arXiv:nucl-th/0502069
26. D. Molnar, S. Voloshin, Phys. Rev. Lett. **91**, 092301 (2003)
27. R.J. Fries, J. Phys. G **31**, 379 (2005), nucl-th/0410085 and references therein
28. R. Hwa, C.B. Yang, Phys. Rev. C **70**, 024904 (2004) and reference therein
29. H. Sorge, Phys. Rev. C **52**, 3291 (1995)
30. M. Bleicher et al., J. Phys. G **25**, 1859 (1999) [arXiv:hep-ph/9909407]
31. S.A. Bass et al., Prog. Part. Nucl. Phys. **41**, 225 (1998)
32. M. Bleicher et al., in preparation (2006)
33. H. van Hecke, H. Sorge, N. Xu, Phys. Rev. Lett. **81**, 5764 (1998)
34. STAR Collaboration, K.H. Ackermann et al., Phys. Rev. Lett. **86**, 402 (2001)
35. STAR Collaboration, J. Adams et al., Phys. Rev. C **72**, 014904 (2005)
36. G. Baur, J. Bleibel, C. Fuchs, A. Faessler, L.V. Bravina, E.E. Zabrodin, Phys. Rev. C **71**, 054905 (2005) [arXiv:nucl-th/0411117]
37. Z. Xu, C. Greiner, in Proc. Quark Matter 2005, August 4–9, 2005, Budapest [arXiv:hep-ph/0509324]
38. E.L. Bratkovskaya et al., Phys. Rev. C **69**, 054907 (2004) [arXiv:nucl-th/0402026]
39. S.A. Bass, C. Hartnack, H. Stoecker, W. Greiner, Phys. Rev. C **50**, 2167 (1994)
40. C.M. Hung, E.V. Shuryak, Phys. Rev. C **57**, 1891 (1998) [arXiv:hep-ph/9709264]
41. C. Anderlik et al., Phys. Rev. C **59**, 3309 (1999) [arXiv:nucl-th/9806004]
42. V.K. Magas et al., Heavy Ion Phys. **9**, 193 (1999) [arXiv:nucl-th/9903045]
43. T. Hirano, M. Gyulassy, nucl-th/0506049
44. S.A. Bass, A. Dumitru, M. Bleicher, L. Bravina, E. Zabrodin, H. Stöcker, W. Greiner, Phys. Rev. C **60**, 021902 (1999) [arXiv:nucl-th/9902062]
45. A. Dumitru, S.A. Bass, M. Bleicher, H. Stöcker, W. Greiner, Phys. Lett. B **460**, 411 (1999) [arXiv:nucl-th/9901046]
46. S.A. Bass, A. Dumitru, Phys. Rev. C **61**, 064909 (2000) [arXiv:nucl-th/0001033]
47. D. Teaney, J. Lauret, E. Shuryak, nucl-th/0110037
48. C. Nonaka, S.A. Bass, arXiv:nucl-th/0510038
49. T. Hirano, Proc. Quark Matter 2005, August 4–9, 2005, Budapest [arXiv:nucl-th/0510005]
50. P. Huovinen, private communications (2003)
51. STAR Collaboration, M. Oldenburg et al., Proc. Quark Matter 2005, August 4–9, 2005, Budapest [arXiv:nucl-ex/0510026]
52. STAR Collaboration, J. Adam et al., Phys. Lett. B **612**, 181 (2005)
53. T.S. Biro, H.B. Nielsen, J. Knoll, Nucl. Phys. B **245**, 449 (1984)
54. H.J. Möring, J. Ranft, C. Merino, C. Pajares, Phys. Rev. D **47**, 4142 (1993)
55. H. Sorge, Phys. Rev. C **52**, 3291 (1995) [arXiv:nucl-th/9509007]
56. H. Sorge, M. Berenguer, H. Stöcker, W. Greiner, Proceedings of Hot and Dense Nuclear Matter, 621–633, 1993, Bodrum
57. N. Xu, Prog. Part. Nucl. Phys. **53**, 165 (2004)