

Form factors in the radiative pion decay

V. Mateu, J. Portolés^a

IFIC, Universitat de València – CSIC, Apt. Correus 22085, 46071 València, Spain

Received: 13 June 2007 /

Published online: 5 September 2007 – © Springer-Verlag / Società Italiana di Fisica 2007

Abstract. We perform an analysis of the form factors that rule the structure-dependent amplitude in radiative pion decay. The resonance contributions to $\pi \rightarrow e\nu_e\gamma$ decays are computed through the proper construction of the vector and axial-vector form factors by setting the QCD driven asymptotic properties of the three-point Green functions $\langle VVP \rangle$ and $\langle VAP \rangle$, and by demanding the smoothening of the form factors at high transfer of momentum. A comparison between theoretical and experimental determination of the form factors is also carried out. We also consider and evaluate the role played by a non-standard tensor form factor. We conclude that, at present and due to the hadronic uncertainties, the search for new physics in this process is not feasible.

PACS. 11.15.Pg; 12.38.-t; 12.39.Fe

1 Introduction

The radiative decay of the pion is a suitable process to be analysed within chiral perturbation theory (χ PT) [1–4], the effective field theory of QCD in the very low-energy region. This framework provides the structure of the relevant form factors through i) a polynomial expansion in momenta, essentially driven by the contributions of heavier degrees of freedom that have been integrated out, and ii) the required chiral logarithms generated by the loop expansion and compelled by unitarity. Both contributions correspond to the chiral expansion in p^2/M_V^2 and p^2/Λ_χ^2 , respectively, where M_V is the mass of the lightest vector resonance, $\Lambda_\chi \sim 4\pi F$ and F is the decay constant of the pion. Hence their magnitude is, in principle, comparable. The chiral logarithms have thoroughly been studied in later years up to $\mathcal{O}(p^6)$ both in SU(2) [5] and SU(3) [6, 7].

However, the size of the polynomial contributions is more controversial. They involve short-distance dynamics through the chiral low-energy constants (LECs) of the χ PT Lagrangian and their determination from QCD is a difficult non-perturbative problem. Phenomenology and theoretical arguments suggest that the main role is played by the physics at the scale M_V , i.e. the physics of low-lying resonances, and a lot of effort has been dedicated to pursue this goal [8, 9]. This assumption, widely known as resonance saturation of the LECs in χ PT, implies that the structure of the form factors is given by the pole dynamics of resonances and this hint works well in all known cases in the allowed range of energies as, e.g. scalar, vector and axial-vector form factors in hadronic decays of the tau lepton [10–14].

Because $M_\pi \ll M_V$, it is expected that the structure of the form factors in pion decays should be less relevant and, accordingly, the approach provided by χ PT should be good enough even when only the first terms of the chiral expansion are included. This is the case of the radiative pion decay, namely $\pi \rightarrow \ell\nu_\ell\gamma$, $\ell = \mu, e$, where constant form factors (which correspond to the leading order contribution in the chiral expansion) have widely been employed in the analyses of data. However, the PIBETA collaboration [15] showed that a strong discrepancy between theory and experiment arises for the branching ratio of the process in a specific region of the electron and photon energies. Lately the same collaboration, after their 2004 analysis, concludes that the discrepancy has faded away [16]. Curiously enough this decay has a persistent story of deceptive comparisons between theory and experiment [17], which have prompted the publication of proposals beyond the standard model (SM) to account for the variance [18–25]. Of these the possibility of allowing a tensor contribution has received particular attention, which could explain the discrepancy by interfering destructively with the standard model prescription though showing some inconsistency with the corresponding tensor contribution in nuclear β decay [26]. Related with this issue it is essential, seeking to discern the presence of a new physics contribution to the radiative decay of the pion, to provide an accurate profile of the involved form factors within QCD.

In order to settle the standard model description of the vector and axial-vector form factors participating in $\pi \rightarrow \ell\nu_\ell\gamma$ decays we study, in this article, the structure provided by the lightest meson resonances. This is very much relevant on the experimental side because high-statistics experiments as PIBETA [16] already are able to deter-

^a e-mail: Jorge.Portoles@ific.uv.es

mine, for instance, the slopes of the form factors involved in these decays. The procedure consists, essentially, in the construction of the relevant three-point Green functions (GF) in the resonance energy region with a finite spectrum of states at the poles of the corresponding meromorphic functions. This can be carried out by employing a parametric *ansatz* or a Lagrangian theory (we work it out in both cases). In a second stage several constraints are imposed on the parameters or coupling constants, namely chiral symmetry at $q^2 \ll M_R^2$ and the asymptotic behavior ruled by the operator product expansion of the GF for $q^2 \gg M_R^2$, where M_R is short for the typical masses of the resonances in the poles and q^2 an indicative squared transfer of momenta of this energy region. This method, guided by the large- N_C limit of QCD [27, 28] (N_C is short for the number of colors), has proven to be efficient [29–41] in order to collect the constraints that drive form factors of QCD currents.

Section 2 is devoted to the construction of the vector and axial-vector form factors using the procedure outlined above. The study of the beyond the SM contribution of a tensor current to the radiative pion decay is performed in Sect. 3 and, finally, Sect. 4 describes the analysis of the photon spectrum in this process. After the Conclusions in Sect. 5, three appendices complete the setting of this article.

2 Radiative pion decay: vector and axial-vector form-factors

The amplitude that describes the $\pi^+ \rightarrow \ell^+ \nu_\ell \gamma$ process can be split into two different contributions:

$$M(\pi^+ \rightarrow \ell^+ \nu_\ell \gamma) = M_{\text{IB}} + M_{\text{SD}}. \quad (1)$$

Here M_{IB} is the inner bremsstrahlung (IB) amplitude, where the photon is radiated by the electrically charged external legs, either pion or lepton; consequently the interaction is driven by the axial-vector current. M_{SD} is the structure-dependent (SD) contribution, where the photon is emitted from intermediate states generated by strong interactions. In the latter case both vector and axial-vector form factors arise from the hadronization of the QCD currents within the standard model.

Because $\pi^+ \rightarrow e^+ \nu_e$ is helicity suppressed, the IB contribution to its radiative counterpart suffers the same inhibition and, consequently, the electron case is the appropriate channel to uncover the non-perturbative SD amplitude. On the contrary, the $\pi^+ \rightarrow \mu^+ \nu_\mu \gamma$ decay is fairly dominated by IB. As a consequence the $\pi^+ \rightarrow e^+ \nu_e \gamma$ is of great interest to investigate the hadronization of the currents contributing to the SD amplitudes that are driven, within the standard model, by the vector ($F_V(q^2)$) and axial-vector ($F_A(q^2)$) form factors defined by¹

$$\langle \gamma | \bar{u} \gamma_\alpha d | \pi^- \rangle = -\frac{e}{M_{\pi^+}} \varepsilon^{\beta*} F_V(q^2) \varepsilon_{\alpha\beta\mu\nu} r^\mu p^\nu,$$

¹ We use the convention $\varepsilon_{0123} = +1$ for the Levi-Civita tensor $\varepsilon_{\mu\nu\alpha\beta}$ throughout this paper.

$$\langle \gamma | \bar{u} \gamma_\alpha \gamma^5 d | \pi^- \rangle = i \frac{e}{M_{\pi^+}} \varepsilon^{\beta*} F_A(q^2) [(r \cdot p) g_{\alpha\beta} - p_\alpha r_\beta] + i e \varepsilon_\alpha^* \sqrt{2} F, \quad (2)$$

where r and p are the pion and photon momenta, respectively, $q^2 = (r - p)^2$ and e is the electric charge of the electron. The second term in the matrix element of the axial-vector current corresponds to the pion pole contribution (which coupling is given by the decay constant of the pion F) to the IB amplitude.

For the full set of expressions for the differential decay rate of $\pi \rightarrow \ell \nu_\ell \gamma$ in terms of the vector and axial-vector form factors see, for instance, [42].

Form factors drive the hadronization of QCD currents and embed non-perturbative aspects that we still do not know how to evaluate from the underlying strong interaction theory. Their determination is all-important in order to disentangle those aspects. It is reasonable to assume, as has been common lore in the literature on this topic, that hadronic resonance states should dominate the structure of form factors and, accordingly, meromorphic functions with poles in the relevant resonances coupled to the corresponding channels have been extensively proposed in order to fit hadronic data. This procedure by itself is, however, not fully satisfactory because it does not impose known QCD constraints.

On the one side, chiral symmetry of massless QCD drives the very low-energy region of form factors [1–3]. Hence the latter have to satisfy its constraints in this energy domain. On the other side, one can also demand that form factors in the resonance energy region should match short-distance QCD properties. This idea was pioneered by [36] and has been used extensively in the last years [29–35, 37–41].

In the following we apply these techniques in order to determine the standard model description of the vector and axial-vector form factors in the radiative pion decay. Their definition, given by (2), illustrates the fact that they follow from three-point Green functions of the corresponding QCD currents. The proper GF in this case, namely $\langle VVP \rangle$ and $\langle VAP \rangle$, happen to be order parameters of the spontaneous breaking of chiral symmetry, and they hence are free of perturbative contributions in the chiral limit. This is a key aspect required by our procedure. Hence we consider in this article their study in the chiral limit, which otherwise should provide the dominant features. In the following we handle the GF in order to provide a description constrained by QCD, and then we will work out the form factors.

2.1 Vector form factor

The relevant GF is the $\langle VVP \rangle$ defined by

$$\begin{aligned} & (\Pi_{VVP})_{\mu\nu}^{abc}(p, q) \\ &= i^2 \int d^4x d^4y e^{i(px+qy)} \langle 0 | T \{ V_\mu^a(x) V_\nu^b(y) P^c(0) \} | 0 \rangle, \end{aligned} \quad (3)$$

where

$$V_\mu^a(x) = \left(\bar{\psi} \gamma_\mu \frac{\lambda^a}{2} \psi \right) (x), \quad P^a(x) = \left(\bar{\psi} i \gamma_5 \lambda^a \psi \right) (x) \quad (4)$$

are the octets of the vector and pseudoscalar currents, with $a = 1, \dots, 8$. Moreover, $SU(3)_V$ symmetry, parity and time reversal demand that

$$(II_{VVP})_{\mu\nu}^{abc}(p, q) = d^{abc} \varepsilon_{\mu\nu\alpha\beta} p^\alpha q^\beta II_{VVP}(p^2, q^2, r^2), \quad (5)$$

with $r_\mu = (p+q)_\mu$. In addition Bose symmetry requires $II_{VVP}(p^2, q^2, r^2) = II_{VVP}(q^2, p^2, r^2)$.

The vector form factor $F_V(q^2)$, defined by (2), derives from $II_{VVP}(p^2, q^2, r^2)$ in the chiral limit by

$$F_V(q^2) = \frac{\sqrt{2} M_{\pi^+}}{6FB_0} \lim_{p^2, r^2 \rightarrow 0} r^2 II_{VVP}(p^2, q^2, r^2), \quad (6)$$

with $B_0 = -\langle \bar{\psi} \psi \rangle_0 / F^2$.

We do not know how to determine $II_{VVP}(p^2, q^2, r^2)$ (II_{VVP} for short) from QCD in all energy domains but our knowledge of the strong interaction theory allows us to know this function in two specific limits as we will comment now [32, 33, 43].

1. *Very low-energy region.* The GF has to satisfy the constraints of chiral symmetry encoded in χ PT. The leading $\mathcal{O}(p^4)$ contribution in the chiral expansion is provided by the Wess–Zumino anomaly [44, 45] and gives

$$II_{VVP}(p^2, q^2, r^2) \simeq \frac{B_0}{r^2} \left(\frac{N_C}{4\pi^2} + \mathcal{O}(p^2, q^2, r^2) \right), \quad (7)$$

2. *Asymptotic energy region.* The perturbative QCD determination of the GF, within the operator product expansion (OPE) framework gives, in the chiral limit and at $\mathcal{O}(\alpha_S^0)^2$,

$$\begin{aligned} & \lim_{\lambda \rightarrow \infty} II_{VVP}((\lambda p)^2, (\lambda q)^2, (\lambda p + \lambda q)^2) \\ &= -\frac{B_0 F^2}{\lambda^4} \frac{p^2 + q^2 + r^2}{p^2 q^2 r^2} + \mathcal{O}\left(\frac{1}{\lambda^6}\right), \end{aligned} \quad (8)$$

$$\begin{aligned} & \lim_{\lambda \rightarrow \infty} II_{VVP}((\lambda p)^2, (q - \lambda p)^2, q^2) \\ &= -\frac{2B_0 F^2}{\lambda^2} \frac{1}{p^2 q^2} + \mathcal{O}\left(\frac{1}{\lambda^3}\right), \end{aligned} \quad (9)$$

$$\begin{aligned} & \lim_{\lambda \rightarrow \infty} II_{VVP}((\lambda p)^2, q^2, (q + \lambda p)^2) \\ &= i \frac{2}{\lambda^2} \frac{1}{p^2} II_{VT}(q^2) + \mathcal{O}\left(\frac{1}{\lambda^3}\right), \end{aligned} \quad (10)$$

² Although this result is completely symmetric in the three momenta, this does not longer hold when gluon corrections are included [46, 47]. Hence we do not expect that this symmetry will be sustained when constructing the GF in the resonance energy region.

where

$$\begin{aligned} & \delta^{ab} (II_{VT})_{\mu\rho\sigma}(p) \\ &= i \int d^4x e^{ip \cdot x} \langle 0 | T \{ V_\mu^a(x) T_{\rho\sigma}^b(0) \} | 0 \rangle, \\ & (II_{VT})_{\mu\rho\sigma}(p) = (p_\rho g_{\mu\sigma} - p_\sigma g_{\mu\rho}) II_{VT}(p^2), \\ & \lim_{\lambda \rightarrow \infty} II_{VT}((\lambda p)^2) = i \frac{B_0 F^2}{\lambda^2} \frac{1}{p^2} + \mathcal{O}\left(\frac{1}{\lambda^4}\right), \\ & T_{\rho\sigma}^a(x) = \left(\bar{q} \sigma_{\rho\sigma} \frac{\lambda^a}{2} q \right) (x), \end{aligned} \quad (11)$$

where $(II_{VT})_{\mu\rho\sigma}(p)$ stands for the vector tensor GF.³

In addition to these constraints on the $\langle VVP \rangle$ GF there is also a requirement that we will enforce in any hadronic form factor of vector or axial-vector QCD currents. It is known [48] that the leading perturbative contribution, within QCD, to the spectral functions of both vector and axial-vector correlators is constant. Then it turns out, as a heuristic deduction, that any of the infinite hadron contributions to the spectral functions should vanish at high transfer of momentum. This implies, in order, that hadron form factors of those currents should behave smoothly at high energy [49, 50]. Incidentally this feature coincides with the known Brodsky–Lepage condition on form factors (derived within a partonic framework) [51]. Specifically the condition, in our case, from (6) reads

$$\lim_{\substack{p^2, r^2 \rightarrow 0 \\ q^2 \rightarrow \infty}} r^2 II_{VVP}(p^2, q^2, r^2) = 0. \quad (12)$$

Our task is to construct a function for $\langle VVP \rangle$ that satisfies, at least, the set of conditions in (7)–(12).

To proceed we use the ideas of the large- N_C limit of QCD [27, 28] that, essentially, tell us that GF of currents should be described, in the $N_C \rightarrow \infty$ limit, by meromorphic functions emerging from a theory with an infinite hadron spectrum of stable states. This setting is difficult to handle and, in practice, one cuts the spectrum reducing it to the lightest Goldstone or resonance mesons that obey specific hints⁴.

In [33] a Lagrangian theory, including one multiplet of vector resonances only, was designed in order to obtain an expression for II_{VVP} that satisfied all conditions but for the one in (12). Indeed the fact that only one multiplet of vector resonances was not enough in order to satisfy all short-distance constraints for this GF was already previously noticed [32] with the use of a parametric ansatz. It is already well known [32, 34] that the MHA is more involved if we want that our representation of the GF satisfies both OPE and the Brodsky–Lepage requirements.

The obvious extension is to extend our spectrum by including also a multiplet of pseudoscalar resonances in the construction of the II_{VVP} function. Although it can be

³ We use $\sigma_{\mu\nu} = (i/2) [\gamma_\mu, \gamma_\nu]$.

⁴ This method is known as minimal hadronic ansatz (MHA) [29–31].

shown that indeed this parameterization satisfies the conditions in (7), (8), (12), it fails to meet the OPE condition in (9)⁵. It is not difficult to relate this problem to the fact that the $\langle AP \rangle$ correlator in the chiral limit is saturated by one pion exchange [46, 47].

It can be seen that all conditions are met if we consider in the spectrum of the $\langle VVP \rangle$ GF two non-degenerated multiplets of vector resonances, together with the Goldstone pseudoscalar mesons. Then the ansatz would read

$$\Pi_{VVP}^{\text{res}}(p^2, q^2, r^2) = \frac{P_0 + P_1 + P_2 + P_3}{(M_{V_1}^2 - q^2)(M_{V_1}^2 - p^2)(M_{V_2}^2 - q^2)(M_{V_2}^2 - p^2)r^2}, \quad (13)$$

with

$$P_n = \sum_{k=0}^n \sum_{l=0}^k c_{n-k, k-l, l} (r^2)^{n-k} (q^2)^{k-l} (p^2)^l, \quad (14)$$

where the coefficients are symmetric under interchange of the last two indices $c_{kml} = c_{klm}$ and M_{V_i} ($i = 1, 2$) are the masses of two nonets of vector resonances in the $N_C \rightarrow \infty$ limit⁶. In order to satisfy the short-distance constraints several conditions on the c_{ijk} coefficients arise. Hence the chiral symmetry behavior (7) gives

$$c_{000} = B_0 M_{V_1}^4 M_{V_2}^4 \frac{N_C}{4\pi^2}. \quad (15)$$

The OPE condition in (8) is satisfied for

$$\begin{aligned} c_{300} = c_{030} = c_{120} = c_{210} &= 0, \\ c_{021} = c_{111} &= -B_0 F^2. \end{aligned} \quad (16)$$

Finally the Brodsky–Lepage behavior on the vector form factor, defined by condition (12), fixes one additional parameter, namely

$$c_{020} = 0. \quad (17)$$

Our ansatz, with all these constraints, satisfies also the OPE conditions in (9) and (10).

If we evaluate now the vector form factor from (6) we get

$$F_V(q^2) = \frac{M_{\pi^+}}{3\sqrt{2}B_0 F M_{V_1}^2 M_{V_2}^2} \frac{c_{000} + c_{010}q^2}{(M_{V_1}^2 - q^2)(M_{V_2}^2 - q^2)}, \quad (18)$$

and we observe that only one parameter, c_{010} , has not been fixed by our procedure. The expression for the vector form factor in the radiative pion decay given by (18)

is the most general one that satisfies the short-distance constraints specified above. As the transferred momenta in the $\pi^+ \rightarrow e^+ \nu_e \gamma$ process is small by comparison with the mass of the lightest vector meson resonance, $q^2 \ll M_V^2$, it is appropriate to perform the relevant expansion until first order in q^2 . Using the result for c_{000} given by (15), it gives

$$F_V(q^2) = F_V(0) \left[1 + \lambda_V \frac{q^2}{M_{\pi^+}^2} + \mathcal{O}(q^4) \right], \quad (19)$$

where $\lambda_V = A_{N_C \rightarrow \infty}^V + A_{1/N_C}^V + \dots$ admits an expansion in $1/N_C$ and

$$\begin{aligned} F_V(0) &= \frac{\sqrt{2}N_C M_{\pi^+}}{24\pi^2 F}, \\ A_{N_C \rightarrow \infty}^V &= \frac{M_{\pi^+}^2}{M_{V_1}^2} + \frac{M_{\pi^+}^2}{M_{V_2}^2} + M_{\pi^+}^2 \frac{c_{010}}{c_{000}}. \end{aligned} \quad (20)$$

We must compare our value⁷ for $F_V(0) \simeq 0.0271$ with the result coming from $\Gamma(\pi^0 \rightarrow \gamma\gamma)$ and CVC, $F_V(0) = 0.0261(9)$, and with the recent experimental fit by the PIBETA collaboration, $F_V(0) = 0.0259(18)$ [52, 53]. We recall that our result for the vector form factor (18) arises from a large- N_C procedure in which a model of the $N_C \rightarrow \infty$ limit has been implemented, namely the cut in the spectrum. At $q^2 \ll M_V^2$ this form factor has been studied up to $\mathcal{O}(p^6)$ in χ PT [5, 6]. At $\mathcal{O}(p^4)$ the Wess–Zumino Lagrangian determines $F_V(0)$ as given in (20). Higher chiral order corrections to this result vanish in the chiral limit; accordingly their size is suppressed over the leading order by powers of M_π^2/M_V^2 or M_π^2/Λ_χ^2 that are tiny. Indeed, using the $\mathcal{O}(p^6)$ odd-intrinsic parity Lagrangian \mathcal{L}_6^W worked out in [54], this modification to $F_V(0)$ is proportional to a low-energy constant as $M_{\pi^+}^2 C_7^W$, which also contributes to the $\pi^0 \rightarrow \gamma\gamma$ decay. From the latter one obtains [55] $C_7^W \simeq (0.013 \pm 1.17) \times 10^{-3} \text{ GeV}^{-2}$, i.e. a value compatible with zero.

The slope λ_V arises at $\mathcal{O}(p^6)$ with the usual two features: the local operator O_{22}^W in \mathcal{L}_6^W provides the $N_C \rightarrow \infty$ contribution:

$$A_{N_C \rightarrow \infty}^{\chi\text{PT}} = \frac{64\pi^2}{N_C} M_{\pi^+}^2 C_{22}^W r(\mu), \quad (21)$$

that is $\mathcal{O}(1)$ in the large- N_C expansion, and a one-loop calculation that provides the chiral logs corresponding to the next-to-leading order [6]:

$$A_{1/N_C}^{\chi\text{PT}} = -\frac{M_{\pi^+}^2}{48\pi^2 F^2} \left[1 + \log \left(\frac{M_\pi^2}{\mu^2} \right) \right]. \quad (22)$$

There is another process directly related with the $\langle VVP \rangle$ GF, namely $\pi \rightarrow \gamma\gamma^*$; hence it should be related with radiative pion decay. Indeed within the assumptions that carried us to $F_V(q^2)$ in (18), the momenta structure for the

⁵ The constraints in (10) and (11) are, in this case, undetermined.

⁶ Hadron phenomenology of the pion vector form factor suggests that the mass of the lightest vector multiplet in this limit is very well approximated by the $\rho(770)$ mass. Hence we will take $M_{V_1} = M_{\rho(770)}$.

⁷ In the following numerical determinations we will use $F = 0.0924 \text{ GeV}$, $M_\pi = 0.138 \text{ GeV}$, $M_K = 0.496 \text{ GeV}$, $M_{\pi^0} = 0.135 \text{ GeV}$, $M_{\pi^+} = 0.140 \text{ GeV}$ and $M_{V_1} = M_{\rho(770)} = 0.775 \text{ GeV}$.

$\pi \rightarrow \gamma\gamma^*$ decay should be the same, though with different normalization. In consequence the $A_{N_C \rightarrow \infty}^V$ slope in (20) is the same for both processes.

The $\pi^0 \rightarrow \gamma e^+ e^-$ amplitude can be expressed by

$$\mathcal{M}_{\pi \rightarrow \gamma\gamma^*} = \mathcal{M}_{\pi \rightarrow \gamma\gamma} \left[1 + \lambda_\gamma \frac{q^2}{M_{\pi^0}^2} + \dots \right], \quad (23)$$

where $q^2 = (p_{e^+} + p_{e^-})^2$. The slope arises at $\mathcal{O}(p^6)$ in χ PT and it is [56]⁸

$$\lambda_\gamma = \frac{64\pi^2}{N_C} M_{\pi^0}^2 C_{22}^{Wr}(\mu) - \frac{M_{\pi^0}^2}{96\pi^2 F^2} \left[2 + \log \left(\frac{M_\pi^2 M_K^2}{\mu^4} \right) \right]. \quad (24)$$

Fortunately it has been measured rather accurately [57] to be $\lambda_\gamma = 0.032 \pm 0.004$, and we can input this measure to determine the LEC $C_{22}^W(\mu)$, obtaining

$$C_{22}^{Wr}(M_\rho) = 7.0_{-1.5}^{+1.0} \times 10^{-3} \text{ GeV}^{-2}, \quad (25)$$

where the error includes also the uncertainty of the renormalization point μ between M_ρ and 1 GeV. Coming back to the slope of $F_V(q^2)$, we get

$$\lambda_V = 0.041_{-0.007}^{+0.004}, \quad (26)$$

which compares well with the recent PIBETA measurement $\lambda_V = 0.070 \pm 0.058$ [52, 53].

By comparing now $A_{N_C \rightarrow \infty}^V$ in (20) and $A_{N_C \rightarrow \infty}^{\chi\text{PT}}$ in (21) we can provide a determination for the undetermined parameter in the GF and then give a full prescription for the $F_V(q^2)$ form factor in (18). For the mass of the first multiplet of vector resonances we take $M_{\rho(770)}$ and for the second $M_{\rho(1450)} = 1.459 \text{ GeV}$:

$$\frac{c_{010}}{c_{000}} = (-0.7 \pm 0.3) \text{ GeV}^{-2}. \quad (27)$$

Notice that the size of this parameter is of the same order as the other two terms in $A_{N_C \rightarrow \infty}^V$. With this result we end the construction of the vector form factor in radiative pion decays in the large- N_C limit given by (18).

It is interesting to compare our results with those in [33, 58]. As commented above the construction of the $\langle VVP \rangle$ GF in those references was carried out using only one multiplet of vector resonances; hence the vector form factor in radiative pion decay did not satisfy the constraint in (12). With this setting we were able to give a full prediction for the leading contribution to the slope λ_V , namely,

$$A_{N_C \rightarrow \infty}^{1R} = \frac{M_{\pi^+}^2}{M_V^2} \left[1 - \frac{4\pi^2 F^2}{N_C M_V^2} \right]. \quad (28)$$

Using $M_V = M_\rho$ we got $A_{N_C \rightarrow \infty}^{1R} \simeq 0.027$, to be compared with $A_{N_C \rightarrow \infty}^V = 0.028 \pm 0.006$ from our analysis above.

⁸ Notice that one-loop $\mathcal{O}(p^6)$ χ PT contributions, encoded in $A_{1/N_C}^{\chi\text{PT}}$, coincide in $\pi \rightarrow e\nu_e\gamma$ and $\pi \rightarrow \gamma\gamma^*$ in the $\text{SU}(3)_V$ limit, i.e. for $M_K = M_\pi$.

The study of the $\langle VVP \rangle$ GF along the lines outlined in this section can also be carried out within a resonance Lagrange theory instead of a parametric representation as given by (13). We collect this procedure in Appendix A.

2.2 Axial-vector form factor

We now come back to the axial-vector form factor defined by (2). In order to determine the $F_A(q^2)$ form factor we follow an analogous procedure to the one outlined before for the vector form factor. The relevant GF is, in this case, the $\langle VAP \rangle$ defined by

$$\begin{aligned} (II_{VAP})_{\mu\nu}^{abc}(p, q) = \\ i^2 \int d^4x d^4y e^{i(px+qy)} \langle 0 | T \{ V_\mu^a(x) A_\nu^b(y) P^c(0) \} | 0 \rangle, \end{aligned} \quad (29)$$

where

$$A_\mu^a(x) = \left(\bar{\psi} \gamma_\mu \gamma_5 \frac{\lambda^a}{2} \psi \right) (x), \quad (30)$$

and the other currents have been defined in (4). The structure of this GF is slightly more complicated than $\langle VVP \rangle$ as it involves two scalar functions:

$$\begin{aligned} (II_{VAP})_{\mu\nu}^{abc}(p, q) = 2f^{abc} \left\{ B_0 F^2 \left[\frac{(p+2q)_\mu q_\nu}{q^2 r^2} - \frac{g_{\mu\nu}}{r^2} \right] \right. \\ \left. + P_{\mu\nu}(p, q) \mathcal{F}(p^2, q^2, r^2) \right. \\ \left. + Q_{\mu\nu}(p, q) \mathcal{G}(p^2, q^2, r^2) \right\}, \end{aligned} \quad (31)$$

with $r_\mu = (p+q)_\mu$ and

$$\begin{aligned} P_{\mu\nu}(p, q) &= q_\mu p_\nu - (p \cdot q) g_{\mu\nu}, \\ Q_{\mu\nu}(p, q) &= p^2 q_\mu q_\nu + q^2 p_\mu p_\nu - (p \cdot q) p_\mu q_\nu - p^2 q^2 g_{\mu\nu}. \end{aligned} \quad (32)$$

Then the axial-vector form factor is obtained:

$$F_A(q^2) = \frac{\sqrt{2} M_{\pi^+}}{B_0 F} \lim_{p^2, r^2 \rightarrow 0} r^2 \mathcal{F}(p^2, q^2, r^2). \quad (33)$$

A detailed study of this GF along the line we have performed above for the $\langle VVP \rangle$ function was performed in [35]. One of the conclusions achieved was that the inclusion of one multiplet of vector, axial-vector and pseudoscalar resonances (together with the pseudoscalar mesons) was enough to satisfy the matching to the OPE expansion of the $\langle VAP \rangle$ GF at leading order. Moreover the analogous to the Brodsky–Lepage condition (12), in this case, was also satisfied, i.e. the resulting axial-vector form factor $F_A(q^2)$ behaves smoothly at high q^2 . We refer the reader to that reference for details. Hence we obtain for $N_C \rightarrow \infty$ with a cut spectrum

$$F_A(q^2) = \frac{\sqrt{2} F M_{\pi^+}}{M_A^2 - q^2} \left(\frac{M_A^2}{M_V^2} - 1 \right), \quad (34)$$

Table 1. Comparison of theoretical and experimental determinations for the low-energy expansion of vector and axial-vector form factors. The PIBETA determination assumes that the axial-vector form factor is constant, i.e. it does not consider a slope

	Experiment [16]	SU(2) [5]	SU(3) [7]	Our work
$F_V(0)$	0.0258(18)	0.0271	0.0272	0.0271
λ_V	0.070(58)	0.044	0.045	0.041
$F_A(0)$	0.0121(18)	0.0091	0.0112	exp. input
λ_A	not measured	0.0034	~ 0	0.0197(19)

where M_A is the mass of the lightest axial-vector multiplet of resonances in the $N_C \rightarrow \infty$ limit.

At $q^2 \ll M_A^2$ we may resort again to χ PT [5, 7] with the expansion

$$F_A(q^2) = F_A(0) \left[1 + \lambda_A \frac{q^2}{M_{\pi^+}^2} + \dots \right]. \quad (35)$$

Both terms, $F_A(0)$ and slope, satisfy an expansion in $1/N_C$, for instance $\lambda_A = A_{N_C \rightarrow \infty}^A + A_{1/N_C}^A + \dots$. From our result above we get

$$F_A(0) = \sqrt{2} F M_{\pi^+} \left(\frac{1}{M_V^2} - \frac{1}{M_A^2} \right),$$

$$A_{N_C \rightarrow \infty}^A = \frac{M_{\pi^+}^2}{M_A^2}. \quad (36)$$

$F_A(q^2)$ arises first at $\mathcal{O}(p^4)$ with a constant local contribution from the χ PT Lagrangian, namely

$$F_A^{(4)}(q^2) = 4\sqrt{2} \frac{M_{\pi^+}}{F} (L_9^r + L_{10}^r). \quad (37)$$

The next corrections appear at $\mathcal{O}(p^6)$ in the chiral expansion [7]. One of them results from local operators of the $\mathcal{O}(p^6)$ chiral Lagrangian that, in the chiral limit, only contribute to λ_A :

$$\lambda_A^{(6)} \Big|_{N_C \rightarrow \infty} = \frac{M_{\pi^+}^2}{L_9^r + L_{10}^r} \left[C_{78}^r - 2C_{87}^r + \frac{1}{2}C_{88}^r \right]. \quad (38)$$

There is also a subleading term, in the large- N_C expansion, that comes from one-loop diagrams involving the $\mathcal{O}(p^4)$ chiral Lagrangian. However, it only affects $F_A(0)$ and it is zero in the chiral limit. The third correction is sub-subleading and results from two-loop diagrams evaluated with the $\mathcal{O}(p^2)$ chiral Lagrangian. The latter contributes both to $F_A(0)$ and λ_A . All local additions, $F_A^{(4)}(q^2)$ and $\lambda_A^{(6)} \Big|_{N_C \rightarrow \infty}$, correspond to our result in (36), i.e. $N_C \rightarrow \infty$, when LECs are saturated by resonance contributions [35]. Though the full $\mathcal{O}(p^6)$ chiral result is rather cumbersome, the authors of [7] have provided a numerical expression for the renormalization scale $\mu = M_\rho$. The conclusion is that, in the chiral limit, subleading contributions to the slope are negligible. This notwithstanding, it is relevant to emphasize that both χ PT results of [5, 7] use models to evaluate the resonance contributions to the $\mathcal{O}(p^6)$ local terms and, accordingly, their final conclusion is tamed by this estimate.

We now turn to our numerical results. Contrarily to what happens in the vector case, where the lightest vector resonance mass in the $N_C \rightarrow \infty$ limit is well approximated by the $\rho(770)$ mass, the axial-vector mass in that limit (M_A) differs appreciably from the lightest multiplet of these resonances, namely $a_1(1260)$. The result $M_A = \sqrt{2}M_V$ was obtained in [36] by imposing several short-distance constraints on the couplings of the resonance Lagrangian. Lately [49, 50] it has been noticed that the inclusion of NLO effects in the large- N_C expansion is a pointer to $M_A \leq \sqrt{2}M_V$. These results are rather different from the mass of the lightest axial-vector meson determined experimentally $M_A \simeq 1.230$ GeV $\simeq M_{a_1(1260)}$ [14] but it is important to remind that this resonance is rather wide.

Our strategy is the following: we will use the experimental value of $F_A(0)$ as given in Table 1 to determine M_A through (36); then we provide a prediction for λ_A . We find⁹

$$M_A = 998(49)\text{MeV},$$

$$\lambda_A = 0.0197(19), \quad (39)$$

where the error stems only from the experimental uncertainty in $F_A(0)$. Notice that this result satisfies $M_A \leq \sqrt{2}M_V \simeq 1096$ MeV.

2.3 Theory versus experiment

We are now ready to compare our results with other theoretical settings and experimental determinations. In Table 1 we compare our outcome for the low-energy expansion of the form factors with the one provided by $\mathcal{O}(p^6)$ χ PT and the recent PIBETA published values.

As $F_V(0)$ is ruled by the Wess–Zumino anomaly all the theoretical results agree for this parameter. Leading corrections to this value are driven by the pion mass and as a result happen to be tiny [33]. This is also reflected in the excellent comparison with the experimental determination. The agreement is also good for the slope of the vector form factor, considering the large error of the experimental value.

For the axial-vector form factor a similar consensus does not arise. As indicated above χ PT can only predict

⁹ It is important to notice that the value of $F_A(0)$ measured by the PIBETA experiment assumes no slope for the axial-vector form factor. We should repeat this exercise when λ_A is included.

reliably all loop contributions (up to $\mathcal{O}(p^4)$ in the even-intrinsic parity and $\mathcal{O}(p^6)$ in the odd-intrinsic parity sectors), while higher order loops involve the couplings of local operators. Moreover tree-level $\mathcal{O}(p^4)$ (37) and $\mathcal{O}(p^6)$ (38) terms can only be determined in different models for resonance saturation contributions. The excellent agreement between the χ PT results and the experimental determination of $F_A(0)$ is, indeed, not a major issue as the axial-vector form factor in radiative pion decay is the main phenomenological source¹⁰ to fix the value of L_{10}^r . It happens that $F_A^{(4)}(0)$ arises from a strong cancellation between the L_9^r and L_{10}^r LECs and, in consequence, it is very sensitive to the chosen value for L_{10}^r . In terms of resonance saturation this sensitivity moves to the value of the axial-vector mass M_A input in the numerical determination. The value of $L_{10}^r \simeq -5.5 \times 10^{-3}$, used by [7], arises for $M_A \simeq 1$ GeV.

Our model of large N_C gives the leading result for the axial-vector form factor parameters and there are leading Goldstone mass driven contributions that we have not considered. In the χ PT framework these $\mathcal{O}(p^6)$ corrections arise from the LECs and, a priori, it is difficult to estimate their contribution due to our lack of reliable knowledge on those low-energy couplings. However, it has been pointed out [7] that the role of LECs is unimportant in the $\mathcal{O}(p^6)$ corrections. As the subleading loop contributions are also tiny, it is concluded that λ_A is not sizeable, and $F_A(0)$ is ruled by the leading $\mathcal{O}(p^4)$ contribution by far.

However, using as input the experimental value of $F_A(0)$ we find a large value for λ_A . As subleading $1/N_C$ loop contributions seem to be tiny our leading result shows a clear discrepancy with the estimates of tree-level contributions performed in the chiral framework [5, 7]. It would be very interesting to have an experimental determination of λ_A in order to disentangle the different resonance models.

3 Beyond SM: tensor form factor

As pointed out in Sect. 1, the history of the radiative decay of the pion accumulates a few clashes between theory and experiment. It seems though that, after the latest analysis by the PIBETA collaboration, the landscape has very much been smoothed. However, it has become customary to investigate possible contributions beyond the standard model in order to appease alleged discrepancies. Between the latter the possible role played by a tensor form factor has thoroughly been studied [18–25].

The new short-distance interaction can be written in terms of quark and lepton currents, and it reads

$$\mathcal{L}_T = \frac{G_F}{2\sqrt{2}} V_{ud} F_T [\bar{q} \sigma_{\mu\nu} (1 - \gamma_5) q] [\bar{\ell} \sigma^{\mu\nu} (1 - \gamma_5) \nu_\ell], \quad (40)$$

where F_T is an adimensional parameter measuring the strength of the new interaction. As the product $\sigma^{\mu\nu} \gamma^5$

is not an independent Dirac matrix (due to the identity $\sigma^{\mu\nu} \gamma^5 = -\frac{i}{2} \varepsilon^{\mu\nu\alpha\beta} \sigma_{\alpha\beta}$) we can write (40) as

$$\mathcal{L}_T = -\frac{G_F}{\sqrt{2}} V_{ud} F_T [\bar{q} \sigma_{\mu\nu} \gamma_5 q] [\bar{\ell} \sigma^{\mu\nu} (1 - \gamma_5) \nu_\ell]. \quad (41)$$

In the standard model the latter structure, a tensor-like quark–lepton interaction, arises from loop corrections to the tree-level amplitudes and gives a tiny value for $F_T \sim 10^{-8}$ [19, 20]. More sizeable contributions could come from new physics models. Leptoquark exchanges, for instance, could give $F_T \sim 10^{-3}$ [21], while SUSY contributions provide $F_T \sim 10^{-4} - 10^{-5}$ [19, 20] for light supersymmetric partners.

The hadronization of the tensor current, at very low transfer of momenta, is driven by the constant f_T defined by

$$\langle \gamma | \bar{u} \sigma_{\mu\nu} \gamma_5 d | \pi^- \rangle = -\frac{e}{2} f_T (p_\mu \epsilon_\nu - p_\nu \epsilon_\mu), \quad (42)$$

where p is the photon momentum¹¹. The determination of f_T involves QCD in its non-perturbative regime and, consequently, is a non-trivial task. We will come back to this issue in the next subsection.

It is possible to obtain the product $\mathcal{T} = F_T f_T$ from the analyses of different processes. Hence from some previous discrepancy in the $\pi \rightarrow e \nu_e \gamma$ process it is found that $\mathcal{T}_\pi = -(5.6 \pm 1.7) \times 10^{-3}$ [18], while from the introduction of a Gamow–Teller term in the amplitude of nuclear β -decay [26] gives $\mathcal{T}_N = (1.8 \pm 1.7) \times 10^{-3}$.

3.1 $\langle VT \rangle$ Green function: the tensor form factor

If we want to extract information on the value of F_T from experimental data, we need a reliable QCD-based determination of the hadronic tensor form factor. Using LSZ and working at leading order in the pion mass we can express the matrix element (42) as follows¹²:

$$\begin{aligned} \langle \gamma | \bar{u} \sigma_{\mu\nu} \gamma_5 d | \pi^- \rangle &= \frac{i}{\sqrt{2}F} \langle \gamma | \bar{u} \sigma_{\mu\nu} u + \bar{d} \sigma_{\mu\nu} d | 0 \rangle \\ &= -i \frac{\sqrt{2}e}{3F} \Pi_{VT}(0) (p_\mu \epsilon_\nu - p_\nu \epsilon_\mu), \end{aligned} \quad (43)$$

where in the last step we have used again the LSZ reduction formula applied to the $\langle VT \rangle$ correlator defined in (11). Then we have

$$f_T = i \frac{2\sqrt{2}}{3F} \Pi_{VT}(0). \quad (44)$$

¹⁰ L_9^r is rather well determined from the phenomenology (squared charge radius of the pion) and its numerical value agrees nicely with resonance saturation.

¹¹ There is in fact another Lorentz structure contributing to this matrix element, but it carries higher orders in the momenta. If the latter is included, f_T acquires a dependence on the square of the transferred momenta, i.e. $f_T(q^2)$. See Appendix B for a detailed evaluation of both form factors.

¹² See Appendix C for a derivation of this expression.

To determine the $\langle VT \rangle$ correlator it suffices to employ (10) with our ansatz in (13). It gives

$$\Pi_{VT}(q^2) = \frac{i}{2} \frac{c_{110} + c_{200} + 2c_{111}q^2}{(M_{V_1}^2 - q^2)(M_{V_2}^2 - q^2)}, \quad (45)$$

which satisfies the proper high-energy behavior ruled by (10). Unfortunately $\Pi_{VT}(0)$ is not fully specified by short-distance constraints within our approach as we have only fixed the c_{111} parameter.

In order to provide an estimate for f_T we consider this correlator including one multiplet of vector resonances only. To proceed we can use the results for the $\langle VVP \rangle$ GF as in [33] (using (10) for instance) or, equivalently, from the $\langle VAP \rangle$ GF [32, 35]. Both procedures yield the same result, namely

$$\Pi_{VT}(q^2) = -i \frac{B_0 F^2}{M_V^2 - q^2}, \quad (46)$$

which matches the OPE result [61]:

$$\lim_{\lambda \rightarrow \infty} \Pi_{VT}(\lambda^2 q^2) = i \frac{B_0 F^2}{\lambda^2 q^2} + \mathcal{O}\left(\frac{1}{\lambda^4}\right), \quad (47)$$

when the large momentum limit is taken. Notice that the result (46) can be recovered from (45) by taking the limit $M_{V_2} \rightarrow \infty$, demanding $c_{111} = 0$ and identifying the rest of the constants. Using (46) in (44) yields:

$$f_T = \frac{2\sqrt{2}B_0 F}{3M_V^2}. \quad (48)$$

An educated guess can be obtained by writing $B_0 F = -\langle \bar{\psi}\psi \rangle_0 / F$ and the estimate $\langle \bar{\psi}\psi \rangle_0 (1 \text{ GeV}) = -(242 \pm 15 \text{ MeV})^3$ [62]. We obtain $f_T = 0.24 \pm 0.04$.

Another parameter of interest is the susceptibility of the quark condensate χ_z defined by the vacuum expectation value of the tensor current in the presence of an external source $Z_{\mu\nu}$ [63–65]:

$$\langle 0 | \bar{\psi} \sigma_{\mu\nu} \psi | 0 \rangle_Z = g_\psi \chi_z \langle \bar{\psi}\psi \rangle_0 Z_{\mu\nu}. \quad (49)$$

In our case we consider the magnetic susceptibility χ given by an external electromagnetic field by

$$\langle \gamma | \bar{u} \sigma_{\mu\nu} u + \bar{d} \sigma_{\mu\nu} d | 0 \rangle = -ie (e_u + e_d) \chi \langle \bar{\psi}\psi \rangle_0 F_{\mu\nu}, \quad (50)$$

with $e_u = 2/3$ and $e_d = -1/3$. Using the first equality of (43), we get

$$f_T = -\frac{\sqrt{2}}{3} \chi B_0 F, \quad (51)$$

and comparing with (48) we obtain

$$\chi = -\frac{2}{M_V^2} \simeq -3.3 \text{ GeV}^{-2}. \quad (52)$$

There are several determinations of the magnetic susceptibility that provide a range that runs from $\chi = -(8.16 \pm 0.41) \text{ GeV}^{-2}$ [63] up to $\chi \simeq -2.7 \text{ GeV}^{-2}$ [66, 67].

3.2 Lattice data and sum rules

The last years have witnessed increasing attention to the determination of matrix elements of tensor quark currents. For instance, together with the QCD sum rules technique [66–68], by lattice methods one has also performed evaluations of the amplitudes involving the tensor current and a vector resonance [69, 70]:

$$\begin{aligned} \langle 0 | V_\mu^a | \rho^b(p, \lambda) \rangle &= -\frac{1}{\sqrt{2}} \delta^{ab} M_V f_V \epsilon_\mu^\lambda, \\ \langle 0 | T_{\mu\nu}^a | \rho^b(p, \lambda) \rangle &= -\frac{i}{\sqrt{2}} \delta^{ab} f_V^\perp(\mu) (\epsilon_\mu^\lambda p_\nu - \epsilon_\nu^\lambda p_\mu), \end{aligned} \quad (53)$$

where $\rho^b(p, \lambda)$ is a vector resonance with momentum p , helicity λ and polarization vector ϵ_μ^λ . The vector and tensor quark currents have been defined in (4) and (11). The scale dependence of the $f_V^\perp(\mu)$ in (53) reflects the fact that the tensor current has a non-vanishing anomalous dimension.

Within the large- N_C framework it can be shown [61, 71] that if we consider a single multiplet of vector mesons we get

$$\Pi_{VT}(q^2) = -\frac{i f_V f_V^\perp M_V}{2 M_V^2 - q^2}. \quad (54)$$

Comparing with (46) we obtain the relation

$$f_V f_V^\perp = \frac{2B_0 F^2}{M_V}. \quad (55)$$

The f_V coupling can be obtained from the measured $\Gamma(\rho^0 \rightarrow e^+ e^-)$ [57]. We obtain $f_V \simeq 221 \text{ MeV}$ with an expected tiny error¹³. Then from (55) and using the value of the quark condensate quoted above we get

$$f_V^\perp (1 \text{ GeV}) = 165 \pm 31 \text{ MeV}, \quad (56)$$

where the error collects only the uncertainty in the value of $\langle \bar{\psi}\psi \rangle_0$. Our result is in excellent agreement with those coming from QCD sum rules: $f_\rho^\perp = 160(10) \text{ MeV}$ [66, 67] and $f_\rho^\perp = 157(5) \text{ MeV}$ [68].

Lattice evaluations determine the ratio with the vector coupling. From our results we get

$$\frac{f_V^\perp}{f_V} (1 \text{ GeV}) = 0.75 \pm 0.14. \quad (57)$$

This is to be compared with the quenched value [69, 70], run down to $\mu = 1 \text{ GeV}$:

$$\frac{f_\rho^\perp}{f_\rho} (1 \text{ GeV}) = 0.74 \pm 0.03. \quad (58)$$

¹³ The vector coupling can also be determined from short-distance analyses within resonance theory [35], giving $f_V^2 = 2 \frac{F^2 M_A^2}{M_A^2 - M_V^2}$, which translates into $f_V = 207(15) \text{ MeV}$ for $M_A = 998(49) \text{ MeV}$ (39), in excellent agreement with the quoted phenomenological result.

Finally from the latter result and the phenomenological value of f_V , the lattice method provides the determination

$$f_\rho^\perp(1 \text{ GeV}) = 164 \pm 7 \text{ MeV}, \quad (59)$$

to compare with our figure in (56).

4 Analysis of the photon spectrum in the radiative pion decay

The PIBETA experiment has thoroughly measured the photon spectrum in the radiative decay of the pion [15]. Though the results of that reference seemed to confirm a serious discrepancy with theoretical determinations, an ensuing analysis of more data and the refinement of systematic errors [16, 52, 53] has brought a close agreement between theory and experiment.

The experimental available data amount to the branching ratio of the radiative pion decay integrated in different subregions (Q) of the final state phase space:

$$R_Q = \frac{1}{\Gamma_{\pi \rightarrow e\nu}} \int_Q dQ_3 \sum_\lambda |\mathcal{M}(E_e, E_\nu)|^2, \quad (60)$$

where the sum runs over the polarizations of the final particles. The three regions and the experimental results are shown in Table 2.

We test the predictions ruled by our determination for the hadronic form factors with the experimental data, ignoring first a possible tensor interaction, and compare them with other theoretical settings. In order to achieve the accuracy required by the experimental information, higher order radiative corrections to the decay [72] must be included, and they have been implemented in our analysis. The numerical input for vector and axial-vector form factors is given in Table 1.

In the fourth column of Table 2 the latest experimental data are given; in the fifth and sixth ones we show the results provided by our analysis. We study the numerical impact of the momenta dependence of the form factors by setting the slopes to zero and we conclude that it is tiny: the q^2 dependence tends to increase the central value of R but the modification is by far within the errors. The last two columns bring the results yielded by two- and three-flavor two-loop χ PT calculations. The evaluation of the errors for the theoretical predictions is ruled by those in the form factors. The estimate of the latter has been done in the following way: we assume no error coming from the

slopes (since their numerical impact is very poor); to the vector form factor we assign the same error as that of the experimental determination, $\sim 7\%$, and to the axial-vector form factor we attach the error of the experimental input. Finally the error given for the χ PT calculations only considers the scale dependence, which is tiny, 5% .

We conclude then that the corrections induced by the q^2 dependence of vector and axial-vector form factors are numerically negligible unless the theoretical error is reduced. For this we would need a better determination of vector and, specially, axial-vector form factors at $q^2 = 0$. When comparing our results with the experimental data, we see that our predictions are in agreement with previous estimates.

As a final exercise we use the experimental data to fit the value of $\mathcal{T} = F_T f_T$ defined above. In order to reach this purpose we use the experimentally fitted values for the hadronic inputs $F_V(0)$ and $F_A(0)$, and our results for the slopes λ_V and λ_A . Finally, to extract the value of the F_T coupling from the fit, we use our determination for the tensor form factor f_T . The value that we obtain is compatible with zero and its order of magnitude is compatible with that dictated by SUSY:

$$F_T = (1 \pm 14) \times 10^{-4}. \quad (61)$$

5 Conclusions

Radiative pion decay has been a continuous source of debate between theoretical predictions and experimental determinations. Nevertheless the latest analysis by the PIBETA Collaboration seems to bring about close agreement between both sides.

In this article we have performed a detailed analysis of the structure-dependent amplitudes contributing to $\pi \rightarrow e\nu_e\gamma$. The q^2 dependence of vector and axial-vector form factors, driven by the standard model, has been rigorously constructed through the study of the $\langle VVP \rangle$ and $\langle VAP \rangle$ Green functions, by matching meromorphic ansätze with their leading OPE contributions. Moreover, we have also required that our form factors are soft at high transfer of momenta. Hence we obtain the most general (and simple) functions that satisfy all those constraints. The appropriate structure of the form factors requires a double vector resonance pole for the vector form factor and a single axial-vector resonance pole for the axial-vector form factor. After a small momenta expansion we compare our results with those of χ PT and while in the vector sector we find complete agreement, our slope for the axial-vector form

Table 2. Comparison of the theoretical predictions and the experimental data for $\mathcal{R}_Q = 10^8 R_Q$ for constant form-factors and different predictions of the q^2 dependence

$E_{e^+}^{\min}$ (MeV)	E_γ^{\min} (MeV)	$\theta_{e\gamma}^{\min}$	\mathcal{R}_{exp} [52, 53]	\mathcal{R}_{th} (without slopes)	\mathcal{R}_{th} (with slopes)	\mathcal{R}_{th} SU(2) [5]	\mathcal{R}_{th} SU(3) [7]
50	50	–	2.614(21)	2.78(38)	2.81(38)	2.46(35)	2.72(38)
10	50	40°	14.46(22)	14.81(54)	15.08(58)	14.73(53)	15.00(57)
50	10	40°	37.69(46)	38.08(98)	38.41(103)	37.51(94)	38.17(103)

factor is much larger than the one provided by models of local terms in the chiral framework.

The role of a tensor contribution to the radiative pion decay has customarily been taken into account in order to analyse the experimental results. We use those in order to fix the size of the contribution and we find that it is compatible with zero. Incidentally we have given a prediction for f_V^\perp that measures the coupling of a vector resonance $J^{PC} = 1^{--}$ to the tensor current. Our results agree well with determinations from QCD sum rules and quenched lattice.

We conclude that the standard model is able to embody the experimentally known features of the radiative pion decay. As happens with other decays involving non-perturbative strong effects, the rather large size of the numerical uncertainties generated by our lack of knowledge of this QCD regime shows that this process is, at present, unsuitable for the search of new physics.

Acknowledgements. We thank M. Bychkov and D. Počanić for useful discussions and for providing us with the latest experimental results on radiative pion decay. We also thank A. Pich for a thorough reading of the manuscript and his suggestions. The work of V. Mateu is supported by a FPU contract (MEC). This work has been supported in part by the EU MRTN-CT-2006-035482 (FLAVIANet), by MEC (Spain) under grant FPA2004-00996 and by Generalitat Valenciana under grant GVACOMP2007-156.

Appendix A: $\langle VVP \rangle$ from a Lagrangian

We used the meromorphic ansatz in (13) in order to determine the vector form factor in radiative pion decay. To reach the same result, one could also proceed starting with a Lagrangian like the one given by resonance chiral theory (R χ T). This study, with one multiplet of vector resonances, was already carried out in [33]. However, we have concluded that QCD constraints seem to indicate the need of including a second vector multiplet. In this appendix we construct the short-distance constrained $\langle VVP \rangle$ function within R χ T.

In order to proceed we need to build the odd-intrinsic parity R χ T Lagrangian with two multiplets of vector resonances. The interaction pieces are

$$\mathcal{L}_V^{\text{even}} = \frac{F_V}{2\sqrt{2}} \langle V_1^{\mu\nu} f_{+\mu\nu} \rangle + \frac{F_{V'}}{2\sqrt{2}} \langle V_2^{\mu\nu} f_{+\mu\nu} \rangle, \quad (\text{A.1})$$

$$\begin{aligned} \mathcal{L}_V^{\text{odd}} = & \mathcal{L}_{\text{WZ}} + i\varepsilon_{\mu\nu\alpha\beta} \left\{ \tilde{C}_7^W \langle \chi_- f_+^{\mu\nu} f_+^{\alpha\beta} \rangle + i\tilde{C}_{22}^W \right. \\ & \times \left. \langle \nabla_\lambda f_+^{\lambda\mu} \{ f_+^{\alpha\beta}, u^\nu \} \rangle \right\} \\ & + \sum_{i=1}^7 \frac{c_i}{M_{V_1}} \mathcal{O}_{V_1 JP}^i + \sum_{i=1}^7 \frac{c'_i}{M_{V_2}} \mathcal{O}_{V_2 JP}^i \\ & + \sum_{i=1}^4 d_i \mathcal{O}_{V_1 V_1 P}^i + \sum_{i=1}^4 d'_i \mathcal{O}_{V_2 V_2 P}^i \\ & + \sum_{n=a,b,c,d,e} d_n \mathcal{O}_{V_1 V_2 P}^n + d_f \mathcal{O}_{V_1 V_2 J}^f, \quad (\text{A.2}) \end{aligned}$$

where \mathcal{L}_{WZ} is the Wess–Zumino Lagrangian [44, 45] that arises at $\mathcal{O}(p^4)$ in χ PT. As specified above only two operators contribute at $\mathcal{O}(p^6)$ [54]. The operators $\mathcal{O}_{V_i JP}$ and $\mathcal{O}_{V_i V_i P}$ were already given in [33] and will not be repeated here. For the last part of the Lagrangian there are two subsets of pieces [59]:

- $V_1 V_2 P$ terms, which contain vertices with Goldstone and two vector resonances from different multiplets:

$$\begin{aligned} \mathcal{O}_{V_1 V_2 P}^a &= \varepsilon_{\mu\nu\rho\sigma} \langle \{ V_1^{\mu\nu}, V_2^{\rho\sigma} \} \nabla_\alpha u^\sigma \rangle, \\ \mathcal{O}_{V_1 V_2 P}^b &= \varepsilon_{\mu\nu\rho\sigma} \langle \{ V_1^{\mu\alpha}, V_2^{\rho\sigma} \} \nabla_\alpha u^\nu \rangle, \\ \mathcal{O}_{V_1 V_2 P}^c &= \varepsilon_{\mu\nu\rho\sigma} \langle \{ \nabla_\alpha V_1^{\mu\nu}, V_2^{\rho\sigma} \} u^\sigma \rangle, \\ \mathcal{O}_{V_1 V_2 P}^d &= \varepsilon_{\mu\nu\rho\sigma} \langle \{ \nabla_\alpha V_1^{\mu\alpha}, V_2^{\rho\sigma} \} u^\nu \rangle, \\ \mathcal{O}_{V_1 V_2 P}^e &= \varepsilon_{\mu\nu\rho\sigma} \langle \{ \nabla^\sigma V_1^{\mu\nu}, V_2^{\rho\alpha} \} u_\alpha \rangle; \quad (\text{A.3}) \end{aligned}$$

- $V_1 V_2 J$ terms, with two vector resonances from different multiplets and one pseudoscalar external source:

$$\mathcal{O}_{V_1 V_2 J}^f = i\varepsilon_{\mu\nu\rho\sigma} \langle \{ V_1^{\mu\nu}, V_2^{\rho\sigma} \} \chi_- \rangle. \quad (\text{A.4})$$

The result for the $\langle VVP \rangle$ Green function defined in (3) is

$$\begin{aligned} \Pi_{VVP}^{R\chi T} = & -B_0 \left\{ 64\tilde{C}_7^W - 16\tilde{C}_{22}^W \frac{p^2 + q^2}{r^2} \right. \\ & + \frac{Ar^2 + B(p^2 + q^2)}{(M_{V_1}^2 - p^2)(M_{V_1}^2 - q^2)r^2} \\ & + C \frac{1}{(M_{V_1}^2 - p^2)(M_{V_1}^2 - q^2)} - \frac{N_C}{4\pi^2 r^2} \\ & + D \left(\frac{1}{M_{V_1}^2 - p^2} + \frac{1}{M_{V_1}^2 - q^2} \right) \\ & + \frac{1}{r^2} \left(\frac{Er^2 + Kp^2 + Gq^2}{M_{V_1}^2 - p^2} + \frac{Er^2 + Kq^2 + Gp^2}{M_{V_1}^2 - q^2} \right) \\ & + \frac{A'r^2 + B'(p^2 + q^2)}{(M_{V_2}^2 - p^2)(M_{V_2}^2 - q^2)r^2} \\ & + C' \frac{1}{(M_{V_2}^2 - p^2)(M_{V_2}^2 - q^2)} \\ & + D' \left(\frac{1}{M_{V_2}^2 - p^2} + \frac{1}{M_{V_2}^2 - q^2} \right) \\ & + \frac{1}{r^2} \left(\frac{E'r^2 + K'p^2 + G'q^2}{M_{V_2}^2 - p^2} \right. \\ & \left. + \frac{E'r^2 + K'q^2 + G'p^2}{M_{V_2}^2 - q^2} \right) \\ & + \frac{1}{r^2} \left[\frac{A''r^2 + B''p^2 + Hq^2}{(M_{V_1}^2 - p^2)(M_{V_2}^2 - q^2)} \right. \\ & \left. + \frac{A''r^2 + B''q^2 + Hp^2}{(M_{V_2}^2 - p^2)(M_{V_1}^2 - q^2)} \right] \end{aligned}$$

$$+ C'' \left[\frac{1}{(M_{V_1}^2 - p^2)(M_{V_2}^2 - q^2)} + \frac{1}{(M_{V_1}^2 - q^2)(M_{V_2}^2 - p^2)} \right], \quad (\text{A.5})$$

where

$$\begin{aligned} A &= 8F_V^2 (d_1 - d_3), \\ A' &= 8F_{V'}^2 (d'_1 - d'_3), \\ B &= 8F_V^2 d_3, \\ B' &= 8F_{V'}^2 d'_3, \\ C &= 64F_V^2 d_2, \\ C' &= 64F_{V'}^2 d'_2, \\ D &= -\frac{32\sqrt{2}F_V c_3}{M_{V_1}}, \\ D' &= -\frac{32\sqrt{2}F_{V'} c'_3}{M_{V_2}}, \\ E &= -\frac{4\sqrt{2}F_V}{M_{V_1}} (c_1 + c_2 - c_5), \\ E' &= -\frac{4\sqrt{2}F_{V'}}{M_{V_2}} (c'_1 + c'_2 - c'_5), \\ K &= -\frac{4\sqrt{2}F_V}{M_{V_1}} (-c_1 + c_2 + c_5 - 2c_6), \\ K' &= -\frac{4\sqrt{2}F_{V'}}{M_{V_2}} (-c'_1 + c'_2 + c'_5 - 2c'_6), \\ G &= -\frac{4\sqrt{2}F_V}{M_{V_1}} (c_1 - c_2 + c_5), \\ G' &= -\frac{4\sqrt{2}F_{V'}}{M_{V_2}} (c'_1 - c'_2 + c'_5), \\ C'' &= 32F_V F_{V'} d_f, \\ B'' &= 4F_V F_{V'} (d_b + d_c - d_a - 2d_d), \\ A'' &= 4F_V F_{V'} (d_a + d_b - d_c), \\ H &= 4F_V F_{V'} (d_a + d_c - d_b). \end{aligned} \quad (\text{A.6})$$

In terms of the parameters of our ansatz in (13) we obtain

$$\begin{aligned} c_{031} &= -G - G', \\ c_{022} &= -2(K + K') - \frac{N_C}{4\pi^2}, \\ c_{121} &= -D - D' - E - E', \\ c_{120} &= (D + E)M_{V_2}^2 + (D' + E')M_{V_1}^2, \\ c_{111} &= A + A' + 2A'' + C + C' + 2C'' \\ &\quad + 2(M_{V_1}^2 + M_{V_2}^2)(D + D' + E + E'), \\ c_{021} &= B + B' + B'' + H + K(M_{V_1}^2 + 2M_{V_2}^2) \\ &\quad + K'(2M_{V_1}^2 + M_{V_2}^2) \\ &\quad + (M_{V_1}^2 + M_{V_2}^2) \left(G + G' + \frac{N_C}{4\pi^2} \right), \\ c_{030} &= GM_{V_2}^2 + G'M_{V_1}^2, \end{aligned} \quad (\text{A.7})$$

$$\begin{aligned} c_{110} &= -(D + E)M_{V_2}^4 - (D' + E')M_{V_1}^4 \\ &\quad - (A + A'' + C + C'')M_{V_2}^2 \\ &\quad - (A' + A'' + C' + C'')M_{V_1}^2 \\ &\quad - 2(D + D' + E + E')M_{V_1}^2 M_{V_2}^2, \\ c_{011} &= -2KM_{V_2}^4 - 2K'M_{V_1}^4 - (M_{V_1}^2 + M_{V_2}^2)^2 \frac{N_C}{4\pi^2} \\ &\quad - 2(H + B')M_{V_1}^2 - 2(B + B'')M_{V_2}^2 \\ &\quad - 2(K + K' + G + G')M_{V_1}^2 M_{V_2}^2, \\ c_{020} &= -G'M_{V_1}^4 - GM_{V_2}^4 \\ &\quad - \left(K + K' + G + G' + \frac{N_C}{4\pi^2} \right) M_{V_1}^2 M_{V_2}^2 \\ &\quad - (B' + B'')M_{V_1}^2 - (B + H)M_{V_2}^2, \\ c_{010} &= BM_{V_2}^4 + B'M_{V_1}^4 + (K' + G')M_{V_2}^2 M_{V_1}^4 \\ &\quad + (K + G)M_{V_1}^2 M_{V_2}^4 \\ &\quad + (B'' + H)M_{V_1}^2 M_{V_2}^2 + (M_{V_1}^2 + M_{V_2}^2) M_{V_1}^2 M_{V_2}^2 \frac{N_C}{4\pi^2}, \\ c_{100} &= (A' + C')M_{V_1}^4 + (A + C)M_{V_2}^4 \\ &\quad + 2(A'' + C'')M_{V_1}^2 M_{V_2}^2 \\ &\quad + 2(E' + D')M_{V_1}^4 M_{V_2}^2 + 2(E + D)M_{V_2}^4 M_{V_1}^2, \\ c_{000} &= -M_{V_2}^4 M_{V_1}^4 \frac{N_C}{4\pi^2}, \end{aligned} \quad (\text{A.8})$$

in units of $-B_0$. Chiral symmetry, implemented in our Lagrangian, brings about features that with the ansatz had to be forced by hand. In this way we immediately find that $c_{300} = 0$ and $c_{210} = 0$. Moreover, as a bonus we also find $c_{200} = 0$.

The rest of constraints are given in (16) and (17). In addition we find five more relations:

$$\begin{aligned} \tilde{C}_7^W &= 0, \\ \tilde{C}_{22}^W &= 0, \\ G + G' &= 0, \\ D + D' + E + E' &= 0, \\ 2(K + K') &= -\frac{N_C}{4\pi^2}. \end{aligned} \quad (\text{A.9})$$

After applying all the constraints coming from the OPE expansion and Brodsky–Lepage asymptotic condition we obtain the following relations among the Lagrangian couplings:

$$\begin{aligned} 4c_3 + c_1 &= 0, \\ 4c'_3 + c'_1 &= 0, \\ c_1 - c_2 + c_5 &= 0, \\ c'_1 - c'_2 + c'_5 &= 0, \\ c_5 - c_6 + \frac{F_{V'} M_{V_1}}{F_V M_{V_2}} (c'_5 - c'_6) &= \frac{M_{V_1}}{F_V} \frac{N_C}{64\sqrt{2}\pi^2}, \\ 8F_V^2 d_3 + 8F_{V'}^2 d'_3 + 8F_V F_{V'} (d_c - d_a) \\ &\quad + 8\sqrt{2}F_{V'} \frac{M_{V_2}^2 - M_{V_1}^2}{M_{V_2}} (c'_5 - c'_6) \\ &\quad + M_{V_1}^2 \frac{N_C}{8\pi^2} = F^2, \end{aligned}$$

$$\begin{aligned}
& 4F_V^2(d_1 + 8d_2) + 4F_{V'}^2(d'_1 + 8d'_2) \\
& + 4F_V F_{V'}(d_a + d_b - d_d + 8d_f) \\
& + 4\sqrt{2}F_{V'} \frac{M_{V_2}^2 - M_{V_1}^2}{M_{V_2}}(c'_5 - c'_6) \\
& + M_{V_1}^2 \frac{N_C}{16\pi^2} = F^2, \quad (\text{A.10}) \\
& 8M_{V_2}^2 F_V^2 d_3 + 8F_{V'}^2 M_{V_1}^2 d'_3 \\
& + 4F_V F_{V'} [M_{V_1}^2(d_b + d_c - d_a - 2d_d) \\
& + M_{V_2}^2(d_a + d_c - d_b)] = -M_{V_1}^2 M_{V_2}^2 \frac{N_C}{8\pi^2}.
\end{aligned}$$

Appendix B: Chiral Lagrangians with external tensor sources

In this appendix we determine the matrix element in (42) by incorporating tensor sources to the resonance Lagrangian. The inclusion of those external fields in χ PT has been studied in [60].

We extend the QCD Lagrangian to accommodate an external tensor source:

$$\begin{aligned}
\mathcal{L}_{\text{QCD}} &= \mathcal{L}_{\text{QCD}}^0 + \mathcal{L}_{\text{ext}}, \\
\mathcal{L}_{\text{ext}} &= \bar{\psi} \gamma_\mu (v^\mu + \gamma_5 a^\mu) \psi - \bar{\psi} (s - i\gamma_5 p) \psi + \bar{\psi} \sigma_{\mu\nu} \bar{t}^{\mu\nu} \psi. \quad (\text{B.1})
\end{aligned}$$

The tensor source includes both octet and singlet currents

$$\bar{t}^{\mu\nu} = \sum_{a=0}^8 \frac{\lambda^a}{2} \bar{t}_a^{\mu\nu}. \quad (\text{B.2})$$

One finds that

$$\bar{\psi} \sigma_{\mu\nu} \bar{t}^{\mu\nu} \psi = \bar{\psi}_L \sigma^{\mu\nu} t_{\mu\nu}^\dagger \psi_R + \bar{\psi}_R \sigma^{\mu\nu} t_{\mu\nu} \psi_L, \quad (\text{B.3})$$

and the change of basis reads

$$\begin{aligned}
\bar{t}^{\mu\nu} &= P_L^{\mu\nu\lambda\rho} t_{\lambda\rho} + P_R^{\mu\nu\lambda\rho} t_{\lambda\rho}^\dagger, \\
t^{\mu\nu} &= P_L^{\mu\nu\lambda\rho} \bar{t}_{\lambda\rho}, \quad (\text{B.4})
\end{aligned}$$

where $P_{L,R}^{\mu\nu\lambda\rho}$ are the chiral projectors for the tensor fields, given by

$$\begin{aligned}
P_R^{\mu\nu\lambda\rho} &= \frac{1}{4}(g^{\mu\lambda} g^{\nu\rho} - g^{\nu\lambda} g^{\mu\rho} - i\epsilon^{\mu\nu\lambda\rho}), \\
P_L^{\mu\nu\lambda\rho} &= \left(P_R^{\mu\nu\lambda\rho}\right)^\dagger. \quad (\text{B.5})
\end{aligned}$$

In order to maintain the chiral invariance of the extended QCD Lagrangian (B.1) the tensor source must transform as

$$t_{\mu\nu} \rightarrow g_R t_{\mu\nu} g_L^\dagger. \quad (\text{B.6})$$

It is convenient, in order to build an effective Lagrangian invariant under the chiral group, to define the tensor operators

$$t_\pm^{\mu\nu} = u^\dagger t^{\mu\nu} u^\dagger \pm u t^{\mu\nu} u, \quad (\text{B.7})$$

transforming with the compensating field

$$t_\pm^{\mu\nu} \rightarrow h(g, x) t_\pm^{\mu\nu} h(g, x)^\dagger. \quad (\text{B.8})$$

To the lowest order in the chiral expansion we have one operator that contributes to both the radiative pion decay and the $\langle VT \rangle$ function,

$$\mathcal{L}_4^{\chi\text{PT}} \doteq A_1 \langle t_+^{\mu\nu} f_{+\mu\nu} \rangle. \quad (\text{B.9})$$

When explicitly including resonances the corresponding interacting Lagrangian of interest reads

$$\begin{aligned}
\mathcal{L}^{\text{R}\chi\text{T}} &\doteq \tilde{A}_1 \langle t_+^{\mu\nu} f_{+\mu\nu} \rangle + \frac{F_V}{2\sqrt{2}} \langle V_{\mu\nu} f_+^{\mu\nu} \rangle \\
&+ \sqrt{2} F_{VT} M_V \langle V_{\mu\nu} t_+^{\mu\nu} \rangle. \quad (\text{B.10})
\end{aligned}$$

The couplings F_V and F_{VT} are related with those defined in (53) by $f_V = \sqrt{2}F_V$ and $f_V^\perp = \sqrt{2}F_{VT}$. Upon integration of the vector meson we can relate the couplings of the two Lagrangians:

$$A_1 = \tilde{A}_1 - \frac{F_V F_{VT}}{M_V}. \quad (\text{B.11})$$

If resonance saturation of the chiral LECs would hold when including external tensor sources, A_1 should be given only by the second term in (B.11). In fact this is the case when we enforce short-distance constraints. If we evaluate the $\langle VT \rangle$ Green function defined in (11) we obtain

$$\Pi_{VT}(p^2) = i \left(\tilde{A}_1 - \frac{F_V F_{VT} M_V}{M_V^2 - p^2} \right), \quad (\text{B.12})$$

which satisfies the OPE constraint in (11) provided that

$$\begin{aligned}
\tilde{A}_1 &= 0, \\
F_V F_{VT} &= \frac{B_0 F^2}{M_V}. \quad (\text{B.13})
\end{aligned}$$

Using now the relation in (44) we get again the result in (48) for f_T .

As mentioned in Sect. 3 there are two form factors involved in the hadronic tensor matrix element

$$\begin{aligned}
\langle \gamma | \bar{u} \sigma_{\mu\nu} \gamma_5 d | \pi^- \rangle &= -\frac{e}{2} f_T(q^2) (p_\mu \epsilon_\nu - p_\nu \epsilon_\mu) \\
&- \frac{e}{2} g_T(q^2) [\epsilon q (p^\mu q^\nu - p^\nu q^\mu) + q p (q^\mu \epsilon^\nu - q^\nu \epsilon^\mu)], \quad (\text{B.14})
\end{aligned}$$

where p is the photon momentum and q the transferred momentum by the tensor current. To generate the second Lorentz structure one needs operators of higher order in the chiral expansion, such as

$$Y_{93} = \langle \nabla_\mu t_+^{\mu\nu} \nabla^\alpha f_{+\alpha\nu} \rangle, \quad (\text{B.15})$$

of [60]. In the framework of resonance chiral theory one needs in addition of the operators in (B.10) the basis of

odd-intrinsic parity operators of [33]. These new contributions do not modify result of the $f_T = f_T(0)$. The result reads

$$f_T(q^2) = \frac{\sqrt{2}F_{VT}}{3FM_V} \left[2F_V - (4\sqrt{2}M_V(c_1 - c_2 - c_5 + 2c_6) + 8F_V d_3) \frac{q^2}{M_V^2 - q^2} \right], \quad (\text{B.16})$$

$$g_T(q^2) = \frac{8F_{VT}}{3FM_V^2} \left[\frac{2M_V^2 - q^2}{M_V^2 - q^2} (c_1 - c_2 - c_5) + 2 \frac{M_V^2}{M_V^2 - q^2} c_6 - 2c_7 + \sqrt{2} \frac{F_V}{M_V} \left(\frac{M_V^2}{M_V^2 - q^2} d_3 + d_4 \right) \right]. \quad (\text{B.17})$$

The spectral function of the tensor–tensor currents correlator to which the amplitude in (B.14) contributes behaves as a constant at high q^2 and leading order in α_S [61]. Hence the $f_T(q^2)$ and $g_T(q^2)$ form factor should exhibit a smooth behavior, vanishing at large transferred momentum.

Imposing that the $f_T(q^2)$ form factor vanishes at large momentum we get the constraint

$$c_1 - c_2 - c_5 + 2c_6 = -\frac{\sqrt{2}}{4} \frac{F_V}{M_V} (1 + 4d_3). \quad (\text{B.18})$$

The same procedure with $g_T(q^2)$ gives

$$c_1 - c_2 - c_5 - 2c_7 = -\sqrt{2} \frac{F_V}{M_V} d_4. \quad (\text{B.19})$$

Interestingly enough these constraints fully determine both form factors:

$$f_T(q^2) = \frac{2\sqrt{2}F_{VT}F_V}{3F} \frac{M_V}{M_V^2 - q^2},$$

$$g_T(q^2) = -\frac{f_T(q^2)}{M_V^2}. \quad (\text{B.20})$$

Notice that the contribution of the $g_T(q^2)$ form factor to the matrix element under study is fairly suppressed, typically $\mathcal{O}(q^2/M_V^2)$ over the $f_T(q^2)$ contribution.

Appendix C: LSZ formula for a soft pion

In this appendix we discuss the derivation of (43). Let us start defining three quark currents:

$$\begin{aligned} A_\pi^\mu(x) &= \bar{d}(x)\gamma^\mu\gamma_5 u(x), \\ T_5^{\mu\nu}(x) &= \bar{u}(x)\sigma^{\mu\nu}\gamma_5 d(x), \\ T_\pi^{\mu\nu} &= \bar{u}(x)\sigma^{\mu\nu}u(x) + \bar{d}(x)\sigma^{\mu\nu}d(x). \end{aligned} \quad (\text{C.1})$$

Then we can construct the following GF that after partial integration can be expressed as

$$\langle \gamma(p) | \partial_\alpha A^\alpha T_5^{\mu\nu} | 0 \rangle \quad (\text{C.2})$$

$$\begin{aligned} &\equiv i \int d^4x e^{irx} \langle \gamma(p) | T \{ \partial_\alpha A_\pi^\alpha(x), T_5^{\mu\nu}(0) \} | 0 \rangle \\ &= r_\alpha \int d^4x e^{irx} \langle \gamma(p) | T \{ A_\pi^\alpha(x), T_5^{\mu\nu}(0) \} | 0 \rangle \\ &\quad - i \langle \gamma(p) | T_\pi^{\mu\nu}(0) | 0 \rangle. \end{aligned} \quad (\text{C.3})$$

We can relate this GF to the $\langle \gamma(p) | \bar{u}\sigma_{\mu\nu}\gamma_5 d | \pi^-(r) \rangle$ matrix element (40) through the LSZ formula:

$$\begin{aligned} &\langle \gamma(p) | \bar{u}\sigma^{\mu\nu}\gamma_5 d | \pi^-(r) \rangle \\ &= \lim_{r^2 \rightarrow M_\pi^2} \frac{r^2 - M_\pi^2}{\sqrt{2}FM_\pi^2} \langle \gamma(p) | \partial_\alpha A^\alpha T_5^{\mu\nu} | 0 \rangle. \end{aligned} \quad (\text{C.4})$$

Then the $\langle \gamma(p) | \partial_\alpha A^\alpha T_5^{\mu\nu} | 0 \rangle$ GF must have a pole at $r^2 = M_\pi^2$. Since the second piece in (C.3) has no r dependence it cannot have a pole, and this must be due to the first term:

$$\begin{aligned} &r_\alpha \int d^4x e^{irx} \langle \gamma(p) | T \{ A_\pi^\alpha(x), T_5^{\mu\nu}(0) \} | 0 \rangle \\ &\equiv \frac{1}{r^2 - M_\pi^2} \mathcal{F}^{\mu\nu}(p, r). \end{aligned} \quad (\text{C.5})$$

As the divergence of the axial current is zero in the chiral limit, the left hand side must vanish for $M_\pi = 0$, which implies

$$\mathcal{F}^{\mu\nu}(p, r)|_{M_\pi=0} = ir^2 \langle \gamma(p) | T_\pi^{\mu\nu}(0) | 0 \rangle_{M_\pi=0}. \quad (\text{C.6})$$

Finally, assuming that pion mass corrections are small, (C.4) yields the desired result:

$$\langle \gamma(p) | \bar{u}\sigma^{\mu\nu}\gamma_5 d | \pi^-(r) \rangle = i \frac{1}{\sqrt{2}F} \langle \gamma(p) | T_\pi^{\mu\nu}(0) | 0 \rangle, \quad (\text{C.7})$$

which is also the result that stems from the soft pion theorem.

References

1. S. Weinberg, *Physica A* **96**, 327 (1979)
2. J. Gasser, H. Leutwyler, *Ann. Phys.* **158**, 142 (1984)
3. J. Gasser, H. Leutwyler, *Nucl. Phys. B* **250**, 465 (1985)
4. B.R. Holstein, *Phys. Rev. D* **33**, 3316 (1986)
5. J. Bijnens, P. Talavera, *Nucl. Phys. B* **489**, 387 (1997) [arXiv:hep-ph/9610269]
6. L. Ametller, J. Bijnens, A. Bramon, F. Cornet, *Phys. Lett. B* **303**, 140 (1993) [arXiv:hep-ph/9302219]
7. C.Q. Geng, I.L. Ho, T.H. Wu, *Nucl. Phys. B* **684**, 281 (2004) [arXiv:hep-ph/0306165]
8. G. Ecker, J. Gasser, A. Pich, E. de Rafael, *Nucl. Phys. B* **321**, 311 (1989)
9. V. Cirigliano, G. Ecker, M. Eidemüller, R. Kaiser, A. Pich, J. Portolés, *Nucl. Phys. B* **753**, 139 (2006) [arXiv:hep-ph/0603205]

10. A. Pich, QCD tests from Tau decay data, SLAC Report 343, (1989), p. 416
11. J.H. Kühn, A. Santamaría, Z. Phys. C **48**, 445 (1990)
12. F. Guerrero, A. Pich, Phys. Lett. B **412**, 382 (1997) [arXiv:hep-ph/9707347]
13. M. Jamin, A. Pich, J. Portolés, Phys. Lett. B **640**, 176 (2006) [arXiv:hep-ph/0605096]
14. D. Gómez Dumm, A. Pich, J. Portolés, Phys. Rev. D **69**, 073002 (2004) [arXiv:hep-ph/0312183]
15. E. Frlez et al., Phys. Rev. Lett. **93**, 181804 (2004) [arXiv:hep-ex/0312029]
16. D. Počanić, talk given at the 5th International Workshop on Chiral Dynamics, Theory and Experiment “Chiral Dynamics 2006”, September 18th–22nd (2006), Durham (USA)
17. V.N. Bolotov et al., Phys. Lett. B **243**, 308 (1990)
18. A.A. Poblaguev, Phys. Lett. B **238**, 108 (1990)
19. V.M. Belyaev, I.I. Kogan, Phys. Lett. B **280**, 238 (1992)
20. Y.Y. Komachenko, Sov. J. Nucl. Phys. **55**, 1384 (1992) [Yad. Fiz. **55**, 2487 (1992)]
21. P. Herczeg, Phys. Rev. D **49**, 247 (1994)
22. A.A. Poblaguev, Phys. Rev. D **68**, 054020 (2003) [arXiv:hep-ph/0307166]
23. M.V. Chizhov, Mod. Phys. Lett. A **8**, 2753 (1993) [arXiv:hep-ph/0401217]
24. M.V. Chizhov, arXiv:hep-ph/0310203
25. M.V. Chizhov, Phys. Part. Nucl. **2**, 193 (2005) [Pisma Fiz. Elem. Chast. Atom. Yadra **2N4**, 7 (2005)] [arXiv:hep-ph/0402105]
26. P.A. Quin, T.E. Pickering, J.E. Schewe, P.A. Voytas, J. Deutsch, Phys. Rev. D **47**, 1247 (1993)
27. G. 't Hooft, Nucl. Phys. B **72**, 461 (1974)
28. E. Witten, Nucl. Phys. B **160**, 57 (1979)
29. S. Peris, M. Perrottet, E. de Rafael, JHEP **9805**, 011 (1998) [arXiv:hep-ph/9805442]
30. M. Knecht, S. Peris, M. Perrottet, E. de Rafael, Phys. Rev. Lett. **83**, 5230 (1999) [arXiv:hep-ph/9908283]
31. S. Peris, B. Phily, E. de Rafael, Phys. Rev. Lett. **86**, 14 (2001) [arXiv:hep-ph/0007338]
32. M. Knecht, A. Nyffeler, Eur. Phys. J. C **21**, 659 (2001) [arXiv:hep-ph/0106034]
33. P.D. Ruiz-Femenia, A. Pich, J. Portolés, JHEP **0307**, 003 (2003) [arXiv:hep-ph/0306157]
34. J. Bijnens, E. Gamiz, E. Lipartia, J. Prades, JHEP **0304**, 055 (2003) [arXiv:hep-ph/0304222]
35. V. Cirigliano, G. Ecker, M. Eidemüller, A. Pich, J. Portolés, Phys. Lett. B **596**, 96 (2004) [arXiv:hep-ph/0404004]
36. G. Ecker, J. Gasser, H. Leutwyler, A. Pich, E. de Rafael, Phys. Lett. B **223**, 425 (1989)
37. A. Pich, in: Proc. Phenomenology of Large N_C QCD, ed. by R. Lebed (World Scientific, Singapore, 2002), p. 239 [arXiv:hep-ph/0205030]
38. G. Amorós, S. Noguera, J. Portolés, Eur. Phys. J. C **27**, 243 (2003) [arXiv:hep-ph/0109169]
39. J. Portolés, P.D. Ruiz-Femenía, Nucl. Phys. Proc. Suppl. **131**, 170 (2004) [arXiv:hep-ph/0311251]
40. V. Cirigliano, G. Ecker, M. Eidemüller, R. Kaiser, A. Pich, J. Portolés, JHEP **0504**, 006 (2005) [arXiv:hep-ph/0503108]
41. J. Portolés, Nucl. Phys. Proc. Suppl. **164**, 292 (2007) [arXiv:hep-ph/0509279]
42. D.A. Bryman, P. Depommier, C. Leroy, Phys. Rep. **88**, 151 (1982)
43. B. Moussallam, Phys. Rev. D **51**, 4939 (1995) [arXiv:hep-ph/9407402]
44. J. Wess, B. Zumino, Phys. Lett. B **37**, 95 (1971)
45. E. Witten, Nucl. Phys. B **223**, 433 (1983)
46. V. Mateu, arXiv:hep-ph/0610262
47. V. Mateu, M. Jamin, A. Pich, work in preparation
48. E.G. Floratos, S. Narison, E. de Rafael, Nucl. Phys. B **155**, 115 (1979)
49. I. Rosell, J.J. Sanz-Cillero, A. Pich, JHEP **0701**, 039 (2007) [arXiv:hep-ph/0610290]
50. I. Rosell, arXiv:hep-ph/0701248
51. G.P. Lepage, S.J. Brodsky, Phys. Rev. D **22**, 2157 (1980)
52. M.A. Bychkov, talk given at the APS April Meeting-2007, April 14th–17th (2007), Jacksonville (USA)
53. D. Počanić, talk given at INT, June 4th (2007), Seattle (USA)
54. J. Bijnens, L. Giralanda, P. Talavera, Eur. Phys. J. C **23**, 539 (2002) [arXiv:hep-ph/0110400]
55. O. Strandberg, Master of Science Thesis, Determination of the anomalous chiral coefficients of order p^6 (2003), [arXiv:hep-ph/0302064]
56. J. Bijnens, A. Bramon, F. Cornet, Z. Phys. C **46**, 599 (1990)
57. Particle Data Group, W.M.R. Yao et al., J. Phys. G **33**, 1 (2006)
58. P.D. Ruiz-Femenía, A. Pich, J. Portolés, Nucl. Phys. Proc. Suppl. **133**, 215 (2004) [arXiv:hep-ph/0309345]
59. P.D. Ruiz-Femenía, Effective field theories for heavy and light fermions, PhD Thesis, University of Valencia (2003)
60. O. Catà, V. Mateu, arXiv:0705.2948 [hep-ph]
61. N.S. Craigie, J. Stern, Phys. Rev. D **26**, 2430 (1982)
62. M. Jamin, Phys. Lett. B **538**, 71 (2002) [arXiv:hep-ph/0201174]
63. B.L. Ioffe, A.V. Smilga, Nucl. Phys. B **232**, 109 (1984)
64. H.X. He, X.D. Ji, Phys. Rev. D **54**, 6897 (1996) [arXiv:hep-ph/9607408]
65. L.S. Kisslinger, Phys. Rev. C **59**, 3377 (1999) [arXiv:hep-ph/9804320]
66. P. Ball, V.M. Braun, Phys. Rev. D **54**, 2182 (1996) [arXiv:hep-ph/9602323]
67. P. Ball, V.M. Braun, N. Kivel, Nucl. Phys. B **649**, 263 (2003) [arXiv:hep-ph/0207307]
68. A.P. Bakulev, S.V. Mikhailov, Eur. Phys. J. C **17**, 129 (2000) [arXiv:hep-ph/9908287]
69. V.M. Braun, T. Burch, C. Gatteringer, M. Gockeler, G. Lacagnina, S. Schaefer, A. Schafer, Phys. Rev. D **68**, 054501 (2003) [arXiv:hep-lat/0306006]
70. D. Becirevic, V. Lubicz, F. Mescia, C. Tarantino, JHEP **0305**, 007 (2003) [arXiv:hep-lat/0301020]
71. O. Catà, V. Mateu, work in preparation
72. E.A. Kuraev, Yu.M. Bystritsky, Phys. Rev. D **69**, 114004 (2004) [arXiv:hep-ph/0310275]