

## Emerging Multipotent Aspects of Hepatocyte Growth Factor

Kunio Matsumoto and Toshikazu Nakamura<sup>1</sup>

Division of Biochemistry, Biomedical Research Center, Osaka University Medical School, Suita, Osaka 565

Received for publication, January 24, 1996

Specific tissue interactions between epithelia and mesenchyme (or stroma), e.g., epithelial-mesenchymal (or -stromal) interactions mediate crucial aspects of normal development and tissue regeneration. These events affect tissue induction, organogenesis, cell movement, and morphogenesis of multicellular structures. Extensive and diverse studies have established that hepatocyte growth factor (HGF), a ligand for the *c-met* protooncogene product of receptor tyrosine kinase, is a mesenchymal- or stromal-derived multipotent polypeptide which mediates epithelial-mesenchymal interactions. During embryogenesis, HGF supports organogenesis and morphogenesis of various tissues and organs, including the liver, kidney, lung, gut, mammary gland, tooth, skeletal system, etc. In adult tissues, HGF elicits a potent organotrophic function which supports regeneration of organs including the liver, kidney, and lung. In the brain, HGF is a new member of the family of neurotrophic factors. In neoplastic tissue, HGF is involved in tumor invasion and metastasis, through tumor-stromal interactions. While HGF was originally identified as a potent mitogen for mature hepatocytes, the biological functions of this factor reach far beyond the original identifications. Such being the case, use of HGF for purposes of therapeutics is being given increasing attention.

**Key words:** *c-met*, epithelial-mesenchymal interactions, HGF, morphogenesis, organogenesis, organ regeneration.

Growth factors which share multipotent characteristics regulating proliferation, motility, and differentiation of cells are members of critical molecules responsible for complex biological processes, including embryogenesis, angiogenesis, tissue regeneration, and malignant transformation. Hepatocyte growth factor (HGF) was initially identified in a partially purified form to be a potent mitogen for mature hepatocytes in primary culture (1-4). HGF was thereafter completely purified (5-7) and was molecularly cloned in 1989 (8, 9).

In 1990 to 1991, independent approaches led to isolation of bioactive molecules using different assay method were unexpectedly joined upon molecular cloning of factors. The cloning of cDNAs for scatter factor (10), tumor cytotoxic factor (11), and fibroblast-derived epithelial growth factor (12) revealed these molecules to be identical with HGF. Scatter factor was originally identified as a fibroblast-derived factor which "scatters" tightly growing epithelial cell colonies (13). Tumor cytotoxic factor proved to be fibroblast-derived factor which inhibits growth of certain species of carcinoma cells (14). In 1991, a fibroblast-derived epithelial morphogen which induces branching tubulogenesis in epithelial cells also proved to be HGF (15), and a natural ligand for the receptor-tyrosine kinase, *c-met* protooncogene product was identified as HGF in the same year (16, 17).

While HGF is a potent hepatotrophic factor responsible for vigorous regeneration of the liver, it has become a well

characterized multipotent growth factor which targets a wide variety of cells (reviewed in Refs. 18-22). HGF has "trophic" roles for regeneration and maintenance of various tissues and organs (18, 23). Recent extensive studies on expression and functional analysis of HGF during embryogenesis revealed a distinct aspect of this factor as a mediator in morphogenic epithelial-mesenchymal interactions essential for organogenesis (24-34). Based on its potent motogenic (enhancement of cell motility) and angiogenic activities, it seems clear that HGF is involved in growth, invasion, and metastasis of tumor cells. In this review, we will focus on unique multipotent aspects of HGF as a mediator in specific cell-cell interactions.

### Biochemical characteristics of HGF and c-Met

HGF is a heterodimer with a 69 kDa  $\alpha$ -chain and a 34 kDa  $\beta$ -chain, linked by a single disulfide bridge (5-7) (Fig. 1). The  $\alpha$ -chain contains the N-terminal hairpin structure and four homologous "kringle domains" and the  $\beta$ -chain has serine protease-like motif. Thus, HGF has a structural homology with plasminogen (8, 12, 35-37). But HGF has no serine protease activity, while plasminogen and its active form, plasmin share no biological activities of HGF (8). HGF is translated from a single mRNA, as a single chain preproHGF. Extracellular processing by specific serine protease, HGF-activator or HGF-converting enzyme (38, 39), results in conversion from a biologically inactive form to active two chain mature HGF.

There are two known distinct forms of naturally occurring variant HGF, biosynthesized through alternative splicing of pre-mRNA; one form is deleted with 5 amino acids in the first kringle domain (12, 37) while the other

<sup>1</sup>To whom correspondence should be addressed. Fax: +81-6-879-3789

Abbreviations: HGF, hepatocyte growth factor; HLP/MSP, HGF-like protein/macrophage stimulating protein.

form consists of only the N-terminal hairpin domain and two kringle domains (40, 41). The former has mitogenic and motogenic activities whereas the latter form has no mitogenic activity but does have motogenic activity (42, 43). This smaller variant is likely to be a minimum unit for binding to the c-Met/HGF receptor, with a relatively high-affinity. In accordance with this, deletion of the N-terminal hairpin domain, the first kringle domain, or the second kringle domain in the HGF molecule results in a total loss of biological activities (43, 44).

The tumorigenic met oncogene was initially isolated from chemically transformed human osteosarcoma cells. Although the primary structure of the c-met protooncogene product predicted it to be a receptor-type tyrosine kinase, it remained orphan (or lonesome) receptor until two research groups independently identified its natural ligand to be HGF (16, 17). The c-Met/HGF receptor is heterodimeric molecule composed of a 50 kDa  $\alpha$ -chain and a membrane spanning 145 kDa  $\beta$ -chain which contains the intracellular tyrosine kinase domain (45). The Met/HGF receptor, when autophosphorylated in response to HGF, binds a number of substrata containing the Src homology region 2 (SH2) domains such as phosphatidylinositol 3-kinase, Grb-2(Ash)/Sos complex, Ras GTPase activating protein, pp60<sup>src</sup>, and phospholipase C- $\gamma$  (46, 47). These intracellular signaling

molecules associate with a docking site of the tandemly arranged C-terminal tyrosine residues 1349 and 1354. Mutation of these tyrosine residues results in loss-of-function mutation (46, 48, 49), while mutation of the juxta-membrane tyrosine residue suppresses the loss-of-function mutation of the Met/HGF receptor (49).

While growth factors are often classified into certain families, based on structural similarity, cDNA encoding an unique protein with a similar domain structure to HGF was isolated and the putative protein was termed HGF-like protein (HLP) (50). HLP was later shown to be an molecule identical with macrophage stimulating protein (MSP) (51, 52), originally purified from human serum. On the other hand, the c-Met/HGF receptor has two distinct family members; Ron and Sea (53, 54). Ron tyrosine kinase was identified as a specific receptor for HLP/MSP (55-57), but a ligand for Sea tyrosine kinase remains to be identified (Fig. 2).

### Biological activities

The growth-regulating activity of HGF for various cell types has been well-characterized, as described in Table I. HGF has mitogenic activity for epithelial cells (12, 18, 58-68), endothelial cells (69-71), some stromal cells (72-76), and various species of carcinoma cells (58, 78-82). HGF




Fig. 1. Schematic structure of HGF mRNA, preproHGF, and mature HGF.

also has an angiogenic activity when implanted *in vivo* (70, 71). Recent studies revealed that HGF is also involved in hematopoiesis (72, 74, 76), chondrogenesis, and bone remodeling (75, 77). HGF stimulates proliferation of hematopoietic progenitor cells and enhances the formation of colonies toward erythroid lineage or granulocyte-erythroid-megakaryocyte lineage (72-74, 76). HGF enhances growth and differentiation of osteoclastic cells at the terminal stage (77). Articular chondrocytes are target cells of HGF and HGF mRNA is expressed at presumptive articular regions during development (75). *In vitro* and *in vivo* studies provided that HGF has anti-tumor activity for certain species of carcinoma cells, and in particular, growth of most hepatoma cells is inhibited by HGF (14, 83-85).

Cell movement is an important process during embryogenesis, wound healing, and tumor invasion. Although some growth factors are known to enhance cell motility, HGF is one of most potent motogens to induce dissociation and cell movement in various types of cells (Table I) (13, 30, 59, 70, 78, 79, 81, 86-89). The motogenic activity of HGF is mediated by activation of small GTP-binding proteins, Rho, Ras, and Rac (90-93). Disruption and regulation of cell-cell and cell-matrix interactions are related to the phosphorylation of E-cadherin-associated molecules ( $\beta$ -catenin, plakoglobin, and p120) (94, 95) and focal adhesion kinase (p125<sup>Fak</sup>) (88), respectively. HGF also disrupts intercellular communications mediated by gap junctions (96, 97).

Among the multipotent characteristics of HGF, the morphogenic activity is notable and unique. This activity was initially noted in three-dimensional collagen gel cultures using MDCK cells derived from renal epithelium, wherein HGF induces branching tubular structures (15). Induction of similar branching tubules and gland-like structures in epithelial cells also occurs in other cells, including cell lines derived from hepatic duct and mammary gland (29, 30, 80). Therefore, HGF is an important factor regulating morphogenic processes during development and tissue reconstruction (see below).

Several ligands for receptor-tyrosine kinases have distinct neurotrophic actions in the brain, including members of the nerve growth factor family, basic fibroblast growth factor, and epidermal growth factor. HGF and c-Met/HGF receptor are expressed in various regions of the brain (98,

99). HGF activates Ras in neurons (100) and acts as a potent survival factor for primary cultured neurons and PC12 pheochromocytoma cells (99, 101, 102), all findings to support the thesis that HGF belongs to the family of neurotrophic factors. Likewise, HGF acts as a mitogen for Schwann cells (103).

**HGF in epithelial-mesenchymal interactions and development**

Interactions between epithelium and mesenchyme, *e.g.*, epithelial-mesenchymal interactions mediate crucial aspects of normal development, affecting tissue induction, organogenesis, and morphogenesis of specific multicellular structures. Development and morphogenesis of various organs and tissues, including kidney, lung, liver, pancreas, limb, tooth, mammary gland, hair follicle, *etc.* depend on epithelial-mesenchymal interactions. A conceptual frame-

TABLE I Typical biological activities of HGF and target cells.

Biological activity	Target cells
Mitogenic	Hepatocytes
	Hepatoblast-like cells
	Hepatic ductular epithelial cells
	Renal tubular cells
	Keratinocytes
	Hair cells
	Melanocytes
	Gastric epithelial cells
	Corneal epithelial cells
	Bronchial epithelial cells
	Alveolar type II epithelial cells
	Thyroid cells
	Mammary gland epithelial cells
	Schwann cells
	Pancreatic $\beta$ cells
	Placental cytotrophoblasts
	Prostate epithelial cells
	Osteoclast-like cells
	Vascular endothelial cells
	Articular chondrocytes
	Hematopoietic progenitor cells
	Gallbladder cancer cells, <i>etc.</i>
	Motogenic
Hepatic ductular epithelial cells	
Keratinocytes	
Thyroid cell	
Mammary gland epithelial cells	
Vascular endothelial cells	
Articular chondrocytes	
Myogenic precursor cells	
Oral squamous carcinoma cells	
Gallbladder carcinoma cells	
A431 epidermoid carcinoma, <i>etc.</i>	
Morphogenic	Renal epithelial cells
	Hepatic epithelial cells
	Mammary gland epithelial cells
	Colon carcinoma cells, <i>etc.</i>
Promotion of cell survival	Neurons
	PC12 rat pheochromocytoma cells
Tumor inhibition	Hepatoma cells (HepG2, <i>etc.</i> )
	B6/F1 melanoma cells
	KB squamous carcinoma cells, <i>etc.</i>




Fig. 2. Ligand-receptor relationship in molecules of HGF and c-Met/HGF receptor families.

work of epithelial-mesenchymal interactions was established in the 1950s and 1960s, but molecular mechanisms responsible for these interactions have not been elucidated. Recent extensive works on HGF have established that HGF is a mesenchymal-derived mediator in epithelial-mesenchymal interactions.

During kidney development, the first interaction between epithelial ureteric bud and mesenchymal metanephric blastema is essential for the development of the kidney. Epithelial cells derived from ureteric bud form branching collecting tubules and mesenchymal cells at the tip of collecting tubules convert to epithelial cells that form the nephron. During organogenesis of the kidney, c-Met/HGF receptor mRNA is expressed in epithelial cells, while HGF mRNA is expressed in mesenchymal cells in close proximity to renal epithelial cells (24, 26). A specific antibody against HGF inhibits both morphogenesis of the kidney in organ culture system and differentiation of metanephric mesenchymal cells into epithelial precursors of the nephron (25, 26). Together with *in vitro* induction of branching tubulogenesis by HGF, this factor is a mesenchymal-derived morphogen for renal epithelial cells and is involved in transdifferentiation from mesenchymal to epithelial cells. Likewise, HGF and c-Met/HGF receptor mRNA are expressed in mammary gland tissue, and HGF potently promotes the formation of branching duct-like structures by mammary gland epithelial cells *in vitro* (29-31). Therefore, HGF may mediate inducing effects of mesenchyme (or stroma) on mammary gland development.

The potential participation of HGF in organogenesis was also demonstrated by disruption of the HGF gene (27, 28). In the homozygous mutant mice of HGF gene, embryos are lethal, due to defected development of the placenta (28) or both placenta and liver (27). Likewise, in c-Met homozygous mutant mouse embryos, development of the liver and placenta was defected (89). These defects are in consistent with the finding that HGF is a potent mitogen for placental cytotrophoblasts (64), as well as hepatocytes. The essential role of HGF in liver development has recently been demonstrated using *in vivo* loss-of-function mutation in the *Xenopus* embryo. Overexpression of mutant c-Met/HGF receptor of tyrosine kinase-minus (TK<sup>-</sup>Met) in *Xenopus* embryos resulted in liver defects and impaired development of pronephros, gut, and skeletal morphogenesis in tail regions (Aoki *et al.*, submitted). These results indicate that HGF and the c-Met/HGF receptor are highly conserved molecules, at least with regard to development of the liver, from amphibians to mammals. In contrast, in transgenic mice that express HGF specifically in the liver, a new population of small hepatocytes (presumably blastic hepatocytes) appears in the liver (104), thereby indicating that HGF may be involved in proliferation of hepatoblast-like cells (61, 105).

Localization of HGF and c-Met/HGF receptor mRNA in various tissues indicates that functional coupling between HGF and the c-Met/HGF receptor is important for development, morphogenesis, and migration of cells in other tissues, including limb, branchial arches, lung, tooth, and bone. Figure 3 shows the *in situ* localization of HGF and c-Met/HGF receptor mRNA in developing lung of day 13 of the rat embryo. The c-Met/HGF receptor mRNA is specifically localized in bronchial epithelial cells (Fig. 3, C and

D), while HGF mRNA locates in surrounding mesenchymal cells (Fig. 3, A and B). Antisense HGF oligonucleotide specifically inhibits branching tubulogenesis of the developing lung *in vitro* (our unpublished data). Thus, HGF is a mesenchymal-derived factor for branching morphogenesis during lung development. Similarly, the functional coupling between HGF and c-Met/HGF receptor supports tooth development (32). The c-Met/HGF receptor is expressed in epithelial tissue while HGF is expressed in mesenchymal tissue in tooth germ, and antisense HGF oligonucleotide specifically induces abnormal tooth morphogenesis in organ culture system (32). Expression pattern of HGF and c-Met, and biological activities of HGF implicate HGF in skeletal morphogenesis and chondrogenesis. HGF is mitogenic and motogenic for chondrocytes (75), and HGF and c-Met are expressed in chondrogenic regions, including rib, limb joints, and branchial arches (33, 34). In met<sup>-/-</sup> embryos, migration of myogenic precursor cells into the limb bud, diaphragm, and tip of tongue is impaired, and as a consequence, skeletal muscles of the limb and diaphragm do not form (89). These observations mean that HGF is involved in migration of cells during development. In the chick embryo, HGF is involved in early steps of neural induction, presumably by inducing or maintaining




Fig 3 Expression of HGF and c-Met/HGF receptor mRNA in developing rat lung. Localization of HGF mRNA (A, B) and c-Met/HGF receptor mRNA (C, D) was analyzed by *in situ* hybridization using day 13 rat embryo. B and D indicate bright field views for A and C, respectively.

competence of the epiblast to respond to neural-inducing signals (106–109).

### Organotrophic roles

Regeneration of the liver is one of the most dramatic phenomena in higher animals. When 70% of the liver is resected, the cells in the remaining liver rapidly proliferate and the original liver mass and functions are restored within a week. As partial purification of HGF was originally done using peripheral blood of partially hepatectomized rats, HGF was considered to be a humoral hepatotrophic factor which enhances liver regeneration. During the last 10 years, the hepatotrophic roles of HGF have been well-established. HGF is now seen to have the role of organotrophic factor for regeneration of other tissues and organs (19–23).

Liver injuries can be induced in rats or mice by means of partial hepatectomy, ischemia, liver crush, or administration of hepatotoxins such as CCl<sub>4</sub> and  $\alpha$ -naphthyl-isothiocyanate. Expression of HGF mRNA rapidly increases following the onset of these injuries in the injured liver and distant intact organs such as the lung and spleen. The liver is composed of several types of cells, including parenchymal hepatocytes, sinusoidal endothelial cells, Kupffer cells (liver macrophages), Ito (fat-storing) cells, and bile duct epithelial cells. Cell fractionation and *in situ* hybridization revealed that HGF is expressed in non-parenchymal stromal cells such as Kupffer cells, sinusoidal endothelial cells, and Ito cells, but not in parenchymal hepatocytes, indicating that this factor acts through a paracrine mechanism. Additionally, up-regulation of HGF mRNA in intact organs, together with a marked increase in blood HGF levels, means that an endocrine-related mechanism is likely to be functioning in liver regeneration. Elevated levels of plasma HGF were also well-demonstrated in patients with hepatic diseases (110, 111).

Based on a wide spectrum in target cell specificity of HGF, the involvements of HGF in regeneration of other organs was noted. Expression of HGF is rapidly induced after injuries in the kidney and lung (112, 113). In analogy with the case of liver injury, non-epithelial stromal cells produce HGF. Taken together with *in vitro* and *in vivo* mitogenic actions of HGF for renal (114, 115) and lung epithelial cells (60, 116), HGF seems to trigger regeneration of these organs at least through a paracrine mechanism. Therefore HGF is a stromal-derived mediator responsible for organ regeneration. Changes in blood HGF levels were noted in patients with renal diseases (117) and in patients treated by renal transplantation (118).

Expression of HGF is regulated by various factors. Interleukin-1, platelet-derived growth factor, acidic and basic fibroblast growth factor, epidermal growth factor, prostaglandins, and heparin are potent inducers of HGF expression (119–121). In contrast, transforming growth factor- $\beta$ 1 and glucocorticoids suppress the gene expression of HGF (122, 123). Although these regulatory molecules are likely to have distinct roles, the regulatory network for expression of HGF may be involved not only in organ regeneration but also in epithelial-mesenchymal and tumor-stromal interactions during organogenesis and tumor progression (see below), respectively.

Direct evidence for the organotrophic roles of HGF has been obtained from *in vivo* studies and these studies

suggested potential therapeutic strategies using recombinant HGF. Administration of HGF to experimental animals with liver injury strongly enhanced liver regeneration (124–126), and importantly, HGF suppressed the onset of hepatic dysfunction (124, 127). Likewise, HGF enhances renal regeneration and suppresses the onset of acute renal failure caused by renal toxins, renal ischemia, or unilateral nephrectomy (114, 115). More importantly, HGF prevented the onset of liver fibrosis/cirrhosis and abrogated lethal hepatic dysfunction due to chronic liver injury (128). Mitogenic, motogenic, and morphogenic activities, all of which are required for reconstruction of tissue architecture, no doubt are responsible for the organotrophic functions of HGF.

### HGF in tumor-stromal interactions

Because of its profound effects on cell growth, motility, and angiogenesis, HGF is implicated in the growth, invasion, and metastasis of tumor cells. As establishment of an autocrine loop of growth factors and their receptors is involved in tumorigenic transformation of cells, gene transfer experiments indicate that autonomous activation of c-Met/HGF receptor results in tumorigenic transformation (129–132). Stable transfection of the HGF gene in *met*-expressing epithelial cells (130–132), and the *c-met* gene in HGF-producing fibroblasts (129) both confer *in vivo* tumorigenicity in these cells. Such an autocrine activation of the Met/HGF receptor is found in certain tumor cells derived from cancer patients (87, 133), however, most carcinoma cells derived from epithelial tissues express c-Met/HGF receptor but do not express the HGF gene. This may mean that autocrine activation of the Met/HGF receptor is restricted to certain species of tumor cells.

Studies indicate the particular importance of stromal-derived HGF in invasion and metastasis of carcinoma cells. Growth and invasive potentials of tumor cells are influenced by their interactions with normal stromal fibroblasts (134–136). *In vitro* invasion of carcinoma cells into the collagen gel matrix was induced in co-cultivation with stromal fibroblasts (135), and fibroblasts can produce migration-stimulating factor (137). Although molecular mechanisms underlying these tumor-stromal interactions are of current interest to tumor biologists, one fibroblast-derived invasion factor is known to be HGF (88) and HGF induces invasion of various types of carcinoma cells *in vitro* (78, 79, 81, 86, 87). In addition to stromal fibroblast-derived HGF, we recently found that carcinoma cells secrete inducing factors for HGF expression in fibroblasts (Matsumoto *et al.*, submitted) and the presence of such inducing factor(s) was also noted by other workers (138, 139). Therefore, HGF seems to be a predominant stromal-derived invasion factor for carcinoma cells. The mutual interaction between HGF-expressing stromal cells and Met-expressing carcinoma cells mediated by HGF and its inducers may result in an acquisition of invasive phenotype in tumor cells. The epithelial-mesenchymal (or -stromal) interactions mediated by HGF are likely to be functional in tumor-stromal interactions, as well as in tissue regeneration.

### Perspective and future directions

The biological and physiological functions of HGF have been much greater than expected (Fig. 4). However, much




Fig. 4. Pleiotropic roles of HGF during embryogenesis, organ regeneration, and tumor progression.

work remains to be determined how HGF exerts its highly diversified activities and how HGF is involved in constructing an organized multicellular tissue structures (e.g., branching tubules).

Target cells of HGF are distributed widely and specific biological roles of HGF for the development and homeostasis of each tissue need to be studied. Although localization and *in vitro* analysis indicate that HGF may well play a neurotrophic role in the brain, specific roles of HGF for the maintenance and regeneration of the central nervous system, and also for the inductive processes, development, and network formation of neural cells are of general interest.

Because of its organotrophic functions, HGF may well have therapeutic potential for disorders of the liver and kidney. HGF is highly effective for chronic and often incurable hepatic disease, e.g., liver fibrosis/cirrhosis. Application of the HGF gene as a therapeutic for chronic diseases may be feasible. The generation and application of antagonistic molecules of HGF may prove to be therapeutic in inhibiting tumor invasion and metastasis.

The biological functions of HLP/MSP and Ron are still uncharacterized. Identification of novel members in HGF and the Met family, and elucidation of biological functions of HLP/MSP will shed light on the biological significance of

HGF family molecules in embryogenesis, organogenesis, and tissue regeneration.

Due to space limitation, the papers of some scientists may not have been cited. Nevertheless, we are entirely grateful to all our colleagues for "working on HGF." Gratitude is extended to M. Ohara for helping us to write up this review.

#### REFERENCES

1. Nakamura, T., Nawa, K., and Ichihara, A. (1984) Partial purification and characterization of hepatocyte growth factor from serum of hepatectomized rats. *Biochem. Biophys. Res. Commun.* **122**, 1450-1459
2. Russell, W.E., McGowan, J.A., and Bucher, N.L.R. (1984) Partial characterization of hepatocyte growth factor from rat platelets. *J. Cell. Physiol.* **119**, 183-192
3. Thaler, J. and Michalopoulos, G.K. (1985) Hepatopoietin A: A partial characterization and trypsin activation of a hepatocyte growth factor. *Cancer Res.* **45**, 2545-2549
4. Gohda, E., Tsubouchi, H., Nakayama, H., Hirono, S., Takahashi, K., Koura, M., Hashimoto, S., and Daikuhara, Y. (1986) Human hepatocyte growth factor in plasma from patients with fulminant hepatic failure. *Exp. Cell Res.* **166**, 139-150
5. Nakamura, T., Nawa, K., Ichihara, A., Kaize, N., and Nishino, T. (1987) Purification and subunit structure of hepatocyte growth factor from rat platelets. *FEBS Lett.* **224**, 311-318
6. Gohda, E., Tsubouchi, H., Nakayama, H., Hirono, S., Sakiyama, O., Takahashi, K., Miyazaki, H., Hashimoto, S., and Daikuhara, Y. (1988) Purification and partial characterization of hepatocyte growth factor from plasma of a patient with fulminant hepatic failure. *J. Clin. Invest.* **88**, 414-419
7. Zarnegar, R. and Michalopoulos, G.K. (1989) Purification and biological characterization of human hepatopoietin A, a polypeptide growth factor for hepatocytes. *Cancer Res.* **49**, 3314-3320
8. Nakamura, T., Nishizawa, T., Hagiya, M., Seki, T., Shimonishi, M., Sugimura, A., Tashiro, K., and Shimizu, S. (1989) Molecular cloning and expression of human hepatocyte growth factor. *Nature* **342**, 440-443
9. Miyazawa, K., Tsubouchi, H., Naka, D., Takahashi, K., Okigaki, M., Gohda, E., Daikuhara, Y., and Kitamura, N. (1989) Molecular cloning and sequence analysis of cDNA for human hepatocyte growth factor. *Biochem. Biophys. Res. Commun.* **163**, 967-973
10. Weidner, K.M., Arakaki, N., Hartmann, G., Vanderkerckhove, J., Weingart, S., Rieder, H., Fonatsch, C., Tsubouchi, H., Hishida, T., Daikuhara, Y., and Birchmeier, W. (1991) Evidence for the identity of human scatter factor and human hepatocyte growth factor. *Proc. Natl. Acad. Sci. USA* **88**, 7001-7005
11. Shima, N., Nagao, M., Ogaki, F., Tsuda, E., Murakami, A., and Higashio, K. (1991) Tumor cytotoxic factor-hepatocyte growth factor from human fibroblasts: Cloning of its cDNA, purification and characterization of recombinant protein. *Biochem. Biophys. Res. Commun.* **180**, 1151-1158
12. Rubin, J.S., Chan, A.M.-L., Bottaro, D.P., Burges, W.H., Taylor, W.G., Cech, A.C., Hirschfield, D.W., Wong, J., Miki, T., Finch, P.W., and Aaronson, S.A. (1991) A broad-spectrum human lung fibroblast-derived mitogen is a variant of hepatocyte growth factor. *Proc. Natl. Acad. Sci. USA* **88**, 415-419
13. Gherardi, E., Gray, J., Stoker, M., Perryman, M., and Furlong, R. (1989) Purification of scatter factor, a fibroblast-derived basic protein that modulates epithelial interactions and movement. *Proc. Natl. Acad. Sci. USA* **86**, 5844-5848
14. Higashio, K., Shima, N., Goto, M., Itagaki, Y., Nagao, M., Yasuda, H., and Morinaga, T. (1990) Identity of a tumor cytotoxic factor from human fibroblasts and hepatocyte growth factor. *Biochem. Biophys. Res. Commun.* **170**, 397-404
15. Montesano, R., Matsumoto, K., Nakamura, T., and Orci, L. (1991) Identification of a fibroblast-derived epithelial morphogen as hepatocyte growth factor. *Cell* **67**, 901-908
16. Bottaro, D.P., Rubin, J.S., Faletto, D.L., Chan, A.M.-L., Kmieciak, T.E., Vande Woude, G.F., and Aaronson, S.A. (1991) Identification of the hepatocyte growth factor receptor as the

- c-met* proto-oncogene product. *Science* **251**, 802-804
17. Naldini, L., Vigna, E., Narcinham, R.P., Gaudino, G., Zarnegar, R., Michalopoulos, G.K., and Comoglio, P.M. (1991) Hepatocyte growth factor stimulates the tyrosine kinase activity of the receptor encoded by the proto-oncogene *c-MET*. *Oncogene* **6**, 501-504
  18. Matsumoto, K. and Nakamura, T. (1992) Hepatocyte growth factor: Molecular structure, roles in liver regeneration, and other biological functions. *Crit. Rev. Oncogenesis* **3**, 27-54
  19. Jiang, W.G., Hallett, M.B., and Puntis, M.C.A. (1993) Hepatocyte growth factor/scatter factor, liver regeneration and cancer metastasis. *Br. J. Surg.* **80**, 1368-1373
  20. Zarnegar, R. and Michalopoulos, G.K. (1995) The many faces of hepatocyte growth factor: From hepatopoiesis to hematopoiesis. *J. Cell Biol.* **129**, 1177-1180
  21. Boros, P. and Miller, C.M. (1995) Hepatocyte growth factor: A multifunctional cytokine. *Lancet* **345**, 293-295
  22. Rubin, J.S., Bottaro, D.P., and Aaronson, S.A. (1993) Hepatocyte growth factor/scatter factor and its receptor, the *c-met* proto-oncogene product. *Biochim. Biophys. Acta* **1155**, 357-371
  23. Matsumoto, K. and Nakamura, T. (1994) Pleiotropic roles of HGF in mitogenesis, morphogenesis, and organ regeneration in *Gann Monograph on Cancer Research 42, Growth Factors: Cell Growth, Morphogenesis, and Transformation* (Nakamura, T. and Matsumoto, K., eds.) pp. 91-112, Japan Scientific Press Tokyo and CRC Press, Tokyo
  24. Sonnenberg, E., Meyer, D., Weidner, K.M., and Birchmeier, C. (1993) Scatter factor/hepatocyte growth factor and its receptor, the *c-met* tyrosine kinase can mediate a signal exchange between mesenchymal and epithelia during mouse development. *J. Cell Biol.* **123**, 223-235
  25. Santos, O.F.P., Barros, E.J.G., Yang, X.-M., Matsumoto, K., Nakamura, T., and Nigam, S.K. (1994) Involvement of hepatocyte growth factor in kidney development. *Dev. Biol.* **163**, 525-529
  26. Woolf, A.S., Kolatsi-Joannou, M., Hardman, P., Andermarcher, E., Moorby, C., Fine, L.G., Jat, P.S., Noble, M.D., and Gherardi, E. (1995) Roles of hepatocyte growth factor/scatter factor and the Met receptor in the early development of the metanephros. *J. Cell Biol.* **128**, 171-184
  27. Schmidt, C., Bladt, F., Goedecke, S., Brinkmann, V., Zschiesche, W., Sharpe, M., Gherardi, E., and Birchmeier, C. (1995) Scatter factor/hepatocyte growth factor is essential for liver development. *Nature* **373**, 699-702
  28. Uehara, Y., Minowa, O., Mori, C., Shiota, K., Kuno, J., Noda, T., and Kitamura, N. (1995) Placental defect and embryonic lethality in mice lacking hepatocyte growth factor/scatter factor. *Nature* **373**, 702-705
  29. Soriano, J.V., Pepper, M.S., Nakamura, T., Orci, L., and Montesano, R. (1995) Hepatocyte growth factor stimulates extensive development of branching duct-like structures by cloned mammary gland epithelial cells. *J. Cell Sci.* **108**, 413-430
  30. Niranjan, B., Buluwela, L., Yant, J., Perusinghe, N., Atherton, A., Phippard, D., Dale, T., Gusterson, B., and Kamalati, T. (1995) HGF/SF: A potent cytokine for mammary growth, morphogenesis and development. *Development* **121**, 2897-2908
  31. Yang, Y.M., Spitzer, E., Meyer, D., Sachs, M., Niemann, C., Hartmann, G., Weidner, K.M., Birchmeier, C., and Birchmeier, W. (1995) Sequential requirement of hepatocyte growth factor and neuregulin in the morphogenesis and differentiation of the mammary gland. *J. Cell Biol.* **131**, 215-226
  32. Tabata, M.J., Kim, K., Liu, J., Yamashita, K., Kato, J., Iwamoto, M., Wakisaka, S., Matsumoto, K., Nakamura, T., and Kurisu, K. (1996) Hepatocyte growth factor involves in the morphogenesis of tooth germ in murine molars. *Development*, in press
  33. Myokai, F., Washio, N., Asahara, Y., Yamaai, T., Tanda, N., Aoki, S., Kurihara, H., Murayama, Y., Saito, T., Matsumoto, K., Nakamura, T., and Nohno, T. (1995) Expression of the hepatocyte growth factor gene during chick limb development. *Dev. Dynamics* **202**, 80-90
  34. Thery, C., Sharpe, M.J., Batley, S.J., Stern, C.D., and Gherardi, E. (1995) Expression of HGF/SF, HGF1/MSP, and *c-met* suggests new functions during early chick development. *Dev. Genet.* **17**, 90-101
  35. Tashiro, K., Hagiya, M., Nishizawa, T., Seki, T., Shimonishi, M., Shimizu, S., and Nakamura, T. (1990) Deduced primary structure of rat hepatocyte growth factor and expression of the mRNA in rat tissues. *Proc. Natl. Acad. Sci. USA* **87**, 3200-3204
  36. Okajima, A., Miyazawa, K., and Kitamura, N. (1990) Primary structure of rat hepatocyte growth factor and induction of its mRNA during liver regeneration following hepatic injury. *Eur. J. Biochem.* **193**, 375-381
  37. Seki, T., Ihara, I., Sugimura, A., Shimonishi, M., Nishizawa, T., Asami, O., Hagiya, M., Nakamura, T., and Shimizu, S. (1990) Isolation and expression of cDNA for different forms of hepatocyte growth factor from human leukocytes. *Biochem. Biophys. Res. Commun.* **172**, 321-327
  38. Miyazawa, K., Shimomura, T., Kitamura, A., Kondo, J., Morimoto, Y., and Kitamura, N. (1993) Molecular cloning and sequence analysis of the cDNA for a human serine protease responsible for activation of hepatocyte growth factor. *J. Biol. Chem.* **268**, 10024-10028
  39. Mizuno, K., Tanoue, Y., Okano, I., Harano, T., Takada, K., and Nakamura, T. (1994) Purification and characterization of hepatocyte growth factor (HGF)-converting enzyme: Activation of proHGF. *Biochem. Biophys. Res. Commun.* **198**, 1161-1169
  40. Chan, A.M.-L., Rubin, J.S., Bottaro, D.P., Hirschfeld, D.W., Chedid, M., and Aaronson, S.A. (1991) Identification of a competitive HGF antagonist encoded by an alternative transcript. *Science* **254**, 1382-1385
  41. Miyazawa, K., Kitamura, A., Naka, D., and Kitamura, N. (1991) An alternatively processed mRNA generated from human hepatocyte growth factor gene. *Eur. J. Biochem.* **197**, 15-22
  42. Hartmann, G., Naldini, L., Weidner, M., Sachs, M., Vigna, E., Comoglio, P.M., and Birchmeier, W. (1992) A functional domain in the heavy chain of scatter factor/hepatocyte growth factor binds the c-Met receptor and induces cell dissociation but not mitogenesis. *Proc. Natl. Acad. Sci. USA* **89**, 11574-11578
  43. Lokker, N.A., Mark, M.R., Luis, E.A., Bennett, G.L., Robbins, K.A., Baker, J.B., and Godowski, P.J. (1992) Structure-function analysis of hepatocyte growth factor: Identification of variants that lack mitogenic activity yet retain high affinity receptor binding. *EMBO J.* **11**, 2503-2510
  44. Matsumoto, K., Takehara, T., Inoue, H., Hagiya, M., Shimizu, S., and Nakamura, T. (1991) Deletion of kringle domains or the N-terminal hairpin structure in hepatocyte growth factor results in marked decreases in related biological activities. *Biochem. Biophys. Res. Commun.* **181**, 691-699
  45. Park, M., Dean, M., Kaul, K., Braun, M.J., Gonda, M.A., and Vande Woude, G.F. (1987) Sequence of MET protooncogene cDNA has features characteristic of the tyrosine kinase family of growth factor receptors. *Proc. Natl. Acad. Sci. USA* **84**, 6379-6383
  46. Ponzetto, C., Bardelli, A., Zhen, Z., Maina, F., Zonca, P., Giordano, S., Graziani, A., Panayotou, G., and Comoglio, P.M. (1994) A multifunctional docking site mediates signaling and transformation by the hepatocyte growth factor/scatter factor receptor family. *Cell* **77**, 261-271
  47. Faletto, D.L., Kaplan, D.R., Halverson, D.O., Rosen, E.M., and Vande Woude, G.F. (1993) Signal transduction in *c-met* mediated mitogenesis in *Hepatocyte Growth Factor-Scatter Factor (HGF-SF) and the C-Met Receptor* (Goldberg, I.D. and Rosen, E.M., eds.) pp. 107-130, Birkhauser Verlag, Basel, Switzerland
  48. Zhu, H., Naujokas, M.A., Fixman, E.D., Torossian, K., and Park, M. (1994) Tyrosine 1356 in the carboxy-terminal tail of the HGF/SF receptor is essential for the transduction of signals for cell motility and morphogenesis. *J. Biol. Chem.* **269**, 29943-29948
  49. Weidner, K.M., Sachs, M., Riethmacher, D., and Birchmeier, W. (1995) Mutation of juxtamembrane tyrosine residue 1001 suppresses loss-of-function mutations of the met receptor in epithelial cells. *Proc. Natl. Acad. Sci. USA* **92**, 2597-2601
  50. Han, S., Stuart, L.A., and Degen, S.J.F. (1991) Characterization

- of the DNF15S2 locus on human chromosome 3: Identification of a gene coding for four kringle domains with homology to hepatocyte growth factor. *Biochemistry* **30**, 9768-9780
51. Yoshimura, T., Yuhki, N., Wang, M., Skeel, A., and Leonard, E.J. (1993) Cloning, sequencing, and expression of human macrophage stimulating protein (MSP, MST1) confirms MSP as a member of the family of kringle proteins and locates the MSP gene on chromosome 3. *J. Biol. Chem.* **268**, 15461-15468
  52. Shimamoto, A., Kimura, T., Matsumoto, K., and Nakamura, T. (1993) Hepatocyte growth factor-like protein is identical to macrophage stimulating protein. *FEBS Lett.* **333**, 61-66
  53. Ronsin, C., Muscatelli, F., Mattei, M.-G., and Breathnach, R. (1993) A novel putative receptor tyrosine kinase of the met family. *Oncogene* **8**, 1195-1202
  54. Huff, J.L., Jelinek, M.A., Borgman, C.A., Lansing, T.J., and Parsons, J.T. (1993) The proto-oncogene c-sea encodes a transmembrane protein tyrosine kinase related to the Met/hepatocyte growth factor/scatter factor receptor. *Proc. Natl. Acad. Sci. USA* **90**, 6140-6144
  55. Wang, M.H., Ronsin, C., Gesnel, M.C., Coupey, L., Skeel, A., Leonard, E.J., and Breathnach, R. (1994) Identification of the *ron* gene product as the receptor for the human macrophage stimulating protein. *Science* **266**, 117-119
  56. Wang, M.H., Iwama, A., Skeel, A., Suda, T., and Leonard, E.J. (1995) The murine *stk* gene product, a transmembrane protein tyrosine kinase, is a receptor for macrophage-stimulating protein. *Proc. Natl. Acad. Sci. USA* **92**, 3933-3937
  57. Gaudino, G., Follenzi, A., Naldini, L., Collesi, C., Santoro, M., Gallo, K.A., Godowski, P.J., and Comoglio, P.M. (1994) Ron is heterodimeric tyrosine kinase receptor activated by the HGF homologue MSP. *EMBO J.* **13**, 3524-3532
  58. Kan, M., Zhang, G.H., Zarnegar, R., Michalopoulos, G.K., Myoken, Y., McKeehan, W.L., and Stevens, J. (1991) Hepatocyte growth factor/hepatopoietin A stimulates the growth of rat kidney proximal tubule epithelial cells (RPTE), rat non-parenchymal liver cells, human melanoma cells, mouse keratinocytes and stimulates anchorage-independent growth of SV40 transformed RPTE. *Biochem. Biophys. Res. Commun.* **174**, 331-337
  59. Dremier, S., Taton, M., Coulonval, K., Nakamura, T., Matsumoto, K., and Dumont, J.E. (1994) Mitogenic, dedifferentiating, and scattering effects of hepatocyte growth factor on dog thyroid cells. *Endocrinology* **135**, 135-140
  60. Mason, R.J., Leslie, C.C., McCormickshannon, K., Deterding, R.P., Nakamura, T., Rubin, J.S., and Shannon, J.M. (1994) Hepatocyte growth factor is a growth factor for rat alveolar type II cells. *Am. J. Respir. Cell Mol. Biol.* **11**, 561-567
  61. Mitaka, T., Norioka, K., Nakamura, T., and Mochizuki, Y. (1994) Effect of mitogens and co-mitogens on the formation of small cell colonies in primary cultures of rat hepatocytes. *J. Cell. Physiol.* **157**, 461-468
  62. Otonski, T., Beattie, G.M., Rubin, J.S., Lopez, A.D., Baird, A., and Hayek, A. (1994) Hepatocyte growth factor/scatter factor has insulinotrophic activity in human fetal pancreatic cells. *Diabetes* **43**, 947-953
  63. Wilson, S.E., He, Y.G., Weng, J.A., Zieske, J.D., Jester, J.V., and Shurtz, G.S. (1994) Effect of epidermal growth factor, hepatocyte growth factor, and keratinocyte growth factor, on proliferation, motility and differentiation of human corneal epithelial cells. *Exp. Cell Res.* **59**, 665-678
  64. Saito, S., Sakakura, S., Enomoto, M., Ichijo, M., Matsumoto, K., and Nakamura, T. (1995) Hepatocyte growth factor promotes the growth of cytotrophoblasts by the paracrine mechanism. *J. Biochem.* **117**, 671-676
  65. Shimaoka, S., Tsuboi, R., Jindo, T., Imai, R., Takamori, K., Rubin, J.S., and Ogawa, H. (1995) Hepatocyte growth factor expressed in follicular papilla cells stimulates human hair growth in vitro. *J. Cell. Physiol.* **165**, 333-338
  66. Strain, A.J., Wallace, L., Joplin, R., Daikuhara, Y., Ishii, T., Kelly, D.A., and Neuberger, J.M. (1995) Characterization of biliary epithelial cells isolated from needle biopsies of human liver in the presence of hepatocyte growth factor. *Am. J. Pathol.* **146**, 537-545
  67. Takahashi, M., Ota, S., Shimada, T., Hamada, E., Kawabe, T., Okudaira, T., Matsumura, M., Kaneko, N., Terano, A., Nakamura, T., and Omata, M. (1995) Hepatocyte growth factor is the most potent endogenous stimulant of rabbit gastric epithelial cell proliferation and migration in primary culture. *J. Clin. Invest.* **95**, 1994-2003
  68. Vila, M.R., Nakamura, T., and Real, F.X. (1995) Hepatocyte growth factor is a potent mitogen for normal human pancreas cells in vitro. *Lab. Invest.* **73**, 409-418
  69. Morimoto, A., Okamura, K., Hamanaka, R., Sato, Y., Shima, N., Higashio, K., and Kuwano, M. (1991) Hepatocyte growth factor modulates migration and proliferation of human microvascular endothelial cells in culture. *Biochem. Biophys. Res. Commun.* **179**, 1042-1049
  70. Bussolino, F., DiRenzo, M.F., Ziche, M., Bochietto, E., Olivero, M., Naldini, L., Goudino, G., Tamagnone, L., Coffe, A., and Comoglio, P.M. (1992) Hepatocyte growth factor is a potent angiogenic factor which stimulates endothelial cell motility and growth. *J. Cell Biol.* **119**, 629-641
  71. Grant, D.S., Kleinman, H.K., Goldberg, I.D., Bhargava, M.M., Nickoloff, B.J., Kinsella, J.I., Polverini, P., and Rosen, E.M. (1993) Scatter factor induces blood vessel formation in vivo. *Proc. Natl. Acad. Sci. USA* **90**, 1937-1941
  72. Kmiecik, T.E., Keller, J.R., Rosen, E., and Vande Woude, G.F. (1992) Hepatocyte growth factor is a synergistic factor for the growth of hematopoietic progenitor cells. *Blood* **80**, 2454-2457
  73. Mizuno, K., Higuchi, O., Ihle, J.N., and Nakamura, T. (1993) Hepatocyte growth factor stimulates growth of hematopoietic progenitor cells. *Biochem. Biophys. Res. Commun.* **194**, 178-186
  74. Galimi, F., Bagnara, G.P., Bonsi, L., Cottone, E., Follenzi, A., Simeone, A., and Comoglio, P.M. (1994) Hepatocyte growth factor induces proliferation and differentiation of multipotent and erythroid hematopoietic progenitor cells. *J. Cell Biol.* **127**, 1743-1754
  75. Takebayashi, T., Iwamoto, M., Jikko, A., Matsumura, T., Enomoto-Iwamoto, M., Myoukai, F., Koyama, E., Yamaai, T., Matsumoto, K., Nakamura, T., Kurisu, K., and Noji, S. (1995) Hepatocyte growth factor/scatter factor modulates cell motility, proliferation, and proteoglycan synthesis of chondrocytes. *J. Cell Biol.* **129**, 1411-1419
  76. Nishino, T., Hisha, H., Nishino, N., Adachi, M., and Ikehara, S. (1995) Hepatocyte growth factor as a hematopoietic regulator. *Blood* **85**, 3093-3100
  77. Sato, T., Haketa, Y., Yamaguchi, Y., Mano, H., Tezuka, K., Matsumoto, K., Nakamura, T., Mori, Y., Yoshizawa, K., Sumitani, K., Kodama, H., and Kumegawa, M. (1995) Hepatocyte growth factor is involved in formation of osteoclast-like cells mediated by clonal stromal cells (MC3T3-G2/PA6). *J. Cell. Physiol.* **164**, 197-204
  78. Tajima, H., Matsumoto, K., and Nakamura, T. (1992) Regulation of cell growth and motility in various cell species. *Exp. Cell Res.* **202**, 423-431
  79. Jiang, W.G., Lloyd, D., Puntis, M.C.A., Nakamura, T., and Hallet, M.B. (1993) Regulation of spreading and growth of human colon cancer cells by hepatocyte growth factor. *Clin. Exp. Metastasis* **11**, 235-242
  80. Johnson, M., Koukoulis, G., Matsumoto, K., Nakamura, T., and Iyer, A. (1993) Hepatocyte growth factor induces proliferation and morphogenesis in nonparenchymal epithelial liver cells. *Hepatology* **17**, 1052-1061
  81. Shimura, H., Date, K., Matsumoto, K., Nakamura, T., and Tanaka, M. (1995) Induction of invasive growth in a gallbladder cancer cell line by hepatocyte growth factor in vitro. *Jpn. J. Cancer Res.* **86**, 662-669
  82. Hasegawa, Y., Yamamoto, M., Maeda, S., and Saitoh, Y. (1995) Hepatocyte growth factor and its receptor c-MET regulates both cell growth and invasion of human pancreatic cancer. *Int. J. Oncol.* **7**, 877-881
  83. Tajima, H., Matsumoto, K., and Nakamura, T. (1991) Hepatocyte growth factor has a potent anti-proliferative activity for various tumor cell lines. *FEBS Lett.* **291**, 229-232
  84. Shiota, G., Rhoads, D.B., Wang, T.C., Nakamura, T., and



- Schmidt, E.V. (1992) Hepatocyte growth factor inhibits growth of hepatocellular carcinoma cells. *Proc. Natl. Acad. Sci. USA* **89**, 373-377
85. Liu, M.L., Mars, W.M., and Michalopoulos, G.K. (1995) Hepatocyte growth factor inhibits cell proliferation in vivo of rat hepatocellular carcinomas induced by dimethylnitrosamine. *Carcinogenesis* **16**, 841-843
86. Weidner, K.M., Behrens, J., Vanderkerckhove, J., and Birchmeier, W. (1990) Scatter factor: Molecular characterization and effect on invasiveness of epithelial cells. *J. Cell Biol.* **111**, 2097-2108
87. Itakura, Y., Yamamoto, T., Matsumoto, K., and Nakamura, T. (1994) Autocrine stimulation of motility in SBC-5 small cell lung carcinoma cells by a two-kringle variant of HGF. *Cancer Lett.* **83**, 235-243
88. Matsumoto, K., Matsumoto, K., Nakamura, T., and Kramer, R.H. (1994) Hepatocyte growth factor/scatter factor induces tyrosine phosphorylation of focal adhesion kinase (p125<sup>FAK</sup>) and promotes migration and invasion by oral squamous cell carcinoma cells. *J. Biol. Chem.* **269**, 31807-31813
89. Bladt, F., Riethmacher, Isenmann, Aguzzi, A., and Birchmeier, C. (1995) Essential role for the *c-met* receptor in the migration of myogenic precursor cells into the limb bud. *Nature* **376**, 768-771
90. Takaishi, K., Sasaki, T., Kato, M., Yamori, W., Kuroda, S., Nakamura, T., Takeichi, M., and Takai, Y. (1994) Involvement of rho p21 small GTP-binding protein and its regulator in the HGF-induced cell motility. *Oncogene* **9**, 273-279
91. Nishiyama, T., Sasaki, T., Takaishi, K., Kato, M., Yaku, H., Araki, K., Matsuura, Y., and Takai, Y. (1994) rac p21 is involved in insulin-induced membrane ruffling and rho p21 is involved in hepatocyte growth factor induced and 12-O-tetradecanoylphorbol-13-acetate (TPA)-induced membrane ruffling in KB cells. *Mol. Cell. Biol.* **14**, 2447-2456
92. Hartmann, G., Weidner, K.M., Schwarz, H., and Birchmeier, W. (1994) The motility signal of scatter factor/hepatocyte growth factor mediated through the receptor tyrosine kinase Met requires intracellular action of Ras. *J. Biol. Chem.* **269**, 21936-21939
93. Ridley, A.J., Comoglio, P.M., and Hall, A. (1995) Regulation of scatter factor/hepatocyte growth factor responses by Ras, Rac, and Rho in MDCK cells. *Mol. Cell. Biol.* **15**, 1110-1122
94. Watabe, M., Matsumoto, K., Nakamura, T., and Takeichi, M. (1993) Effect of hepatocyte growth factor on cadherin-mediated cell-cell adhesion. *Cell Struct. Funct.* **18**, 117-124
95. Shibamoto, S., Hayakawa, M., Takeuchi, K., Hori, T., Oku, N., Miyazawa, K., Kitamura, N., Takeichi, M., and Ito, F. (1994) Tyrosine phosphorylation of  $\beta$ -catenin and plakoglobin enhanced by hepatocyte growth factor and epidermal growth factor in human carcinoma cells. *Cell Adhesion Commun.* **1**, 295-305
96. Moorby, C.D., Stoker, M., and Gherardi, E. (1995) HGF/SF inhibits junctional communication. *Exp. Cell Res.* **219**, 657-663
97. Ikejima, K., Watanabe, S., Kitamura, T., Hirose, M., Miyazaki, A., and Sato, N. (1995) Hepatocyte growth factor inhibits intercellular communication via gap junctions in rat hepatocytes. *Biochem. Biophys. Res. Commun.* **214**, 440-414
98. Jung, W., Castren, E., Odenthal, M., Vande Woude, G.F., Ishii, T., Dienes, H.-P., Lindholm, D., and Schirmacher, P. (1994) Expression and functional interaction of hepatocyte growth factor-scatter factor and its receptor *c-met* in mammalian brain. *J. Cell Biol.* **126**, 485-494
99. Honda, S., Kagoshima, M., Wanaka, A., Tohyama, M., Matsumoto, K., and Nakamura, T. (1995) Localization and functional coupling of HGF and *c-Met*/HGF receptor in rat brain: Implication as neurotrophic factor. *Mol. Brain Res.* **32**, 197-210
100. Yamagata, T., Muroya, K., Mukasa, T., Igarashi, H., Momoi, M., Tsukahara, T., Arahata, K., Kumagai, H., and Momoi, T. (1995) Hepatocyte growth factor specifically expressed in microglia and activated Ras in the neurons, similar to the action of neurotrophic factors. *Biochem. Biophys. Res. Commun.* **210**, 231-237
101. Hamanoue, M., Takemoto, N., Matsumoto, K., Nakamura, T., Nakajima, K., and Kohsaka, S. (1996) Neurotrophic effect of hepatocyte growth factor on CNS neurons in vitro. *J. Neurosci. Res.*, in press
102. Matsumoto, K., Kagoshima, M., and Nakamura, T. (1995) Hepatocyte growth factor acts as a potent survival factor for rat pheochromocytoma PC12 cells. *Exp. Cell Res.* **220**, 71-78
103. Krasnoselaky, A., Massay, M.J., Defrances, M.C., Michalopoulos, G.K., Zarnegar, R., and Ratner, N. (1994) Hepatocyte growth factor is a mitogen for Schwann cells and is present in neurofibromas. *J. Neurosci.* **14**, 7284-7290
104. Shiota, G., Wang, T., Monte, S., Nakamura, T., and Schmidt, E.V. (1994) Hepatocyte growth factor in transgenic mice: Effects on hepatocyte growth, liver regeneration and gene expression. *Hepatology* **19**, 962-972
105. Allison, M.R., Poulsom, R., Jeffery, R., Anikumar, T.V., Jagoe, R., and Sarraf, C.E. (1993) Expression of hepatocyte growth factor messenger RNA during oval cell activation in the rat liver. *J. Pathol.* **171**, 291-299
106. Stern, C.D., Ireland, G.W., Herrick, S.E., Gherardi, E., Gray, J., Perryman, M., and Stoker, M. (1990) Epithelial scatter factor and development of the chick embryonic axis. *Development* **110**, 1271-1284
107. Stern, C.D. (1994) The avian embryo: A powerful model system for studying neural induction. *FASEB J.* **8**, 687-691
108. Streit, A., Stern, C., Thery, C., Ireland, G.W., Aparicio, S., Sharpe, M.J., and Gherardi, E. (1995) A role for HGF/SF in neural induction and its expression in Hensen's node during gastrulation. *Development* **121**, 813-824
109. Bronner-Fraser, M. (1995) Hepatocyte growth factor/scatter factor in early development: Evidence for a role in neural induction. *Trends Genet.* **11**, 423-425
110. Tsubouchi, H., Niitani, Y., Hirono, S., Nakayama, H., Gohda, E., Arakaki, N., Sakiyama, O., Takahashi, K., Kimoto, M., Kawakami, S., Setoguchi, M., Tachikawa, T., Shin, S., Arima, T., and Daikuhara, Y. (1991) Levels of human hepatocyte growth factor in serum of patients with liver diseases determined by an enzyme-linked immunosorbent assay. *Hepatology* **13**, 1-5
111. Shiota, G., Okano, J., Kawasaki, H., Kuwamoto, T., and Nakamura, T. (1995) Serum hepatocyte growth factor levels in liver diseases: Clinical implications. *Hepatology* **21**, 106-112
112. Nagaïke, M., Hirao, S., Tajima, H., Noji, S., Taniguchi, S., Matsumoto, K., and Nakamura, T. (1991) Renotropic functions of hepatocyte growth factor in renal regeneration after unilateral nephrectomy. *J. Biol. Chem.* **266**, 22781-22784
113. Yanagita, K., Matsumoto, K., Sekiguchi, K., Ishibashi, H., Niho, Y., and Nakamura, T. (1993) Hepatocyte growth factor may act as a pulmotrophic factor on lung regeneration after acute lung injury. *J. Biol. Chem.* **268**, 21212-21217
114. Kawaida, K., Matsumoto, K., Shimazu, H., and Nakamura, T. (1994) Hepatocyte growth factor prevents acute renal failure and accelerates renal regeneration in mice. *Proc. Natl. Acad. Sci. USA* **91**, 4357-4361
115. Miller, S.B., Martin, D.R., Kissane, J., and Hammermann, M.R. (1994) Hepatocyte growth factor accelerates recovery from acute ischemic renal injury in rats. *Am. J. Physiol.* **266**, F129-F134
116. Ohmichi, H., Matsumoto, K., and Nakamura, T. (1996) In vivo mitogenic action of hepatocyte growth factor on lung epithelial cells: Pulmotrophic role in lung regeneration. *Am. J. Physiol.*, in press
117. Chang, H., Nagao, T., Ichikawa, N., Kawamoto, T., Nakamura, T., Kurokawa, K., Asano, S., and Katoh, T. (1994) Blood hepatocyte growth factor levels in chronic renal failure patients. *Nephron* **67**, 497-498
118. Takada, S., Namiki, M., Takahara, S., Matsumiya, K., Kondoh, N., Kitamura, M., Uchida, K., Matsumoto, K., Nakamura, T., and Okuyama, A. (1996) Early detection and prediction of the prognosis of acute renal rejection using serum HGF after renal transplantation. *Transplant. Int.*, in press
119. Matsumoto, K., Okazaki, H., and Nakamura, T. (1992) Up-regulation of hepatocyte growth factor gene expression by interleukin-1 in human skin fibroblasts. *Biochem. Biophys. Res. Commun.* **188**, 235-243

120. Gohda, E., Matsunaga, T., Kataoka, H., Takebe, T., and Yamamoto, I. (1994) Induction of hepatocyte growth factor in human skin fibroblasts by epidermal growth factor, platelet-derived growth factor and fibroblast growth factor. *Cytokine* **6**, 633-640
121. Matsumoto, K., Okazaki, H., and Nakamura, T. (1995) Novel function of prostaglandins as inducers of gene expression of HGF and putative mediators of tissue regeneration. *J. Biochem.* **117**, 458-464
122. Matsumoto, K., Tajima, H., Okazaki, H., and Nakamura, T. (1992) Negative regulation of hepatocyte growth factor gene expression in human lung fibroblasts and leukemic cells by transforming growth factor- $\beta$ 1 and glucocorticoids. *J. Biol. Chem.* **267**, 24917-24920
123. Gohda, E., Matsunaga, T., Kataoka, H., and Yamamoto, I. (1992) TGF- $\beta$  is a potent inhibitor of hepatocyte growth factor secretion by human fibroblasts. *Cell Biol. Int. Rep.* **16**, 917-926
124. Ishiki, Y., Ohnishi, H., Muto, Y., Matsumoto, K., and Nakamura, T. (1992) Direct evidence that hepatocyte growth factor is a hepatotrophic factor for liver regeneration and has potent antihepatitis effects *in vivo*. *Hepatology* **16**, 1227-1235
125. Fujiwara, K., Nagoshi, S., Ohno, A., Hirata, K., Ohta, Y., Mochida, S., Tomiya, T., Higashio, K., and Kurokawa, K. (1993) Stimulation of liver growth by exogenous human hepatocyte growth factor in normal and partially hepatectomized rats. *Hepatology* **18**, 1443-1449
126. Ishii, T., Sato, M., Sudo, K., Suzuki, M., Nakai, H., Hishida, T., Niwa, T., Umez, K., and Yuasa, S. (1995) Hepatocyte growth factor stimulates liver regeneration and elevates blood protein level in normal and partially hepatectomized rats. *J. Biochem.* **117**, 1105-1112
127. Roos, F., Terrell, T.G., Godowski, P.J., Chamow, S.W., and Schwall, R.H. (1992) Reduction of alpha-naphthylisothiocyanate-induced hepatotoxicity by recombinant human hepatocyte growth factor. *Endocrinology* **131**, 2540-2544
128. Matsuda, Y., Matsumoto, K., Ichida, T., and Nakamura, T. (1995) Hepatocyte growth factor suppresses the onset of liver cirrhosis and abrogates lethal hepatic dysfunction in rats. *J. Biochem.* **118**, 643-649
129. Rong, S., Bodescot, M., Blair, D., Dunn, J., Nakamura, T., Mizuno, K., Park, M., Chan, A.M.-L., Aaronson, S.A., and Vande Woude, G.F. (1992) Tumorigenicity of the met protooncogene and the gene for hepatocyte growth factor. *Mol. Cell. Biol.* **12**, 5152-5158
130. Kanda, H., Tajima, H., Lee, G.-H., Nomura, K., Ohtake, K., Matsumoto, K., Nakamura, T., and Kitagawa, T. (1993) Hepatocyte growth factor immortalized mouse liver epithelial cells. *Oncogene* **8**, 3047-3053
131. Bellusci, S., Moens, G., Gaudino, G., Comoglio, P.M., Nakamura, T., Thiery, J.P., and Joanneau, J. (1994) Creation of hepatocyte growth factor/scatter factor autocrine loop in carcinoma cells induces invasive properties associated with increased tumorigenicity. *Oncogene* **9**, 1091-1099
132. Johnson, M., Koukoulis, G., Kochher, K., Kubo, C., Nakamura, T., and Iyer, A. (1995) Selective tumorigenesis in non-parenchymal liver epithelial cell lines by hepatocyte growth factor transfection. *Cancer Lett.* **98**, 37-48
133. Ferracini, R., Direnzo, M.F., Scotlaudi, K., Baldini, N., Olivero, M., Lollini, P., Cremona, O., Campanacci, M., and Comoglio, P.M. (1995) The Met/HGF receptor is over-expressed in human osteosarcomas and is activated either a paracrine or an autocrine circuit. *Oncogene* **10**, 739-749
134. Picard, O., Rolland, Y., and Poupon, M.F. (1986) Fibroblast-dependent tumorigenicity of cells in nude-mice: Implication for implantation of metastases. *Cancer Cells* **46**, 3290-3294
135. Matsumoto, K., Horikoshi, M., Rikimaru, K., and Enomoto, S. (1989) A study of an *in vitro* model for invasion of oral squamous cell carcinoma. *J. Oral Pathol. Med.* **18**, 498-501
136. Camps, J.L., Chang, S., Hsu, T.C., Freeman, M.R., Hong, S., Zhau, H.E., von Eschenbach, A.C., and Chung, L.W.K. (1990) Fibroblast-mediated acceleration of human epithelial tumor growth *in vivo*. *Proc. Natl. Acad. Sci. USA* **87**, 75-79
137. Grey, A.-M., Schor, A.M., Rushton, G., Ellis, I., and Schor, S. (1989) Purification of the migration stimulating factor produced by fetal and breast cancer patient fibroblasts. *Proc. Natl. Acad. Sci. USA* **86**, 2438-2442
138. Seslar, S., Nakamura, T., and Byers, S. (1993) Regulation of fibroblast hepatocyte growth factor/scatter factor expression by human breast carcinoma cells and peptide growth factors. *Cancer Res.* **53**, 1233-1238
139. Rosen, E.M., Joseph, A., Jin, L., Rockwell, S., Elias, J.A., Knesel, J., Wines, J., McClellan, J., Kluger, M.J., Goldberg, I.D., and Zitnik, R. (1994) Regulation of scatter factor production via a soluble inducing factor. *J. Cell Biol.* **127**, 225-234