

# A Disulfide Bridge Mediated by Cysteine 574 Is Formed in the Dimer of the 70-kDa Heat Shock Protein

Takayuki K. Nemoto<sup>1,\*</sup>, Yutaka Fukuma<sup>1</sup>, Hideaki Itoh<sup>2</sup>, Takashi Takagi<sup>3</sup> and Toshio Ono<sup>1</sup>

<sup>1</sup>Division of Oral Molecular Biology, Department of Developmental and Reconstructive Medicine, Course of Medical and Dental Sciences, Nagasaki University Graduate School of Biomedical Sciences, 1-7-1 Sakamoto, Nagasaki 852-8588; <sup>2</sup>Department of Material-Process Engineering and Applied Chemistry for Environment, Akita University Faculty of Engineering and Resource Science, 1-1 Tegata Gakuen Town, Akita City 010-8502; and <sup>3</sup>Department of Developmental Biology and Neurosciences, Graduate School of Life Sciences, Tohoku University, Sendai 980-8578

Received November 15, 2005; accepted February 6, 2006

**The 70-kDa heat shock protein (Hsp70) is predominantly present intracellularly as a monomer, but a small population is converted to dimers and oligomers under certain conditions. In the present study, we investigated the dimeric structure of human inducible Hsp70. As reported earlier, the C-terminal client-binding domain (amino acids 382–641) was required for the dimerization. A 40-amino acid deletion in the client-binding domain from either the N-terminus or C-terminus greatly enhanced the dimerization potential of Hsp70. Limited proteolysis indicated that the dimer formed through truncation from the C-terminus had a conformation similar to that of the non-truncated form. Truncation experiments demonstrated that the client-binding sub-domain (amino acids 382–520) with its adjacent region up to amino acid 541 was not sufficient for the dimerization but that the region up to amino acid 561 was sufficient. Interestingly, the dimer formed through truncation from the C-terminus acquired a homomeric disulfide bridge at Cys574.**

**Key words:** client-binding, dimer, disulfide bridge, lid, Hsp70.

Abbreviations: Hsp, heat shock protein; Hsp70, inducible 70-kDa heat shock protein; Hsc70, 70-kDa heat shock cognate protein; Hsp90, 90-kDa heat shock protein; Hsp60/GroEL 60-kDa heat shock protein; mAb, monoclonal antibody; PAGE, polyacrylamide gel electrophoresis; SDS, sodium dodecylsulfate; and TPCK, *N*<sup>ε</sup>-acetyl-L-tosyl-L-phenylalanine chloromethyl ketone.

Heat shock proteins (Hsps) essentially exist in all organisms. Reflecting their important and general roles as molecular chaperones, their amino acid sequences are highly conserved in every Hsp family, even among eukaryotic and prokaryotic organisms. Among the three major Hsps in the high-molecular-weight class, *i.e.*, 60-kDa, 70-kDa and 90-kDa heat shock proteins (Hsp60/GroEL, Hsp70 and Hsp90, respectively), the architecture of Hsp60/GroEL oligomers complexed with Hsp10/GroES has been extensively investigated (1). Hsp90 forms a dimer through its C-terminal region (2, 3). Based on the results of X-ray crystallographic studies (4–7), Prodromou *et al.* (4, 5) and Meyer *et al.* (7) proposed the molecular clamp model, in which a dimer of Hsp90 traps a client protein between the N-terminal domains or middle domains. On the other hand, Hsp70 predominantly exists as a monomer (8). Nevertheless, it has also been reported that Hsp70 and DnaK, a bacterial ortholog of Hsp70, are dimerized and further oligomerized in a concentration-dependent manner (9). Moreover, ATP induces the dissociation of dimers and oligomers of Hsp70 (9–12). Therefore, it has been suggested that the monomer-dimer conversion is closely related to the chaperone function of Hsp70.

Hsp70 is composed of at least two structural domains (13). The 44-kDa N-terminal domain of bovine brain Hsp70 contains a nucleotide-binding site and possesses weak ATP-hydrolytic activity (13). The 30-kDa C-terminal domain of DnaK has the capacity to bind unfolded polypeptides (14, 15). Although the three-dimensional structure of the full-length forms of Hsp70 remains unknown, it is obvious that the activities of these domains are co-operative, because the client-binding activity of Hsp70 is relatively high in the ADP-bound state, but is reduced in the ATP-binding state (9, 16), and the binding of client peptides leads to the stimulation of ATP hydrolysis by Hsp70 (17, 18).

There is some evidence that the 30-kDa C-terminal client-binding domain is responsible for the dimerization of 70-kDa heat shock cognate protein Hsc70 (19, 20). It has been reported that the 30-kDa C-terminal domain can be divided into N-terminal 18-kDa and C-terminal 10-kDa sub-domains. The N-terminal sub-domain (designated as the client-binding sub-domain) contains the binding site for client peptides (21), and the C-terminal sub-domain (designated as the lid sub-domain) appears to function as a lid that stabilizes the client peptide bound to the client-binding sub-domain. Fouchaq *et al.* (20) reported that the 17-kDa client-binding sub-domain of Hsc70 could be oligomerized. In contrast, Chou *et al.* (22) proposed that the 10-kDa lid sub-domain of Hsc70 is responsible

\*To whom correspondence should be addressed. Tel: +81-95-849-7640, Fax: +81-95-849-7642, E-mail: tnemoto@net.nagasaki-u.ac.jp

for the dimerization and oligomerization. Therefore, it remains unknown as to whether the client-binding sub-domain or the lid sub-domain is critical for the dimerization.


In the present study, we expressed the full-length form of human Hsp70 (hHsp70) and the C-terminal client-binding domain, and their truncated forms to determine the region responsible for the dimerization and oligomerization of Hsp70. We found that several forms of the truncated C-terminal domain acquired a dimerization potential that was much higher than that of the client-binding domain itself or the full-length form. Unexpectedly, some of the dimeric complexes possessed disulfide bridge(s) connecting the Cys574 residues of the molecules.

#### EXPERIMENTAL PROCEDURES

**Materials**—The materials used and their sources were as follows: Expression vector pTrcHis TOPO and TOP10 competent cells, Invitrogen (Carlsbad, CA, USA); low-molecular-weight markers and peptide markers, Amersham Biosciences (Piscataway, NJ, USA); kaleidoscope prestained molecular standards, Bio-Rad (Richmond, CA, USA); restriction enzymes and DNA-modifying enzymes, Nippon Gene (Tokyo, Japan); Talon metal affinity resin, Clontech Laboratories Inc. (Palo Alto, CA, USA); trypsin (5,200 USP units/mg protein) and *N*<sup>ε</sup>-acetyl-L-tosyl-L-phenylalanine chloromethyl ketone (TPCK), Sigma (St. Louis, MO, USA.); anti-histidine hexamer tag monoclonal antibodies (anti-His tag mAb), Qiagen Inc. (Chatworth, CA, USA); and alkaline-phosphatase-conjugated rabbit anti-mouse Ig(G+A+M) antibodies, Zymed Laboratories Inc. (San Francisco, CA, USA).

**Construction of Bacterial Expression Plasmids**—All recombinant proteins used in this study were encoded by the pTrcHis TOPO vector (Invitrogen), and therefore possessed a 35-amino acid tag carrying a histidine-hexamer at the N-terminus. cDNA encoding the full-length form of hHsp70 was obtained by RT-PCR from *HeLa* cell mRNA with the aid of its cDNA sequence (23, 24). For construction of plasmids carrying the full-length form of hHsp70, the ATP-binding domain (amino acids 1–381), the client-binding domain (amino acids 382–641), and truncated forms of the client-binding domain (Fig. 1), DNA fragments encoding the corresponding regions were amplified by PCR, and then directly inserted into the pTrcHis TOPO expression vector in frame according to the manufacturer's recommendation. TOP10 cells were transformed with the plasmids and selected on Luria-Bertani medium-agar plates containing 50 µg/ml of ampicillin. The insertion and orientation of the DNA fragments were confirmed by conducting Hot Star PCR (Qiagen Inc., Chatsworth, CA, USA).

**Expression and Purification of Recombinant Proteins**—After overnight cultivation of transformed bacteria at 37°C, recombinant proteins were expressed at 30°C for 4 h in the presence of 0.2 mM isopropyl-β-D-thiogalactopyranoside. The proteins were purified by affinity chromatography on a Talon affinity column according to the manufacturer's protocol except that 10 mM imidazole and 10 µg/ml of leupeptin were added to the lysis buffer (20 mM Tris-HCl, pH 8.0, and 0.1 M NaCl containing 0.5 mg/ml lysozyme). After extensive washing with washing



**Fig. 1. Dimerization of the full-length form of hHsp70.** The full-length form of recombinant hHsp70 was subjected to SDS-PAGE at a polyacrylamide concentration of 7.5% under non-reducing and reducing conditions (a) or PAGE at a polyacrylamide concentration of 7.5% under non-denaturing conditions (b). Lane 1, 0.3 µg; lane 2, 1 µg; lane 3, 3 µg; and lane 4, 10 µg. The molecular markers (Mk) used were low-molecular-weight markers and kaleidoscope markers (a), and markers for native PAGE (b). M, monomer; D, dimer; and T, trimer.

buffer (20 mM Tris-HCl, pH 8.0, containing 0.1 M NaCl and 10 mM imidazole) to remove non-adsorbed proteins, bound proteins were eluted with 0.1 M imidazole (pH 8.0) containing 10% (v/v) glycerol. The purified proteins were used immediately or stored at –80°C until used.

**Sodium Dodecyl Sulfate–Polyacrylamide Gel Electrophoresis (SDS-PAGE)**—Electrophoresis was performed in the presence of 0.1% (w/v) SDS according to Laemmli (25). Unless otherwise stated, a polyacrylamide concentration of 12.5% (w/v) was used. For separation of small molecular species such as proteolytic fragments, SDS-PAGE with the Tris-Tricine system at a polyacrylamide concentration of 10% (w/v) was employed (26). Samples were denatured at 94°C for 10 min in the sample buffer in the presence or absence of 10% (v/v) 2-mercaptoethanol. Low-molecular-weight markers (Amersham Biosciences, Piscataway, NJ, USA) and peptide markers (Amersham Biosciences) were used as references.

**Polyacrylamide Gel Electrophoresis under Nondenaturing Conditions**—In order to determine molecular configurations, samples were subjected to PAGE under nondenaturing conditions (27). Unless otherwise stated, a polyacrylamide concentration of 7.5% (w/v) was used. Electrophoresis was performed at a low voltage (50 V) at room temperature to avoid elevation of the gel temperature. The separated proteins were stained with Coomassie brilliant blue. Ovalbumin (45 kDa), bovine serum albumin (66 kDa as monomer and 132 kDa as dimer), and catalase (240 kDa) were used as molecular references.

**Two-Dimensional PAGE**—Purified proteins were separated by two-dimensional PAGE. A polyacrylamide concentration of 7.5% (w/v) was used for the first dimension PAGE under non-denaturing conditions, and one of 11% (w/v) was used for the second dimension SDS-PAGE under reducing or non-reducing conditions.

**Immunoblotting Analysis**—Recombinant proteins in bacterial lysates separated by PAGE under non-denaturing conditions were subjected to immunoblotting analysis as described previously (28). An anti-histidine tag mAb (Qiagen) was used as the first antibody at 1 µg/ml, and the second antibody was used at 1/3,000 dilution. The blotted proteins were finally visualized by incubation with 5-bromo-4-chloro-3-indolyl phosphate and nitro blue tetrazolium (Nakarai, Kyoto, Japan) at room temperature for 10–20 min.

**Size-Exclusion Gel Chromatography**—Size-exclusion gel chromatography was performed on a Sephacryl S100HR (Amersham Biosciences, 1.8 × 36 cm) or Sephacryl S300 (Amersham Biosciences, 1.0 × 90 cm) column equilibrated with 20 mM Tris-HCl (pH 8.0) containing 1 mM EDTA, 50 mM NaCl and 0.05% (w/v) NaN<sub>3</sub>. Blue dextran 2000 (void volume), catalase (232 kDa, Stokes radius = 52.2 Å), aldolase (158 kDa, 48.1 Å), bovine serum albumin (66 kDa, 35.5 Å), and ovalbumin (45 kDa, 30.5 Å) were used as molecular size references. Forty-drop fractions (2 ml) were collected from the Sephacryl S100HR column, and 20-drop ones (1 ml) from the Sephacryl S300 column.

**N-Terminal Amino Acid Sequencing**—In order to determine the difference in molecular configuration, we partially cleaved the client-binding domain and its truncated forms with trypsin treated with TPCK. The recombinant proteins in the elution buffer were incubated at 30°C with 3 ng of TPCK-trypsin for 0–12 h or with 0–300 ng of TPCK-trypsin at 30°C for 12 h. Following electrophoretic separation of the proteolytic fragments by use of a Tris-Tricine system, the major tryptic fragments were transferred to a polyvinylidene difluoride membrane (Bio-Rad). After having been stained with Coomassie brilliant blue, the bands were excised and the respective materials directly sequenced with a model 492A protein sequencer (PE Biosystems).

**Protein Concentrations**—Protein concentrations were determined by the bicinchoninic acid method using bovine serum albumin as a standard (Pierce, Rockford, IL, USA).

## RESULTS

**Dimerization of the Full-Length Form of hHsp70**—Initially we expressed the full-length form of hHsp70 in a bacterial expression system. Upon SDS-PAGE under reducing conditions (Fig. 1a, right, lanes 1–4), the purified protein migrated as a single band to the position corresponding to 75 kDa, which was slightly larger than the value of endogenous hHsp70 (calculated molecular mass, 70,053; Ref. 23) due to the 35-amino acid tag at the N-terminus.

Even if large amounts of the purified sample were loaded onto an SDS-PAGE gel, no additional band appeared under reducing conditions (Fig. 1a, right). In contrast, under non-reducing conditions, additional bands appeared at positions corresponding to high molecular masses


(Fig. 1a, left, lanes 3 and 4). Among them, the most predominant band migrated to the 150-kDa position, indicating the covalently-associated dimeric configuration of 75-kDa hHsp70.

Polyacrylamide gel electrophoresis under non-denaturing conditions gave one major band (Fig. 1b, lanes 1 and 2), and a second, slow-migrating species appeared at higher protein concentrations (lanes 3 and 4). At the highest protein concentration, a third band appeared (lane 4). Taken the migration positions and the results of previous studies (11, 12) into consideration, the three bands may correspond to the monomer, dimer and trimer.

We noticed that the bands presumably corresponding to the monomer, dimer and trimer of hHsp70 migrated slower than expected, compared with the molecular mass references (bovine serum albumin, ovalbumin and catalase) used in this study. This discrepancy seemed to be due to the molecular size references used for PAGE under non-denaturing conditions, because the isoelectric point (pI) of Hsp70 (pI = 5.5–5.6; Ref. 8) is more basic than that of ovalbumin (pI = 4.6) or bovine serum albumin (pI = 4.8; Ref. 27).

**Charge Properties of hHsp70 Affecting Its Electrophoretic Mobility on PAGE under Non-Denaturing Conditions**—The net charges of the N-terminal ATP-binding and C-terminal client-binding domains simply calculated from the amino acid composition are –1 and –10, respectively (24). Hence, we suspected that the slow migration of hHsp70 was primarily caused by the charge of the N-terminal ATP-binding domain. To examine this, we expressed the N-terminal ATP-binding (amino acids 1–381), and the two C-terminal client-binding domains (amino acids 382–641), and the two sub-domains (amino acids 382–519 and 520–641) of the client-binding domain (Fig. 2a). Upon SDS-PAGE (Fig. 2b, left), they each migrated as a single band with the exception of that of the lid sub-domain (hHsp70 520–641), which contained a 22-kDa species as a minor constituent (lane 5). The migration of hHsp70 1–381 was much slower than that of hHsp70 382–641 (Fig. 2b, right, compare lanes 2 and 3). This retardation of hHsp70 1–381 should reflect either the dimerization or the charge. To examine these two possibilities, hHsp70 1–381 was separated by size exclusion gel chromatography on a Sephacryl S100HR column. As a result, hHsp70 1–381 was eluted as a single peak corresponding to a molecular size smaller than that of 45-kDa ovalbumin (Fig. 2c). Therefore, we concluded that hHsp70 was monomeric, but migrated slowly on PAGE under non-denaturing conditions because of its less acidic charge. To the contrary, the main band of hHsp70 382–641 migrated fast on PAGE under non-denaturing conditions, reflecting its acidic property (Fig. 2b, right, lane 3).

**Dimerization of the C-Terminal Client-Binding Domain**—Polyacrylamide gel electrophoresis under non-denaturing conditions revealed the presence of slow-migrating species in the hHsp70 382–641 and 520–641 preparations in addition to the respective fast-migrating major ones (Fig. 2b, right, arrowheads). We suspected that the fast- and slow-migrating species corresponded to the monomer and dimer, respectively. In fact, two-dimensional PAGE of hHsp70 382–641 confirmed the existence of monomeric, dimeric, trimeric and presumably oligomeric



**Fig. 2. Dimerization of the client-binding domain and its truncated forms.** (a) The ATP-binding domain (amino acids 1–381), the client-binding domain (amino acids 382–641), and two sub-domains of the client-binding domain are schematically illustrated. (b) Recombinant proteins were separated by SDS-PAGE under reducing conditions (left) or by PAGE under non-denaturing conditions (right). Protein samples loaded for SDS-PAGE under reducing conditions (0.5  $\mu$ g) and for PAGE under non-denaturing conditions (1.5  $\mu$ g) in lanes 1–5 correspond to numbers 1–5 in panel “a.” Mk, molecular markers. Arrowheads indicate minor species migrating slower than the major species (right, lanes 1, 3 and 5). An asterisk indicates a 22-kDa band (see text). (c) hHsp70 1–381 (1.3 mg/2 ml) was separated by size-exclusion gel chromatography on a Sephacryl S100HR column (upper panel) as described under


“EXPERIMENTAL PROCEDURES.” Ovalbumin (45 kDa, Stokes’ radius = 30.5 Å) was applied to the column separately as a size reference. hHsp70 1–381 (lane P, 1.5  $\mu$ g) or aliquots (10  $\mu$ l) of fractions 27–36 obtained on chromatography were separated by SDS-PAGE under non-reducing conditions (middle panel) or by PAGE under non-denaturing conditions (lower panel). (d and e) Twenty-five micrograms of hHsp70 382–641 (d) and 520–641 (e), respectively, was separated by two-dimensional PAGE as described under “EXPERIMENTAL PROCEDURES.” The second dimension SDS-PAGE was performed under reducing conditions. The apparent molecular masses of recombinant proteins are indicated on the right. Asterisks indicate a 22-kDa band (see text). M, D, T and O represent monomer, dimer, trimer and oligomer, respectively.

species (Fig. 2d). In contrast, the slow-migrating species of hHsp70 520–641 was not a dimer, but corresponded to the 22-kDa band, which was co-purified with the major 24-kDa band (Fig. 2e, asterisks). The 22-kDa species seemed to be a

degradation product of the 24-kDa hHsp70 520–641. These findings confirmed the results of the previous study (19) demonstrating that the dimerization as well as the oligomerization of Hsp70 was mediated through the C-terminal

client-binding domain. Moreover, the results in Fig. 2 show that the two sub-domains of the client-binding domain dimerized no more.

We here produced a series of N- and C-terminally truncated forms of the C-terminal client-binding domain (Fig. 3a). They were purified to near homogeneity, as shown on SDS-PAGE under reducing conditions (Fig. 3b, left). However, upon SDS-PAGE under non-reducing conditions, large-molecular-weight bands (63 or 58 kDa) appeared for hHsp70 382–601 (Fig. 3b, right, lane 3) and hHsp70 382–581 (lane 4). Accompanying this, the 33- and 30-kDa species decreased in amount. Hence, the high-molecular-mass bands appeared to be dimeric forms mediated by a disulfide bond(s).



**Fig. 3. Dimerization of the client-binding domain and its truncated forms.** (a) The client-binding domain (amino acids 382–641) and its truncated forms are presented. The states of the proteins (M, monomer and D, dimer) and the formation of disulfide bridges (S-S) observed in panels “b” and “c” are summarized on the right. Recombinant proteins were separated by SDS-PAGE under reducing and non-reducing conditions (b) or by PAGE under non-denaturing conditions (c). Protein samples loaded for SDS-PAGE (0.5  $\mu$ g) and for PAGE under non-denaturing conditions (1.5  $\mu$ g) in lanes 1–9 in panels “b” and “c,” respectively, correspond to numbers 1–9 in panel “a.” Mk, molecular markers. M, D and T on the left of panel “c” represent monomer, dimer and trimer, respectively.

We also examined the electrophoretic motility of truncated forms of the client-binding domain on PAGE under non-denaturing conditions (Fig. 3c). The C-terminal truncation of 20 amino acids (hHsp70 382–621) produced smear bands in the dimer and trimer regions (lane 2). Successive truncations (hHsp70 382–601 and 382–581) substantially increased the proportion of the slower-migrating species (lanes 3 and 4). Further truncation (hHsp70 382–519) reversed this trend (lane 5).


Next we examined the effect of N-terminal truncation of the client-binding domain on the dimerization. Eighteen-amino acid truncation (hHsp70 400–641) had little effect (Fig. 3c, lane 6), but further truncations (hHsp70 420–641 and hHsp70 440–641) caused a significant increase in the amount of the slow-migrating species (lanes 7 and 8) corresponding to the dimer. Again, further truncation (hHsp70 520–641) reversed this trend (lane 9).

In order to further characterize the molecular configurations, we subjected hHsp70 382–601 to size-exclusion gel chromatography on Sephacryl S100HR. hHsp70 382–601 was split into two peaks (Fig. 4a, top). Polyacrylamide gel electrophoresis under non-denaturing conditions revealed that the dimer and monomer were eluted in the first and second peaks, respectively (Fig. 4a, middle). A trimer was also detected in fractions 24 and 25 (Fig. 4a, middle). Upon SDS-PAGE under non-reducing conditions, the first peak contained the 63-kDa covalently-associated dimeric species (Fig. 4a, low). Interestingly, the 33-kDa species was also found in the first peak, which indicated that this species existed as a dimer, although it did not possess a disulfide bridge.

hHsp70 382–601 was also subjected to two-dimensional PAGE, with the first dimension PAGE under non-denaturing conditions and second dimension SDS-PAGE under non-reducing conditions (Fig. 4b). As a result, five major species were recognized: a monomer with a molecular mass of 33 kDa, two dimers with molecular masses of 33 and 63 kDa, and two trimers molecular masses of 33 and 63 kDa. These results demonstrated that a disulfide bridge was not a prerequisite for the dimerization. Noticeably, the dimer with a disulfide bridge migrated slightly faster than that without the bridge on PAGE under non-denaturing conditions, indicating a difference in their configurations (Fig. 4b).

One might argue that these dimers and trimers were artificially formed through nonspecific interactions of the recombinant proteins. In fact, an X-ray crystallographic study on the C-terminal 10-kDa lid sub-domain of rat Hsc70 revealed the packaging of 48 molecules in a unit cell. To address this issue, we performed a mixing experiment on the lysates containing hHsp70 382–581 and 382–601. If a dimeric interaction randomly occurred between any free sulfhydryl groups, twice as much of a heterodimer of the two proteins as each homo-dimer should be formed. However, the hetero-dimer was scarcely formed (Fig. 5, an arrow), indicating its specific interactions.

We also performed immunoblotting analyses of truncated forms of hHsp70 in the bacterial lysates. As a result, small populations of hHsp70 382–641, 382–621, 382–601 and 382–581 were found to possess a disulfide bridge(s) in the lysates (Fig. 6a, right arrowheads). However, the immunoblotting profile on PAGE under non-denaturing conditions (Fig. 6b) was indistinguishable from that of




**Fig. 4. Size-exclusion gel chromatography and two-dimensional PAGE of hHsp70 382–601.** (a) hHsp70 382–601 (3 mg/2 ml) was separated by size-exclusion gel chromatography on a Sephacryl S100HR column (upper panel) as described under “EXPERIMENTAL PROCEDURES.” Size markers [Blue dextran 2000 (void volume), bovine serum albumin (66 kDa, Stokes’ radius = 35.5 Å), and ovalbumin (45 kDa, Stokes’ radius = 30.5 Å)] were subjected to chromatography separately. An aliquot (7.5  $\mu$ l) of each fraction was subjected to PAGE under non-denaturing conditions (middle panel) or SDS-PAGE under non-reducing conditions (lower panel). P, purified hHsp70 382–601 loaded on the chromatography column. Mk, molecular markers. M, D and T on the left represent monomer, dimer and trimer, respectively. (b) Two-dimensional PAGE of hHsp70 382–601 (20  $\mu$ g). As references for first and second dimensional PAGE, hHsp70 382–601 (2  $\mu$ g) was run in parallel. The apparent molecular masses of recombinant proteins are indicated on the right. Mk, marker proteins. M, D and T represent monomer, dimer and trimer, respectively.

Coomassie-stained purified proteins (Fig. 3c), which clearly demonstrated that dimers of hHsp70 were already formed in the lysates even without a disulfide bridge.

Taking the results obtained in the above studies together, we concluded that (i) the dimer-forming activity of hHsp70 resided in the C-terminal domain, (ii) the activity was significantly enhanced by deletion of either N-terminal or C-terminal amino acids ( $\geq 38$ –40 residues) of the domain, (iii) neither the client-binding sub-domain (amino acids 382–519) nor the lid sub-domain (amino acids 522–641) was sufficient for dimer formation, (iv) the dimer

of the C-terminally truncated forms (hHsp70 382–581 and 382–601) contained disulfide bond(s), and (v) the disulfide bridge was not a prerequisite for dimerization. We suspected that free sulfhydryl groups of cysteines were located close to each other in the dimeric configuration.


*Characterization of the Dimeric Complex Formed with the N- and C-Terminal Truncations*—We investigated the structural integration of the truncated proteins by limited proteolysis. Were the dimeric configurations of the truncated proteins similar to or considerably different from that of the client-binding domain? As shown in Fig. 7a, the



**Fig. 5. Time course of the formation of the disulfide bridge(s).** A bacterial lysate containing hHsp70 382–581 or hHsp70 382–601, or an equal mixture of them was prepared. Immediately after the lysate preparation or after 1 or 5 days of being kept at 4°C, the lysates (1 ml) were mixed with Talon affinity resin [50  $\mu$ l as a 50% (v/v) suspension]. After washing of the resin with washing buffer (1 ml) five times, the bound proteins were extracted with 50  $\mu$ l of 100 mM imidazole (pH 8.0) containing 10% (v/v) glycerol. Then, aliquots (7.5  $\mu$ l) were subjected to SDS-PAGE under non-reducing conditions. M, monomer; and D, dimer. Mk, molecular markers. An arrow indicates a hetero-dimer.

time courses of proteolysis of hHsp70 382–621 and hHsp70 400–641 were similar to each other and to the profile of the client-binding domain (hHsp70 382–641). Further truncations at the C-terminus indicated that the proteolytic profiles of hHsp70 382–601 and 382–581 were similar to each other and relatively well resembled the profile of hHsp70 382–641. In contrast, the N-terminally-truncated forms (hHsp70 420–641 and hHsp70 440–641) were highly susceptible to trypsin and their profiles were considerably different from the hHsp70 382–641 profile (Fig. 7a). This finding was confirmed by the results for the concentration-dependent proteolytic profiles of the three recombinant proteins: The proteolytic profile of hHsp70 382–641 was similar to that of hHsp70 382–601, and that of hHsp70 420–641 was obviously different from that of hHsp70 382–641 or hHsp70 382–601 (Fig. 7, b and c).


The results of limited proteolysis and subsequent N-terminal amino acid sequencing of tryptic fragments (Table 1) are schematically summarized in Fig. 7c. Noticeably, hHsp70 382–641 as well as hHsp70 382–601 was preferentially attacked at sites in the C-terminal lid sub-domain, but hHsp70 420–641 was preferentially attacked at sites in the N-terminal client-binding sub-domain (Table 1 and Fig. 7c). Importantly, the alteration of the proteolytic profiles from hHsp70 382–621 to hHsp70 382–601 and that from hHsp70 400–641 to hHsp70 420–641 (Fig. 7) were accompanied by the acquisition of potent dimer-forming activity (Fig. 3). Accordingly, we tentatively classified the truncated proteins into three types: (i) hHsp70 382–641, 382–621 and 400–641, which were predominantly monomeric; (ii) hHsp70 382–581 and 382–601, which were predominantly dimeric, and had a configuration similar to that of the first type; and (iii) hHsp70 420–641 and 440–641, which were predominantly dimeric,



**Fig. 6. Immunoblotting analysis of the client-binding domain and its truncated forms.** Bacterial lysates containing recombinant proteins (2.5  $\mu$ l) were subjected to SDS-PAGE under reducing and non-reducing conditions (a), or PAGE under non-reducing conditions (b). The separated proteins were immunoblotted with an anti-His tag monoclonal antibodies as described under “EXPERIMENTAL PROCEDURES.” Mk, molecular markers. On PAGE under non-denaturing conditions, the markers in the polyacrylamide gel were stained with Coomassie Brilliant Blue. M, monomer; and D, dimer. The proteins loaded in lanes 1–9 are identical to those in Fig. 3.

and had a structure that was considerably different from those of other types.

**Disulfide Bridge(s) of the Full-Length Form of hHsp70—**The C-terminal truncation of the client-binding domain induced efficient conversion from a monomer to a dimer, and the resulting dimer possessed a disulfide bridge(s). The disulfide bridge(s) appeared to be also present in the full-length form of hHsp70 (Fig. 1). For the full-length form, although the population of the species that possessed the disulfide bond appeared to be limited, the proportion in the dimer should be relatively high if the small population of the dimeric species is considered (Fig. 1b). However, there were additional high molecular-mass bands, the subunit compositions for which could not be readily determined. Hence, we performed a truncation experiment on the full-length form similar to that performed on the client-binding domain. We expressed a series of C-terminal truncation forms of hHsp70 (Fig. 8a). Sodium dodecylsulfate-PAGE under non-reducing conditions revealed the existence of a high molecular species (Fig. 8b, left, lanes 1–3). Moreover, consistent with the result for the client-binding domain (Fig. 3b), C-terminal truncation of 40 or 60 amino acids caused more efficient dimer formation (Fig. 8b, left, lanes 2 and 3). Noticeably, the higher molecular species of the respective proteins migrated slowest for the full-length form (lane 1), moderately for hHsp70 1–601 (lane 2), and fastest for hHsp70 1–581 (lane 3), reflecting their molecular masses. This finding confirms that the high-molecular species of the



**Fig. 7. Limited proteolysis of the client-binding domain and its truncated forms.** (a) The client-binding domain (hHsp70 382–641) and its truncated forms (5 µg/50 µl) were incubated without trypsin (lane 1) or with 0.2 µg TPCK-treated trypsin at 30°C for 1.5 h (lane 2) or 10 h (lane 3). Proteolytic fragments were separated by Tris-Tricine SDS-PAGE under reducing conditions. Mk, molecular markers. (b) hHsp70 382–641, hHsp70 382–601 and hHsp70 420–641 (12.5 µg/50 µl) were incubated at 30°C for 12 h without (lane 1), or with 3 ng (lane 2), 10 ng (lane 3), 30 ng (lane 4), 0.1 µg (lane 5), 0.3 µg (lane 6), or 1 µg of TPCK-trypsin. Aliquots (5 µg) were separated on SDS-PAGE gels under reducing conditions by use of a Tris-Tricine system. Other aliquots were separated in the same way and used for N-terminal amino acid sequencing as described under “EXPERIMENTAL PROCEDURES.” Mk, molecular markers. The apparent molecular masses of the major proteolytic fragments are indicated in panel “b.” (c) Tryptic fragments of the three recombinant proteins are schematically arranged based on their profiles and N-terminal sequences. The molecular masses of the fragments were estimated from the results shown in panel “b.” The N-terminal amino acids are defined in Table 1. The hatched areas represent the 35-amino acid N-terminal tag.

full-length form migrating to the 150-kDa position was a dimer of hHsp70 mediated by a disulfide bridge(s). The recombinant protein with the C-terminal 80 amino acids deleted (hHsp70 1–561) had completely lost the bridge (lane 4).

We subjected the recombinant proteins to size exclusion chromatography on Sephacryl S300 (Fig. 8c). Under the conditions employed, monomeric species were eluted in fractions 46–48. Gradual retardation of migration from fraction 46 for hHsp70 1–641 and hHsp70 1–581 to fraction 48 for hHsp70 1–561 and hHsp70 1–541 should reflect successive C-terminal truncations. The full length form was mainly eluted as a monomer, with a small population around fraction 40, the position of the dimer (Fig. 8c). In

contrast, hHsp70 1–581 and hHsp70 1–561 were each split into two peaks, i.e., dimer (fractions 40–42) and monomer (fractions 46–48). The first peak of hHsp70 1–581 migrated to the 140-kDa position on SDS-PAGE gel under non-reducing conditions, but that of hHsp70 1–561 migrated to the 60-kDa position, indicating that the hHsp70 1–561 dimer did not possess a disulfide bridge. hHsp70 1–541 (Fig. 8c) and hHsp70 1–520 (data not shown) were each eluted as a monomer.

Taken together, these findings indicate that efficient formation of an hHsp70 dimer and disulfide bond(s) induced by C-terminal truncation were not specific to the client-binding domain, but also occurred in the case of the full-length form. Importantly, hHsp70 1–561 formed



Table 1. N-terminal sequences of tryptic fragments of truncated hHsp70.

Origin	Fragment (kDa)	Amino acids detected on sequencing	Corresponding sequence
382–641	35 (intact)	n.d.	
	33	GGSHHHHH	m <sub>-35</sub>   g <sub>-34</sub> gshhhhh
	30	DLYDDDD	r <sub>-13</sub>   g <sub>-12</sub> lydddd
	25	DLYDDDDKDP	r <sub>-13</sub>   g <sub>-12</sub> lydddskdp
	21 <sup>a</sup>	GVPQIEVTFD	R <sub>469</sub>   G <sub>470</sub> QIEVTFD
		MVQEAEKYKA	R <sub>517</sub>   M <sub>518</sub> QEAEKYKA
	15	GVPQIEVT	R <sub>469</sub>   G <sub>470</sub> QIEVTFD
	14	GVPQIEVT (major <sup>b</sup> )	R <sub>469</sub>   G <sub>470</sub> QIEVTFD
382–601	33 (intact)	n.d.	
	27	DYLDKDDK	r <sub>-13</sub>   g <sub>-12</sub> lydddsk
	25	GGSHHHHH	m <sub>-35</sub>   g <sub>-34</sub> gshhhhh
420–641	33 (intact)	n.d.	
	23	GVPQIEVTFD	R <sub>469</sub>   G <sub>470</sub> VPQIEVTFD
	21 <sup>a</sup>	GVPQIEVTFD	R <sub>469</sub>   G <sub>470</sub> VPQIEVTFD
		MVQEAEKYKA	R <sub>517</sub>   M <sub>518</sub> VQEAEKYKA

Deduced sequences given in capital and small letters were derived from hHsp70 and the tagged sequence, respectively. <sup>a</sup>A mixture of two fragments. <sup>b</sup>N-terminal sequence of the major species was determined. n.d., not determined.

a dimer, although it did not contain a disulfide bond. This finding confirmed that the disulfide bridge is not a prerequisite for the dimer formation.

#### DISCUSSION

In the present study, we investigated the dimeric structure of hHsp70. Although a small population of hHsp70 was present as a dimer, the dimer was more efficiently formed when the C-terminal amino acids (40–80 residues) were deleted from the client-binding domain as well as the full-length form. Similarly, Ohno *et al.* (29) reported that mouse Hsp70 1–615, from which the C-terminal 26 residues had been deleted, tended to form a dimer and oligomers more efficiently than the full-length form did. There are only two cysteine residues, *i.e.*, Cys574 and Cys603, in the client-binding domain of hHsp70. Because hHsp70 382–581 as well as hHsp70 382–601, from which Cys603 has been lost, formed a disulfide bridge, it is reasonable to conclude that Cys574 of a molecule was associated with Cys574 of another molecule in a dimer. In fact, hHsp70 1–561, of which a half population still formed a dimer, did not form a disulfide bridge (Fig. 8c).

Although N-terminal deletion of the client-binding domain also induced dimer formation, the configuration of the resulting dimer was considerably distinct from that of the dimer of the client-binding domain, because the client-binding domain with a truncated N-terminus was highly susceptible to proteolysis. N-terminal sequencing analysis of the proteolytic fragments confirmed this notion, because peptide bonds (Arg469–Gly470 and Arg517–Met518) located within the client-binding




Fig. 8. Dimerization of the full-length form and its C-terminally truncated forms. The full-length form and its C-terminally truncated forms expressed are schematically illustrated (a). Cys574 and Cys603 represent the two sole cysteines located within the client-binding domain. The results in terms of disulfide bridge formation (S-S) and the states [monomer (M) or dimer (D)] are summarized on the right. (b) SDS-PAGE under non-reducing (left) and reducing conditions (right). Numbers 1–6 are identical to those in panel “a.” Mk, molecular markers. (c) The recombinant proteins (1 mg/ml, 0.8 ml) were separated on a Sepharyl S300 (1 × 90 cm) column as described under “EXPERIMENTAL PROCEDURES.” Affinity-purified samples (P, 1 μg) and aliquots (6 μl) of fractions 28–50 obtained on the gel chromatography were separated by SDS-PAGE on a 10% polyacrylamide gel under non-reducing conditions. The apparent molecular masses of recombinant proteins are indicated on the right. Mk, molecular markers.

sub-domain became susceptible to trypsin on N-terminal truncation of the domain (Table 1). In contrast, the dimer formed through the truncation from the C-terminus showed proteolytic profiles similar to that of the client-binding domain, indicating their structural integrity. Moreover, we found the presence of a trimer of hHsp70

(Fig. 1). A trimer was also recognized for hHsp70 382–641, 382–601 and 382–581, but not for hHsp70 420–641 or 440–641 (Figs. 3 and 4), which further suggested the similarity of the dimeric structure of the full-length form to those of hHsp70 382–641 and its C-terminally-truncated forms.

Previous studies consistently showed that the C-terminal client-binding domain is responsible for the dimer/oligomer formation of Hsc70 (19). However, there was an apparent discrepancy as to the roles of sub-domains, *i.e.*, whether the client-binding sub-domain or the lid sub-domain is critical for this role. The 10-kDa C-terminal lid sub-domain (amino acids 542–646) of rat Hsc70, which forms an elongated bundle-like structure, dictates self-association (22). In contrast, Foucaq *et al.* (20) reported that the 17-kDa peptide-binding sub-domain (amino acids 385–540 of Hsc70) was involved in the oligomerization process. The results obtained in the present study are not in accord with either of these reports. Figure 3 showed that either of the sub-domains, *i.e.*, the client-binding sub-domain or the lid sub-domain, did not form a dimer, but that the client-binding domain partially truncated from either the N- or C-terminus could form dimers. We also demonstrated substantial changes in the configuration if the N-terminal segment of the client-binding sub-domain was lost, strongly suggesting that the client-binding sub-domain is essential for dimer formation. Hence, we propose that the client-binding sub-domain (amino acids 382–519) is a prerequisite, but that an adjacent region (amino acids 520–561 of the lid sub-domain) is indispensable for the dimerization.

Chou *et al.* (22) reported the crystal structure of the C-terminal 10-kDa lid sub-domain of rat Hsc70. The lid sub-domain formed a package of 48 molecules in a unit cell. In this structure, the association was mediated by coiled coil-like dimers in an anti-parallel manner, and then the dimers formed a cruciform tetrameric structure in an asymmetric manner. However, the lid sub-domain could not form a dimer under our conditions, and deletion of more than 38 amino acid residues from the N-terminus of the client-binding domain (residues 382–641) produced a dimer with a conformation that was distinct from that of the client-binding domain. These findings strongly suggest that the oligomeric configuration formed by the lid sub-domain is distinct from that of the native dimer. According to the structure proposed by Chou *et al.* (22), the lid sub-domain of one molecule associates with that of another molecule in an anti-parallel manner, in which two cysteines are separated from each other and therefore do not seem to form a disulfide bridge. Thus, we stress that the dimeric structure proposed by Chou *et al.* (22) should be carefully re-investigated.

Finally we propose a mechanism for the dimerization and the formation of the disulfide bridge through Cys574, as follows: Hsp70 could form a dimer through the C-terminal client-binding domain. In the dimeric structure, Cys574 of an Hsp70 molecule is located close to Cys574 of another molecule in the dimer, but the distance is not short enough for efficient formation of a disulfide bridge. When the C-terminal 40–60 amino acids are deleted, the two cysteine residues can gain access to each other, presumably because of destabilization of the C-terminal domain. This is likely to be the case, because

the formation of the bridge resulted in conversion of an hHsp70 dimer into a different configuration, as shown in Fig. 4b: The dimer of hHsp70 382–601 that possessed the disulfide bridge had a configuration that migrated faster on PAGE under non-denaturing conditions than that without the bridge did (Fig. 4b). This finding strongly suggests that the dimer with the disulfide bridge had a more compact configuration than that without the bridge. A further study on the dimeric structure of hHsp70 is under way in our laboratory.

We greatly acknowledge Mr. Takeshi Kobayakawa for his technical assistance.

#### REFERENCES

- Xu, Z., Horwich, A.L., and Sigler, P.B. (1997) The crystal structure of the asymmetric GroEL-GroES-(ADP) 7 chaperonin complex. *Nature* **388**, 741–750
- Minami, Y., Kimura, Y., Kawasaki, H., Suzuki, K., and Yahara, I. (1994) The carboxy-terminal region of mammalian HSP90 is required for its dimerization and function *in vivo*. *Mol. Cell. Biol.* **14**, 1459–1464
- Nemoto, T., Ohara-Nemoto, Y., Ota, M., Takagi, T., and Yokoyama, K. (1995) Mechanism of dimer formation of the 90-kDa heat-shock protein. *Eur. J. Biochem* **233**, 1–8
- Prodromou, C., Roe, S.M., O'Brien, R., Ladbury, J.E., Piper, P.W., and Pearl, L.H. (1997) Identification and structural characterization of the ATP/ADP-binding site in the Hsp90 molecular chaperone. *Cell* **90**, 65–75
- Prodromou, C., Roe, S.M., Piper, P.W., and Pearl, L.H. (1997) A molecular clamp in the crystal structure of the N-terminal domain of the yeast Hsp90 chaperone. *Nature Struct. Biol* **4**, 477–482
- Stebbins, C.E., Russo, A.A., Schneider, C., Rosen, N., Hartl, F.U., and Pavletich, N.P. (1997) Crystal structure of an Hsp90-geldanamycin complex: targeting of a protein chaperone by an antitumor agent. *Cell* **89**, 239–250
- Meyer, P., Prodromou, C., Hu, B., Vaughan, C., Roe, S.M., Panaretou, B., Piper, P.W., and Pearl, L.H. (2003) Structural and functional analysis of the middle segment of Hsp90: implications for ATP hydrolysis and client protein cochaperone interactions. *Mol. Cell* **11**, 647–658
- Welch, W.J. and Feramisco, J.R. (1982) Purification of the major mammalian heat shock proteins. *J. Biol. Chem.* **257**, 14949–14959
- Palleros, D.R., Reid, K.L., Shi, L., and Fink, A.L. (1993) DnaK ATPase activity revisited. *FEBS Lett.* **336**, 124–128
- Blond-Elguindi, S., Fourie, A.M., Sambrook, J.F., and Gething, M.-J., H. (1993) Peptide-dependent stimulation of the ATPase activity of the molecular chaperone BiP is the result of conversion of oligomers to active monomers. *J. Biol. Chem.* **268**, 12730–12735
- Azem, A., Oppliger, W., Lustig, A., Jenö, P., Feifel, B., Schatz, G., and Horst, M. (1997) The mitochondrial hsp70 chaperone system: effect of adenine nucleotides, peptide substrate, and mGrpE on the oligomeric state of mhsp70. *J. Biol. Chem.* **272**, 20901–20906
- Kim, D., Lee, Y.J., and Corry, P.M. (1992) Constitutive HSP70: oligomerization and its dependence on ATP binding. *J. Cell Physiol.* **153**, 353–361
- Chappell, T.G., Konforti, B.B., Schmid, S.L., and Rothman, J.E. (1987) The ATPase core of a clathrin uncoating protein. *J. Biol. Chem.* **262**, 746–751
- Stevens, S.Y., Cai S., Pellecchia, M., and Zuideweg, E.R.P. (2003) The solution structure of the bacterial HSP70 chaperone protein domain DnaK (393–507) in

- complex with the peptide NRRLLTG. *Prot. Sci.* **12**, 2588–2596
15. Burkholder, W.F., Zhao, X., Zhu, X., Hendrickson, W.A., Gragerov, A., and Gottesman, M.E. (1996) Mutations in the C-terminal fragment of DnaK affecting peptide binding. *Proc. Natl. Acad. Sci. USA* **93**, 10632–10637
  16. Takeda, S. and McKay, D.B. (1996) Kinetics of peptide binding to the bovine 70 kDa heat shock cognate protein, a molecular chaperone. *Biochemistry* **35**, 4636–4644
  17. Wang, C. and Lee M.R. (1993) High-level expression of soluble rat hsc70 in *Escherichia coli*: purification and characterization of the cloned enzyme. *Biochem. J.* **294**, 69–77
  18. Flynn, G.C., Chappell, T.G., and Rothman, J.E. (1989) Peptide binding and release by proteins implicated as catalysts of protein assembly. *Science* **245**, 385–390
  19. Benaroudj, N., Fouchaq, B., and Ladjimi, M.M. (1997) The COOH-terminal peptide binding domain is essential for self association of the molecular chaperone HSC70. *J. Biol. Chem.* **272**, 8744–8751
  20. Fouchaq, B., Benaroudj, N., Ebel, C., and Ladjimi, M.M. (1999) Oligomerization of the 17-kDa peptide-binding domain of the molecular chaperone HSC70. *Eur. J. Biochem.* **259**, 379–384
  21. Wang, T.F., Chang, J.-H., and Wang, C. (1993) Identification of the peptide binding domain of hsc70: 18-kilodalton fragment located immediately after ATPase domain is sufficient for high affinity binding. *J. Biol. Chem.* **268**, 26049–26051
  22. Chou, C.-C., Forouhar, F., Yeh, Y.-H., Shr, H.-L., Wang, C., and Hsiao, C.-D. (2003) Crystal structure of the C-terminal 10-kDa subdomain of Hsc70. *J. Biol. Chem.* **278**, 30311–30316
  23. Hunt, C. and Morimoto, R.I. (1985) Conserved features of eukaryotic hsp90 genes revealed by comparison with the nucleotide sequence of human hsp70. *Proc. Natl. Acad. Sci. USA* **82**, 6455–6459
  24. Milner, C.M. and Campbell, R.D. (1990) Structure and expression of the three MHC-linked HSP70 genes. *Immunogenetics* **32**, 242–251
  25. Laemmli, U.K. (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680–685
  26. Nishihara, J., Ishibashi, T., Sakai, M., Nishi, S., Kumazaki, T., Hatanaka, Y., Tsuda, S., and Hikichi, K. (1992) Characterization of cysteine residues of glutathione S-transferase P: evidence for steric hindrance of substrate binding by a bulky adduct to cysteine 47. *Biochem. Biophys. Res. Commun.* **188**, 424–432
  27. Nemoto, T., Ota, M., Ohara-Nemoto, Y., and Kaneko, M. (1995) Identification of dimeric structure of proteins by use of the glutathione S-transferase-fusion expression system. *Anal. Biochem.* **227**, 396–399
  28. Nemoto, T. and Sato, N. (1998) Oligomeric forms of the 90-kDa heat shock protein. *Biochem. J.* **330**, 989–995
  29. Ohno, M., Kitabatake, N., and Tani, F. (2004) Role of the C-terminal region of mouse inducible Hsp72 in the recognition of peptide substrate for chaperone activity. *FEBS Lett.* **576**, 281–286