

Palytoxin Acts on Na⁺,K⁺-ATPase but not Nongastric H⁺,K⁺-ATPase

Saida Guennoun-Lehmann · James E. Fonseca ·
Jean-Daniel Horisberger · Robert F. Rakowski

Received: 5 March 2007 / Accepted: 14 May 2007 / Published online: 17 July 2007
© Springer Science+Business Media, LLC 2007

Abstract Palytoxin (PTX) opens a pathway for ions to pass through Na,K-ATPase. We investigate here whether PTX also acts on nongastric H,K-ATPases. The following combinations of cRNA were expressed in *Xenopus laevis* oocytes: *Bufo marinus* bladder H,K-ATPase α_2 - and Na,K-ATPase β_2 -subunits; *Bufo* Na,K-ATPase α_1 - and Na,K-ATPase β_2 -subunits; and *Bufo* Na,K-ATPase β_2 -subunit alone. The response to PTX was measured after blocking endogenous *Xenopus* Na,K-ATPase with 10 μ M ouabain. Functional expression was confirmed by measuring ⁸⁶Rb uptake. PTX (5 nM) produced a large increase of membrane conductance in oocytes expressing *Bufo* Na,K-ATPase, but no significant increase occurred in oocytes expressing *Bufo* H,K-ATPase or in those injected with *Bufo* β_2 -subunit alone. Expression of the following combinations of cDNA was investigated in HeLa cells: rat colonic H,K-ATPase α_1 -subunit and Na,K-ATPase β_1 -subunit; rat Na,K-ATPase α_2 -subunit and Na,K-ATPase β_2 -subunit; and rat Na,K-ATPase β_1 - or Na,K-ATPase β_2 -subunit alone. Measurement of increases in ⁸⁶Rb uptake confirmed that both rat Na,K and H,K pumps were functional in HeLa cells expressing rat colonic HK α_1 /NK β_1 and NK α_2 /NK β_2 . Whole-cell patch-clamp measurements in HeLa cells

expressing rat colonic HK α_1 /NK β_1 exposed to 100 nM PTX showed no significant increase of membrane current, and there was no membrane conductance increase in HeLa cells transfected with rat NK β_1 - or rat NK β_2 -subunit alone. However, in HeLa cells expressing rat NK α_2 /NK β_2 , outward current was observed after pump activation by 20 mM K⁺ and a large membrane conductance increase occurred after 100 nM PTX. We conclude that nongastric H,K-ATPases are not sensitive to PTX when expressed in these cells, whereas PTX does act on Na,K-ATPase.

Keywords Palytoxin · Na⁺,K⁺-ATPase · H⁺,K⁺-ATPase

Introduction

The PII_C-type ion-motive adenosine triphosphatase (ATPase) subgroup (also termed X,K-ATPases) includes the ubiquitous Na,K-ATPase, the gastric H,K-ATPase found in parietal cells and the nongastric (ng) H,K-ATPases expressed in distal colon (Crowson & Shull, 1992). ngH,K-ATPase is overexpressed under pathophysiological conditions such as chronic hypokalemia, NaCl deficiency or renal acidosis (Silver & Soleimani, 1999). Several variants of ngH,K-ATPases have been identified and isolated from the amphibian toad bladder, human skin, guinea pig and rabbit distal colon. X,K-ATPases require another polypeptide, the glycosylated β -subunit, to be expressed and functional within the plasma membrane (Geering, 1998), where they maintain K⁺ homeostasis and intracellular ionic composition. The closely related PII_A-type ATPase subgroup, including the sarcoendoplasmic reticulum Ca²⁺-ATPase (SERCA), has characteristics similar to those of PII_C-type ATPases. The structure of their catalytic α -subunits shows a high level of sequence similarity (Sweadner

S. Guennoun-Lehmann · R. F. Rakowski (✉)
Department of Biological Sciences, Ohio University, Athens,
OH 45701, USA
e-mail: rakowski@ohio.edu

J. E. Fonseca
School of Electrical Engineering and Computer Science, Ohio
University, Athens, OH 45701, USA

J.-D. Horisberger
Department of Pharmacology and Toxicology, University of
Lausanne, Rue du Bugnon 27, CH-1005 Lausanne, Switzerland

& Donnet, 2001). They exchange Ca^{2+} or Na^+ for H^+ or K^+ by using the energy derived from ATP hydrolysis. P_{IIc} - and P_{IIa} -type ATPases are inhibited by vanadate acting at the ATP binding site (Cantley, Resh & Guidotti, 1978). Experimental data have shown that both rat colonic and *Bufo* bladder H,K-ATPases can transport Na^+ rather than H^+ ions (Cougnon et al., 1998; Spicer et al., 2001). Despite these similarities, however, Na,K-ATPase differs in several important aspects from both gastric and ngH,K-ATPases. Na,K-ATPase functions with a transport stoichiometry of $3\text{Na}^+/2\text{K}^+$, resulting in outward transport of one net charge per pump cycle (Post & Jolly, 1957; Rakowski, Gadsby & De Weer, 1989), whereas H,K-ATPases have a $1\text{K}^+/1\text{H}^+$ or $2\text{K}^+/2\text{H}^+$ stoichiometry, resulting in net electroneutral exchange (Sachs et al., 1976; Rabon, McFall & Sachs, 1982; Burnay et al., 2001). In polarized epithelial tissues, Na,K-ATPase is mainly located in the basolateral plasma membrane (Gottardi & Caplan, 1993), whereas H,K-ATPases are present primarily at the apical surfaces (Caplan, 1997; Smolka, Helander & Sachs, 1983; Pestov et al., 2002). *Bufo*, rat and human ngH,K-ATPases are moderately sensitive to ouabain; and both *Bufo* and rat ngH,K-ATPases can be inhibited by high concentrations of SCH-28080 (Del Castillo, Rajendran & Binder, 1991; Codina et al., 1996).

At picomolar concentrations, the highly potent marine toxin palytoxin (PTX) binds to Na,K-ATPase and converts it from an ion pump into an ion channel (Habermann, 1989; Wang & Horisberger 1997; Artigas & Gadsby 2003). This greatly increases the membrane conductance and results in a net inward current carried by Na^+ . The gating of PTX-induced ion channels can be modulated by Na,K-ATPase ligands such as Na^+ or ATP (Artigas & Gadsby 2003, 2004; Hilgemann, 2003). PTX-induced membrane conductance can also be inhibited by ouabain or K^+ ions (Ozaki, Nagase & Urakawa, 1985; Artigas & Gadsby, 2003). Experimental data from cysteine-scanning accessibility studies combined with structural modeling based on the three-dimensional (3-D) structure of SERCA (Guennoun & Horisberger 2000, 2002) have suggested that the PTX-induced ion channel includes at least a part of the Na^+ and K^+ transport pathway. Reports on the effect of PTX on the apical vs. basolateral side of polarized epithelial cells from kidney (LLC-PK1) provide additional evidence that the site of PTX action is Na,K-ATPase (Mullin, Snock & McGinn, 1991). However, it has been reported that PTX acts on both distal and proximal parts of the descending colon (Scheiner-Bobis, Hubschle & Diener, 2002). Extensive pharmacological characterizations of colonic and other H,K-ATPases found in collecting duct have yielded conflicting conclusions regarding its sensitivity to ouabain (Codina et al., 1996; Cougnon et al., 1996; Del Castillo et al., 1991). Hence, definitive conclusions cannot be drawn regarding the site of PTX action based on studies in colon and collecting duct,

and further investigation of PTX action on both Na,K-ATPase and H,K-ATPase in tissues with well-defined expression of these transport proteins is warranted.

The goal of the present study was to determine whether PTX increases the conductance of ngH,K-ATPases or acts only on Na,K-ATPase. We expressed *Bufo* ngH,K-ATPase, Na,K-ATPase or Na,K-ATPase β_2 -subunit alone in *Xenopus* oocytes and rat colonic H,K-ATPase, Na,K-ATPase and Na,K-ATPase β_1 - or β_2 -subunit alone in HeLa cells. *Bufo* and rat Na,K-ATPases are both ouabain-resistant. Both *Xenopus* oocyte and HeLa Na,K-ATPase are ouabain-sensitive. Thus, we can inhibit the endogenous sodium pump with a low dose (e.g., $10 \mu\text{M}$) of ouabain without completely blocking the exogenous Na,K-ATPase in the cells used for these studies. To investigate the action of PTX, we measured conductance changes using the two-microelectrode voltage-clamp technique in *Xenopus* oocytes and whole-cell current in HeLa cells using the patch-clamp technique. We also examined models of Na,K- and ngH,K-ATPase based on the known 3-D structure of SERCA in order to determine if a major difference exists between the two that could be related to their ability to respond to PTX.

Materials and Methods

Plasmid Construction

Full-length cDNAs encoding rat Na,K-ATPase α_1 -subunit ($\text{NK}\alpha_1$), rat Na,K-ATPase β_1 -subunit ($\text{NK}\beta_1$), rat colonic H,K-ATPase α_2 -subunit ($\text{HK}\alpha_2$) and rat Na,K-ATPase β_2 -subunit ($\text{NK}\beta_2$) were digested by restriction enzymes that conserve the 5' Kozak translation initiation sequence, the methionine start codon and the stop codon 3' ends of the genes of interest. Using T4 DNA ligase, we inserted each cDNA fragment into the previously linearized pcDNA3.1⁺. In order to achieve high expression levels in mammalian cells, we used a restriction enzyme that cut within multiple cloning sites conserving the CMV promoter and BGH polyadenylation signal. The coding regions of *Bufo* Na,K-ATPase α_1 -subunit ($\text{NK}\alpha_1$), Na,K-ATPase β_2 -subunit ($\text{NK}\beta_2$) or *Bufo* bladder H,K-ATPase α_2 -subunit ($\text{HK}\alpha_2$) cRNAs were inserted into pSD5 vector with SP6 promoter, allowing high levels of protein expression in *Xenopus* oocytes.

Expression Systems

Xenopus oocytes were microinjected with *Bufo* $\text{NK}\alpha_1$ / $\text{NK}\beta_2$ cRNAs to overexpress *Bufo* Na,K-ATPase, *Bufo* $\text{HK}\alpha_2$ / $\text{NK}\beta_2$ cRNAs to overexpress *Bufo* (bladder) ngH,K-ATPase or *Bufo* β_2 -subunit ($\text{NK}\beta_2$) cRNA alone. HeLa

cells were transiently transfected with rat NK β_1 -subunit cDNA and cotransfected with a total of 2 mg cDNA of rat NK α_1 /NK β_1 to overexpress rat Na,K-ATPase or with a total of 2 mg cDNA of HK α_2 /HK β_2 to overexpress rat (colonic) ngH,K-ATPase, using PolyFect reagent (Qiagen, Valencia, CA) following the protocol described by the vendor.

^{86}Rb -Uptake Measurements

^{86}Rb -uptake measurements were performed to ensure that we achieved high levels of functional expression of both *Bufo* ngH,K- and Na,K-ATPase in *Xenopus* oocytes and rat ngH,K- and Na,K-ATPase in HeLa cells. *Xenopus* oocytes expressing *Bufo* bladder H,K-ATPase, Na,K-ATPase or β_2 -subunit alone were loaded with Na $^+$ by 2-h incubation in a K $^+$ -free/Ca $^{2+}$ -free solution containing (in mM) 90 NaCl and 0.5 ethyleneglycoltetraacetic acid (EGTA). Na $^+$ -loaded oocytes were transferred to a solution containing (in mM) 5 KCl, 90 NaCl, 1 CaCl $_2$, 10 4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid (HEPES, pH 7.4), 0.2 μM ouabain (to inhibit endogenous Na,K-ATPase) and 10 μM bumetanide (to inhibit ^{86}Rb uptake mediated by the Na-K-2Cl cotransporter). Oocytes were incubated 12 min with ^{86}Rb (5 $\mu\text{Ci/ml}$) at room temperature and washed with a solution containing (in mM) 90 NaCl, 1 CaCl $_2$, 1 MgCl $_2$ and 10 HEPES (pH 7.4). Individual oocytes were then dissolved in 0.5% sodium dodecyl sulfate, and ^{86}Rb uptake was determined by scintillation counting. HeLa cells grown in 24-well cluster dishes at 60–80% confluence were transiently transfected as described above. Three days later, ^{86}Rb -uptake measurements were performed using a wash tray according to Sangan et al. (2000). After drilling a hole in each coverwell (~12.5 mm), we inserted a plastic test tube (#2052; Falcon, Lincoln Park, NJ) and then glued it into position. This allowed us to fill each tube with wash solutions and invert the whole assembly on the 24-well culture dish containing transfected HeLa cells. The solution obtained in each tube of the wash tray was transferred into individual test wells.

Steady-State Voltage-Clamp Measurements

Xenopus oocytes were microinjected with *Bufo* NK α_1 /NK β_2 , HK α_2 /NK β_2 and NK β_2 cRNAs encoding for *Bufo* Na,K-ATPase, bladder ngH,K-ATPase and Na,K-ATPase β_2 -subunit, respectively. The steady-state current activated by 10 mM extracellular K $^+$ was measured 3 or 4 days later at a holding potential of -50 mV using the two-electrode voltage-clamp technique. The experimental solution contained (in mM) 100 Na $^+$ -gluconate, 0.82 MgCl $_2$, 0.41 CaCl $_2$, 10 *N*-methyl-D-glucamine/HEPES, 5 BaCl $_2$, 10


tetraethylammonium (TEA)-Cl $_2$ and 0.2 μM ouabain (to block endogenous *Xenopus* Na,K pumps). Ba $^{2+}$ and TEA $^+$ were present to block passive K $^+$ channels (Rakowski & Paxson, 1988) so that the current produced by ouabain-resistant *Bufo* Na,K pumps could be measured upon addition of extracellular K $^+$. An aliquot of 100 μM PTX (Sigma, St. Louis, MO) was thawed just prior to each experiment and diluted to a final concentration of 5 nM in the external K $^+$ -free solution containing 0.002% bovine serum albumin (BSA) to minimize PTX binding to nonglass surfaces. All solutions used in two-microelectrode voltage-clamp experiments had a pH of 7.4 \pm 0.05 and osmolality of approximately 200 mOsm/kg.

Steady-State Patch-Clamp Measurements

HeLa cells were transiently transfected with rat NK α_1 /NK β_1 cDNA encoding for rat Na,K-ATPase, rat HK α_2 /NK β_2 cDNA encoding for rat colonic ngH,K-ATPase or rat NK β_1 cDNA encoding for rat Na,K-ATPase β_1 -subunit alone. Two days later, the cells were seeded on polylysine-coated coverslips and 10 μM ouabain was added to the culture medium in order to inhibit endogenous Na,K pumps. The cells were then incubated at 37°C in a 5% CO $_2$ atmosphere until confluence. The day after achieving confluence, steady-state patch-clamp currents were measured at room temperature (22–25°C). The patch pipette (1–3 M Ω) was filled with an intracellular solution containing (in mM) 85 Na-sulfamate, 20 TEA-Cl, 3 MgCl $_2$, 5.5 dextrose, 10 EGTA, 10 HEPES, 5 Na-pyruvate, 10 MgATP and 7.9 phosphocreatine disodium salt. The K $^+$ -free bathing solution contained (in mM) 145 NaCl, 23 MgCl $_2$, 2 BaCl $_2$, 5.5 dextrose, 10 HEPES/Na $^+$ and 0.2 CdCl $_2$. The solution containing 20 mM K $^+$ to activate the Na,K pump was the same as the K $^+$ -free solution but with equimolar replacement of NaCl by KCl. Ouabain (10 μM) was added to bathing solutions to inhibit endogenous sodium pumps before starting electrophysiological recordings. An aliquot of 100 μM PTX (Sigma) was thawed just prior to each experiment and diluted to a final concentration of 100 nM in the external K $^+$ -free solution containing 0.002% BSA. All internal and external solutions used for patch-clamp measurements had a pH of 7.4 \pm 0.05 and osmolality of 280–300 mOsm/kg.

Direct PTX Application to Confluent HeLa Cells

The effect of PTX on cells overexpressing rat ngH,K-ATPase and Na,K-ATPase was studied in HeLa cells grown in 35-mm Petri dishes to 50–60% confluence and then transiently transfected with rat NK α_1 /NK β_1 cDNAs encoding for Na,K-ATPase, rat ngHK α_2 /NK β_2 cDNAs



encoding for (colonic) ngH,K-ATPase or rat NKβ₁ cDNA encoding for Na,K-ATPase β₁-subunit. We treated all Petri dishes 36 h later with 20 μM ouabain in 1 ml culture medium for 30 min to inhibit endogenous HeLa Na,K-ATPase. We then added 1 μM PTX to each Petri dish, and the cells were incubated for 90 min at 37°C in a 5% CO₂ atmosphere. Photographs were taken with a digital camera under phase-contrast illumination at magnifications of x100, x250 and x400. Petri dishes were photographed with phase-contrast illumination at x250 and x400.

Modeling of Rat Na,K-ATPase and Rat Colonic H,K-ATPase

Modeller version 8.2 (Marti-Renom et al., 2000) was used to create structural models of rat colonic ngH,K-ATPase and rat Na,K-ATPase based on a template of SERCA (Toyoshima, Numora & Tsuda, 2004) in the E2-P conformation (PDB ID: 1WPG). Modeller's SALIGN command was used to construct a global, multiple alignment that included rat Na,K and rat gastric H,K sequences to provide

Fig. 1 a Measurements of ⁸⁶Rb uptake by HeLa cells. The four groups of three bars represent data obtained from HeLa cells transiently transfected with rat NKα₁/NKβ₁ cDNAs encoding for rat Na,K-ATPase (group 1 from the left), NKβ₁ cDNA encoding for rat Na,K-ATPase β₁-subunit (group 2), HKα₂/NKβ₂ cDNAs encoding for rat ngH,K-ATPase (group 3) and NKβ₂ cDNA encoding for rat Na,K-ATPase β₂-subunit (group 4). Assays for ⁸⁶Rb uptake (nmol/mg protein/10 min) were performed on each category of cells exposed to 10 μM ouabain (black bars), 10 mM ouabain (light gray) and 10 μM ouabain plus 10 mM SCH-28080 (gray). The mean ± standard error of the mean is shown for six assays under each condition. HeLa cells expressing rat Na,K-ATPase or colonic ngH,K-ATPase exhibited increases of ⁸⁶Rb uptake of 8.8 ± 0.2-fold and 6.9 ± 0.3-fold, respectively, over the ⁸⁶Rb uptake background observed with cells transfected with rat Na,K-ATPase β₁- or β₂- subunit alone. **b** Measurements of ⁸⁶Rb uptake by *Xenopus* oocytes. The mean ± standard error of the mean of ⁸⁶Rb uptake (pmol/oocyte/12 min, 8–10 oocytes in each group) was measured in oocytes microinjected with cDNA encoding for *Bufo* Na,K-ATPase β₂-subunit alone (β₂) (left), cDNA encoding for *Bufo* bladder H,K-ATPase and β₂ (middle) and cDNA encoding for *Bufo* Na,K-ATPase and β₂ (right). Oocytes expressing *Bufo* bladder H,K-ATPase and *bufo* Na,K-ATPase exhibited an increase of ⁸⁶Rb uptake of 4.09 ± 1.04-fold and 4.35 ± 0.57-fold, respectively, over that measured in oocytes microinjected with β₂-subunit alone. Oocytes were Na⁺-loaded prior to uptake measurements. The solutions for uptake measurements in both **a** and **b** contained 10 μM ouabain to inhibit endogenous Na,K pumps and 10 μM bumetanide to block ⁸⁶Rb uptake mediated by the Na-K-2Cl cotransporter

a consensus alignment in regions of lower identity. SERCA and rat ngH,K-ATPase share about 31% identity (Wang & Takeyasu, 1997), but there is a much higher similarity in the intracellular domains as well as in the transmembrane sections, where ion binding and permeation occur. All images were prepared with the PyMOL program (<http://www.pymol.org>), which is a molecular graphics system with a Python interpreter designed for real-time visualization and generation of high-quality molecular graphics images.

Results

Functional Expression of Na,K-ATPase and ngH,K-ATPase

We carried out ⁸⁶Rb-uptake measurements to test whether Na,K- and ngH,K-ATPase proteins expressed in HeLa cells and in *Xenopus* oocytes are functional. As shown in Figure 1a, coexpression of rat Na,K-ATPase α₁ and β₁ or rat colonic ngH,K-ATPase α₂ and rat Na,K-ATPase β₂ in HeLa cells resulted in significant increases of ⁸⁶Rb uptake compared to cells expressing rat Na,K-ATPase β₁- or β₂-subunits alone. HeLa cells (n = 6) expressing rat Na,K-ATPase gave an increase of 8.8 ± 0.2-fold over the background ⁸⁶Rb uptake measured in cells transfected with rat β₁-subunit alone. Inhibition of Na,K-ATPase by 10 mM




Fig. 2 Effect of PTX on confluent HeLa cell morphology. PTX, 1 μM , was applied for 90 min to 35-mm Petri dishes containing 1 ml of culture medium previously grown to confluence of HeLa cells transfected with rat NK α_1 /NK β_1 cDNA encoding for rat Na,K-ATPase (**a,b**), rat ngHK α_2 /NK β_2 cDNA encoding for rat colonic ngH,K-ATPase (**c,d**) and NK β_1 cDNA encoding for rat Na,K-ATPase β_1 -subunit (**e,f**). Large clusters of cells expressing Na,K-ATPase (**a,b**) have swollen and detached from their neighboring cells and the substrate. They eventually become small and round and are found freely floating in the medium. Cells expressing rat Na,K-ATPase (**b**)

show cytoplasmic granulations, whereas no granulations are seen within cells expressing ngH,K-ATPase (**d**) or β_1 -subunit alone (**f**). **a, c** and **e** were photographed with phase-contrast illumination at $\times 250$ magnification. **b, d** and **f** were photographed with phase-contrast illumination at $\times 400$ magnification. Petri dishes were treated with 20 μM ouabain for 30 min prior to PTX application. Four such experiments all showed similar morphological changes in cells expressing rat Na,K-ATPase, and no similar changes were observed in cells expressing rat colonic ngH,K-ATPase or those transfected with rat Na,K-ATPase β_1 -subunit cDNA alone

ouabain gave background ^{86}Rb uptake similar to the controls with β_1 - and β_2 -subunits alone. Neither the ^{86}Rb uptake mediated by Na,K-ATPase nor that mediated by ngH,K-ATPase was affected by application of 10 mM SCH-28080, suggesting that neither is sensitive to this high dose of this compound. HeLa cells expressing rat ngH,K-ATPase displayed an ^{86}Rb uptake increase of 6.9 ± 0.3 -fold over the ^{86}Rb uptake background measured in HeLa cells transfected with β_2 -subunit alone. Application of 10 mM ouabain reduced ^{86}Rb uptake mediated by rat ngH,K-ATPase by about one-fifth, consistent with a moderate sensitivity of rat ngH,K-ATPase to ouabain. Results of ^{86}Rb -

uptake studies in Na^+ -loaded oocytes are shown in Figure 1b. Oocytes expressing either *Bufo* bladder ngH,K-ATPase or *Bufo* Na,K-ATPase exhibited increases of 4.09 ± 1.04 -fold and 4.35 ± 0.57 -fold over ^{86}Rb uptake in oocytes injected with *Bufo* β_2 -subunit alone. These results confirm that ngH,K- or Na,K-ATPase expressed in HeLa cells and in *Xenopus* oocytes are capable of significant ^{86}Rb uptake and, therefore, are expressed in the surface membrane as functional pumps. Additionally, the data in Figure 1 are consistent with a moderate sensitivity of rat colonic ngH,K-ATPase to ouabain and its resistance to SCH-28080.




Fig. 3 Conductance changes produced by PTX in *Xenopus* oocytes. **a** Representative traces of currents recorded using the two-microelectrode voltage-clamp technique in oocytes expressing *Bufo* Na,K-ATPase (top), *Bufo* bladder H,K-ATPase (middle) and *Bufo* Na,K-ATPase β_2 -subunit alone (bottom). After clamping the oocyte at -50 mV in K^+ -free solution (flow rate of 1 ml/min), solution containing 10 mM K^+ was applied for 30–60 s. Oocytes expressing Na,K-ATPase and exposed to 10 mM K^+ produced a small, rapid increase in outward current ($\sim 0.2 \mu A$), while oocytes expressing H,K-ATPase or those injected by β_2 -subunit alone did not produce a response. PTX (5 nM) generated a large inward current that continued to increase for several minutes after PTX removal in oocytes expressing *Bufo* Na,K-ATPase. No similar inward current was produced in oocytes expressing *Bufo* bladder ngH,K-ATPase or those injected with β_2 -subunit alone. **b** PTX-induced conductance, G_m (μS), measured from oocytes injected with cRNA coding for *Bufo* Na,K-ATPase β_2 -subunit (left bar), *Bufo* Na,K-ATPase/ β_2 (middle bar) and *Bufo* bladder ngH,K-ATPase/ β_2 (right bar). A large increase of membrane conductance occurred in oocytes expressing *Bufo* Na,K-ATPase after exposure to 5 nM PTX (middle bar). No significant membrane conductance changes were produced by 5 nM PTX in oocytes expressing ngH,K-ATPase or oocytes injected with β_2 -subunit cRNA alone. The oocytes were exposed to 10 μM ouabain prior to electrophysiological measurements. Values are means \pm standard error of the mean of 8–10 measurements

PTX Produces Morphological Changes on Confluent HeLa Cells Expressing Na,K-ATPase

The effect of PTX could be directly observed as morphological changes produced in HeLa cells expressing Na,K-ATPase but not in those expressing ngH,K-ATPase. HeLa cells were treated with 20 μM ouabain (to inhibit endogenous Na,K pumps) 30 min prior to PTX application. To visualize the changes in morphology that occurred after application of PTX, 1 μM PTX was added to the culture medium. After exposure to PTX (90 min), the medium was replaced by the solution used for electrophysiological measurements. Examination with phase-contrast microscopy of cells expressing rat Na,K-ATPase showed that many clusters of cells had detached from the substrate and were freely floating in the medium (Fig. 2a). These detached cells were small and round compared to attached cells that were spread flat against the substrate. Examination of the cells that remained attached to the surface (at x400) revealed granulations within their cytoplasm (Fig. 2b). The middle and lower rows in Figure 2 show photographs of Petri dishes of cells expressing rat ngH,K-ATPase (Fig. 2c,d) or transfected with rat Na,K-ATPase β_1 -subunit alone (Fig. 2e,f). Only a few small, rounded cells were freely floating in the medium. Most cells were flat and adherent to the substrate. Examination of these cells at x400 magnification showed that they were confluent and did not have cytoplasmic granulations (Fig. 2d,f).

Effect of PTX on Oocytes and HeLa Cells Expressing Na,K and ngH,K Pumps

The two-microelectrode voltage-clamp technique was used to measure currents generated by K^+ activation and PTX application in *Xenopus* oocytes expressing *Bufo* Na,K-ATPase or ngH,K-ATPase or those injected with *Bufo* Na,K-ATPase β_2 -subunit alone (Fig. 3a). An oocyte expressing *Bufo* Na,K-ATPase (top trace) was activated by 10 mM K^+ and generated a small outward Na,K pump current ($\sim 0.2 \mu A$). Exposure of oocytes expressing ngH,K-ATPase (middle trace) or those injected with β_2 -subunit alone (bottom trace) to 10 mM K^+ did not produce this small initial outward current. A small inward current was generated after 10 mM K^+ activation of *Bufo* bladder H,K-ATPase. After returning to K^+ -free solution, measurements of membrane conductance were performed by making 50-mV depolarizing voltage steps from the holding potential of -50 mV at intervals of 30 s (vertical current excursions). Application of 5 nM PTX to oocytes expressing *Bufo* Na,K-ATPase resulted in an inward current and an increase of membrane conductance. The conductance increase was very large, on average up to 30 times the baseline membrane conductance

(Fig. 3b, middle column). Oocytes expressing *Bufo* ngH,K-ATPase or those injected with β_2 -subunit alone did not produce an inward current after 1-min exposure of 5 nM PTX, and the membrane conductance remained at baseline levels. Similar results were obtained at 10 nM PTX (*data not shown*). Results at 5 and 10 nM PTX from 8–10 oocytes were combined and are summarized in Figure 3b.

We also measured K^+ -activated currents and the effect of PTX in HeLa cells using the whole-cell patch-clamp technique at -40 mV (Fig. 4). The current traces in Figure 4 illustrate activation of an outward Na,K pump current on application of 20 mM K^+ in cells expressing rat Na,K-ATPase (Fig. 4a, top trace) but not in cells either expressing ngH,K-ATPase (Fig. 4a, middle trace) or transfected with β_1 -subunit alone (Fig. 4a, lower trace). After returning to K^+ -free solution, all three categories of cells were exposed to 100 nM PTX, and the membrane conductance was measured after about 3 min of exposure. The membrane conductance of HeLa cells expressing rat Na,K-ATPase and exposed to PTX averaged 2.69 ± 0.24 nS, ~ 30 -fold greater than the membrane conductance measured in HeLa cells exposed to PTX and expressing rat ngH,K-ATPase (0.095 ± 0.03 nS, $n = 6$) or those transfected with β_1 -subunit alone (0.05 ± 0.01 nS, $n = 4$). These results are summarized in Figure 4b.

Rat ngH,K-ATPase and Na,K-ATPase Models

If the failure of PTX to increase the conductance of ngH,K-ATPase is a result of failure to bind to the protein, we would expect to find significant structural differences in the extracellular domains of ngH,K-ATPase and Na,K-ATPase that could account for this difference. In order to determine if there are structural differences between ngH,K-ATPase and Na,K-ATPase that could be related to PTX binding, we constructed structural models of rat Na,K- and rat (colonic) ngH,K-ATPase by alignment with SERCA. The alignments obtained correspond well with other alignments of type II_C-ATPases with SERCA (Munson, Garcia & Sachs, 2005; Rakowski & Sagar, 2003). These models are based on the known crystal structure of SERCA and were constructed for the E2-P conformation (PDB ID: 1WPG) (Toyoshima et al., 2004) as described in Materials and Methods. The N termini in type II_C ATPases are, in general, regions of low sequence identity. Indeed, Figure 5 shows a clear difference between the long N terminus (green) of the rat Na,K-ATPase model and the short one (also green) of rat ngH,K-ATPase. The Na,K-ATPase N terminus is in close proximity to the actuator domain, whereas the ngH,K-ATPase N terminus is hardly in contact with the actuator domain. Figure 5 also shows differences of the shape of the protruding M1–2 loop that may account for differences in ouabain and/or PTX binding affinity.





Fig. 4 Measurements of conductance changes produced by PTX in HeLa cells. **a** Typical current traces obtained in whole-cell patch-clamp experiments (-40 mV holding potential) in HeLa cells expressing rat Na,K-ATPase (*top trace*), ngH,K-ATPase (*middle*) or rat Na,K-ATPase β_1 -subunit *cDNA* alone (*bottom*). Cells expressing Na,K-ATPase produced an outward current, presumably mediated by the Na,K pump, whereas cells expressing ngH,K-ATPase or rat Na,K-ATPase β_1 -subunit alone did not show an immediate response to K^+ (*middle and lower traces*, respectively). PTX application produced a large inward current in cells expressing rat Na,K-ATPase, while no significant inward current was seen after 3-min application of PTX to cells expressing rat ngH,K-ATPase or those transfected with rat Na,K-ATPase β_1 -subunit *cDNA*. **b** Mean values of PTX-induced membrane conductance changes (G_m). Vertical bars indicate mean values \pm standard error of the mean of conductance changes measured from cells expressing rat Na,K-ATPase (*left bar*) or rat ngH,K-ATPase (*middle bar*) and cells transfected with rat Na,K-ATPase β_2 -subunit (*right bar*). A large conductance was produced by PTX application on cells expressing rat Na,K-ATPase, but no significant membrane conductance increase occurred after 3-min application of 100 nM PTX to cells expressing rat ngH,K-ATPase or transfected with rat Na,K-ATPase β -subunit

Discussion

The present study was designed to determine if PTX has an effect on ngH,K-ATPases (rat colonic and *Bufo marinus* bladder) in cells in which they can be functionally

Fig. 5 Views of rat Na,K-ATPase (*left*) and rat ngH,K-ATPase (*right*) models based on E2 (PDB ID: 1WPG) crystal structures of SERCA. Transmembrane segments M1 are shown in *red*, actuator domains in *blue* and N termini in *green*. Boxes indicate the M1–2 loops, and *arrows* indicate N termini. The remaining parts of the models are quite similar and are shown in gray



expressed (*Xenopus* oocytes and HeLa cells) under experimental conditions in which the participation of endogenous Na,K-ATPase can be prevented by prior application of a low dose of ouabain. The data in Figure 1 demonstrate that exogenous ngH,K-ATPase and ouabain-resistant Na,K-ATPase expressed in HeLa cells and oocytes are functional as measured by ^{86}Rb uptake in cells in which endogenous NaK-ATPase activity is blocked by $10\ \mu\text{M}$ ouabain. A clear and simple demonstration of the effect of PTX on the morphology of confluent HeLa cells overexpressing ouabain-resistant rat Na,K-ATPase is shown in Figure 2a and b. After exposure to PTX, the cells can no longer maintain their normal gradients of electrolytes. The cells develop intracellular granulations, become more rounded as they swell, detach from the substrate and adjacent cells and eventually shrink to small spheres floating freely in the medium, presumably because they lose their cellular contents across leaky surface membranes. HeLa cells expressing ngH,K-ATPase (Fig. 2c,d) or Na,K-ATPase β_1 -subunit alone (Fig. 2e,f) that have their endogenous Na,K-ATPase blocked by $20\ \mu\text{M}$ ouabain were not similarly affected.

We also performed measurements of membrane conductance on *Xenopus* oocytes expressing *Bufo* Na,K-ATPase or *Bufo* (bladder) ngH,K-ATPase. The results in Figure 3 show that PTX produced a large increase of membrane conductance in oocytes expressing Na,K-ATPase but not in those expressing either ngH,K-ATPase or H,K-ATPase β_2 alone. Patch-clamp experiments in HeLa cells (Fig. 4) showed that PTX produced a very large increase in conductance (~ 30 -fold) in cells expressing Na,K-ATPase but no significant increase in conductance in cells expressing (rat colonic) ngH,K-ATPase or rat Na,K-ATPase β_1 -subunit alone. We conclude from these studies that Na,K-ATPase, but not ngH,K-ATPase, is the target of PTX action.

Rat ngH,K-ATPase and Na,K-ATPase Structural Models

Figure 5 shows a marked difference between the N termini of rat Na,K-ATPase and rat ngH,K-ATPase models. This difference is due to a 40-residue shorter N terminus in ngH,K-ATPase than the one in Na,K-ATPase. The Na,K-ATPase N terminus is situated close to the actuator domain, and that is thought to tilt the M1 helix by rotation of the A domain (Toyoshima et al., 2004). This change is thought to play a key role in the E1 to E2 conformational change of the enzyme. The absence of 40 residues in the H,K-ATPase N terminus appears to reduce the interaction between the A domain and TM1. This is evident since the short α helix and strand that loop around the A domain in the Na,K-ATPase model are absent from the H,K-ATPase model. The Na,K-ATPase N terminus has been shown to play a role in PTX-induced channel inactivation (Wu et al., 2003). The absence of these 40 residues in the ngH,K-ATPase N terminus may, therefore, account for the absence of a PTX effect on ngH,K-ATPase. Additionally, the TM1–2 extracellular loop, which is critical for the high-affinity binding of ouabain (Sweadner & Donnet, 2001), protrudes toward the extracellular region in a different way in the ngH,K-ATPase and Na,K-ATPase models. We suggest that this structural difference may account for the difference in sensitivity of ngH,K-ATPase and Na,K-ATPase to ouabain and/or to PTX. Experiments that directly measure PTX binding to the two proteins are warranted to test if this is the explanation of the difference in their responsiveness. If differences in binding affinity are found, it should be possible to test which regions are involved in PTX and ouabain binding by constructing chimeras of the two ATPases.

It is well established that PTX binds to Na,K-ATPase and opens a conducting pathway through it (Scheiner-

Bobis, 1998; Wang & Horisberger, 1997; Guennoun & Horisberger, 2000, 2002; Hilgemann, 2003; Artigas & Gadsby, 2003, 2004). ngH,K-ATPase has been extensively studied since the successful cloning of its α -subunit in 1992 (Crowson & Shull, 1992). There is controversy regarding its pharmacological sensitivity to ouabain (Codina et al., 1996; Coughon et al., 1996, 1998; Sangan et al., 2000), and this complicates the interpretation of experiments in which both PTX and ouabain are present. It has been reported that PTX has an effect on both distal and proximal colon, and it has been suggested that this action of PTX is mediated by interaction with ngH,K-ATPase (Scheiner-Bobis et al., 2002). However, since the conductance increase produced by PTX action on Na,K-ATPase is so large, the presence of only a small amount of Na, K-ATPase in distal and proximal colon would be sufficient to explain the results obtained in those tissues. For example, in oocytes in which endogenous Na,K-ATPase was blocked by 10 μ M ouabain, even very low levels of expression of cysteine mutants of Na,K-ATPase resulted in a large increase of membrane conductance upon exposure to 2–4 nM PTX (Guennoun & Horisberger 2000, 2002; Horisberger et al., 2004). Low levels of Na,K-ATPase expression have been reported within the apical membranes of nongastric cells (Gottardi & Caplan 1993), whereas H,K-ATPases are present primarily at the apical surfaces (Caplan, 1997; Smolka et al., 1983). We suggest therefore that the reported action of PTX on proximal and distal colon is due to the presence of Na,K-ATPase in those tissues, even though mucosal tissue was treated with 1 mM ouabain prior PTX application. This apparently did not prevent the effect of PTX on the apical membranes. The effect of PTX on *Bufo* bladder H,K-ATPase (Guennoun, Rakowski & Horisberger, 2005) and on ATP1A1, the human ngH,K-ATPase (Grishen et al., 1994), was tested by electrophysiological measurements and no increase of membrane conductance was found (*unpublished data*). These results support the conclusion that PTX does not increase the conductance of ngH,K-ATPases and that the conductance increase produced by PTX in various tissues is due to the presence of Na,K-ATPase.

Acknowledgment We thank Gary Shull for providing the rat colonic H,K-ATPase α -subunit and the Na,K-ATPase α_1 -subunit cDNAs. We also thank Robert Levenson for providing the rat Na,K-ATPase β_1 - and β_2 -subunit cDNAs. We also thank Joshua Berlin for assistance with the mammalian expression system and for providing facilities to start this project. We thank Kaethi Geering for providing suggestions for this work. This investigation was supported in part by funds from an American Heart Association postdoctoral fellowship (to S. G.-L.), National Institutes of Health grant NS-22979, National Science Foundation grant CCF-0622158 and Swiss National Fund grant 31-65441.01.

References

- Artigas P, Gadsby DC (2003) Na⁺/K⁺-pump ligands modulate gating of palytoxin-induced ion channels. *Proc Natl Acad Sci USA* 100:501–505
- Artigas P, Gadsby DC (2004) Large diameter of palytoxin-induced Na/K pump channels and modulation of palytoxin interaction by Na/K pump ligands. *J Gen Physiol* 123:357–376
- Burnay M, Crambert G, Geering K, Horisberger J-D (2001) *Bufo marinus* bladder H,K-ATPase carries out electroneutral ion transport. *Am J Physiol* 281:F869–F874
- Caplan MJ (1997) Ion pumps in epithelial cells: sorting, stabilization, and polarity. *Am J Physiol* 272:G1304–G1313
- Cantley LG, Resh MD, Guidotti G (1978) Vanadate inhibits the red cell Na,K-ATPase from the cytoplasmic side. *Nature* 272:552–554
- Codina J, Kone BC, Delmas-Mata JT, Dubose TD Jr (1996) Functional expression of the H⁺,K⁺-ATPase α subunit. *J Biol Chem* 271:29759–29763
- Coughon M, Planelles G, Crowson MS, Shull GE, Rossier B, Jaisser F (1996) The rat distal colon P-ATPase α -subunit encodes a ouabain-sensitive H,K-ATPase. *J Biol Chem* 271:7277–7280
- Coughon M, Bouyer P, Planelles G, Jaisser F (1998) Does the colonic H,K-ATPase also act as an Na,K-ATPase? *Proc Natl Acad Sci USA* 95:6516–6520
- Crowson MS, Shull GE (1992) Isolation and characterization of a cDNA encoding the putative distal colon H⁺,K⁺-ATPase. *J Biol Chem* 267:13740–13748
- Del Castillo JR, Rajendran VM, Binder HJ (1991) Apical membrane localization of ouabain-sensitive K⁺-activated ATPase activities in rat distal colon. *Am J Physiol* 261:G1005–G1011
- Geering K (1998) Molecular mechanisms involved in the regulation of Na,K-ATPase expression. *Adv Mol Cell Biol* 23B:275–309
- Gottardi CJ, Caplan MJ (1993) Delivery of Na,K-ATPase in polarized epithelial cells. *Science* 260:552–554
- Grishen AV, Sverdlov VE, Costina MB, Modyanov NN (1994) Cloning and characterization of the entire cDNA encoded by ATP1A1—a member of the human Na,K/H,K-ATPase gene family. *FEBS Lett* 349:144–150
- Guennoun S, Horisberger J-D (2000) Structure of the 5th transmembrane segment of the Na,K-ATPase alpha subunit: a cysteine-scanning mutagenesis study. *FEBS Lett* 482:144–148
- Guennoun S, Horisberger J-D (2002) Cysteine-scanning mutagenesis study of the sixth transmembrane segment of the Na,K-ATPase alpha subunit. *FEBS Lett* 513:277–281
- Guennoun S, Rakowski RF, Horisberger J-D (2005) *Bufo marinus* H,K-ATPase is not sensitive to palytoxin. *J Gen Physiol* 126:58a
- Habermann E (1989) Palytoxin acts through Na⁺,K⁺ ATPase. *Toxicol* 27:1171–1187
- Hilgemann DW (2003) From a pump to a pore: how palytoxin opens the gates. *Proc Natl Acad Sci USA* 100:386–388
- Horisberger J-D, Kharoubi-Hess S, Guennoun S, Michielin O (2004) The fourth transmembrane segment of the Na,K-ATPase α subunit. *J Biol Chem* 279:29542–29550
- Marti-Renom MA, Stuart AC, Fiser A, Sanchez R, Melo F, Sali A (2000) Comparative protein structure modeling of genes and genomes. *Annu Rev Biophys Biomol Struct* 29:291–325
- Mullin JM, Snock KV, McGinn MT (1991) Effects of apical vs. basolateral palytoxin on LLC-PK1 renal epithelia. *Am J Physiol* 260:C1201–C1211
- Munson KR, Garcia G, Sachs G (2005) Inhibitor and ion binding sites on the gastric H,K-ATPase. *Biochemistry* 44:5267–5284
- Ozaki H, Nagase H, Urakawa N (1985) Interaction of palytoxin and cardiac glycosides on erythrocytes membrane and (Na⁺,K⁺)ATPase. *Eur J Biochem* 152:475–480

- Pestov NB, Korneenko TV, Adams G, Tillekeratne M, Shakhparonov MI, Modyanov NN (2002) Nongastric H-K-ATPase in rodent prostate: lobe-specific expression and apical localization. *Am J Physiol* 282:C907–C916
- Post RL, Jolly PC (1957) The linkage of sodium, potassium, and ammonium active transport across the human erythrocyte membrane. *Biochim Biophys Acta* 25:118–128
- Rabon EC, McFall TL, Sachs G (1982) The gastric [H,K]ATPase: H⁺/ATP stoichiometry. *J Biol Chem* 257:6296–6299
- Rakowski RF, Gadsby DC, De Weer P (1989) Stoichiometry and voltage dependence of the sodium pump in voltage-clamped, internally dialyzed squid giant axon. *J Gen Physiol* 93:903–941
- Rakowski RF, Paxson CL (1988) Voltage dependence of Na/K pump current in *Xenopus* oocytes. *J Membr Biol* 106:173–182
- Rakowski RF, Sagar S (2003) Found: Na and K binding sites of the sodium pump. *News Physiol Sci* 18:164–168
- Sachs G, Chang HH, Rabon E, Schackman R, Lewin M, Saccomani G (1976) A nonelectrogenic H⁺ pump in plasma membranes of hog stomach. *J Biol Chem* 251:7690–7698
- Sangan P, Thevananther S, Sangan S, Rajendran VM, Binder HJ (2000) Colonic H,K-ATPase alpha- and beta-subunits express ouabain-insensitive H,K-ATPase. *Am J Physiol* 278:C182–C189
- Scheiner-Bobis G (1998) Ion-transporting ATPases as ion channels. *Naunyn Schmiedebergs Arch Pharmacol* 357:477–482
- Scheiner-Bobis G, Hubschle T, Diener M (2002) Action of palytoxin on apical H⁺/K⁺-ATPase in rat colon. *Eur J Biochem* 269:3905–3911
- Silver RB, Soleimani M (1999) H⁺,K⁺-ATPases: regulation and role in pathophysiological states. *Am J Physiol* 276:F799–F811
- Smolka M, Helander HF, Sachs G (1983) Monoclonal antibodies against gastric H⁺/K⁺ ATPase. *Am J Physiol* 245:G589–G596
- Spicer Z, Clarke LL, Gawenis LR, Shull GE (2001) Colonic H⁺-K⁺-ATPase in K⁺ conservation and electrogenic Na⁺ absorption during Na⁺ restriction. *Am J Physiol* 281:G1369–G1377
- Sweadner KJ, Donnet C (2001) Structural similarities of Na,K-ATPase and SERCA, the Ca²⁺-ATPase of the sarcoplasmic reticulum. *Biochem J* 356:685–704
- Toyoshima C, Numora H, Tsuda T (2004) Luminal gating mechanism revealed in calcium pump crystal structures with phosphate analogues. *Nature* 432:361–368
- Wang X, Horisberger J-D (1997) Palytoxin effects through interaction with the Na,K-ATPase in *Xenopus* oocyte. *FEBS Lett* 409:391–395
- Wang S, Takeyasu K (1997) Primary structure and evolution of the ATP-binding domains of the P-type ATPases in *Tetrahymena thermophila*. *Am J Physiol* 272:C715–C728
- Wu CH, Vasilets LA, Takeda K, Kawamura M, Schwarz W (2003) Functional role of the N-terminus of Na⁺,K⁺-ATPase α -subunit as an inactivation gate of palytoxin-induced pump channel. *Biochim Biophys Acta* 1609:55–62