

## Unrestricted Diffusion of Exogenous and Endogenous PIP<sub>2</sub> in Baby Hamster Kidney and Chinese Hamster Ovary Cell Plasmalemma

Alp Yaradanakul · Donald W. Hilgemann

Published online: 8 February 2008  
© Springer Science+Business Media, LLC 2008

**Erratum to: J Membrane Biol (2007) 220(1–3):53–67**  
**DOI 10.1007/s00232-007-9074-4**

In the above-mentioned article, published in SpringerLink on November 16, 2007, and in print in vol 220/no 1–3, December 2007, pp 53–67, a Note Added in Proof was missing from the article (see below):

### Note Added in Proof

At the time of publication, the authors became aware that the group of Stuart McLaughlin (Stony Brook) had determined similar diffusion coefficients for fluorescent PIP<sub>2</sub> in intact cells using Fluorescence Correlation Spectroscopy technique.

---

The online version of the original article can be found under doi:  
[10.1007/s00232-007-9074-4](https://doi.org/10.1007/s00232-007-9074-4)

---

A. Yaradanakul (✉) · D. W. Hilgemann  
Department of Physiology, University of Texas Southwestern  
Medical Center at Dallas, Dallas, TX 75390-9040, USA  
e-mail: alp.yaradanakul@utsouthwestern.edu