

Erratum to: Investigation on Stability of Transporter Protein, Glucuronide Transporter from *Escherichia coli*

Noriyuki Ishii

Published online: 22 March 2011
© Springer Science+Business Media, LLC 2011

Erratum to: J Membrane Biol (2010) 235:63–72 DOI 10.1007/s00232-010-9256-3

Original Article has an error in the caption of Figure 2. In the last sentence of the caption, “heptameric” should be read as “pentameric”. Corrected caption for figure 2 is given below.

Fig. 2 Size-exclusion HPLC elution profile of GusB with molecular size marker standards. The GusB–DDM

complex was found to have a Stokes radius of 57 Å. Six soluble proteins of known Stokes radius were used for calibration: ferritin (63 Å), catalase (52 Å), aldorase (46 Å), bovine serum albumin (35 Å), ovalbumin (28 Å) and chymotrypsinogen (22 Å). Inset is a negatively stained transmission electron microscopic image of the peak fraction. Scale bar, 500 nm. Based on the elution profile and TEM image, GusB in the presence of DDM most probably stays as a pentameric oligomer.

The online version of the original article can be found under doi:[10.1007/s00232-010-9256-3](https://doi.org/10.1007/s00232-010-9256-3).

N. Ishii
Biological Information Research Center, National Institute of Advanced Industrial Science and Technology (AIST), Tsukuba Central-6, 1-1-1 Higashi, Tsukuba, Ibaraki 305-8566, Japan

N. Ishii
Skirball Institute of Biomolecular Medicine, Department of Cell Biology, New York University School of Medicine, 540 First Avenue, New York, NY 10016, USA

N. Ishii
Life Science Division, Donner Laboratory, Lawrence Berkeley National Laboratory, Molecular and Cell Biology Department, University of California, Berkeley, CA 94720, USA

N. Ishii (✉)
Biomedical Research Institute, National Institute of Advanced Industrial Science and Technology (AIST), Tsukuba Central-6, 1-1-1 Higashi, Tsukuba, Ibaraki 305-8566, Japan
e-mail: ishii@ni.aist.go.jp