

21-OH steroid hydroxylase (21-OH-ase) uses both I and II in corticoid biosynthesis in other species, it was considered of interest to study the comparative interactions which could exist between these two precursors and the rat adrenal 21-OH-ase, determining enzymatic constants for I and II (usual and unusual substrates, respectively). Homogenized adrenals from normal rats were incubated with various combinations of concentrations of I- ^3H and/or II- ^{14}C , acting as substrates and/or inhibitors of 21-OH-ase. The results showed that 21-OH-ase uses II almost as efficiently as I. The K_m values were about the same for both I and II (13.9 and 14.2×10^{-6} M/L), respectively, however, the V_{max} values were 54.6 and 26.0×10^{-7} M/L/h for I and II, respectively. The amounts of I required to saturate the 21-OH-ase was double than that of II. Further kinetic studies showed that both I and II inhibit the 21-hydroxylation of the other in a reciprocal fashion. While II inhibits the 21-hydroxylation of I by competitive inhibition, I inhibits the 21-hydroxylation of II through a mixed type of inhibition. The results suggest that, rather than the existence of two different specific enzymes (one for I and another for II) as it has been postulated by others, it seems that we are dealing with a 21-hydroxylating system with two active sites. One site uses only I and the other site uses I and/or II indistinctively.

59. The 11β -hydroxylase activity of cell-free adrenal preparations from *Echidnas* (*Tachyglossus aculeatus*) in various physiological states

WEISS, M. and MALECKI, J., Department of Physiology, Monash University, Clayton, Victoria, Australia

The echidna has a much lower rate of corticosteroid secretion than eutherian mammals and it survives adrenalectomy. Adrenal 11β -hydroxylase activity was found to be extremely low, although 17α - and 21-hydroxylase activities were comparable to eutherians. In echidnas treated for 1 week with frusemide (Lasix), which caused Na^+ depletion and dehydration, there was a marked increase in 11β -hydroxylation of the adrenal homogenates, 60% of the end products from progesterone being 11β -hydroxylated, compared with less than 1% in controls. This activity was also enhanced by dehydration or treatment of the animal with ACTH. Using purified mitochondrial preparations and deoxycorticosterone substrate the yield of corticosterone was 8.3% in an echidna treated with frusemide, less than 0.1% in a control echidna and 25.5% in a normal rat. The K_m values from Lineweaver-Burke plots for 11β -hydroxylase for the treated echidna and the rat in these experiments were 0.24 and 0.30 μM and the V_{max} values 8.2×10^{-6} and 6.7×10^{-4} $\mu\text{M min}^{-1} \text{mg}^{-1}$, respectively. Thus the efficiencies of 11β -hydroxylase in the two animals were comparable, but the amount of active enzyme in the echidna was approximately 1% of that in the normal rat. The low corticosteroid secretion rate in the echidna may be a consequence of a relatively meagre mitochondrial enzyme system.

60. A lasting effect of ACTH on adrenal 11β -hydroxylation in guinea-pig

KOLANOWSKI, J. and CRABBÉ, J., Endocrine Unit, Dept. Physiology, University of Louvain (Belgium)

The day following i.v. ACTH administration to guinea-pigs, the steroidogenic response to further ACTH stimulation is enhanced, as evidenced by plasma and adrenal tissue levels of cortisol (F), while 11-deoxycortisol (S) is not modified. This suggests a prolonged ACTH effect on late steps of

F biosynthesis. This hypothesis was evaluated on isolated guinea-pig adrenal cells by studying ACTH action on F, S, cyclic-AMP production and on 11β -hydroxylation of ^3H -S. Adrenal cells were harvested from 26 control adult male guinea-pigs and from 18 animals treated with ACTH ($75 \mu\text{g ACTH}_{1-24}$ i.v. over 3 h, 24 h previously). The 11β -hydroxylation index [(F/F+S) \times 100] determined from F and S adrenal tissue content before cell dispersion, was 82.7 ± 1.1 in controls and 99.5 ± 0.2 24 h after ACTH ($p < 0.001$). Incubation of isolated cells with tracer amounts of ^3H -S resulted in ^3H -F formation, itself undergoing conversion into cortisone (E). The 11β -hydroxylation index, calculated therefore as (F+E/F+E+S) \times 100, increased in cells from ACTH-treated animals, averaging after 30 min incubation 88 ± 1 vs. 77 ± 2 in controls ($p < 0.01$). The cells from both groups were then challenged with ACTH *in vitro* for 2 h, at doses ranging from 1 to 1000 pg/ml cell suspension. The F secretory response of isolated adrenal cells from ACTH-treated animals was enhanced when compared to controls, maximal F production (at 1000 pg ACTH/ml) averaging 1236 and 836 ng F/10⁵ cells, respectively ($p < 0.05$), while net S and cyclic-AMP production did not differ in both groups. An acute exposure of cells to ACTH during incubation does not influence any further the activity of the 11β -hydroxylation enzyme system, as judged from ^3H -S conversion into ^3H -F and ^3H -E. In conclusion: (1) No change in 11β -hydroxylase activity of guinea-pig adrenocortical cells results from acute exposure to ACTH; (2) Activity of this enzyme system increases as a delayed effect of ACTH stimulation, demonstrable the day after infusion with this hormone; (3) The increase in 11β -hydroxylase activity could account, at least in part, for the enhanced F secretory response obtained upon repeated ACTH stimulation. In contrast, generation of cyclic-AMP does not seem to be modified under these circumstances.

61. Regulation of androgen synthesis in the human adrenal gland *in vitro*

DESHPANDE, N., JENSEN, V. and YATES, J., Imperial Cancer Research Fund, Lincoln's Inn Fields, London WC2A 3PX, England

The possibility that local factors at a cellular level might influence androgen synthesis by the human adrenal gland was investigated in an incubation system *in vitro*. The synthesis of DHA was controlled by the availability of its major precursor 17α -hydroxypregnenolone, the obligatory cofactor for this reaction NADPH, and by the ratio of NADP^+ to NADPH. 17α -hydroxyprogesterone and DHA itself exerted non-competitive inhibition on the conversion. The synthesis of androstenedione from its two immediate precursors, 17α -hydroxyprogesterone and DHA, was also investigated. Both reactions were dependent upon the availability of the substrate and of the obligatory cofactors, NADP and NAD^+ , respectively. The conversion of 17α -hydroxyprogesterone to androstenedione was competitively inhibited by pregnenolone, progesterone and 17α -hydroxypregnenolone, while the synthesis of androstenedione from DHA was found to be non-competitively inhibited by oestrone and oestradiol- 17β . It is suggested that androgen synthesis is regulated by feedback inhibition at the cellular level.

62. Bovine adrenal cortex 3β -hydroxysteroid dehydrogenase and 3-oxosteroid- Δ^5 -4-isomerase: phospholipid requirement?

GEYNET, P., DE PAILLEREIS, C., NICOT, C. and