

E > 90% within 3 days, presumably by reducing endogenous D/DS. E (mainly estrone) was determined by RIA after enzyme hydrolysis of conjugates. Increments in E, estimated from the difference in E during 5 days pre- and post-treatment, were [ $\mu\text{g}$  mean (range), number of expt.]: Control A (no treatment), -47 (-462/+472), 8; Control B (5 mg estradiol-17 $\beta$ ), +2465 (+1358/+3537), 7; 100 mg D, +3206 (+1610/+5321), 7; 100 mg D + betamethazone, +2244 (+1655/+2824), 2; 100 mg DS ( $\equiv$  74 mg D), +35 (-1587/+659), 7; 100 mg DS + betamethazone, +107 (+46/+169), 2. Thus, preferential use of tracer D is unrelated to pool size, and variations in D availability and placental sulfatase activity may both contribute to regulation of estrogen production.

#### 84. The biochemical basis of placental sulfatase deficiency

FRANCE, J. T. and McNAUGHT, R. W., Postgraduate School of Obstetrics and Gynaecology, University of Auckland, Auckland, New Zealand

The human placenta is normally a rich source of the enzymes  $3\beta$ -steroid sulfatase and arylsulfatase but pregnancies have been recognized recently with deficiency in the activity of these enzymes. Fetal growth and development are normal but estrogen levels are extremely low since the placenta is unable to metabolise the estrogen precursors androst-5-en-17-one-3 $\beta$ -yl sulfate (DHEAS) and 16 $\alpha$ -hydroxyandrost-5-en-17-one-3 $\beta$ -yl sulfate. In attempting to establish the biochemical mechanism of the enzyme defect we have investigated the possibility that lack of enzyme activity was due to the presence of a sulfatase inhibitor. Sulfatase activity was determined kinetically from the hydrolysis of DHEAS-7 $\alpha$ - $^3\text{H}$  by microsomal preparations under optimum conditions. Microsomal preparations from affected placentas exhibited sulfatase activity. Ultrasonic (100 W, 24 kHz) treatment of the microsomes directed at disrupting a possible enzyme-inhibitor complex failed to increase the sulfatase activity of the preparation. Addition of 105,000  $\mu\text{g}$  supernatant from affected placental tissue to the incubate did not inhibit sulfatase activity of normal placental microsomes. It is concluded that placental sulfatase deficiency is likely to be due to deficiency of enzyme rather than the presence of a sulfatase inhibitor.

#### 85. Human placental $\Delta_5$ -3 $\beta$ -hydroxysteroid dehydrogenase: intracellular distribution, substrate specificity and inhibition

FERRE F., BREUILLER, M., DUCHESNE, M. J., SAINTOT, M. and DESCOMPS, B., Maternité de Port-Royal (E.R. CNRS No. 123) Paris Institut de Biologie (U 58 INSERM) Montpellier, France

$\Delta_5$ -3 $\beta$ -hydroxysteroid dehydrogenase ( $\Delta_5$ -3 $\beta$  HSD) activity has been assayed either by spectrophotometric method or by use of radioactive substrates. The enzymatic activity is equally distributed between mitochondrial and microsomal fractions verified by electronic microscopy. The specific activity is similar in both fractions, as well as the  $K_m$  for NAD ( $5 \times 10^{-5}\text{M}$ ), for the substrate, and the optimal pH. But the enzymatic activity is different with  $\text{C}_{21}$  or  $\text{C}_{19}$  natural or synthetic substrates; the optimal pH is 7.4-8 for  $\text{C}_{21}$  steroids and 10-10.2 for  $\text{C}_{19}$  steroids. The  $\Delta_5$ -3 $\beta$  HSD is strongly inhibited by  $\Delta_4$ -3-oxosteroids (progesterone and 4-androstenedione) and the inhibition seems to be of the

competitive type. Oestrogens and cyclic AMP have also an inhibitory action. These results suggest that human placental  $\Delta_5$ -3 $\beta$  HSD which has identical characteristics in mitochondrial and microsomal fractions has different properties with  $\text{C}_{21}$  or  $\text{C}_{19}$  steroid substrates, and that endogenous steroids play a physiological role in the regulation of the enzymatic activity.

### 3F. Steroid biosynthesis: Non endocrine tissues

#### 86. Testosterone biosynthesis by human skin

GARZON, P., CORONA, E. and TOPETE, F., Departamento de Bioquímica, Facultad de Medicina, Universidad de Guadalajara, Guadalajara, Jalisco, Mexico

Steroid metabolism in the human skin was studied using purified progesterone-4- $^{14}\text{C}$  as a testosterone and 17 $\alpha$ -hydroxyprogesterone precursor. Skin minces were placed in Eagle's medium, with penicillin and streptomycin, and thereafter incubated for a five days period at 37 C. Neither PPLO or other bacteria were detected as contributing factors to steroid metabolism. Organic extracts were paper chromatographed and labelled steroid detected by a gas-flow strip scanner. Separation, elution and radioactivity quantification was performed until constant specific activities of the compounds were obtained. 17 $\alpha$ -Hydroxy-4-pregnene-3,20-dione and 17 $\alpha$ -hydroxy-4-androsten-3-one yields accounted for less than 1% of the incubated precursor. The remaining metabolites have been previously described. Skin steroid metabolic pathways are becoming established. 17 $\alpha$ -Hydroxy-4-pregnene-3,20-dione might constitute an intermediate step to testosterone: both may be formed either as a compensatory route or just as a tissue response to form adaptive enzymes.

#### 87. *In vitro* synthesis of active sex steroids in normal and inflamed human gingiva

ELATTAR, T. M. A., Department of Biochemistry, School of Dentistry, University of Missouri, Kansas City, Mo., U.S.A.

This study employed 4 groups (each of 12 samples) of normal and inflamed human gingival tissue of male and female subjects. 300 mg of tissue slices were incubated, separately, in 3 ml of 0.1 M potassium phosphate buffer (pH 7.4) with 50  $\mu\text{Ci}$  of androstenedione-1,2- $^3\text{H}$  ( $\Delta^4\text{A}$ ) or estrone-2,4,6,7- $^3\text{H}$  ( $\text{E}_1$ ) in the presence of  $\text{NADPH}_2$  at 37 for 3 h. Organic solvent extracts of the incubated tissue slices were separated by paper and silica gel thin layer chromatography. Radioactive testosterone (T) and estradiol-17 $\beta$  ( $\text{E}_2$ ) were identified as the only conversion products of ( $\Delta^4\text{A}$ ) and ( $\text{E}_1$ ). The mean rates of conversion of ( $\Delta^4\text{A}$ ) to (T) and ( $\text{E}_1$ ) to ( $\text{E}_2$ ) in normal and inflamed tissue samples were 3.5 and 7.3  $\times 10^{-7}$   $\mu\text{M/g/min}$  in the males and 6.8 and 19.4  $\times 10^{-7}$   $\mu\text{M/g/min}$  in the females. These results reveal the presence of 17 $\beta$ -ol-dehydrogenase in human gingival tissue. The significant increase in the biosynthesis of active sex steroids, (T) and ( $\text{E}_2$ ), in inflamed as compared to normal gingiva of both sexes might be a factor in the aggravation of gingival inflammation due to the known hyperemic effects of these hormones. (Supported by USPH DE-03031-03.)