

significant increases occur in the theca and interstitium between days 10 and 25 of age whereas the granulosa exhibits an enzyme decline after day 15. Dehydroepiandrosterone (DHA), pregnenolone, 17-hydroxy-pregnenolone and 16-dehydropregnenolone served as substrates. Administration of 5  $\mu\text{g}$  of estradiol benzoate at 5 days of age delayed the development of the enzyme to day 20 in the interstitium and to day 30 in the theca regardless of the substrate but at age 40 days substrate differences were noted. When estradiol was injected on day 10, enzyme development in the interstitium was actually enhanced through day 40.

Prior studies have indicated that 10  $\mu\text{g}$  of estradiol dipropionate administered to the neonate will invoke infertility if injected at 5 days of age and that this dosage is partially effective at 10 days of age when studied in mice 100 days old. Administering 5 or 10  $\mu\text{g}$  of estradiol benzoate at 5 days of age resulted in 29 of 30 mice being infertile, although cohabitated with males for 90 days. Littermate controls proved fertile in 23 or 24 cases with litters being delivered in 24–31 days after admission of the male. When comparable steroid treatment was given at 10 days of age, 50% of the mice in each group proved infertile. Furthermore, litter size was subnormal and cohabitation time with the male was extended from 24 days (control) to 41 days (5  $\mu\text{g}$ ) to 57 days (10  $\mu\text{g}$ ) with each time period being significantly different. Finally, the influence of aging and neonatal steroid effects were examined 16 to 17 months after 5  $\mu\text{g}$  of estradiol benzoate was administered on day 5 (7 expt., 9 controls) or day 10 (6 expt., 8 controls). After 100–150 days in breeding n4 pregnancies resulted in neonatal steroid treated rats whereas 3 controls produced small litters. The impact of neonatal steroid treatment on aging is unknown but does deserve consideration.

51. **Physiologically available cortisol in the human fetus and mother**, B. E. PEARSON MURPHY and A. C. CAMPBELL, Reproductive Physiology Unit, Montreal General Hospital, and Department of Experimental Medicine, McGill University, Canada

In order to interpret the relationships between total cortisol levels in maternal serum, cord arterial and venous serum and amniotic fluid, it is necessary to consider the fraction of cortisol which is physiologically available to the tissues, i.e. that which is not bound to transcortin. This was measured at 37°C by equilibrium microdialysis of 0.1 ml sample against 0.1 ml saline containing a concentration of albumin equal to that of the sample. Total cortisol was measured by a specific non-chromatographic radiotransinassay. Physiologically available cortisol was then calculated as % unbound  $\times$  total cortisol. Subjects were studied at 12 to 20 weeks gestation. (6 elective hysterectomies) and at term (8 elective Caesarean sections, 14 induced vaginal deliveries, and 18 spontaneous-onset vaginal deliveries. The two vaginal delivery groups were matched for gestational age and duration of labour. The % unbound cortisol in maternal serum ranged from 17 to 29% while that in cord serum ranged from 34 to 76%. Cord arterial and venous values were similar in all instances. Lowest percentages were found at 12 to 20 weeks and highest at spontaneous vaginal delivery. Levels of physiologically available cortisol at term were all higher than at 12 to 20 weeks gestation. Cord arterial levels were consistently higher than venous levels in all groups, but especially in the spontaneous group ( $P < 0.01$ ). Maternal levels were higher than cord levels but the amount crossing the placenta, corrected for 85% conversion of cortisol to cortisone, as estimated previously in our laboratory (*Am. J. Obstet. Gynec.* 118 (1974) 538), was lower than cord levels in all instances ( $P < 0.01$ ).

Spontaneous-onset cord arterial levels ( $61 \pm 3$  ng/ml) were higher ( $P < 0.05$ ) than induced cord arterial levels ( $43 \pm 5$  ng/ml) or Caesarean section cord arterial levels ( $36 \pm 8$  ng/ml). Amniotic fluid levels were about half those of cord levels and correlated well with cord arterial levels but poorly with maternal venous levels. These studies provide evidence that (1) the placenta, by converting maternal cortisol to cortisone, acts as a barrier to prevent fetal pituitary-adrenal axis suppression, and (2) there is a surge of fetal cortisol which precedes the onset of spontaneous-onset labour which may be important in triggering parturition in man.

52. **Urinary steroid metabolites in the human newborn**, R. A. ANDERSON, G. DEFAYE, C. MADANI, E. M. CHAMBAZ, C.H.U. Grenoble (France), and C. J. W. BROOKS, Chemistry Department, University of Glasgow, Scotland

Urinary steroid separations using gas phase analysis (g.l.c. and GC-MS) have shown that the main metabolites excreted in the newborn period might have a different biological significance, according to their mode of conjugation. Steroid sulfates were of the 5-ene-3 $\beta$ -hydroxy type, increased after ACTH stimulation and were absent in a case of anencephaly. By contrast, the glucuronide fraction contained mainly saturated pregnane structures, were not influenced by ACTH and disappeared within the first 10 days of life.

Methods were developed to obtain satisfactory group separations of steroid metabolites using lipophilic substituted dextran gels, either by direct or reversed phase elution (Anderson *et al.*: *J. Chromatog.* 99 (1974) 485).

A number of polyhydroxylated 5-ene-3 $\beta$ -hydroxy- $\text{C}_{19}$  compounds were synthesized, either by biological or chemical routes: 3 $\beta$ ,15 $\alpha$ -dihydroxy-5-androsten-17-one; 5-androstene-3 $\beta$ ,7 $\alpha$  (and 7 $\beta$ ), 17 $\beta$ -triol; 3 $\beta$ ,16 $\beta$ ,17 $\beta$ -trihydroxy-5-androsten-7-one; 5-androstene-3 $\beta$ ,15 $\alpha$ ,16 $\alpha$ ,17 $\beta$ -tetrol; 5-androstene-3 $\beta$ ,15 $\beta$ ,16 $\beta$ ,17 $\beta$ -tetrol; 3 $\beta$ , 18-dihydroxy-5-androsten-17-one; 5-androstene-3 $\beta$ ,17 $\beta$ ,19-triol.

After preliminary group fractionation, 15 $\alpha$ -hydroxy-DHA (trace) and 5-androstene-3 $\beta$ ,15 $\alpha$ ,16 $\alpha$ ,17 $\beta$ -tetrol (average 110  $\mu\text{g}/24$  h) could be identified in the sulfate fraction in the newborn period.

The same methodology could be applied to the study of steroid in amniotic fluid; in a case of sulfatase defect, values within the normal range were obtained for the major 5-ene-3 $\beta$ -hydroxy steroids which could be measured.

53. **Pattern of plasma concentration of dehydroepiandrosterone during the neonatal period and the first year of life in human**, EVELINE DE PERETTI and MAGUELONE G. FOREST, Unité de Recherches Endocriniennes et Métaboliques chez l'Enfant, INSERM-U.34 Hôpital Debrousse, 29, rue Soeur Bouvier, 69322 Lyon Cedex 1, France

A specific and sensitive radioimmunoassay (RIA) for measuring unconjugated plasma dehydroepiandrosterone (DHA) has been developed. Specific antibodies have been obtained in rabbit immunized with a DHA-17 (O-carboxymethyl) oxime - BSA complex. Plasma was extracted by diethyl ether and the dry extract purified on a celite column. At the end of the RIA, bound and free fractions were separated using a Dextran-charcoal solution. In mixed cord blood, the mean values were 593.3 ng/100 ml  $\pm$  186.5 (range 248–1493) in 21 females, and 712.7 ng/100 ml  $\pm$  190.9 in 18 males (range 179–1367). These values were within adult range ( $642 \pm 112$  ng/100 ml in male and  $515 \pm 107$  ng/100 ml in female). During the first day of life the peripheral