

## THE SEXUALLY DIFFERENTIATED METABOLISM OF [6,7-<sup>3</sup>H]17 $\beta$ -OESTRADIOL IN RATS: MALE-SPECIFIC 15 $\alpha$ - AND MALE-SELECTIVE 16 $\alpha$ -HYDROXYLATION AND FEMALE-SELECTIVE CATECHOL FORMATION

J. L. MAGGS, P. MORGAN and B. K. PARK\*

Department of Pharmacology and Therapeutics, University of Liverpool, Liverpool L69 3BX, England

(Received 10 August 1991)

**Summary**—The oxygenated-metabolite profiles of exogenous 17 $\beta$ -oestradiol (E<sub>2</sub>) in adult male and female Wistar rats have been characterized and major sex-dependent biotransformations observed which correlate with the regioselectivities of known sexually differentiated hepatic P450. [6,7-<sup>3</sup>H]E<sub>2</sub> (27  $\mu$ g/kg) was given i.v. The metabolites of E<sub>2</sub> were rapidly and extensively excreted in bile (46 and 78% of the dose over 1 and 6 h, respectively). Female rats metabolized E<sub>2</sub> by one major pathway: oxidation to oestrone (E<sub>1</sub>) followed by C-2 hydroxylation and O-methylation; the principal aglycones (0–1 h bile collections) were E<sub>1</sub> (14%), 2-hydroxyE<sub>1</sub> (2-OHE<sub>1</sub>) (42%) and 2-methoxyE<sub>1</sub> (24%). Male rats metabolized E<sub>2</sub> principally by two parallel composite pathways of E<sub>1</sub> hydroxylation which yielded a complex mixture of mono- and di-oxygenated compounds: 15 $\alpha$ -OHE<sub>1</sub> (33%), 2,15 $\alpha$ -diOHE<sub>1</sub> (7%), and 2-methoxy-15 $\alpha$ OHE<sub>1</sub> (14%); 16 $\alpha$ -OHE<sub>1</sub> (13%), 2,16 $\alpha$ -diOHE<sub>1</sub> (4%) and 2-methoxy-16 $\alpha$ -OHE<sub>1</sub> (2%). 15 $\alpha$ -Hydroxylation was unique to males. The balance of aromatic and alkyl hydroxylation in males was dose-dependent: at 3 mg/kg, 15 $\alpha$ -hydroxylation was decreased approx. 50% in favour of 2-hydroxylation whilst 16 $\alpha$ -hydroxylation was largely unaffected.

The male-specific 15 $\alpha$ -hydroxylation and male-predominant 16 $\alpha$ -hydroxylation of E<sub>1</sub> derived from E<sub>2</sub> *in vivo* may be ascribable to the male-specific isoforms P450IIC13 and P450IIC11, respectively.

### INTRODUCTION

The multiplicity of regioselective and stereospecific steroid hydroxylations in adult rat liver are catalysed by microsomal P450 from several gene sub-families [1–7]. Amongst these enzymes are male-specific [3–7] and female-specific [6–8] forms with broad, yet selective, substrate specificities. Additional but non-constitutive P450 steroid hydroxylases can be induced in both sexes [1, 6, 9]. Finally, certain constitutive steroid hydroxylases are common to both sexes [6, 10]; though they may be preferentially expressed in one sex [11]. The expression of a sexual dimorphism in steroid hydroxylation could be further complicated by P450 polymorphism [2, 12–14].

The androgen and progesterone specificities of rat sex-specific and sex-predominant P450 are known in detail [1–7, 11–15]; a male-specific hepatic form determines the sexual dimorphism of androgen 16 $\alpha$ -hydroxylation [1, 4]. Less

is known about the sexual differentiation of oestrogen hydroxylation—especially alkyl hydroxylation—and the involvement of sex-specific P450. The metabolism of 17 $\beta$ -oestradiol (E<sub>2</sub>) is dominated by dehydrogenation to oestrone (E<sub>1</sub>) and subsequent hydroxylation [16]. C-2 hydroxylation, predominant in adult female rats [17–19], is the principal pathway in rat liver microsomes [20, 21]. Several P450 catalyse oestrogen 2-hydroxylation [20, 22, 23]; including male-specific forms considered responsible [20] for the sexual dimorphism of the microsomal activity [24, 26].

Male rat hepatic microsomes contain a male-specific P450 [1, 4, 22] with oestrogen 16 $\alpha$ -hydroxylase activity [15, 22]. Though another major form possessing this activity is present in both sexes [10], male microsomes are more active [27]. Nevertheless, E<sub>2</sub> 16 $\alpha$ -hydroxylation is a minor [21, 27] and strain-dependent [27] microsomal reaction. Various 16 $\alpha$ -hydroxylated metabolites of exogenous oestrogens have been isolated from rat bile [28, 29] but with incomplete qualitative and quantitative

\*To whom correspondence should be addressed.

characterization. A sexual dimorphism of oestrogen 16 $\alpha$ -hydroxylation *in vivo* has not been reported hitherto. There appears to be no previous evidence that any of the other oestrogen alkyl hydroxylations [30–33] display sexual dimorphism. However, one of the major male-specific, though strain-dependent, hepatic P450 [12–14, 22] has oestrogen 15 $\alpha$ -hydroxylase activity [15, 22] and thus represents an enzymic potential for a metabolic disparity.

In the present study, the pathways of E<sub>2</sub> hydroxylation in rats, but especially the incompletely described alkyl hydroxylations, have been characterized to determine whether they are corollaries of the regioselectivities of known sex-specific P450.

## EXPERIMENTAL

### Materials

[6,7-<sup>3</sup>H]17 $\beta$ -oestradiol ([<sup>3</sup>H]E<sub>2</sub>; 48Ci/mmol; radiochemical purity, 99%; NEN Research Products, Fed. Rep. Germany) was re-purified, as a methanol solution (128  $\mu$ Ci/ $\mu$ mol), on a Techopak column (C<sub>18</sub>, 10  $\mu$ m; 30  $\times$  0.39 cm i.d.; HPLC Technology, Macclesfield, England) by elution with acetonitrile (20 to 75% over 40 min) in 43 mM ammonium dihydrogen orthophosphate, pH 3.0, at 1.5 ml/min. It was recovered by extraction into ether following evaporation of the acetonitrile under nitrogen. Unlabelled E<sub>2</sub>, other oestrogen standards, oestrogen conjugates,  $\beta$ -glucuronidase (Glucurase) and  $\beta$ -glucuronidase-arylsulphohydrolase (Type H-2; 4.2  $\times$  10<sup>3</sup> U/ml arylsulphohydrolase) were from Sigma Chemical Co. (Poole, England). 15 $\alpha$ -Hydroxyoestrone (15 $\alpha$ -OHE<sub>1</sub>) was generously provided by Dr M. Levitz, New York University Medical Centre (NY, USA). Boron tribromide (1 M in dichloromethane) was from Aldrich Chemical Co. (Gillingham, England).

### Animal experiments

Male (220  $\pm$  18 g body wt; mean  $\pm$  SD,  $n$  = 4) and female (220  $\pm$  20 g;  $n$  = 4) Wistar rats were anaesthetized with urethane (1.4 g/kg body wt, in isotonic saline given *i.p.*) and cannulated via the jugular vein and common bile duct. [<sup>3</sup>H]E<sub>2</sub> (1–3  $\mu$ Ci) was administered *i.v.* in saline-polyethylene glycol 200 (1:1, *v/v*) at a dose of 0.1  $\mu$ mol/kg (27.2  $\mu$ g/kg). Bile was collected hourly for 6 h, and assayed for radioactivity [34].

### Chromatographic analysis of biliary metabolites

Bile (10–20  $\mu$ l), deconjugated biliary metabolites (50–80  $\mu$ l methanol solutions) and co-injected unlabelled standards were analysed on a Techopak column linked to an LKB 2150-2152 pump-controller and a Gilson IIIB u.v. (280 nm) detector. Biliary conjugates (15–50  $\times$  10<sup>3</sup> dpm) were resolved with methanol (20 to 60% over 40 min) in 10 mM ammonium acetate, pH 6.9, and aglycones (5–50  $\times$  10<sup>3</sup> dpm) with acetonitrile (20 to 75% over 40 min) in 43 mM ammonium dihydrogen orthophosphate, pH 3.0. The flow rate was 1.5 ml/min. Pairs of polar ketols and methoxyketols, irresolvable on the C<sub>18</sub> column, were separated on a diol column (10  $\mu$ m, 25  $\times$  0.46 cm i.d.; HPLC Technology) with propan-2-ol in hexane (10% for 10 min, then 10–20% over 20 min; flow rate, 1.5 ml/min) after isolation by C<sub>18</sub>-HPLC. They were recovered from the reversed-phase eluate by ether extraction, and reconstituted in methanol for diol-phase HPLC. Metabolites were quantified by scintillation counting: 30 s eluate fractions were dissolved in 4 ml of scintillant (Aqua Luma Plus; Lumac bv, The Netherlands). Recoveries of chromatographed radioactivity were 90–100%.

### Enzymic hydrolysis

Bile (20–50  $\mu$ l; 20–80  $\times$  10<sup>3</sup> dpm; 50–200 ng equivalent E<sub>2</sub>) was incubated with either  $\beta$ -glucuronidase (5  $\times$  10<sup>3</sup> U) or  $\beta$ -glucuronidase-arylsulphohydrolase (approx. 1.0  $\times$  10<sup>3</sup> and 42 U, respectively) in 1.0 ml 0.1 M sodium acetate, pH 5.0, containing 10 mM ascorbate at 37°C for 16 h; the ascorbate prevented oxidative loss of catechol metabolites [35]. Combined ether extracts (5 ml  $\times$  2) of the incubations were evaporated under nitrogen, and reconstituted in 100  $\mu$ l methanol for HPLC.

### Isolation and identification of metabolites

*Isolation by reversed-phase HPLC.* Anaesthetized and cannulated male Wistar rats (350 g) were administered [<sup>3</sup>H]E<sub>2</sub> (3.0 mg/kg; 5.0  $\mu$ Ci) in 0.3 ml polyethylene glycol 200–isotonic saline (2:1, *v/v*) given *i.v.* Bile (0.4–2.7 ml) was collected for 3 h; 68–72% of the dose was recovered. Radiochromatographic analysis of the aglycones revealed substantial dose-related changes [Fig. 2(B and C)]. The metabolite profile combined the male and female low-dose profiles, plus a small amount of E<sub>2</sub>, with elevated proportions of the most polar male-specific metabolites [M- $\alpha$ , R, (<sup>3</sup>H peak) 7.5 min, and

M- $\beta$ , R, 9.0 min]. Three steroids—2-OHE<sub>1</sub> (17.5 min), E<sub>1</sub> (22.5 min) and 2-methoxyE<sub>1</sub> (23.5 min)—were ascribed by co-elution with standards. The remaining fractions, i.e. male-specific M-A (12.0 min) and male-predominant M-B (13.5 min), co-eluted with authentic  $15\alpha$ -OHE<sub>1</sub> and  $16\alpha$ -OHE<sub>1</sub>, respectively, but mass spectrometry (MS) and HPLC showed them to be heterogeneous.

Bile (250  $\mu$ l; 100  $\mu$ g equivalent) was incubated with  $\beta$ -glucuronidase-arylsulphohydrolase ( $2.5 \times 10^3$  and 105 U, respectively) in 2.5 ml of acetate buffer containing 10 mM ascorbate at 37°C for 16 h. The ether (7 ml  $\times$  2)-extracted aglycones, representing 87% of incubated radioactivity, were reconstituted in methanol

(500  $\mu$ l). Portions (70–120  $\mu$ l; 12–21  $\mu$ g equivalent) were chromatographed on a Techopak (C<sub>18</sub>) column. E<sub>2</sub>, five metabolites and two heterogeneous fractions (83% of chromatographed radioactivity) were eluted with acetonitrile-phosphate buffer, and recovered by ether extraction (2 vol  $\times$  2). The isolated metabolites and fractions (1–18  $\mu$ g) were radiochromatographically homogeneous on the C<sub>18</sub> column. Aliquots were analysed by electron impact (EI) and chemical ionization (CI) MS.

Detailed chromatographic analysis of M-B (16  $\mu$ g) revealed two sub-fractions separated by 0.7 min; base-line resolution of u.v. absorbance but not of radioactivity by fraction collection was achieved. The first-eluted component




Fig. 1. High-performance liquid radiochromatograms of the biliary metabolites of [ $^3$ H]E<sub>2</sub> (27  $\mu$ g/kg) in (A) female (B) male rats at 0–1 h. The metabolites and unlabelled standards were resolved on a C<sub>18</sub> column using a gradient of methanol in ammonium acetate.

exactly co-chromatographed with authentic  $16\alpha$ -OHE<sub>1</sub>,  $16\alpha$ -OHE<sub>1</sub> and its methoxylated co-fraction ( $m/z$  316 in spectrum of M-B) were isolated (ratio of radioactivity, 0.95:1.0) and characterized.

*Isolation by diol-phase HPLC.* EI spectra of M-A indicated  $15\alpha$ -OHE<sub>1</sub> [ $M$ ]<sup>+</sup> at  $m/z$  286, relative intensity (RI) 72%—acquired spectrum was computer-matched with standard's spectrum—and a methoxylated derivative ([ $M$ ]<sup>+</sup> at  $m/z$  316; RI 100%). These were irresolvable on the C<sub>18</sub> column but could be separated ( $R$ , 19.5 and 21.5 min; ratio of radioactivity, 0.75:1.0) on a diol column with propan-2-ol in hexane (10% for 10 min, then 10–20% over 20 min) at 1.5 ml/min; the second-eluted component co-chromatographed with authentic  $15\alpha$ -OHE<sub>1</sub>. Isolated M-A (21  $\mu$ g) was fractionated on the diol column, and yielded radiochromatographically homogeneous  $15\alpha$ -OHE<sub>1</sub> and methoxylated  $15\alpha$ -OHE<sub>1</sub> which was 86% pure. The sub-fractions were analysed by MS.

*Chemical demethylation of methoxylated metabolites.* Characterization of the methoxylated, i.e. methylated catechol, metabolites and their putative catechol precursors was aided by demethylation with boron tribromide [36]. Authentic 2-methoxylated E<sub>1</sub>, E<sub>2</sub> and  $16\alpha$ -OHE<sub>2</sub> (50–100  $\mu$ g; approx. 0.16–0.33  $\mu$ mol) were dissolved in 1 ml dichloromethane in a conical glass tube, the head-space flushed with nitrogen, and boron tribromide (10–20  $\mu$ l; 10–20  $\mu$ mol) injected via a septum cap. After 3–4 h at 20°C, the reaction was terminated by adding water (2 ml) and the products extracted into ether (6 ml  $\times$  2). Extensive (75–80%) or quantitative formation of catechol was established by C<sub>18</sub>-HPLC and MS. 2-methoxyE<sub>1</sub>, M-A and M-B (20–28  $\mu$ g), isolated from the bile hydrolysates by C<sub>18</sub>-HPLC, were reacted with boron tribromide (50  $\mu$ l) for 1.5–2 h. Essentially quantitative demethylation of 2-methoxyE<sub>1</sub> and the methoxylated sub-fractions of M-A and M-B was confirmed by HPLC and MS; only traces of (non-polar) side products were formed. 2-OHE<sub>1</sub> co-eluted with its standard, and the ketol sub-fractions of M-A and M-B with  $15\alpha$ -OHE<sub>1</sub> and  $16\alpha$ -OHE<sub>1</sub>, respectively. The ketols and the catechols liberated from their co-fractions were isolated (3–6  $\mu$ g) by C<sub>18</sub>-HPLC for MS.

#### MS

Standards and metabolites were analysed by direct-probe EI and CI (isobutane) MS using a VG TS-250: electron energy, 70 eV (EI) or 50 eV

(CI); filament emission, 500  $\mu$ A (EI) or 300  $\mu$ A (CI); source temperature, 180°C; accelerating voltage,  $4 \times 10^3$  V. Spectra were acquired via a VG 11/250J data system over  $m/z$  50 (EI)/75 (CI)–600 (scan time, 1 s) at resolution 800.

## RESULTS

### *Excretion of radiolabelled metabolites*

The radiolabelled metabolites of i.v.-administered [<sup>3</sup>H]E<sub>2</sub> (27  $\mu$ g/kg) were rapidly and extensively excreted in bile by both sexes:  $51.6 \pm 3.2\%$  (mean  $\pm$  SD,  $n = 4$ ) and  $40.6 \pm 8.5\%$  of the dose during the first hour by males and females, respectively;  $73.8 \pm 1.7$  and  $65.9 \pm 4.6\%$  over 3 h;  $79.2 \pm 2.4$  and  $76.8 \pm 9.0\%$  over 6 h.

### *Biliary metabolites*

Female bile contained three major radiolabelled conjugates ( $R$ , 19, 20.5 and 23.5 min) and several minor ones [Fig. 1(A)]. Collectively, the three represented 72–78% of the radiolabel excreted during the first hour; with the principal metabolite ( $R$ , 23.5 min) representing 43–47% (mean, 46%). The conjugate of  $R$ , 19 min co-eluted with authentic E<sub>1</sub>  $\beta$ -glucuronide. Male rats formed a greater number of more polar conjugates [Fig. 1 (B)]. Since 75–86 and 81–89% of the biliary radiolabel in both sexes was ether-extractable after incubations with  $\beta$ -glucuronidase and  $\beta$ -glucuronidase-arylsulphohydrolase, respectively, whereas only 4–6% was immediately extractable, it is evident that the conjugates were predominantly glucuronides.

### *Identification of steroidal metabolites*

E<sub>2</sub>, E<sub>1</sub>, 2-OHE<sub>1</sub> ( $M$ <sup>+</sup>  $m/z$  286, RI 100%) and 2-methoxyE<sub>1</sub> ( $M$ <sup>+</sup>  $m/z$  300, RI 100%) isolated from enzymic hydrolysates of bile [Fig. 2(C)] were identified from their EI spectra; which matched those of unlabelled standards (data not shown).

*Ketol metabolites.*  $15\alpha$ -OHE<sub>1</sub> and  $16\alpha$ -OHE<sub>1</sub> were identified by EI [Table 1; Figs 3(A) and 4(A), respectively] and CI MS. Their EI spectra matched library spectra [37] and acquired spectra of standards [Fig. 3(A and B)]. Ring-D fragmentation yielded diagnostic ions at characteristic RI: metabolic and authentic  $15\alpha$ -OHE<sub>1</sub> gave  $m/z$  214 ([ $M-C_3O_2H_4$ ]<sup>+</sup>) and  $m/z$  213 ([ $M-C_3O_2H_3$ ]<sup>+</sup>) in the RI ratio 5:22 and 7:20, respectively, whereas the corresponding ratios for  $16\alpha$ -OHE<sub>1</sub> were 34:100 and 42:98. The




Fig. 2. High-performance liquid radiochromatograms of the phase I metabolites of  $[^3\text{H}]E_2$  liberated from conjugates in rat bile: (A) females,  $27 \mu\text{g}/\text{kg}$ , 0-1 h; (B) males,  $27 \mu\text{g}/\text{kg}$ , 0-1 h; (C) males,  $3 \text{ mg}/\text{kg}$ , 0-3 h. The metabolites and unlabelled standards were resolved on a  $C_{18}$  column using a gradient of acetonitrile in ammonium dihydrogen orthophosphate.

metabolites' spectra were distinct from those of 14-OHE<sub>1</sub>, 12β-OHE<sub>1</sub>, 11β-OHE<sub>1</sub>, 6β-OHE<sub>1</sub>, 16-keto-17βE<sub>2</sub>, 16-keto-17αE<sub>2</sub> and 6-keto-17βE<sub>2</sub> [37]. These identifications were supported by CI MS. Both metabolites and their standards gave base-peak [M + 1]<sup>+</sup> at *m/z* 287. The pseudomolecular ions underwent regioselective dehydration [39]: the RI ratio [M + 1-18]: [M + 1] for 15α- and 16α-OHE<sub>1</sub> was 68:100 and 21:100, respectively.

*Hydroxyketol (catechol) metabolites.* M-α and M-β [Fig. 2(B)] gave dioxygenated M<sup>+</sup> at *m/z* 302 [Figs 3(C) and 4(B)]. Under CI conditions, *m/z* 303 ([M + 1]<sup>+</sup>) and *m/z* 285 ([M + 1-18]<sup>+</sup>) were obtained; as with the ketols, the more polar metabolite, i.e. M-α, underwent greater dehydration. M-α and M-β were deduced to be C-2 catechol, rather than cycloalkanol, derivatives of 15α- and 16α-OHE<sub>1</sub>, respectively, from EI spectra and by comparison with the catechols liberated from the methoxylated ketols. Thus the peaks at *m/z* 227, 229 and 230 (Table 1) are ring-D fragmentation ions for catechols [37, 38], i.e. analogues of *m/z* 211, 213 and 214 in the ketols' spectra, and the greater fragmentation of M-β is typical of a 16-OH ketol. Those at *m/z* 162, 173, 175 and 188 are catechol fragments containing rings A and B [37] whilst the peaks at *m/z* 123, 131 and 136 appear to be oxygenated ring-A fragments [28, 37]. A structural connection between the dioxygenated polar metabolites and the methoxyketols was conclusively established by demethylation; boron tribromide effects quantitative demethylation of aryl methylethers under mild conditions [36]. The catechols obtained from 2-methoxy-15αOHE<sub>1</sub> and 2-methoxy-16αOHE<sub>1</sub> (*vide infra*) co-eluted with M-α and M-β, respectively, and yielded corresponding EI and CI spectra.

*Methoxyketol (methylated catechol) metabolites.* Putative 2-methoxy-15αOHE<sub>1</sub> and 2-methoxy-16αOHE<sub>1</sub> were isolated by diol- and C<sub>18</sub>-HPLC, respectively; the site of methoxylation was indicated by the predominance of C-2 hydroxylation. Their EI spectra [Table 1; Figs 3(D) and 4(C)] contained the fragments diagnostic of 2-methoxyoestrogens [37, 40], including 2-methoxy-16αOHE<sub>1</sub> [28, 29], e.g. *m/z* 137, 176, 189, 202, 215, 229, 243 and 244. The peaks at *m/z* 243 (16α-OH isomer only) and 244, analogues of *m/z* 213 and 214, respectively, in the ketols' spectra, had RI indicative of 15α- and 16α-hydroxylated isomers. CI spectra contained pseudomolecular base peaks at *m/z* 317 and dehydration fragments at *m/z* 299. The RI ratio [M-H<sub>2</sub>O]:[M + 1] for the methoxylated co-fractions of 15α-OHE<sub>1</sub> and 16α-OHE<sub>1</sub>—28:100 and 11:100, respectively—conformed with those for the corresponding ketols.

Schematic metabolic pathways based upon the above identifications are shown in Fig. 5. However, they represent the minimum number of reactions considered necessary to explain the sexual differentiation of metabolism rather than an attempt to incorporate every possible transformation.

#### Quantitation of steroidal metabolites

The phase EI metabolites of low-dose [<sup>3</sup>H]E<sub>2</sub> (27 μg/kg) displayed an unambiguous sexual dimorphism [Table 1; Fig. 2 (A and B)]. Unchanged E<sub>2</sub> was either absent or present in only trace amounts (approx. 1% deconjugated radioactivity). The major metabolites formed by females, i.e. E<sub>1</sub>, 2-OHE<sub>1</sub> and 2-methoxyE<sub>1</sub>, represented 80% of the radiolabel liberated from conjugates in 0–1 h bile collections. Whilst

Table 1. Molecular ions and principal fragments of the polar metabolites of [<sup>3</sup>H]E<sub>2</sub> isolated from rat bile

Metabolite	Ions (RI)
15α-OHE <sub>1</sub>	286 (M <sup>+</sup> , 100), 268 (9), 240 (8), 225 (7), 214 (22), 213 (5), 211 (7), 199 (8), 186 (36), 172 (17), 159 (67), 157 (18), 146 (45), 133 (21), 120 (13).
2-OH-15αOHE <sub>1</sub>	302 (M <sup>+</sup> , 100), 284 (32), 242 (6), 230 (11), 227 (16), 215 (8), 213 (9), 202 (12), 188 (11), 175 (24), 173 (28), 162 (24), 136 (11).
2-Methoxy-15αOHE <sub>1</sub>	316 (M <sup>+</sup> , 100), 244 (21), 230 (4), 216 (12), 202 (4), 189 (19), 187 (12), 176 (27), 159 (17), 150 (11), 137 (17).
16α-OHE <sub>1</sub>	286 (M <sup>+</sup> , 100), 258 (8), 214 (100), 213 (34), 201 (18), 199 (20), 186 (11), 172 (54), 159 (46), 157 (23), 146 (33), 133 (22), 115 (15).
2-OH-16αOHE <sub>1</sub>	302 (M <sup>+</sup> , 100), 230 (54), 229 (46), 215 (18), 202 (16), 188 (35), 175 (33), 173 (52), 162 (21), 161 (32), 131 (29), 123 (23).
2-Methoxy-16αOHE <sub>1</sub>	316 (M <sup>+</sup> , 100), 288 (3), 244 (36), 243 (14), 229 (8), 215 (4), 202 (15), 189 (14), 187 (12), 176 (11), 163 (8), 145 (7), 137 (26).

Spectra were obtained by direct-probe EI.




Fig. 3. Direct-probe electron impact mass spectra of  $15\alpha$ -hydroxylated metabolites of  $[^3\text{H}]E_2$  formed in male rats: (A)  $15\alpha$ -OHE<sub>1</sub>; (B) authentic  $15\alpha$ -OHE<sub>1</sub>; (C) M- $\alpha$ [2-OH- $15\alpha$ OHE<sub>1</sub>]; and (D) 2-methoxy- $15\alpha$ OHE<sub>1</sub>. Metabolites were isolated from enzymic hydrolysates of bile by HPLC.

E<sub>1</sub> and 2-OHE<sub>1</sub> were consistent though minor products in males, 2-methoxyE<sub>1</sub> was only a trace metabolite (0–1.5% of deconjugated radiolabel).  $16\alpha$ -OHE<sub>1</sub> and, by chromatographic implication, 2-methoxy- $16\alpha$ OHE<sub>1</sub>, were minor female metabolites but  $15\alpha$ -OHE<sub>1</sub> and 2-methoxy- $15\alpha$ OHE<sub>1</sub>, and both the catechol intermediates, appeared to be unique to males.

Male-specific, i.e.  $15\alpha$ -oxygenated, and male-predominant, i.e.  $16\alpha$ -oxygenated, products collectively represented an estimated 54 and 19.5%, respectively, of the males' phase I metabolites (Table 2). Chromatographic ( $C_{18}$ -HPLC) comparison of the metabolites with unlabelled standards indicated that the following were not present in appreciable quantities:

2,16 $\alpha$ -diOHE<sub>2</sub> (2-OH oestriol; *R*<sub>t</sub> 6.5 min), 16 $\alpha$ -OHE<sub>2</sub> (9.5 min), 2-methoxy-16 $\alpha$ OHE<sub>2</sub> (10.0 min) 6- and 16-ketoE<sub>2</sub> (12.5 min; standards not resolved), 4-OHE<sub>1</sub> (18.5 min) and 2-methoxyE<sub>2</sub> (20.5 min). 6 $\beta$ -OHE<sub>2</sub> co-chromatographed with *M*- $\beta$  [2-OH-16 $\alpha$ OHE<sub>1</sub>; Fig. 2(B)] but the latter's mass spectra clearly indicated a molecular weight of 302. 2-OHE<sub>2</sub> (15.5 min) may have been formed in trace amounts by female rats [Fig. 2(A)].

The male and female metabolite profiles remained qualitatively unchanged throughout the 3 h after dosing and the proportion of each metabolite in males was essentially constant. All the quantitative changes observed in females were slight and inconsistent.

The regioselective hydroxylation of E<sub>1</sub> formed from E<sub>2</sub> in males was dose-dependent between 27  $\mu$ g/kg and 3 mg/kg [Fig. 2(B and C)]: the

proportions of the hydroxylations at C-2, C-15 and C-16, estimated from a combination of diol- and C<sub>18</sub>-HPLC analyses, changed from approx. 29:53:18 (Table 2), respectively, to 59:24:17; 15 $\alpha$ -hydroxylation declined approx. 50% in favour of 2-hydroxylation whilst 16 $\alpha$ -hydroxylation was largely unaffected. Consequently, the major metabolites at 3 mg/kg were 2-OHE<sub>1</sub> (20% radiolabel), 2-methoxyE<sub>1</sub> (15%), 15 $\alpha$ -OHE<sub>1</sub> (12%), 2-methoxy-15 $\alpha$ OHE<sub>1</sub> (10%) and 16 $\alpha$ -OHE<sub>1</sub> (10%).

## DISCUSSION

The present studies have identified and quantified the hitherto poorly characterized components of the "polar" fraction of E<sub>2</sub>s biliary metabolites in rats [28]. A sexual




Fig. 4. Direct-probe electron impact mass spectra of 16 $\alpha$ -hydroxylated metabolites of [<sup>3</sup>H]E<sub>2</sub> formed in male rats: (A) 16 $\alpha$ -OHE<sub>1</sub>; (B) *M*- $\beta$ [2-OH-16 $\alpha$ OHE<sub>1</sub>]; and (C) 2-methoxy-16 $\alpha$ OHE<sub>1</sub>. Metabolites were isolated from enzymic hydrolysates of bile by HPLC.






Fig. 5. Scheme of proposed major pathways of  $E_2$  metabolism in male (---) and female (—) rats at a dose of 27  $\mu\text{g}/\text{kg}$ .  $E_1$ , 2-OHE $_1$  and 2-methoxy $E_1$  were the major metabolites in females; 15 $\alpha$ -OHE $_1$ , 16 $\alpha$ -OHE $_1$  and 2-methoxy-15 $\alpha$ OHE $_1$  the major metabolites in males.

differentiation of alkyl hydroxylations—predictable in essence from the known regioselectivities of male-specific hepatic  $P450$  [15, 22]—was established. They, rather than the previously emphasized aromatic hydroxylations [16], were the basis of a sexual dimorphism of  $E_2$  metabolism.

The high ratio of aromatic (C-2) to alkyl (C-16 $\alpha$ ) hydroxylation in female rats approximately reflects the relative enzyme activities of rat hepatic microsomes [21, 24, 27], and yielded a profile of C-2 oxygenated metabolites similar to those described previously [18, 19]. In contrast, and despite the 2.5- to 7-fold higher oestrogen C-2 hydroxylase activity of male microsomes [20, 26], alkyl hydroxylations, i.e.

C-15 $\alpha$  and C-16 $\alpha$ , either alone or in combination with C-2 hydroxylation, were dominant in males: the ratio of ring-D to ring-A oxygenated products was approx. 71:29 and 5:67 in males and females, respectively.

16 $\alpha$ -OHE $_1$  and 2-methoxy-16 $\alpha$ OHE $_1$  were known but previously unquantified metabolites of exogenous  $E_2$  in rats [29]. The male-predominant expression of 16 $\alpha$ -hydroxylation *in vivo* accords with the 2-fold greater 16 $\alpha$ -hydroxylase activity of hepatic microsomes from male Wistar rats [27]. Although partial sexual dimorphisms could arise from sex-predominant expression of a single  $P450$  form [11], the present case can be explained by the combined activities of a male-specific  $P450$  and a non-specific form. The latter,  $P450_{\text{CM/F}}$  [10], is constitutively present in the livers of both male and female Sprague-Dawley rats and actively catalyses 2- and 16 $\alpha$ -hydroxylation of  $E_2$ ; it represents approx. 10–20% of the  $P450$  in hepatic microsomes from both sexes [10]. Male-predominant 16 $\alpha$ -hydroxylation would result from the greater oestrogen 16 $\alpha$ -hydroxylase activity [15, 22] of male-specific  $P450_{\text{IIC11}}$  (equivalent preparations have been termed RLM5 [15], UT-2 [22],  $P450_{\text{h}}$ , UT-A, or  $P450_{\text{2c}}$  [1–4]), a multi-functional steroid hydroxylase [3–6, 22], which is induced at puberty exclusively in males [1] and represents approx. 50% of spectrally-detectable  $P450$  in male rat liver microsomes [5].  $P450_{\text{IIC11}}$  additionally catalyses androgen 16 $\alpha$ -hydroxylation [4] and oestrogen 2-hydroxylation [15, 22], and is

Table 2. The metabolites of [ $^3\text{H}$ ] $E_2$  (27  $\mu\text{g}/\text{kg}$ ) liberated from conjugates in the bile of male and female rats

Metabolite	% Radioactivity (mean $\pm$ SD, $n = 4$ )	
	Males	Females
Oestrone ( $E_1$ )	6.7 $\pm$ 1.0	13.5 $\pm$ 3.4
2-OHE $_1$	3.0 $\pm$ 0.7	42.4 $\pm$ 3.6
2-Methoxy $E_1$	0–1.5	24.3 $\pm$ 6.1
Fraction M-A	47.2 $\pm$ 0.3	0
15 $\alpha$ -OHE $_1$ <sup>a</sup>	33	
2-OH-15 $\alpha$ OHE $_1$ (M- $\alpha$ )	6.7 $\pm$ 0.9	0
2-Methoxy-15 $\alpha$ OHE $_1$ <sup>a</sup>	14	0
Fraction M-B	17.1 $\pm$ 1.3 <sup>b</sup>	5.2 $\pm$ 2.5
16 $\alpha$ -OHE $_1$ <sup>a</sup>	13	ND
2-OH-16 $\alpha$ OHE $_1$ (M- $\beta$ )	4.4 $\pm$ 0.6	0
2-Methoxy-16 $\alpha$ OHE $_1$ <sup>a</sup>	2	ND

Biliary conjugates (0–1 h collections) were hydrolyzed with enzymes. Ether-extracted metabolites were quantified by radiometric  $C_{18}$ -HPLC except as indicated. ND = not determined.

<sup>a</sup>Metabolite quantified by diol-phase radiometric HPLC after isolation of relevant heterogeneous metabolite fraction from hydrolysate of pooled bile by reversed-phase HPLC (mean 2/3 analyses).

<sup>b</sup>Comprised of two minor, more polar, sub-fractions in addition to 16 $\alpha$ -OHE $_1$  and 2-methoxy-16 $\alpha$ OHE $_1$ .

thought to determine the sexual differentiation not only of these reactions [4, 20] but also of transformations of certain non-steroidal substrates [41].

Though it was the principal male pathway, and the only sex-specific transformation, observed in this study, oestrogen 15-hydroxylation is a poorly characterized reaction.  $15\alpha$ -OHE<sub>1</sub> is a minor biliary metabolite of endogenous oestrogens in man [42], and oestrogen 15 $\alpha$ -hydroxylation has been detected in human foetal liver [43], in bovine [30] and human adrenals [44], and in porcine ovary [33]. 11 $\beta$ -Methoxy-17 $\alpha$ -ethinyloestradiol undergoes substantial 15 $\alpha$ -hydroxylation in man but none in rats [45], and 15-hydroxylated oestrogens were apparently otherwise unknown as metabolites of exogenous substrates *in vivo*. The activity in porcine blastocysts is *P450*-mediated though the isoform(s) has not been characterized [33]. However, hepatic microsomes from female 129/J mice have no detectable oestrogen 15 $\alpha$ -hydroxylase activity even though a *P450* purified from them catalyses this activity in a reconstructed system [46]. Hepatic microsomes from male Sprague-Dawley rats likewise have little [22] or no detectable [15] activity and those from male Wistar rats no apparent activity [32]. The cause of the ineffective catalysis of 15 $\alpha$ -hydroxylation by microsomal *P450* *in situ* is unknown.

The only rat hepatic *P450* known to catalyse significant oestrogen 15 $\alpha$ -hydroxylation [22], *P450*IIC13 [14] (equivalent preparations have been termed RLM3 [15], *P450g* [3, 12] or UT-5 [22]), is a male-specific and multi-functional steroid hydroxylase [3, 15, 47] subject to complex sexual dimorphism [12–14]. RLM3 and UT-5 also possess oestrogen C-2 hydroxylase activity [15, 22], and the UT-5 preparation, at least, hydroxylates E<sub>2</sub> at C-16 $\alpha$  [22], but their contributions to microsomal catalysis of these activities are likely to be insignificant [22]. The expression of *P450g* is complicated by a strain- and colony-dependent polymorphism (10- to 20-fold variation) in outbred rats [13, 14] and either very low or, uncommonly, high levels of *P450g* in inbred strains [12, 13]. The *P450g*-phenotype status of the male Wistars used in this study is unknown but, at least, a high frequency of the (+g)-phenotype [12] is predicted. Nevertheless, *P450g* polymorphism clearly implies polymorphic expression of E<sub>2</sub> 15 $\alpha$ -hydroxylation in certain strains. Although 15 $\alpha$ -hydroxylated oestrogens are minor prod-

ucts of purified *P450*<sub>C-M/F</sub> [10], their complete absence from female bile indicates effective non-expression of this enzyme's 15 $\alpha$ -hydroxylase activity *in vivo*.

The 2-hydroxyketol metabolites of E<sub>2</sub> could arise from either concerted or sequential hydroxylations. A concerted mechanism for metabolism of progesterone to dioxygenated products by *P450g* has indeed been described [48]. However, secondary metabolites were seemingly not observed during a study of the products formed by RLM3 and RLM5 [15] whilst 2-OHE<sub>2</sub> is known to be a good substrate for 16 $\alpha$ -hydroxylation in Wistar rats [28]. The probability that sequential hydroxylations at C-2 and ring-D predominate *in vivo* might be inferred from the dominant formation by isolated male rat livers of C-2 monooxygenated E<sub>2</sub> and E<sub>1</sub> metabolites during single-passage perfusion [49].

Despite the emphasis placed upon the role of hepatic *P450*, the metabolism of E<sub>2</sub> to 15 $\alpha$ -OHE<sub>2</sub> by adrenal glands [30] and the presence of male-predominant microsomal *P450* in guinea pig adrenals [50] indicate a potential extra-hepatic contribution to 15-hydroxylation.

In conclusion, it would appear that the regioselective oestrogen hydroxylase activities of two major male-specific *P450*s, as determined in reconstructed systems, are predictive of E<sub>2</sub>'s sexually differentiated metabolism in Wistar rats whereas the activities manifested by intact hepatic microsomes and isolated livers are not. Subsequent publications will describe the complex blocking, impeding and re-directing of oestrogen biotransformations by ring-A halogen substituents.

*Acknowledgements*—P. Morgan was supported by the Department of Education, Northern Ireland. B. K. Park is a Wellcome Principal Research Fellow. The TS-250 mass spectrometer in the Department of Pharmacology and Therapeutics was purchased with generous grants from the Wellcome Trust and the University of Liverpool. We thank Miss S. Newby for her assistance and Miss S. Oliphant for typing the manuscript.

## REFERENCES

1. Waxman D. J.: Interactions of hepatic cytochromes *P450* with steroid hormones. Regioselectivity and stereospecificity of steroid hydroxylation and hormonal regulation of rat *P450* enzyme expression. *Biochem. Pharmac.* **37** (1988) 71–84.
2. Gonzalez F.: The molecular biology of cytochrome *P450*s. *Pharmac. Rev.* **40** (1989) 243–288.
3. Ryan D. E., Iida S., Wood A. W., Thomas P. E., Lieber C. S. and Levin W.: Characterization of three highly

- purified cytochromes P450 from hepatic microsomes of adult male rats. *J. Biol. Chem.* **259** (1984) 1239–1250.
4. Waxman D. J.: Rat hepatic cytochrome P450 isoenzyme 2c: identification as a male-specific developmentally induced steroid 16 $\alpha$ -hydroxylase and comparison to a female-specific cytochrome P450 isoenzyme. *J. Biol. Chem.* **259** (1984) 15481–15490.
  5. Morgan E. T., MacGeoch C. and Gustafsson J.-A.: Hormonal and developmental regulation of expression of the hepatic microsomal 16 alpha-hydroxylase cytochrome P450 apoprotein in the rat. *J. Biol. Chem.* **260** (1985) 11895–11898.
  6. Waxman D. J., Dannan G. A. and Guengerich F. P.: Regulation of rat hepatic cytochrome P450: age-dependent expression, hormonal imprinting and xenobiotic inducibility of sex-specific isozymes. *Biochemistry* **24** (1985) 4409–4417.
  7. Matsunaga T., Nagata K., Holsztyńska E. J., Lapenson D. P., Smith A., Kato R., Gelboin H. V., Waxman D. J. and Gonzalez F. J.: Gene conversion and differential regulation in the rat P450IIA gene subfamily: purification, catalytic activity, cDNA and deduced amino acid sequence, and regulation of an adult male-specific hepatic testosterone 15 $\alpha$ -hydroxylase. *J. Biol. Chem.* **263** (1988) 17995–18002.
  8. MacGeoch C., Morgan E. T., Halpert J. and Gustafsson J.-A.: Purification, characterization and pituitary regulation of the sex-specific cytochrome P450 15 $\beta$ -hydroxylase from liver microsomes of untreated female rats. *J. Biol. Chem.* **259** (1984) 15433–15439.
  9. Namkung M. J., Yang H. L., Hulla J. E. and Juchau M. R.: On the substrate specificity of cytochrome P450III A1. *Molec. Pharmacol.* **34** (1988) 628–637.
  10. Sugita O., Sassa S., Miyairi S., Fishman J., Kubota I., Noguchi T. and Kappas A.: Cytochrome P450<sub>C-M/F</sub>, a new constitutive form of microsomal cytochrome P450 in male and female rat liver with estrogen 2- and 16 $\alpha$ -hydroxylase activity. *Biochemistry* **27** (1988) 678–686.
  11. Waxman D. J., Morrissey J. J. and Le Blanc G. A.: Female-predominant rat hepatic P450 forms j (IIE1) and 3 (IIA1) are under hormonal regulatory controls distinct from those of the sex-specific P450 forms. *Endocrinology* **124** (1989) 2954–2966.
  12. Rampersand A., Bandiera S., Ryan D. E., Levin W., Thomas P. E. and Walz F. G.: A regulatory polymorphism for rat hepatic cytochrome P450g. *Archs Biochem. Biophys.* **252** (1987) 145–151.
  13. McClellan-Green P., Waxman D. J., Caveness M. and Goldstein J. A.: Phenotypic differences in expression of cytochrome P450g but not its mRNA in outbred male Sprague-Dawley rats. *Archs Biochem. Biophys.* **253** (1987) 13–25.
  14. Yeowell H. N., McClellan-Green P. D., Negishi M. and Goldstein J. A.: Characterization of a cDNA for the unexpressed form of cytochrome P450 from the (–g) rat and differentiation of its messenger RNA from that of the (+g) phenotype using specific oligoprobes. *Biochemistry* **29** (1990) 713–719.
  15. Cheng K.-C. and Schenkman J. B.: Metabolism of progesterone and estradiol by microsomes and purified cytochrome P450 RLM<sub>3</sub> and RLM<sub>5</sub>. *Drug Metab. Dispos.* **12** (1984) 222–234.
  16. Fishman J.: Aromatic hydroxylation of estrogens. *A. Rev. Physiol.* **45** (1983) 61–72.
  17. Bolt H. M., Kappas H. and Remmer H.: Studies on the metabolism of ethynylestradiol *in vitro* and *in vivo*: the significance of 2-hydroxylation and the formation of polar products. *Xenobiotica* **3** (1973) 773–785.
  18. Bartke A., Steele R. E., Williams J. G. and Williams K. I. H.: Biliary metabolites of <sup>14</sup>C-estrone and <sup>14</sup>C-estradiol from the rat. *Steroids* **18** (1971) 303–311.
  19. Watanabe H.: Effect of 17 $\alpha$ -ethynylestradiol on the biliary metabolites of 17 $\beta$ -[6,7-<sup>3</sup>H<sub>2</sub>]estradiol in the rat. *Biochim. Biophys. Acta* **231** (1971) 399–405.
  20. Dannan G. A., Porubek D. J., Nelson S. D., Waxman D. J. and Guengerich F. P.: 17 $\beta$ -estradiol 2- and 4-hydroxylation catalysed by rat hepatic cytochrome P450: role of individual forms, inductive effects, developmental patterns, and alterations by gonadectomy and hormone replacement. *Endocrinology* **118** (1986) 1952–1960.
  21. Numazawa M. and Satoh S.: 2-bromo- and 16-bromo-estrogens and related compounds: potential inhibitors of both estradiol 2- and 16 $\alpha$ -hydroxylases in rat liver microsomes. *J. Steroid Biochem.* **33** (1989) 111–117.
  22. Watanabe K., Takahashi K., Imaoka S., Funae Y., Kawano S., Inoue K., Kamataki T., Takagi H. and Yoshizawa I.: Comparison of cytochrome P-450 species which catalyze the hydroxylations of the aromatic ring of estradiol and estradiol 17-sulfate. *J. Steroid Biochem. Molec. Biol.* **38** (1991) 737–743.
  23. Ball S. E., Forrester L. M., Wolf C. R. and Back D. J.: Differences in the cytochrome P450 isoenzymes involved in the 2-hydroxylation of oestradiol and 17 $\alpha$ -ethynylestradiol. Relative activities of rat and human liver enzymes. *Biochem. J.* **267** (1990) 221–226.
  24. Theron C. N., Neethling A. C. and Taljaard J. J. F.: Ontogeny and sexual dimorphism of estradiol 2-hydroxylase activity in rat liver microsomes. *J. Steroid Biochem.* **19** (1983) 1185–1190.
  25. Jellinck P. H., Quail J. A. and Crowley C. A.: Normal and recombinant human growth hormone administered by constant infusion feminize catechol estrogen formation by rat liver microsomes. *Endocrinology* **117** (1985) 2274–2278.
  26. Bui Q. D., Weisz J. and Wrighton S. A.: Hepatic catecholesterogen synthesis: differential effect of sex, inducers of cytochromes P450 and of antibody to the glucocorticoid inducible cytochrome P450 on NADPH-dependent estrogen-2-hydroxylase and of organic hydroperoxidase-dependent estrogen-2/4-hydroxylase activity of rat hepatic microsomes. *J. Steroid Biochem. Molec. Biol.* **37** (1990) 285–293.
  27. Arts C. J. M., Wilmer J. W. G. M., De Bie A. Th. H. J. and Van den Berg H.: *In vitro* synthesis of 16 $\alpha$ -hydroxyestrone by female rat liver microsomes: its possible role in the etiology of breast cancer. *J. Steroid Biochem.* **36** (1990) 527–531.
  28. Watanabe H. and Menzies J. A.: Isolation of 16 $\alpha$ -hydroxy-2-methoxyestrone from rat bile. *Steroids* **21** (1973) 123–132.
  29. Honma S. and Nambara T.: Isolation and characterization of biliary metabolites of estrone in the rat. *Chem. Pharm. Bull.* **22** (1974) 687–695.
  30. Levy H., Hood B., Cha C. H. and Carlo J. J.: The monohydroxylation of 17 $\beta$ -estradiol at the 15 $\alpha$ -, 16 $\alpha$ - and 16 $\beta$ -positions by bovine adrenal perfusion. *Steroids* **5** (1965) 677–686.
  31. Adlercreutz H.: Oestrogen metabolism in liver disease. *J. Endocr.* **46** (1970) 129–163.
  32. Hoppen H.-O., Siekmann L. and Breuer H.: Stoffwechsel von oestron in der mikrosomenfraktion der rattenleber. *Hoppe-Seyler's Z. Physiol. Chem.* **355** (1974) 1305–1315.
  33. Chakraborty C., Davis D. L. and Dey S. K.: Estradiol-15 $\alpha$ -hydroxylation: a new avenue of estrogen metabolism in peri-implantation pig blastocysts. *J. Steroid Biochem.* **35** (1990) 209–218.
  34. Maggs J. L., Morgan P., Hussain F., Page P. C. B. and Park B. K.: The metabolism of 2,4-dibromo-17 $\alpha$ -ethynyl[6,7-<sup>3</sup>H]oestradiol in the rat. *Xenobiotica* **20** (1990) 45–54.

35. Maggs J. L., Grabowski P. S. and Park B. K.: Protection of catechol oestrogen from oxidation during enzymic hydrolysis of biliary conjugates. *J. Steroid Biochem.* **19** (1983) 1235–1237.
36. McOmie J. F. W., Watts M. L. and West D. E.: Demethylation of aryl methyl ethers by boron tribromide. *Tetrahedron* **24** (1968) 2289–2292.
37. McLafferty F. W. and Stauffer D. B. In *The Wiley/NBS Registry of Mass Spectral Data*. John Wiley & Sons, New York (1988).
38. Djerassi C., Wilson J. M., Budzikiewicz H. and Chamberlain J. W.: Mass spectrometry in structural and stereochemical problems. XIV. Steroids with one or two aromatic rings. *J. Am. Chem. Soc.* **84** (1962) 4544–4552.
39. Michnowicz J. and Munson B.: Studies in chemical ionization mass spectrometry: steroidal ketones. *Org. Mass Spectrom.* **8** (1974) 49–60.
40. Nambara T., Ishiguro J., Kawarada Y. and Tajima H.: Isolation and characterization of biliary metabolites of estriol in the rat. *Chem. Pharm. Bull.* **22** (1974) 889–893.
41. Williams D. E., Reed R. L., Kedzierski B., Dannan G. A., Guengerich F. P. and Buhler D. R.: Bioactivation and detoxication of the pyrrolizine alkaloid senecionine by cytochrome P450 enzymes in rat liver. *Drug Metab. Dispos.* **17** (1989) 387–392.
42. Adlercreutz H., Tikkanen M. J. and Hunneman D. H.: Mass fragmentographic determination of eleven estrogens in the body fluids of pregnant and nonpregnant subjects. *J. Steroid Biochem.* **5** (1974) 211–217.
43. Cantineau R., Kremers P., De Graeve J., Gieben J. E. and Lambotte R.: 15- and 16 $\alpha$ -hydroxylations of androgens and estrogens in the human fetal liver: a critical step in estetrol biosynthesis. *J. Steroid Biochem.* **22** (1985) 195–201.
44. Knuppen R., Haupt M. and Breuer H.: Formation of 15 $\alpha$ -hydroxyoestradiol-17 $\beta$  and 1 $\alpha$ -hydroxyestrone by human adrenal tissue. *J. Endocr.* **33** (1965) 529–530.
45. Salmon J., Coussediere D., Cousty C. and Raynaud J. P.: Pharmacokinetics and metabolism of moxestrol in animals—rat, dog and monkey. *J. Steroid Biochem.* **19** (1983) 1223–1234.
46. Harada N. and Negishi M.: Substrate specificities of cytochrome P450, C-P450<sub>16 $\alpha$</sub>  and P450<sub>15 $\alpha$</sub> , and contribution to steroid hydroxylase activities in mouse liver microsomes. *Biochem. Pharmac.* **37** (1988) 4778–4780.
47. Jansson I., Mole J. and Schenkman J. B.: Purification and characterization of a new form (RLM2) of liver microsomal cytochrome P450 from untreated rat. *J. Biol. Chem.* **260** (1985) 7084–7093.
48. Swinney D. C., Ryan D. E., Thomas P. E. and Levin W.: Evidence for concerted kinetic oxidation of progesterone by purified rat hepatic cytochrome P450g. *Biochemistry* **27** (1988) 5461–5470.
49. Hoppen H.-O., Bohlig H.-J., Werner M. and Knuppen R.: Metabolism of oestrone and oestradiol in the isolated perfused rat liver. *Hoppe-Seyler's Z. Physiol. Chem.* **357** (1976) 67–74.
50. Black V. H.: Immunodetectable cytochromes P450I, II, and III in guinea pig adrenal—hormone responsiveness and relationship to capacity for xenobiotic metabolism. *Endocrinology* **127** (1990) 1153–1159.