



## Synthesis of 2 $\beta$ -substituted-14-epi-previtamin D<sub>3</sub> and testing of its genomic activity<sup>☆</sup>

Daisuke Sawada<sup>a</sup>, Yuya Tsukuda<sup>a</sup>, Hiroshi Saito<sup>b</sup>, Ken-ichiro Takagi<sup>b</sup>, Eiji Ochiai<sup>b</sup>, Seiichi Ishizuka<sup>b</sup>, Kazuya Takenouchi<sup>b</sup>, Atsushi Kittaka<sup>a,\*</sup>

<sup>a</sup> Faculty of Pharmaceutical Sciences, Teikyo University, 1091-1, Suwarashi, Sagamiko, Sagamihara, Kanagawa 229-0195, Japan

<sup>b</sup> Teijin Institute for Bio-medical Research, Teijin Pharma Ltd., Tokyo 191-8512, Japan

### ARTICLE INFO

#### Article history:

Received 19 October 2009

Received in revised form 10 February 2010

Accepted 25 February 2010

#### Keywords:

Vitamin D<sub>3</sub>

14-epi-Previtamin D<sub>3</sub>

Osteocalcin

Vitamin D receptor

### ABSTRACT

2 $\beta$ -Substituted analogs of 14-epi-previtamin D<sub>3</sub> were synthesized for the first time by the thermal isomerization of the corresponding 14-epi-vitamin D<sub>3</sub> that were available using coupling reaction between the A-ring phosphine oxide derived from a chiral epoxide and CD-ring *cis*-hydrindanone. The VDR binding affinity and transactivation activity of osteocalcin promoter in HOS cells were evaluated, and the new analogs were found to be less active, 0.01–0.18% of VDR binding affinity compared with the natural hormone and EC<sub>50</sub> 1.0–9.1 nM for transactivation activity, than 14-epi-previtamin D<sub>3</sub> with 0.5% (VDR) and EC<sub>50</sub> 0.46 nM, respectively.

© 2010 Elsevier Ltd. All rights reserved.

### 1. Introduction

It is well established that vitamin D<sub>3</sub> is present in thermal equilibrium with previtamin D<sub>3</sub> via [1,7]-sigmatropic rearrangement. In this equilibrium, the vitamin D form (**A**) with the 6-*s-trans* triene structure is more stable and dominant than the 6-*cis* isomer of the previtamin D form (**B**) (Scheme 1). The biologically most active metabolite of vitamin D<sub>3</sub>, 1 $\alpha$ ,25(OH)<sub>2</sub>D<sub>3</sub> (**1**), also contains 5–10% of its previtamin D form, 1 $\alpha$ ,25(OH)<sub>2</sub>preD<sub>3</sub> (**pre-1**) at 37 °C in similar equilibrium. The major isomer, the vitamin D form (**A**), has been the focus of therapeutic evaluation rather than the previtamin D form, because previtamin D is easily transformed to vitamin D through thermal equilibrium and is almost impossible to isolate in pure form [1]. While **1** is a ligand of the nuclear vitamin D receptor (VDR), regulates gene transcription, and exhibits various biological responses as a hormone [2], **pre-1** is thought to be a weak ligand of VDR and a poor activator of the above genomic actions [3]; however, **pre-1** has been studied as a responsible compound for rapid responses [4], such as stimulation of intestinal Ca<sup>2+</sup> transport (transcaltachia), activation of PKC and MAP kinases, and so on, which are called non-genomic actions [5].


Okamura and coworkers reported that the thermal equilibrium ratio between the vitamin D form (**A**) and previtamin D form (**B**) at 80 °C was reversed by epimerizing the CD-ring bridgehead hydrogen of C14 [6]. Briefly, 14-epi-1 $\alpha$ ,25(OH)<sub>2</sub>preD<sub>3</sub> (**14-epi-pre-1**) was major and dominant to 14-epi-1 $\alpha$ ,25(OH)<sub>2</sub>D<sub>3</sub> (**14-epi-1**), and the former was isolated as a stable single isomer at room temperature. Therefore, we focused on the synthesis of **14-epi-pre-1** analogs with A-ring modification to investigate their more detailed biological properties and potential as therapeutic agents of the previtamin D<sub>3</sub> skeleton.

Previously, we found that 2 $\alpha$ -alkyl and 2 $\alpha$ -( $\omega$ -hydroxyalkyl) substitution afforded great enhancements for VDR binding affinity and the subsequent genomic actions [7]. In the preceding paper, we reported the synthesis and biological evaluations of 2 $\alpha$ -substituted **14-epi-pre-1** [8]. Here, we prepared analogs with 2 $\beta$ -substitution (**14-epi-pre-1a-c**), because 2 $\beta$ -substitution is known as an important modification for vitamin D derivatization (Scheme 2) [9].

**14-epi-pre-1** could be prepared from **14-epi-1** by thermal isomerization; therefore, we synthesized **14-epi-1** analogs as temporary first targets. The **14-epi-1** analogs were divided into two fragments, CD-ring and A-ring, which were coupled by the Roche coupling method [10]. The CD-ring fragment **2** [6,8] is the known compound, which was obtained by epimerization at H14 in Grundmann's ketone derivative derived from vitamin D<sub>3</sub> [11,12]. The A-ring fragments, the phosphine oxides **3a-c**, could be synthesized from dimethyl D-tartrate, and we could introduce various alkyl groups at the 2 $\beta$ -position by nucleophilic epoxide ring-opening reactions [13].

<sup>☆</sup> Special issue selected article from the 14th Vitamin D Workshop held at Brugge, Belgium on October 4–8, 2009.



\* Corresponding author. Tel.: +81 42 685 3713; fax: +81 42 685 3713.  
E-mail address: [akittaka@pharm.teikyo-u.ac.jp](mailto:akittaka@pharm.teikyo-u.ac.jp) (A. Kittaka).

Scheme 1. Equilibrium between vitamin D<sub>3</sub> and previtamin D<sub>3</sub>.


## 2. Results and discussion

2 $\beta$ -Substituted A-ring fragments (**3a–c**) were prepared from the known epoxide **4** derived from dimethyl D-tartrate (Scheme 3) [13,14]. Using the nucleophilic epoxide ring-opening reaction of **4**, three substitutions were introduced as follows: (1) methyl cuprate gave a methyl substitution, (2) an allyl group brought

by Grignard reagent was treated with 9-borabicyclo[3,3,1]nonane (9-BBN), and then with H<sub>2</sub>O<sub>2</sub> to afford a hydroxypropyl substitution, and (3) propylene glycol gave a hydroxypropoxy substitution. After their primary hydroxyls were protected as TBDPS ether, **5a–c** were converted into bromide **6a–c** by the known procedure [15]. Methanolysis of both acetyl groups under basic conditions led to epoxide formation, and the resultant hydroxyl group was

Scheme 2. Retrosynthetic analysis of 2 $\beta$ -substituted 14-epi-1 $\alpha$ ,25(OH)<sub>2</sub>preD<sub>3</sub>.

**Scheme 3.** Synthesis of the A-ring fragments. Conditions: (a) for **5a** MeLi, CuI, Et<sub>2</sub>O, 98%; (b) for **5b** (i) allylmagnesium chloride, toluene, (ii) 9-BBN, THF, H<sub>2</sub>O<sub>2</sub>, NaOH, (iii) TBDPSCI, imidazole, DMF, 90% (3 steps); (c) for **5c** (i) propylene glycol, KOtBu, (ii) TBDPSCI, imidazole, DMF, 92% (2 steps); (d) Pd/C, H<sub>2</sub>, MeOH; (e) MeC(OMe)<sub>3</sub>, PPTS, CH<sub>2</sub>Cl<sub>2</sub>; (f) AcBr, CH<sub>2</sub>Cl<sub>2</sub>, 60% for **6a**, 52% for **6b**, 55% for **6c** (3 steps); (g) K<sub>2</sub>CO<sub>3</sub>, MeOH; (h) BzCl, Et<sub>3</sub>N, CH<sub>2</sub>Cl<sub>2</sub>, 80% for **7a**, 95% for **7b**, 83% for **7c** (2 steps); (i) (trimethylsilyl)acetylene, nBuLi, BF<sub>3</sub>•OEt<sub>2</sub>, THF; (j) TBSOTf, iPr<sub>2</sub>EtN, CH<sub>2</sub>Cl<sub>2</sub>; (k) K<sub>2</sub>CO<sub>3</sub>, MeOH, 68% for **8a**, 83% for **8b**, 95% for **8c** (3 steps); (l) SO<sub>3</sub>•Py, Et<sub>3</sub>N, DMSO, 77% for **9a**, 99% for **9b**, 95% for **9c**; (m) vinylmagnesium chloride, THF, 93% ( $\alpha/\beta$  46/47) for **10a**, 95% ( $\alpha/\beta$  37/58) for **10b**, 73% ( $\alpha/\beta$  15/58) for **10c** (2 steps); (n) TBSOTf, iPr<sub>2</sub>EtN, CH<sub>2</sub>Cl<sub>2</sub>, 100% for **11a $\alpha$** , 97% for **11a $\beta$** , 99% for **11b $\beta$** , 99% for **11c $\beta$** ; (o) nBuLi, (CH<sub>2</sub>O)<sub>n</sub>, THF; (p) Red-Al, Et<sub>2</sub>O, then I<sub>2</sub>, THF; (q) Pd(PPh)<sub>4</sub>, Et<sub>3</sub>N, MeCN; (r) (i) NCS, Me<sub>2</sub>S, CH<sub>2</sub>Cl<sub>2</sub>, (ii) nBuLi, PPh<sub>2</sub>, THF, then 30% H<sub>2</sub>O<sub>2</sub>, 49% for **3a $\alpha$** , 57% for **3a $\beta$** , 27% for **3b $\beta$** , 28% for **3c $\beta$**  (4 steps).




**Scheme 4.** Determination of the stereochemistry of the 1-hydroxy group of **10a–c**. Conditions: (a) for **10a $\beta$** , TBAF, THF, 100%; (b) for **10b $\beta$** , PPTS, EtOH, 60%; (c) dimethoxypropane, PPTS, DMF, 70% for **13a $\beta$** , 90% for **13b $\beta$** .

transformed into benzyl ester **7a–c**. The addition of (trimethylsilyl)acetylene to the epoxide using *n*BuLi was straightforward, and the generated secondary alcohol was protected as TBS ether, and removal of the terminal TMS group and the benzoyl group gave alkyne **8a–c**. The primary alcohol was oxidized to aldehyde by DMSO and SO<sub>3</sub>•pyridine complex (**9a–c**), to which the vinyl group was introduced to give a diastereomixture of alcohol **10a–c**. The stereochemistry of the new hydroxy groups is discussed below (Scheme 4), and both isomers of **10a** and the major isomer of **10b** and **10c** were used for further transformation after column chromatography. The hydroxy group of **10a–c** was protected by the TBS group to obtain 2 $\beta$ -substituted enyne **11a–c**. According to the known procedure, enyne **11a–c** were transformed into phosphine oxide in four steps to give **3a–c**, respectively [8,13]. As above, we were able to prepare four A-ring fragments.

As shown in Scheme 4, the minor diastereomer of **10c** (**10c $\alpha$** ) was converted to the phosphine oxide **3c $\alpha$**  by the same strategy as in Scheme 3, and it was identical to the known compound reported by Hatakeyama et al. [13]. Therefore, the stereochemistry of its 1-hydroxy group (steroidal numbering) was found to be  $\alpha$ -configuration, and the major diastereomer of **10c** was determined to have the 1 $\beta$ -hydroxy group (**10c $\beta$** ). For determination of the stereochemistry in **10a** and **10b**, the TBS groups of the major diastereomers (**10a $\beta$**  and **10b $\beta$** ) were removed, and the resultant 1,3-dihydroxy groups of **12a $\beta$**  and **12b $\beta$**  were converted into acetonide **13a $\beta$**  and **13b $\beta$** , respectively. NOE analysis is described in Scheme 4, and the stereochemistry of 1,3-dihydroxy groups was determined as *syn*, that is, **10a $\beta$**  and **10b $\beta$**  had 1 $\beta$ ,3 $\beta$ -dihydroxy groups. As above, we found that all of the major diastereomer of **10a–c** had 1 $\beta$ -hydroxy groups.

Using the CD- and A-ring fragments prepared as above, we examined the coupling reaction under basic conditions with *n*BuLi (Scheme 5) [6,10]. Small excess amounts of the A-ring fragment worked well and we obtained the coupled products **14a–c** in moderate yields. At this point, isomerization to the previtamin D form was seldom observed, probably because TBS groups at the A-ring should have steric hindrance to prevent from reaching the transition state for the [1,7]-sigmatropic hydrogen shift between the vitamin D form and the previtamin D form. Then, all silyl groups in **14a–c** were removed in one step with excess TBAF, and most of the deprotected compounds remained in the vitamin D form (**14-epi-1a–c**), and small amounts of the previtamin D form (**14-epi-pre-1a–c**) were produced under these reaction conditions. However, once they were heated at 80 °C in benzene, isomerization was found to proceed smoothly by <sup>1</sup>H NMR observation. After 2 h, a large proportion of the vitamin D form had been converted into the previtamin D form, and the isomerization seemed to reach thermal equilibrium, at which the ratio of the compounds was about 5/95 (vitamin D/previtamin D) based on <sup>1</sup>H NMR studies. Using HPLC, the mixture of both forms was separated, and we were able to obtain **14-epi-pre-1a–c** as pure forms, which were used for further biological studies.

The VDR binding affinity and the osteocalcin promoter trans-activation activity of the new compounds were evaluated using chick intestinal VDR and HOS cells, respectively. The results are summarized in Table 1 in comparison with the natural hormone **1** and 14-epi-1 $\alpha$ ,25(OH)<sub>2</sub>preD<sub>3</sub> (**14-epi-pre-1**), which was synthesized in a similar manner in our laboratory. The new compounds showed lower activity than the natural hormone **1**, and also than



**Scheme 5.** Coupling reaction and synthesis of 2 $\beta$ -substituted 14-epi-1 $\alpha$ ,25(OH)<sub>2</sub>preD<sub>3</sub>.

**Table 1**

Relative binding affinity for chick intestinal VDR and osteocalcin promoter transactivation activity in HOS cells of 2 $\beta$ -substituted 14-epi-1 $\alpha$ ,25(OH)<sub>2</sub>preD<sub>3</sub>.

Compound	VDR <sup>a</sup>	Osteocalcin transactivation activity (EC <sub>50</sub> , nM)
<b>1</b>	100	0.03
<b>14-epi-pre-1</b>	0.5	0.46
<b>14-epi-pre-1a<math>\alpha</math></b>	0.08	1.34
<b>14-epi-pre-1a<math>\beta</math></b>	0.08	9.12
<b>14-epi-pre-1b<math>\beta</math></b>	0.18	1.01
<b>14-epi-pre-1c<math>\beta</math></b>	0.01	1.24

<sup>a</sup> The potency of **1** is normalized to 100.

**14-epi-pre-1** regardless of the stereochemistry at the 1-hydroxy group.

### 3. Conclusion

We synthesized 2 $\beta$ -substituted analogs of **14-epi-1** for the first time and were able to isolate these new analogs (**14-epi-pre-1a-c**) after thermal isomerization at 80 °C. We evaluated their VDR binding affinity and transactivation activity of osteocalcin promoter in HOS cells. It was found that 2 $\beta$ -modified analogs of 14-epi-1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> were considerably less active than the natural hormone (**1**) and than **14-epi-pre-1**, although 2 $\beta$ -modification of **1** afforded important knowledge to the vitamin D SAR studies.

### Acknowledgments

We are grateful to Ms. Junko Shimode and Ms. Ayako Kawaji (Teikyo University) for the spectroscopic measurements. This work

was supported in part by a Grant-in-Aid from the Ministry of Education, Culture, Sports, Science and Technology, Japan (to D.S.) and by a Grant-in-Aid from the Japan Society for the Promotion of Science (to A.K.).

### References

- [1] M.L. Curtin, W.H. Okamura, 1 $\alpha$ ,25-Dihydroxyvitamin D<sub>3</sub>: synthesis of the 9,14,19,19-pentadeuterio derivatives and a kinetic study of its [1,7]-sigmatropic shift to 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub>, J. Am. Chem. Soc. 113 (1991) 6958–6966.
- [2] D. Feldman, J.W. Pike, F.H. Glorieux (Eds.), Vitamin D, 2nd ed., Elsevier Academic Press, New York, 2005.
- [3] (a) H.F. DeLuca, Evolution of our understanding of vitamin D, Nutr. Rev. 66 (Suppl. 2) (2008) S73–S87; (b) A.J. Brown, E. Slatopolsky, Vitamin D analogs: therapeutic applications and mechanisms for selectivity, Mol. Aspects Med. 29 (2008) 433–452.
- [4] (a) M.T. Mizwicki, D. Keidel, C.M. Bula, J.E. Bishop, L.P. Zanella, J.-M. Wurtz, D. Moras, A.W. Norman, Identification of an alternative ligand-binding pocket in the nuclear vitamin D receptor and its functional importance in 1 $\alpha$ ,25(OH)<sub>2</sub>-vitamin D<sub>3</sub> signaling, PNAS 101 (2004) 12876–12881; (b) I. Nemere, M.C. Dormanen, M.W. Hammond, W.H. Okamura, A.W. Norman, Identification of a specific binding protein for 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> in basal-lateral membranes of chick intestinal epithelium and relationship to transcaltachia, J. Biol. Chem. 269 (1994) 23750–23756.
- [5] A.W. Norman, M.T. Mizwicki, D.P.G. Norman, Steroid-hormone rapid actions, membrane receptors and a conformational ensemble model, Nat. Rev. Drug Discov. 3 (2004) 27–41.
- [6] D.F. Maynard, W.G. Trankle, A.W. Norman, W.H. Okamura, 14-Epi stereoisomers of 25-hydroxy- and 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub>: synthesis, isomerization to previtamins, and biological studies, J. Med. Chem. 37 (1994) 2387–2393.
- [7] For our review articles, see: (a) N. Saito, S. Honzawa, A. Kittaka, Recent results on A-ring modification of 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub>: design and synthesis of VDR-agonists and antagonists with high biological activity, Curr. Top. Med. Chem. 6 (2006) 1273–1288; (b) N. Saito, A. Kittaka, Highly potent vitamin D receptor antagonists; design, synthesis, and biological evaluation, ChemBiochem 7 (2006) 1478–1490; (c) N. Saito, A. Kittaka, Design and synthesis of highly potent vita-

- min D receptor antagonists based on the structural development of vitamin D<sub>3</sub>-26,23-lactone, *J. Synth. Org. Chem. Jpn.* 65 (2007) 947–958.
- [8] D. Sawada, T. Katayama, Y. Tsukuda, N. Saito, M. Takano, H. Saito, K. Takagi, E. Ochiai, S. Ishizuka, K. Takenouchi, A. Kittaka, Synthesis of 2 $\alpha$ -substituted-14-*epi*-previtamin D<sub>3</sub> and its genomic activity, *Bioorg. Med. Chem. Lett.* 19 (2009) 5397–5400.
- [9] (a) N. Tsugawa, K. Nakagawa, M. Kurobe, Y. Ono, N. Kubodera, K. Ozono, T. Okano, *In vitro* biological activities of a series of 2 $\beta$ -substituted analogues of 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub>, *Biol. Pharm. Bull.* 23 (2000) 66–71;  
(b) Y. Ono, H. Watanabe, A. Shiraishi, S. Takeda, Y. Higuchi, K. Sato, N. Tsugawa, T. Okano, T. Kobayashi, N. Kubodera, Synthetic studies of vitamin D analogs. XXIV. Synthesis of active vitamin D<sub>3</sub> analogs substituted at 2 $\beta$ -position and their preventive effects on bone mineral loss in ovariectomized rats, *Chem. Pharm. Bull.* 45 (1997) 1626–1630;  
(c) K. Miyamoto, E. Murayama, K. Ochi, H. Watanabe, N. Kubodera, Synthetic studies of vitamin D analogs. XIV. Synthesis and calcium regulation activity of vitamin D<sub>3</sub> analogues bearing a hydroxyalkoxy group at the 2 $\beta$ -position, *Chem. Pharm. Bull.* 41 (1993) 1111–1113.
- [10] B. Lythgoe, Simonsen lecture. Synthetic approaches to vitamin D and its relatives, *Chem. Soc. Rev.* 9 (1980) 449–475.
- [11] (a) H.H. Inhoffen, G. Quinkert, S. Siegismund, D. Kampe, G.F. Domagk, Vitamin D series. XXI. Hydrindan compounds derived from vitamin D<sub>3</sub>, *Chem. Ber.* 90 (1957) 664–673;  
(b) J. Kiegiel, P.M. Wovkulich, M.R. Uskokovic, Chemical conversion of vitamin D<sub>3</sub> to its 1,25-dihydroxy metabolite, *Tetrahedron Lett.* 32 (1991) 6057–6060.
- [12] R.R. Sicinski, K.L. Perlman, H.F. DeLuca, Synthesis and biological activity of 2-hydroxy and 2-alkoxy analogs of 1 $\alpha$ ,25-dihydroxy-19-norvitamin D<sub>3</sub>, *J. Med. Chem.* 37 (1994) 3730–3738.
- [13] (a) J. Maeyama, H. Hiyamizu, K. Takahashi, J. Ishihara, S. Hatakeyama, N. Kubodera, Two convergent approaches to the synthesis of 1 $\alpha$ ,25-dihydroxy-2 $\beta$ -(3-hydroxypropoxy)vitamin D<sub>3</sub> (ED-71) by the Lythgoe and the Trost coupling reactions, *Heterocycles* 70 (2006) 295–307;  
(b) S. Hatakeyama, H. Irie, T. Shintani, Y. Noguchi, H. Yamada, M. Nishizawa, An efficient route to a key A-ring synthon for 1 $\alpha$ ,25-dihydroxyvitamin D<sub>3</sub> and its analogs, *Tetrahedron* 50 (1994) 13369–13376.
- [14] K.C. Nicolaou, D.P. Papahatjis, D.A. Claremon, R.L. Magolda, R.E. Dolle, Total synthesis of ionophore antibiotic X-14547A, *J. Org. Chem.* 50 (1985) 1440–1456.
- [15] T. Hanazawa, M. Koiwa, G.P.-J. Hareau, F. Sato, Optically active *trans*-4-(*tert*-butyldimethylsilyloxymethyl)-5-(*tert*-butyldimethylsiloxy)-2-cyclohexenone as a useful chiral building block for preparation of substituted cyclohexane rings: synthesis and its highly stereoselective reaction with RCu(CN)Li, *Tetrahedron Lett.* 41 (2000) 2659–2662.