

PROTEIN BIOSYNTHESIS '99

Human Mitochondrial Complex I in Health and Disease

Jan Smeitink and Lambert van den Heuvel

Department of Pediatrics, Nijmegen Center for Mitochondrial Disorders, University Hospital Nijmegen, Nijmegen, The Netherlands

The oxidative phosphorylation (OXPHOS) system of the mitochondrion uses the products of ~60 nuclear genes and 13 mitochondrial genes to generate cellular ATP. These proteins are organized into five large complexes—electron transport chain complexes I–IV and ATP synthetase (complex V) (Hatefi 1985)—of which complex I, or nicotinamide adenine dinucleotide (NADH):ubiquinone oxidoreductase, is the largest. Studies of the bovine heart have identified at least 35 complex I nuclear gene products, together with 7 mitochondrially encoded proteins (Walker 1992; Skehel et al. 1998). In the past year, our knowledge of the human complex I genes and gene products has made substantial progress. Here we review recent insights into these components and their roles in health and disease.

Complex I, which is embedded in the inner mitochondrial membrane (IMM), serves to dehydrogenate NADH and to shuttle electrons to coenzyme Q. This electron transport generates a proton gradient across the IMM, which provides the proton-motive force that is used in ATP synthesis. The total human complex consists of ≥ 42 subunits, 7 encoded by the mitochondrial genome and the remainder by the nuclear genome (Smeitink et al. 1998b). The coordinated interactions between the nucleus and the mitochondrion that build and maintain this vital multiple-subunit protein complex remain poorly understood. As of last year, however, all 35 known human nuclear-encoded cDNAs for complex I proteins had been characterized (Loeffen et al. 1998b; Smeitink et al. 1998b), thanks, in part, to the work of Walker (1992), who characterized the homologous complex from cattle, and also, in part, to the increasing power of genomic methods that are readily available through public databases. On the basis of the genetic information available from prokaryotes and lower eukaryotes, we expect that knowledge about this intriguing complex will expand rapidly in coming years. For the

many individuals with inherited defects in complex I function, these advances offer the prospect of efficient diagnosis, reliable genetic counseling and prenatal diagnostics, better understanding of the cellular consequences of complex I deficiency, and, ultimately, treatment strategies more rational than those currently available.

Nuclear Genes of Human Complex I

The exact protein composition of complex I has not been studied in detail in mitochondria from whole human organs; however, work with bovine heart tissue has thus far identified 35 nuclear-encoded subunits (Walker 1992; Skehel et al. 1998), including 3 flavoproteins (FPs), 7 iron-sulfur proteins (IPs), and 24 hydrophobic proteins (HPs) (Galante and Hatefi 1979); a recently identified 17.2-kD subunit (Skehel et al. 1998) has not yet been placed in any of these three groups. FP and IP proteins from cattle, bacteria, and fungi protrude from the IMM into the mitochondrial matrix (Grigorieff 1998; Guenebaut et al. 1998). This protruding “arm” includes the NDUFV1 protein, which binds and transfers electrons to NADH, as well as many proteins of the IP group, which carry out intra- and intersubunit electron transfer. The final electron acceptor for complex I, ubiquinone, is predicted to bind one or more of these proteins, but no such interaction has yet been identified. The HP fraction, embedded in the lipid bilayer of the IMM, contains most of the nuclear-encoded proteins and all seven mitochondrially-encoded proteins. This fraction seems to mediate proton translocation from the mitochondrial matrix into the intermembrane space (Belogradov and Hafeti 1994).

Table 1 summarizes the present molecular data regarding the human nuclear-encoded complex I subunits. Presently, the complete gene structure is known for 6 of the 35 genes and chromosomal localization is known for 26 of 35 (Emahazion and Brookes 1998; Emahazion et al. 1998). A small form of complex I, consisting of 14 subunits, is found in *Escherichia coli* (Weidner et al. 1993). In this bacterium, all complex I genes are organized as an operon, called “*Nuo*” after the NADH:ubiquinone oxidase. In most eukaryotes, homologues of *NuoB*–*NuoG* and *NuoI* are nuclear genes. Comparative

Received January 25, 1999; accepted for publication April 15, 1999; electronically published May 10, 1999.

Address for correspondence and reprints: Dr. Jan Smeitink, Department of Pediatrics, Nijmegen Center for Mitochondrial Disorders, University Hospital Nijmegen, P.O. Box 9101, 6500 HB Nijmegen, The Netherlands. E-mail: j.smeitink@ckskg.azn.nl

© 1999 by The American Society of Human Genetics. All rights reserved. 0002-9297/99/6406-0004\$02.00

Table 1**Human Complex I Nuclear Genes: Present Knowledge**

Gene Group and Gene	cDNA Sequence	nDNA Sequence	Chromosomal Localization	Leader Sequence	Biochemical Function
Flavoprotein:					
<i>NDUFV1</i> (<i>NuoF</i>)	+	+	11q13	+	NADH binding, electron transfer
<i>NDUFV2</i> (<i>NuoE</i>)	+	+	18p11.2-p11.21	+	Electron transfer
<i>NDUFV3</i>	+	+	21q22.3	+	
Iron-sulfur:					
<i>NDUFA5</i>	+	–	7q31.33		
<i>NDUFS1</i> (<i>NuoG</i>)	+	–	2q33-34	+	Electron transfer
<i>NDUFS2</i> (<i>NuoD</i>)	+	–	...	+	
<i>NDUFS3</i> (<i>NuoC</i>)	+	–	11p11.11	+	
<i>NDUFS4</i>	+	–	5q11.1	+	Phosphorylation
<i>NDUFS5</i>	+	–	1p34.2-p33		
<i>NDUFS6</i>	+	–	5pter-p15.33	+	
Hydrophobic:					
<i>NDUFA1</i>	+	+	Xq24-25		
<i>NDUFA2</i>	+	–	5q31.2		
<i>NDUFA3</i>	+	–	...		
<i>NDUFA4</i>	+	–	...		
<i>NDUFA6</i>	+	–	21q13.1		
<i>NDUFA7</i>	+	–	19p13.2		Ubiquinone binding?
<i>NDUFA8</i>	+	–	9q33.2-34.11		
<i>NDUFA9</i>	+	–	...	+	
<i>NDUFA10</i>	+	–	12p	+	
<i>NDUFAB1</i>	+	–	16p12.3-12.1		Acylcarrier protein motif
<i>NDUFB1</i>	+	–	14q31.3		
<i>NDUFB2</i>	+	–	7q34-35	+	
<i>NDUFB3</i>	+	–	...		
<i>NDUFB4</i>	+	–	...		
<i>NDUFB5</i>	+	–	...	+	
<i>NDUFB6</i>	+	–	9p13.2		
<i>NDUFB7</i>	+	–	19p13.12-13.11		
<i>NDUFB8</i>	+	–	10q23.2-23.33	+	
<i>NDUFB9</i>	+	+	8p24.21		
<i>NDUFB10</i>	+	–	16p13.3		
<i>NDUFS7</i> (<i>NuoB</i>)	+	–	19p13	+	Electron transfer
<i>NDUFS8</i> (<i>NuoI</i>)	+	+	11q13.1-13.3	+	Electron transfer
<i>NDUFC1</i>	+	–	4q28-28.3		
<i>NDUFC2</i>	+	–	...		
Unknown:					
17.2 kD	+		...		

NOTE.—*E. coli* homologues of human complex I genes are indicated in parentheses in the first column. For primary data, see Walker (1992), Emahazion and Brookes (1998), Emahazion et al. (1998), Loeffen et al. (1998b), Skehel et al. (1998), Smeitink et al. (1998b), and Lin et al. (1999). Ellipses (...) = unknown.

studies show that the number of nuclear-encoded complex I subunits increases with the evolutionary complexity of the organism. The seven human nuclear-encoded counterparts of the *E. coli* Nuo proteins, namely, *NDUFV1* and 2, *NDUFS1*–3, and *NDUFS7* and 8, would be predicted to carry out essential aspects of complex I function, and we began our mutational analysis of human patients with the genes for these subunits.

Nuclear-Mitochondrial Interactions

Except for complex II, in which all four known structural genes are nuclear, all components of the OXPHOS system are under the control of both the mitochondrial

and nuclear genomes. This poses a regulatory challenge that is unique in human cell biology. The control of mitochondrial gene expression is relatively well known (Shadel and Clayton 1997; Clayton 1998), but regulation of the nuclear-encoded genes and coordination of the two genomes are not. Fortunately, work in model systems, including the fungus *Neurospora crassa* (Belogrudov and Hatefi 1994), has begun to address the regulation of transcription and translation of these genes, as well as intracellular signaling, complex assembly, and stoichiometry.

The regulatory regions of several genes encoding complex I proteins have been characterized. We recently found a consensus motif in the human *NDUFV1* gene

for binding the transcription factor nuclear respiratory factor II (Schuelke et al. 1998). This binding sequence motif has also been detected in other nuclear genes for mitochondrial proteins, including those for cytochrome *c* oxidase and for the β subunits of ATP-synthetase and of the mitochondrial transcription factor. Moyes et al. (1998) speculate that these genes respond in a modular fashion to physiological change.

The transport of nuclear-encoded complex I gene products to and through the IMM is poorly understood. Transport of the complex I preproteins across the mitochondrial outer membrane is presumed to occur via a dedicated import machinery, called "Tom" for translocase of the outer membrane (Dekker et al. 1998). This multiple-subunit machinery contains receptors and a general import pore. Recently, Hill et al. (1998) showed that Tom40, one of the eight Tom proteins, forms a cation-selective conductance channel that binds to mitochondrial-targeting sequences on the cytoplasmic face of the mitochondrion and transports them across the membrane. At least 15 of the known nuclear-encoded complex I gene products, including all of the subunits shared with the *E. coli* complex, contain classic mitochondrial-targeting leader sequences, but *cis*-acting targeting sequences have not been defined for the remaining proteins. Alternatively, internal targeting sequences may be present in these proteins; Sirrenberg et al. (1998) recently discovered a pathway in yeast for the mitochondrial transport of proteins containing internal signals.

The matrix arm and the membrane portion of complex I form independently and are joined in the course of assembly. In *N. crassa*, the membrane arm itself is formed from two smaller modules, which assemble through the action of two specific chaperones (Kuffner et al. 1998). The functions of the components of the OXPHOS system, particularly human and yeast cytochrome *c* oxidase, also appear to be mediated by nuclear-encoded factors, namely, the SURF1 protein and its yeast homologue Shy1 (Duhig et al. 1998; Tiranti et al. 1998; Zhu et al. 1998). No such chaperones or accessory factors are yet known for human complex I.

Complex I subunit mRNA expression appears to be ubiquitous in human tissues, and the brain, heart, skeletal muscles, and kidneys accumulate these mRNAs at high levels. Among these mRNAs, only the *NDUFB6* message is expressed most strongly in the kidney (Smeitink et al. 1998a), suggesting that mutations in *NDUFB6* may cause distinctive phenotypes. So far, no mutations in *NDUFB6* have been found in people with isolated complex I deficiency.

Isolated Human Complex I Deficiency

Mitochondriocytopathies occur with an estimated incidence of 1 per 10,000 live births, and isolated complex I deficiency is one of those most frequently encountered

(Robinson 1998). The first clinical symptoms of complex I deficiency, presenting either at birth or in early childhood, seem to result from brain dysfunction, sometimes combined with defects in other energy-consuming organs, such as the skeletal muscle and the heart. For this reason, complex I deficiencies are grouped among the mitochondrial encephalomyopathies. Robinson (1998) categorized complex I-deficient patients into three major clinical groups. The most common presentation is Leigh syndrome (Leigh 1951), with cardiomyopathy occurring in ~40% of Leigh syndrome patients (Morris et al. 1996; Rahman et al. 1996). A second category often seen is fatal neonatal lactic acidosis. A relatively uncommon third group comprises patients who present with hepatopathy and tubulopathy with very mild symptoms, such as exercise intolerance, or with cardiomyopathy and cataracts. In our experience, the majority of patients with complex I deficiency die before the age of 5 years of a multisystem disorder, usually Leigh syndrome or a Leigh-like syndrome. In contrast to the patients described by Robinson (1998), none of our patients with cardiomyopathy or cataract showed isolated complex I deficiency (Sengers et al. 1975; Smeitink et al. 1989). Furthermore, none of the 40 patients with whom we have worked who exhibited isolated complex I deficiency in their cultured fibroblasts or myocytes appear to fit into Robinson's third category (J. L. C. M. Loeffen, J. A. M. Smeitink, J. M. Trijbels, A. J. Janssen, R. T. P. Triepels, R. C. A. Sengers, and L. P. van den Heuvel, unpublished data).

Since the elucidation of the complete human mitochondrial genome, mutational analysis of mtDNA has attracted much attention as a way to explore the underlying cause of isolated complex I deficiency. The most frequently observed pathological mtDNA mutations are found in the genes for mitochondrial tRNAs for leucine (*T3271C*, *A3243G*) and lysine (*A8344G*, *T8356C*) and in the protein-encoding subunits ND1 (*T4160A*, *G3460A*), ND4 (*G11778A*), and ND6 (*T14484C*, *G14459A*) (Wallace 1992; Zeviani et al. 1998). Since the underlying genetic defect can be ascribed to mutations in mtDNA in only ~5% of complex I-deficient patients (Liang and Wong 1998), we focused our attention on mutational analysis of nuclear complex I genes and selected a group of 20 patients with isolated complex I deficiency in whom the enzyme deficiency was confirmed in more than one tissue. Since maternal inheritance was not evident in this group, the known mitochondrial DNA mutations associated with complex I deficiency could be excluded. We started our search for mutations in subunits highly conserved during evolution. Mutational analysis of eight nuclear genes encoding proteins of complex I revealed mutations in five of our patients and two additional siblings (Loeffen et al. 1998a; van den Heuvel et al. 1998; Schuelke et al. 1999; Triepels

Table 2**Nuclear Gene Mutations in Patients with Isolated Complex I Deficiency**

AFFECTED GENE (MUTATION[S])	SEX/AGE (MO) ^a	SYMPTOMS AT PRESENTATION	COURSE OF ILLNESS	LACTIC ACID CONCENTRATION			MRI FINDINGS ^c	AGE AT DEATH/ POSTMORTEM FINDINGS ^d
				Blood	Urine	CSF ^b		
<i>NDUFS4</i> (5-bp duplication)	M/8	Vomiting, failure to thrive, hypotonia	Progressive	Normal	Normal	Normal	Atrophy, basal ganglia abnormalities	16 mo
<i>NDUFS8</i> (P79L, R102H)	M/<1	Feeding difficulties, hypotonia, episodic apnea and cyanosis	Progressive, moderate hypertrophic cardiopathy	Increased	Increased	Increased	Atrophy, symmetrical hypodensities	11 weeks (LS confirmed)
<i>NDUFS7</i> (V122M)	M/26	Feeding problems, dysarthria, ataxia	Progressive	Normal	Normal	Sl. incr.	Symmetrical hypodensities	3.5 years (LS confirmed)
<i>NDUFS7</i> (V122M)	M/11	Vomiting	Progressive after infection (3.4 years)	Normal	Normal	Normal	Symmetrical hypodensities	5 years (LS confirmed)
<i>NDUFV1</i> (R59X, T423M)	M/5	Vomiting, strabismus, hypotonia	Progressive, myoclonic epilepsy	Increased	Normal	Increased	Atrophy	14 mo
<i>NDUFV1</i> (R59X, T423M)	M/7	Vomiting, strabismus, hypotonia	Progressive, myoclonic epilepsy	Increased	Normal	Increased	Atrophy	17 mo
<i>NDUFV1</i> (A341V)	F/6	Infantile myoclonic epilepsy	Progressive	Normal	Normal	Increased	Atrophy, progressive macro-cystic leucodystrophy	Alive at age 10 years

^a Age at presentation.

^b CSF = cerebrospinal fluid; Sl. incr. = slightly increased.

^c MRI = magnetic resonance imaging.

^d LS = Leigh syndrome.

et al., in press). Interestingly, none of our subjects carry mutations in any of the 18 complex I subunit genes we have studied that are not conserved with *E. coli*. We are continuing to search for mutations in the remaining complex I structural genes in this group of patients. On the basis of precedent from *N. crassa* (Kuffner et al. 1998), it is possible that some of these individuals will carry mutations in genes that are required for the folding or assembly of complex I proteins.

The currently known mutations and clinical features of our complex I-deficient patients are summarized in table 2. Six patients had a progressive and ultimately fatal multisystemic disorder in which cerebral dysfunction was the most prominent clinical sign; one is alive at 10 years of age but is severely handicapped. Increased lactic acid concentration, considered to be one of the hallmarks of a mitochondriocytopathy, was certainly not present in all body fluids routinely investigated. Strikingly, one patient showed no lactic acid elevation, not even after a series of provocative tests. We assume, on the basis of past experience with mtDNA studies, that the inheritance pattern of most children with enzymatic complex I deficiency is autosomal recessive.

Outstanding Questions

The cloning of the nuclear genes and cDNAs that encode the human complex I components should make prenatal and postnatal diagnosis of complex I deficiency possible, although the large number of nuclear and mitochondrial genes involved is still somewhat daunting. Chip technology will certainly increase our diagnostic abilities in the near future. At present, extensive mutational detection studies of nuclear genes are performed only after the common mtDNA mutations have been excluded.

Before the emerging molecular data can be put to use to benefit families with complex I deficiency, we badly need answers to several fundamental questions. First, until detailed protein studies of the complex have been performed with human cells, the total number of the human complex I nuclear genes will remain uncertain. Quite recently, Walker's group, which had just reported the identity of the 42d complex I subunit (Skehel et al. 1998), announced the presence of a 43d subunit in bovine heart. Furthermore, the stoichiometry of this multiprotein subunit complex has not been explored in humans, nor have any studies addressed possible differences between cell types in the composition of the complex, which could account for variable clinical presentation of this mitochondrial encephalomyopathy. In addition, very limited information is available about the expression, intracellular transport, and assembly of the subunits of the complex, all steps that might be impaired in individuals deficient in OXPHOS. Insights into these medically relevant biosynthesis issues may come from

detailed studies of bacteria and fungi, as was seen recently with the identification of the *SURF1* gene in cytochrome c oxidase-deficient patients with Leigh syndrome (Tiranti et al. 1998; Zhu et al. 1998). Finally, the metabolic and general physiological effects of aberrant complex I function will need to be explored if rational therapeutic strategies are to be developed.

Acknowledgments

The authors are very grateful to Prof. Rob Sengers and Prof. Frans Trijbels, who founded the Nijmegen Center for Mitochondrial Disorders >20 years ago and who supported us throughout this large-scale project. We cordially thank all the Ph.D. students (Jan Loeffen, Ralf Triepels, Markus Schuelke, Sandy Budde, and Marieke Coenen) and technicians (Roel Smeets, Carin Buskens, Antoon Janssen, and Frans van den Brand) for their enthusiasm and the work they performed in achieving this first part of our research program. The Prinses Beatrix Fonds, the Stichting voor Kinderen die willen maar niet kunnen, and the Fonds Beoefening Wetenschap of the Nijmegen Children's Hospital are gratefully acknowledged for their financial support.

References

- Belogradov G, Hatefi Y (1994) Catalytic sector of complex I (NADH:ubiquinone oxidoreductase): subunit stoichiometry and substrate-induced conformation changes. *Biochemistry* 33:4571-4576
- Clayton DA (1998) Nuclear-mitochondrial intergenomic communication. *Biofactors* 7:203-205
- Dekker PJ, Ryan MT, Brix J, Muller H, Honlinger A, Pfanner N (1998) Preprotein translocase of the outer mitochondrial membrane: molecular dissection and assembly of the general import pore complex. *Mol Cell Biol* 18:6515-6524
- Duhig T, Ruhrberg C, Mor O, Fried M (1998) The human surfeit locus [in process citation]. *Genomics* 52:72-78
- Emahazion T, Beskow A, Gyllenstein U, Brookes AJ (1998) Intron based radiation hybrid mapping of 15 complex I genes of the human electron transport chain. *Cytogenet Cell Genet* 82:115-119
- Emahazion T, Brookes AJ (1998) Mapping of the NDUFA2, NDUFA6, NDUFA7, NDUFAB8, and NDUFS8 electron transport chain genes by intron based radiation hybrid mapping. *Cytogenet Cell Genet* 82:112-114
- Galante YM, Hatefi Y (1979) Purification and molecular and enzymic properties of mitochondrial NADH dehydrogenase. *Arch Biochem Biophys* 192:559-568
- Grigorieff N (1998) Three-dimensional structure of bovine NADH:ubiquinone oxidoreductase (complex I) at 2.2 Å in ice. *J Mol Biol* 277:1033-1046
- Guenebaut V, Schlitt A, Weiss H, Leonard K, Friedrich T (1998) Consistent structure between bacterial and mitochondrial NADH:ubiquinone oxidoreductase (complex I). *J Mol Biol* 276:105-112
- Hatefi Y (1985) The mitochondrial electron transport and oxidative phosphorylation system. *Annu Rev Biochem* 54:1015-1069
- Hill K, Model K, Ryan MT, Dietmeier K, Martin F, Wagner

- R, Pfanner N (1998) Tom40 forms the hydrophilic channel of the mitochondrial import pore for preproteins. *Nature* 395:516–521
- Kuffner R, Rohr A, Schmiede A, Krull C, Schulte U (1998) Involvement of two novel chaperones in the assembly of mitochondrial NADH:ubiquinone oxidoreductase (complex I). *J Mol Biol* 283:409–417
- Leigh D (1951) Subacute necrotizing encephalomyelopathy in an infant. *J Neurol Neurosurg Psychiatr* 14:216–221
- Liang MH, Wong LJ (1998) Yield of mtDNA mutation analysis in 2,000 patients. *Am J Med Genet* 77:395–400
- Lin X, Wells DE, Kimberling WJ, Kumar S (1999) Human NDUFB9 gene: genomic organization and a possible candidate gene associated with deafness disorder mapped to chromosome 8q13. *Hum Hered* 49:75–80
- Loeffen J, Smeitink J, Triepels R, Smeets R, Schuelke M, Sengers R, Trijbels F, et al (1998a) The first nuclear-encoded complex I mutation in a patient with Leigh syndrome. *Am J Hum Genet* 63:1598–1608
- Loeffen JL, Triepels RH, van den Heuvel LP, Schuelke M, Buskens CA, Smeets RJ, Trijbels JM, et al (1998b) cDNA of eight nuclear encoded subunits of NADH:ubiquinone oxidoreductase: human complex I cDNA characterization completed. *Biochem Biophys Res Commun* 253:415–422
- Morris AA, Leonard JV, Brown GK, Bidouki SK, Bindoff LA, Woodward CE, Harding AE, et al (1996) Deficiency of respiratory chain complex I is a common cause of Leigh disease. *Ann Neurol* 40:25–30
- Moyes CD, Battersby BJ, Learly SC (1998) Regulation of muscle mitochondrial design. *J Exp Biol* 201:299–307
- Rahman S, Blok RB, Dahl HH, Danks DM, Kirby DM, Chow CW, Christodoulou J, et al (1996) Leigh syndrome: clinical features and biochemical and DNA abnormalities. *Ann Neurol* 39:343–351
- Robinson BH (1998) Human complex I deficiency: clinical spectrum and involvement of oxygen free radicals in the pathogenicity of the defect. *Biochim Biophys Acta* 1364:271–286
- Schuelke M, Loeffen J, Mariman E, Smeitink J, van den Heuvel L (1998) Cloning of the human mitochondrial 51 kDa subunit (NDUFV1) reveals a 100% antisense homology of its 3'UTR with the 5'UTR of the gamma- interferon inducible protein (IP-30) precursor: is this a link between mitochondrial myopathy and inflammation? *Biochem Biophys Res Commun* 245:599–606
- Schuelke M, Smeitink J, Mariman E, Plecko B, Trijbels F, Stöckler-Ipsiroglu S, van den Heuvel L, et al (1999) Mutant NDUFV1 subunit of mitochondrial complex I causes leukodystrophy and myoclonic epilepsy. *Nat Genet* 21:260–261
- Sengers RC, Trijbels JM, Willems JL, Daniels O, Stadhouders AM (1975) Congenital cataract and mitochondrial myopathy of skeletal and heart muscle associated with lactic acidosis after exercise. *J Pediatr* 86:873–880
- Shadel GS, Clayton DA (1997) Mitochondrial DNA maintenance in vertebrates. *Annu Rev Biochem* 66:409–435
- Sirrenberg C, Endres M, Folsch H, Stuart R, Neupert W, Brunner M (1998) Carrier protein import into mitochondria mediated by the intermembrane proteins Tim10/Mrs11 and Tim12/Mrs5. *Nature* 391:912–915
- Skehel JM, Fearnley IM, Walker JE (1998) NADH:ubiquinone oxidoreductase from bovine heart mitochondria: sequence of a novel 17.2-kDa subunit. *FEBS Lett* 438:301–305
- Smeitink J, Loeffen J, Smeets R, Triepels R, Ruitenbeek W, Trijbels F, van den Heuvel L (1998a) Molecular characterization and mutational analysis in the human B17 subunit of the mitochondrial respiratory chain complex I. *Hum Genet* 103:245–250
- Smeitink JAM, Loeffen JLCM, Triepels RH, Smeets RJP, Trijbels JMF, van den Heuvel LP (1998b) Nuclear genes of human complex I of the mitochondrial electron transport chain: state of the art. *Hum Mol Genet* 7:1573–1579
- Smeitink JA, Sengers RC, Trijbels JM, Ruitenbeek W, Daniels O, Stadhouders AM, Kock-Jansen MJ (1989) Fatal neonatal cardiomyopathy associated with cataract and mitochondrial myopathy. *Eur J Pediatr* 148:656–659
- Tiranti V, Hoertnagel K, Carrozzo R, Galimberti C, Munaro M, Granatiero M, Zelante L, et al (1998) Mutations of SURF-1 in Leigh disease associated with cytochrome c oxidase deficiency. *Am J Hum Genet* 63:1609–1621
- Triepels RH, van den Heuvel LP, Loeffen JLCM, Buskens CAE, Smeets RJP, Rubio Gozalbo ME, Budde SMS, et al. Leigh syndrome associated with a mutation in the NDUF57 (PSST) nuclear encoded subunit of complex I. *Ann Neurol* (in press)
- van den Heuvel L, Ruitenbeek W, Smeets R, Gelman-Kohan Z, Elpeleg O, Loeffen J, Trijbels F, et al. (1998) Demonstration of a new pathogenic mutation in human complex I deficiency: a 5-bp duplication in the nuclear gene encoding the 18-kD (AQDQ) subunit. *Am J Hum Genet* 62:262–268
- Walker JE (1992) The NADH:ubiquinone oxidoreductase (complex I) of respiratory chains. *Q Rev Biophys* 25:253–324
- Wallace DC (1992) Diseases of the mitochondrial DNA. *Annu Rev Biochem* 61:1175–1212
- Weidner U, Geier S, Ptock A, Friedrich T, Leif H, Weiss H (1993) The gene locus of the proton-translocating NADH:ubiquinone oxidoreductase in *Escherichia coli*: organization of the 14 genes and relationship between the derived proteins and subunits of mitochondrial complex I. *J Mol Biol* 233:109–122
- Zeviani M, Tiranti V, Piantadosi C (1998) Mitochondrial disorders. *Medicine (Baltimore)* 77:59–72
- Zhu Z, Yao J, Johns T, Fu K, De Bie I, Macmillan C, Cuthbert AP, et al (1998) SURF1, encoding a factor involved in the biogenesis of cytochrome c oxidase, is mutated in Leigh syndrome. *Nat Genet* 20:337–343