

of the discrepancy between results of our study and those of the report by Ahmad et al. (1998) would be the existence of modifier genes. However, this is very unlikely, given the fact that none of the reported molecular studies of families with APL has observed a case of incomplete penetrance.

Interestingly, although the individual who is homozygous for Gln620 is not affected with papular atrichia, he displays AGA. We therefore examined the frequency of Gln620 in a sample of 103 males with severe AGA and in 58 males (age > 60 years) without AGA. Allele frequencies were similar in both groups (2.91% and 3.45%, respectively; $P = .79$), which suggests that Arg620Gln does not play a role in the development of AGA.

Acknowledgments

We thank all individuals who consented to the study and gave blood samples. This study was supported by a grant from the Deutsche Forschungsgemeinschaft (Forschergruppe "Keratinocyten-Proliferation und differenzierte Leistung in der Epidermis" and Forschergruppe "Genetische Epidemiologie und Medizinische Genetik komplexer Erkrankungen"). R.C.B. is a research fellow from the Deutsche Forschungsgemeinschaft.

AXEL M. HILLMER,¹ ROLAND KRUSE,²
REGINA C. BETZ,¹ JOHANNES SCHUMACHER,¹
UWE HEYN,¹ PETER PROPPING,¹
MARKUS M. NÖTHEN,³ AND SVEN CICHON¹

¹Institute of Human Genetics, University of Bonn, Bonn; ²Department of Dermatology, University of Düsseldorf, Düsseldorf, Germany; ³Department of Medical Genetics, University of Antwerp, Antwerp

Electronic-Database Information

Accession numbers and the URL for data in this article are as follows:

Online Mendelian Inheritance in Man (OMIM), <http://www.ncbi.nlm.nih.gov/Omim/> (for ALUNC [MIM 203655] and APL [MIM 209500])

References

- Ahmad W, Irvine AD, Lam H, Buckley C, Bingham EA, Pan-teleyev AA, Ahmad M, McGrath JA, Christiano AM (1998) A missense mutation in the zinc-finger domain of the human hairless gene underlies congenital atrichia in a family of Irish Travellers. *Am J Hum Genet* 63:984–991
- Kruse R, Cichon S, Anker M, Hillmer AM, Barros-Nunez P, Cantu JM, Leal E, Weinlich G, Schmuth M, Fritsch P, Ruzicka T, Propping P, Nöthen MM (1999) Novel *hairless* mu-

tations in two kindreds with autosomal recessive papular atrichia. *J Invest Dermatol* 113:954–959

Address for correspondence and reprints: Dr. Sven Cichon, Institute of Human Genetics, University of Bonn, Wilhelmstrasse 31, 53111 Bonn, Germany. E-mail: sven.cichon@ukb.uni-bonn.de

© 2001 by The American Society of Human Genetics. All rights reserved. 0002-9297/2001/6901-0026\$02.00

Am. J. Hum. Genet. 69:237–241, 2001

The Presence of Mitochondrial Haplogroup X in Altaians from South Siberia

To the Editor:

For American Indians, extensive RFLP and HVSI sequence analysis has unambiguously identified four major founding mtDNA haplogroups (A, B, C, D), which account together for ~97% of modern American Indian mtDNAs (Wallace 1995). Examination of the distribution of the four founding lineage haplotypes (A, B, C, and D) in American Indian populations (both contemporary and ancient) shows that all four lineages were present in the New World prior to European contact (Wallace 1995; Lalueza et al. 1997; Stone and Stoneking 1998), thus indicating that all American Indian mtDNAs are apparently descended from these four founding lineages. mtDNAs apparently not from haplogroups A–D may result from recent admixture with non-American Indians or may represent additional American Indian founding mtDNA lineages.

A striking example of the presence in American Indians of genotypes not from haplogroups A–D is haplogroup X. This haplogroup represents a minor founding lineage that is restricted in distribution to northern Amerindian groups, including the Ojibwa, the Nu-Chah-Nulth, the Sioux, and the Yakima, as well as the Na Dene-speaking Navajo (Brown et al. 1998). Unlike haplogroups A–D, haplogroup X is also found at low frequencies of ~4% in western Eurasian populations. Despite a shared consensus RFLP haplotype, substantial genetic differences exist between the American Indian and European haplogroup X mtDNAs. Phylogenetic analysis and coalescence estimates for American Indian and European haplogroup X mtDNAs exclude the possibility that the occurrence of haplogroup X in American Indians is due to recent European admixture. They also clearly indicate that the two branches/subgroups are distantly related to each other and that considerable genetic substructure exists within both groups (Brown et al. 1998).

Haplogroup X is remarkable in that it has not been found in Asians, including Siberians, suggesting that it

Table 1**RFLP and HVSI, HVSII Sequence Variation of Altaian Haplogroup X mtDNAs**

SAMPLE	ORIGIN	RFLP	HVSI, HVSII
		HAPLOTYPE ^a	SEQUENCE ^b
		111	111111
		1046	666666
		7345	11112201122333
		1961	88892775916001
		5457	23933833553995
		ccse	a aba
CRS		+++	AAT-CCAATAA—
ALT16	Northern Altaians	-.++	CCCCTTGGCGGCC
ALT43	Northern Altaians	-.++	CCC.TTGGC.G..C
ALT81	Southern Altaians	-.++	CCCCTTGGC.GCCC
ALT161	Southern Altaians	-.++	CCC.TTGGC.GCCC
ALT171	Southern Altaians	-.++	CCC.TTGGC.GCCC
ALT188	Southern Altaians	-.++	CCC.TTGGC.GCCC
ALT208	Southern Altaians	-.++	CCC.TTGGC.GCCC

^a Restriction-endonuclease sites are indicated as follows: c = *DdeI*, e = *HaeIII*, s = *AccI*. A dot (.) denotes identity with the Cambridge reference sequence (Anderson et al. 1981); a plus sign (+), which denotes a site gain, or a minus sign (-), which denotes a site loss, indicates deviation from the CRS.

^b HVSI (nucleotide positions 15991–16400) and HVSII (nucleotide positions 20–420) were sequenced. A dot (.) indicates identity with the CRS, and a dash (—) indicates nucleotide insertion.

may have come to the Americas via a Eurasian migration. The virtual absence of haplogroup X in eastern and northern Asia raises the possibility that some American Indian founders were of European ancestry. In that case, as it has been proposed, haplogroup X was brought to America by the eastward migration of an ancestral white population, of which no trace has so far been found in the mtDNA gene pool of modern Siberian/eastern Asian populations (Brown et al. 1998).

However, it should be stressed that mtDNA-variability studies of the populations living in this major geographic area were performed on a limited number of populations. Some regions remain poorly sampled, and more extensive sampling is required. Moreover, some key markers, including those defining the X-haplogroup sequences, have not been typed for many different populations. These limitations do not allow correct definition of the phylogenetic status of mtDNA lineages.

To extend the survey of Asian mtDNAs for the presence of haplogroup X, we screened the mtDNAs of a total of 790 individuals for the RFLP markers (-1715 *DdeI*, -10394 *DdeI*, +14465 *AccI*, and +16517 *HaeIII*) that define this lineage. These individuals comprised 10 aboriginal Siberian populations: Buryats ($n = 105$), Tuvinians ($n = 111$), Koryaks ($n = 35$), Evens ($n = 65$), Yakuts ($n = 62$), Khakassians ($n = 54$), Shors ($n = 42$), Sojots ($n = 34$), Altaians ($n = 202$), and Evenks ($n = 80$). All individuals belonged to the indigenous population of the regions studied, were unrelated, and

stated that their maternal grandmother had been born in the area considered for this study.

Haplogroup X mtDNAs were detected only in Altaians, at a frequency of 3.5%. The haplogroup X status of these haplotypes was confirmed through HVSI and HVSII mtDNA sequencing (table 1). All Altaian X mtDNAs harbored the consensus haplogroup X motif: -1715 *DdeI*, +14465 *AccI*, +16517 *HaeIII*, 16189C, 16223T, 16278T, 73G, 153G, 195C, 263G, relative to the Cambridge reference sequence (Anderson et al. 1981) and differed from each other by length-polymorphism mutations at nucleotide positions 16193, 309, and 315. One of these X mtDNAs (ALT16) also harbored a 215G variant (table 1) that has not been observed in either American Indian or European X haplotypes. It should also be noted that none of the Altaian X mtDNAs harbored the 225A variant, which is a marker for a major part of haplogroup X (Brown et al. 1998).

Analysis of published data on European HVSI and HVSII mtDNA sequences (Piercy et al. 1993; Calafell et al. 1996; Torroni et al. 1996; Brown et al. 1998; Lutz et al. 1998; Parson et al. 1998; Rousselet and Mangin 1998; Helgason et al. 2000) demonstrates that the overwhelming majority of X haplotypes (23 of 25 X sequences) harbor the 225A variant. In contrast, the X haplotypes without 225A have been observed mostly in American Indians (11 of 14 Ojibwa; see table 1 in Brown et al. 1998). Nevertheless, the X mtDNAs that we detected in the Altaian sample do not bear the 16213A and 200G variants that are characteristic of most American Indian haplogroup X mtDNAs (Brown et al. 1998).

Figure 1 illustrates the reduced median network, constructed by means of the median algorithm of Bandelt et al. (1995), encompassing the HVSI and HVSII variation observed in the American Indian, European, and Altaian haplogroup X mtDNAs. The network suggests that European and American Indian haplogroup X mtDNAs are separated into two major branches, whereas the majority of Altaian X mtDNAs appear to be very similar to the root of haplogroup X phylogeny, differing from it by one step (loss of 225A). The network further suggests that the Altaian X haplotypes occupy the intermediate position between European and American Indian haplogroup X mtDNA lineages (fig. 1).

The Altaians, the native people of Altai Republic (south Siberia) number up to 60,000 persons. "Altaians" is the common denomination for seven formerly distinct Turkic-speaking groups: the Altai-Kizhi, Teleuts, and Telenghits, who are southern Altaians, and the Chelkans, Kumandins, Tubalars, and Maimalars, who are northern Altaians. The differences between southern and northern Altaians are well established, on the basis of anthropological, linguistic, and classical genetic-marker studies (Potapov 1969; Alexeev and Gohman 1984; Luzina 1987). Anthropologically, southern Altaians are typical




Figure 1 Reduced median network of haplogroup X mtDNAs, as defined by HVSI and HVSII variation. RFLP status of the sequences is known for American Indians (NA; Brown et al. 1998); for Altaians (ALT) from the present study; and for some Europeans (CE; Brown et al. 1998). The other European sequences—GER (Lutz et al. 1998), ENG (Piercy et al. 1993), TUS (Torrioni et al. 1996), FRA (Rousselet and Mangin 1998), BUL and TUR (Calafell et al. 1996), AUS (Parson et al. 1998), and ICE (Helgason et al. 2000)—were assumed, on the basis of their HVSI and HVSII sequences, to belong to haplogroup X. Sample origin is indicated by the scheme (see key). mtDNA types are represented by circles, with areas proportional to number of individuals. Lines are labeled by HVSI and HVSII mutations. Nucleotide variants correspond to transitions, whereas transversions are further specified. The insertions/deletions, as well as variations of C's in the unstable regions between nucleotide positions 16182–16193 and 309–315 were not included. The node marked with an asterisk (*) matches the haplogroup X basic motif: –1715c, +14465s, 16189C, 16223T, 16278T, +16517e/16519C, 153A, 195C, and 225A. The restriction site changes indicated by their enzyme letter code listed in table 1. Underlining indicates nucleotide positions that have mutated more than once, and reticulations indicate ambiguity in the topology. Parallel lines in a reticulation represent the same mutation.

central Asian Mongoloids (like Mongolians, Yakuts, and Buryats), whereas northern Altaians exhibit some Caucasoid anthropological features, similar to those of Ugric and Samoyedic groups.

The Altai region was populated during the Lower Paleolithic, and there is ample evidence of settlement during the Middle Paleolithic. It was proposed by anthropologists that, at least from the Neolithic, the territories of Altai and Sayan region were populated by mixed tribes with Caucasoid and Mongoloid anthropological features, but later they were replaced by Mongoloid populations of central Asian origin (Alexeev and Gohman 1984). The analysis of the tribal structure of Southern Altaians has shown that the present-day Altaians have retained their native language and ethnic identity. They have begun to mix with other ethnic groups (mostly Russians and Kazakhs) only recently, so the interethnic admixture is estimated to be <5% (Luzina 1987; Osipova et al. 1997). The haplogroup X mtDNAs have not been found in populations of central Asia, including Kazakhs, Uighurs, and Kirghiz (Comas et al. 1998). Since the frequency of haplogroup X in Russians is extremely low (3 of 336; Orekhov et al. 1999; Malyarchuk and Derenko 2000; authors' unpublished data), the recent European admixture cannot explain the presence of haplogroup X in the Altaians. Hence, the results of the present study allow us to suggest that haplogroup X was the part of the ancestral gene pool for Altaian populations, being found both in northern and southern Altaians.

Recently, the mtDNA studies have shown that both northern and southern Altaians exhibit all four Asian and American Indian-specific haplogroups (A–D) with frequencies of 57.2% (Sukernik et al. 1996) and 46.8% (Derenko et al. 2000a), respectively, exceeding those reported previously for Mongolians, Chinese, and Tibetans. Therefore, they may represent the populations which are most closely related to New World indigenous groups. Since the detection of all four haplogroups (A–D) in an Asian population is thought to be a first criterion in the identification of a possible New World founder, the candidate source population for American Indian mtDNA haplotypes therefore may include the populations originating in the regions to the southwest and southeast of Lake Baikal, including the Altai Mountain region (Derenko et al. 2000b). The presence of X mtDNAs in Altaians is generally consonant with the latter conclusion.

Because the location and identification of the population that was the source of the founding lineages for the New World is a question of considerable interest, several studies on Y-chromosomal DNA polymorphism were performed recently to investigate Pleistocene male migrations to the American continent (Underhill et al. 1996; Lell et al. 1997; Karafet et al. 1999; Santos et al.

1999). It has been shown that the major Y haplotype present in most American Indians could be traced back to recent ancestors they have in common with Siberians: namely, the Kets and Altaians, from the Yenisey River Basin and the Altai Mountains, respectively (Santos et al. 1999). Similarly, based on a comprehensive analysis of worldwide Y-chromosome variation, it has been proposed that populations occupying the general area including Lake Baikal (eastward to the Trans-Baikal and southward into Northern Mongolia), the Lena River headwaters, the Angara and Yenisey River basins, the Altai Mountain foothills, and the region south of the Sayan Mountains (including Tuva and western Mongolia) was the source for dispersals of New World Y-chromosome founders (Karafet et al. 1999). It is obvious that we have now the genetic evidence that will allow closer determination of which Siberian population was the source of the population expansion leading to modern American Indians and will allow relation of the studies of migrations from Siberia to the Americas that are based on paternally inherited genetic systems with those based on maternally inherited ones.

Acknowledgments

The authors would like to thank the following individuals for providing samples for this study: Drs. E. Lotosh, F. Luzina, I. Dambueva, C. Dorzhu, V. Kakpakov, and O. Karamchakova. This work was supported by the Russian Fund for Basic Research (grant 99-06-80430), the State Program Frontiers in Genetics (grant 99-04-30), and by a grant from the Ludwik Rydygier Medical University in Bydgoszcz (BW 90/01).

MIROSLAVA V. DERENKO,¹ TOMASZ GRZYBOWSKI,²
BORIS A. MALYARCHUK,¹ JAKUB CZARNY,²
DANUTA MIŚCICKA-ŚLIWKA,² ILIA A. ZAKHAROV³

¹Genetics Laboratory, Institute of Biological Problems of the North, Magadan, Russia; ²The Ludwik Rydygier Medical University in Bydgoszcz, Forensic Medicine Institute, Bydgoszcz, Poland; and ³Animal Comparative Genetics Laboratory, Vavilov Institute of General Genetics, Moscow

References

- Alexeev VP, Gohman II (1984) Anthropology of Asiatic part of USSR. Nauka, Moscow
- Anderson S, Bankier AT, Barrell BG, de Bruijn MHL, Coulson AR, Drouin J, Eperon IC, Nierlich DP, Roe BA, Sanger F, Schreier PH, Smith AJM, Staden R, Young IG (1981) Sequence and organization of the human mitochondrial genome. *Nature* 290:457–465
- Bandelt HJ, Forster P, Sykes BC, Richards MB (1995) Mitochondrial portraits of human populations using median networks. *Genetics* 141:743–753
- Brown MD, Hosseini SH, Torroni A, Bandelt HJ, Allen JC, Schurr TG, Scozzari R, Cruciani F, Wallace DC (1998)

- mtDNA haplogroup X: an ancient link between Europe/Western Asia and North America? *Am J Hum Genet* 63:1852–1861
- Calafell F, Underhill P, Tolun A, Angelicheva D, Kalaydjieva L (1996) From Asia to Europe: mitochondrial DNA sequence variability in Bulgarians and Turks. *Ann Hum Genet* 60:35–49
- Comas D, Calafell F, Mateu E, Perez-Lezaun A, Bosch E, Martinez-Arias R, Clarimon J, Facchini F, Fiori G, Luiselli D, Pettener D, Bertranpetit J (1998) Trading genes along the Silk Road: mtDNA sequences and the origin of Central Asian populations. *Am J Hum Genet* 63:1824–1838
- Derenko MV, Denisova GA, Malyarchuk BA, Dambueva IK, Dorzhu CM, Stolpovski YM, Lotosh EA, Luzina FA, Zakharov IA (2000a) Mitochondrial DNA variability in Turkic-speaking populations of the Altai and Sayan region from South Siberia. *Am J Hum Genet Suppl* 67:A1161
- Derenko MV, Malyarchuk BA, Dambueva IK, Shaikhaev GO, Dorzhu CM, Nimaev DD, Zakharov IA (2000b) Mitochondrial DNA variation in two South Siberian aboriginal populations: implications for the genetic history of North Asia. *Hum Biol* 72:945–973
- Helgason A, Sigurdardóttir S, Gulcher JR, Ward R, Stefánsson K (2000) mtDNA and the origin of the Icelanders: deciphering signals of recent population history. *Am J Hum Genet* 66:999–1016
- Karafet TM, Zegura SL, Posukh O, Osipova L, Bergen A, Long J, Goldman D, Klitz W, Harihara S, de Kniff P, Wiebe V, Griffiths RC, Templeton AR, Hammer MF (1999) Ancestral Asian source(s) of New World Y-chromosome founder haplotypes. *Am J Hum Genet* 64:817–831
- Lalueza C, Perez-Perez A, Prats E, Cornudella L, Turbon D (1997) Lack of founding Amerindian mitochondrial DNA lineages in extinct Aborigines from Tierra del Fuego-Patagonia. *Hum Mol Genet* 6:41–46
- Lell JT, Brown MD, Schurr TG, Sukernik RI, Starikovskaya YB, Torroni A, Moore LG, Troup GM, Wallace DC (1997) Y chromosome polymorphisms in Native American and Siberian populations: identification of Native American Y chromosome haplotypes. *Hum Genet* 100:536–543
- Lutz S, Weisser HJ, Heizmann J, Pollak S (1998) Location and frequency of polymorphic positions in the mtDNA control region of individuals from Germany. *Int J Legal Med* 111:67–77
- Luzina FA (1987) Hereditary polymorphism and genetic processes in indigenous populations of Altai region. PhD thesis, Moscow State University, Moscow
- Malyarchuk BA, Derenko MV (2000) Mitochondrial DNA variability in Eastern Slavs. In: Renfrew C, Boyle K (eds) *Archaeogenetics: DNA and the population prehistory of Europe*. Cambridge, McDonald Institute for Archaeological Research, pp 249–254
- Orekhov V, Poltoraus A, Zhivotovsky LA, Spitsyn V, Ivanov P, Yankovsky N (1999) Mitochondrial DNA sequence diversity in Russians. *FEBS Letters* 445:197–201
- Osipova LP, Kashinskaya YO, Posukh OL, Ivakin EA, Kryukov YA (1997) A genetic demographic study of the South Altaian Population of the Mendur-Sokkon Village, Altai Republic. *Genetika* 33:1559–1564 (in Russian)
- Parson W, Parsons TJ, Scheithauer R, Holland MM (1998) Population data for 101 Austrian Caucasian mitochondrial DNA d-loop sequences: application of mtDNA sequence analysis to a forensic case. *Int J Legal Med* 111:124–132
- Piercy R, Sullivan KM, Benson N, Gill P (1993) The application of mitochondrial DNA typing to the study of white Caucasian genetic identification. *Int J Leg Med* 106:85–90
- Potapov LP (1969) Ethnical structure and origin of Altaians. Nauka, Leningrad
- Rousselet F, Mangin P (1998) Mitochondrial DNA polymorphisms: a study of 50 French Caucasian individuals and application to forensic casework. *Int J Legal Med* 111:292–298
- Santos FR, Pandya A, Tyler-Smith C, Pena SDJ, Schanfield M, Leonard WR, Osipova L, Crawford MH, Mitchell RJ (1999) The Central Siberian origin for Native American Y chromosomes. *Am J Hum Genet* 64:619–628
- Stone AC, Stoneking M (1998) mtDNA analysis of a prehistoric Oneota population: implications for the peopling of the New World. *Am J Hum Genet* 62:1153–1170
- Sukernik RI, Schurr TG, Starikovskaya YB, Wallace DC (1996) Mitochondrial DNA variation in native Siberians, with special reference to the evolutionary history of American Indians: studies on restriction endonuclease polymorphism. *Genetika* 32:432–439 (in Russian)
- Torroni A, Huoponen K, Francalacci P, Petrozzi M, Morelli L, Scozzari R, Obinu D, Savontaus ML, Wallace DC (1996) Classification of European mtDNAs from an analysis of three European populations. *Genetics* 144:1835–1850
- Underhill PA, Jin L, Zemans R, Oefner PJ, CavalliSforza LL (1996) A preColumbian Y chromosomespecific transition and its implications for human evolutionary history. *Proc Natl Acad Sci USA* 93:196–200
- Wallace DC (1995) Mitochondrial DNA variation in human evolution, degenerative disease and aging. *Am J Hum Genet* 57:201–223

Address for correspondence and reprints: Dr. Miroslava V. Derenko, Genetics Laboratory, Institute of Biological Problems of the North, Portovaya Street 18, Magadan, 685000, Russia. E-mail: ibpn@online.magadan.su

© 2001 by The American Society of Human Genetics. All rights reserved.
0002-9297/2001/6901-0027\$02.00