

B. I. G. Haussmann · D. E. Hess · N. Seetharama  
H. G. Welz · H. H. Geiger

## Construction of a combined sorghum linkage map from two recombinant inbred populations using AFLP, SSR, RFLP, and RAPD markers, and comparison with other sorghum maps

Received: 18 July 2001 / Accepted: 6 August 2001 / Published online: 30 July 2002  
© Springer-Verlag 2002

**Abstract** Sorghum [*Sorghum bicolor* (L.) Moench] is an important crop in the semi-arid tropics that also receives growing attention in genetic research. A comprehensive reference map of the sorghum genome would be an essential research tool. Here, a combined sorghum linkage map from two recombinant inbred populations was constructed using AFLP, SSR, RFLP and RAPD markers. The map was aligned with other published sorghum maps which are briefly reviewed. The two recombinant inbred populations (RIPs) analyzed in this study consisted of 225 (RIP 1) and 226 (RIP 2) F<sub>3,5</sub> lines, developed from the crosses IS 9830 × E 36-1 (RIP 1) and N 13 × E 36-1 (RIP 2), respectively. The genetic map of RIP 1 had a total length of 1,265 cM (Haldane), with 187 markers (125 AFLPs, 45 SSRs, 14 RFLPs, 3 RAPDs) distributed over ten linkage groups. The map of RIP 2 spanned 1,410 cM and contained 228 markers (158 AFLPs, 54 SSRs, 16 RFLPs) in 12 linkage groups. The combined map of the two RIPs contained 339 markers (249 AFLPs, 63 SSRs, 24 RFLPs, 3 RAPDs) on 11 linkage groups and had a length of 1,424 cM. It was in good agreement with other sorghum linkage maps, from which it deviated by a few apparent inversions, deletions, and additional distal regions.

**Keywords** *Sorghum bicolor* · Genetic linkage map · AFLP · RFLP · SSR · RAPD

Communicated by G. Wenzel

B.I.G. Haussmann · H.H. Geiger (✉)  
University of Hohenheim, 350 Institute of Plant Breeding,  
Seed Science, and Population Genetics, 70593 Stuttgart, Germany  
e-mail: geigerhh@uni-hohenheim.de

D.E. Hess  
International Crops Research Institute for the Semi-Arid Tropics  
(ICRISAT), B.P. 320, Bamako, Mali

N. Seetharama  
ICRISAT, Patancheru 502 324, Andhra Pradesh, India

H.G. Welz  
Aventis CropScience, Champaign, IL 61820, USA

### Introduction

Sorghum [*Sorghum bicolor* (L.) Moench], an important crop in the diet of millions of people in the semi-arid tropics, is receiving growing attention in genetic research. A number of genetic linkage maps of sorghum have been published in the last decade (see Table 1). Several are purely based on RFLP (restriction fragment length polymorphism) markers (Hulbert et al. 1990; Binelli et al. 1992; Whitkus et al. 1992; Berhan et al. 1993; Chittenden et al. 1994; Pereira et al. 1994; Ragab et al. 1994; Xu et al. 1994; Dufour et al. 1997; Ming et al. 1998; Crasta et al. 1999; Peng et al. 1999; Bowers et al. 2000; Xu et al. 2000); others include also AFLP (amplified fragment length polymorphism; Boivin et al. 1999; Klein et al. 2001), SSR (simple-sequence repeats; Taramino et al. 1997; Tao et al. 1998, 2000; Bhatramakki et al. 2000; Kong et al. 2000; Bennetzen et al. 2001; Klein et al. 2001), RAPD (randomly amplified polymorphic DNA; Tuinstra et al. 1996, 1997), or morphological markers (Bennetzen et al. 2001). But all of these genetic maps still have deficiencies, including the absence of mapped telomeres and centromeres, and a lack of connections to the physical chromosomal structures. Moreover, these maps have relatively few markers in common which would allow a comparison across sorghum populations or between sorghum and other plant species (Bennetzen et al. 2001). Ideally, all genetic and physical maps of sorghum should be integrated to form a comprehensive reference map. This would facilitate the choice of markers that are either evenly distributed across the genome, or specific to certain genomic regions. It would also ease the comparison of sorghum maps with maps of other *Gramineae* species and several aspects of gene discovery and isolation. Our objective here was to construct a combined sorghum linkage map from two recombinant inbred populations using AFLP, SSR, RFLP, and RAPD markers. Employing previously mapped markers as anchors, we aligned it with other recently published sorghum maps.

## Materials and methods

### Plant material

Two recombinant inbred populations (RIPs) of sorghum, consisting of 225 (RIP 1) and 226 (RIP 2)  $F_3$ -derived  $F_5$  lines ( $F_{3,5}$  lines), were developed from the crosses IS 9830  $\times$  E 36-1 (RIP 1) and N 13  $\times$  E 36-1 (RIP 2). Line IS 9830 is a tall Sudanese feterita belonging to the caudatum race. Line N 13 from India is a durra sorghum. Line E 36-1 is assigned to the guinea/caudatum hybrid race with Ethiopian origin. The crosses were selfed and 226  $F_2$  plants per population advanced by single-seed descent to the  $F_4$  generation. The  $F_4$  lines were multiplied by selfing 40 panicles per line, and the resulting  $F_5$  seed was bulked. These  $F_3$ -plant-derived bulks in  $F_5$  are called  $F_{3,5}$  lines here. Each  $F_{3,5}$  line represents the gene content of one  $F_3$  plant. One  $F_{3,5}$  line in RIP 1 proved to be an off-type and was therefore removed from the data set.

### Marker analyses

Marker analyses were performed with bulked DNA from 20 seedlings per  $F_{3,5}$  line. The genomic DNA was extracted following a modified CTAB (cetyl trimethyl ammonium bromide) extraction protocol (Saghai Maroof et al. 1984). RIP 1 was genotyped at 225 marker loci (131 codominantly and 20 dominantly scored AFLPs, 51 SSRs, 17 RFLPs, 6 RAPDs) and RIP 2 at 292 marker loci (122 codominantly and 75 dominantly scored AFLPs, 58 SSRs, 20 RFLPs, 17 RAPDs). The lower number of markers in RIP 1 was due to a lower genetic distance between IS 9830 and E 36-1 as compared to N 13 and E 36-1.

The AFLP markers were scored by Keygene Inc. (The Netherlands), following the procedures described by Vos et al. (1995). The totals of 151 and 197 AFLP markers in RIPs 1 and 2, respectively, were obtained from ten *EcoRI/Mse* primer combinations. Seven of the ten primer combinations were common to both RIPs: E11/M60, E12/M47, E13/M61, E14/M48, E14/M50, E14/M60 and E33/M50; three were applied to RIP 1 only: E11/M55, E12/M61 and E13/M59; and three to RIP 2 only: E11/M48, E11/M49 and E14/M61. Percentages of codominantly scored AFLP markers were 87% and 62% in RIPs 1 and 2, respectively. The seven primer combinations common for both RIPs yielded 34 common markers.

RFLP analyses were run by Biogenetic Services (Brookings, S.D., USA) according to standard procedures. The BNL (Burr and Burr 1991), CSU (Gardiner et al. 1993), PHP (Beavis and Grant 1991), UMC (Coe et al. 1990) and ISU (cDNA) maize probes were selected from sorghum genetic maps published by Pereira et al. (1994), Dufour et al. (1997), and Boivin et al. (1999). Using three restriction enzymes (*EcoRI*, *EcoRV* and *HindIII*), 201 probes were tested for polymorphism among the parent lines. The RIPs were then genotyped with one to three anchor RFLP markers per linkage group.

SSR analyses were done by Celera AgGen (Davis, Calif., USA) using microsatellites developed by Brown et al. (1996), Taramino et al. (1997), Kong et al. (2000) and Bhatramakki et al. (2000). PCR conditions were standard, following the recommendations of the aforementioned authors. The data was generated on an ABI Prism 377 DNA sequencer (Perkin Elmer Biosystems), and presented as alleles scored as estimated fragment sizes in base pairs compared to size standards or controls. The accuracy of scoring was approximately 0.67 base pairs (G. Hookstra, personal communication), but only markers with at least a two base pair difference between the parent lines were considered polymorphic. The base pair data were converted to letters corresponding to the respective parents, or 'H' when both bands were detected. A total of 241 SSRs was tested for polymorphism among the parent lines. RIPs 1 and 2 were genotyped with 51 and 58 selected SSRs, respectively. Most of these SSRs had been selected from the map of Bhatramakki et al. (2000), either because of their telomeric position or because of their uniform distribution across the genome.

The RAPD analyses were performed by ICRISAT-India. A total of 221 primers (decamers) were used. Of these, 125 were from Operon technologies, Incorporated, and 96 from Genosys, USA. A total of 40 cycles were used for amplification with the PE 9600 GeneAmp system. PCR conditions applied for the first 39 cycles were; denaturation: 93 °C for 1 min; annealing: 40 °C for 1 min; extension: 72 °C for 2 min. In the final cycle the extension period was 10 min. Useful primers were identified using a three-phase screening. In primary screening (stage I), the three parents were screened with all the primers. In secondary screening (stage II), along with each pair of mapping parents, a subset of progeny ( $F_{3,5}$  lines) were screened with only those primers showing polymorphism between parents at stage I. Finally, in the third phase of screening (stage III), primers selected in stage II were used and the whole set of mapping populations was screened along with the parents. Presence or absence of bands was scored for mapping at this stage.

### Statistical analysis

The chi-square test was used to test the markers for a 3:2:3 segregation ratio expected in the  $F_3$  with codominant markers, or for a 3:5 segregation ratio expected with dominantly scored markers.

The computer software Joinmap 2.0 (Stam and Van Ooijen 1995) was employed for map construction. Maps were first computed separately for each RIP, then combined. Initially, markers were assigned to linkage groups, based on modified LOD scores of marker pairs. These modified LOD scores are based on a chi-square test for independence of segregation. The test of independence assures that the LOD scores are not affected by a contingent distortion of segregation, i.e., spurious linkages of markers with segregation distortion are avoided (Stam and van Ooijen 1995). The LOD grouping threshold was set to 5.0 in both RIPs. However, in RIP 2, linkage groups B and J stuck together and could only be separated at a LOD grouping threshold of 11.0. In a second step, recombination frequencies were computed for all marker pairs within each linkage group. Third, the maps for each linkage group were constructed using the "JMAP" module. Data points with a LOD score below 0.1 or a recombination fraction above 0.49 were neglected in the final computations. These non-stringent thresholds were used to retain rather distantly linked markers in the data set. Most linkage groups reacted to the parameter settings ("LOD" and "REC" values) with changes in marker order. More stringent parameter settings resulted in erroneous marker orders as judged from the "list of top-linked markers" in the Joinmap output (see Stam and van Ooijen 1995) and compared to the map of Bhatramakki et al. (2000). Recombination frequencies were converted to centiMorgans (cM) with Haldane's mapping function (Haldane 1919). A "ripple" was performed after three marker additions/insertions. The "jump" and "triple" thresholds were set to 4 and 9, respectively. The linkage groups were named according to common SSR markers selected from the map of Bhatramakki et al. (2000).

The goodness-of-fit of constructed maps, expressed as the chi-square value (Stam and van Ooijen 1995), was calculated as:

$$\chi^2 = -2.0 \ln \{ \text{likelihood}[(r_{\text{joint}})] / \text{likelihood}[(r_{\text{direct}})] \},$$

where ( $r_{\text{joint}}$ ) is the set of pairwise recombination rates corresponding to the calculated map, and ( $r_{\text{direct}}$ ) are the estimates read from the input file. The calculated likelihood covers all pairwise distances in the map for which direct estimates are available.

The two maps had 67 loci in common (30 AFLPs, 33 SSRs and 4 RFLPs). Heterogeneity of recombination frequencies between common marker pairs was tested using the chi-square test in the "JMHE" module. The final combined map was constructed employing the same parameter settings described above.

## Results

### Polymorphism among parent lines

The proportion of polymorphic RFLP probes (polymorphic with at least one of the three restriction enzymes used) was 39.3% for the parent lines of RIP 1 (IS 9830 and E 36-1), and 48.8% for the parent lines of RIP 2 (N 13 and E 36-1). The degree of polymorphism for the 241 tested SSRs was 53.2% and 65.2%, respectively. The average number of polymorphic AFLP markers per primer combination amounted to 15.1 in RIP 1 and 19.7 in RIP 2. Only 2.7% and 7.2% of the tested RAPD markers were finally selected for genotyping the two mapping populations.

### Map of RIP 1 (IS 9830 × E 36-1)

Out of the 225 marker loci used for genotyping RIP 1, 84 (37%) showed significantly ( $P \leq 0.01$ ) distorted segregation. The genetic map of RIP 1 had a total length of 1,264.6 cM, with 187 markers (125 AFLPs, 45 SSRs, 14 RFLPs and 3 RAPDs) distributed over ten linkage groups. Thirty eight marker loci were unlinked or largely disturbed the map construction, thereby decreasing the goodness-of-fit of the map. These markers (mostly RAPDs, dominantly scored AFLPs, or markers with highly distorted segregation) were removed from the data set. The average and maximal distances between two individual markers were 6.8 and 38.8 cM. There was some clustering of (mainly AFLP) markers, i.e., 17 positions in the genome were marked two-fold, and three positions three-fold. Large gaps (>25 cM) occurred in linkage groups B (two gaps), D (one), E (two), F (one) and G (two). A high goodness-of-fit (mean  $\chi^2$  value <2.0 in the map construction of all linkage groups) indicated reliability of the linkage map. Loci with distorted segregation were found in all linkage groups, and mostly appeared in clusters. Linkage group D was exceptional in that 14 out of its 16 loci, covering 64.7 cM, showed significantly distorted segregation in favor of parent line IS 9830.

### Map of RIP 2 (N 13 × E 36-1)

Out of the 292 marker loci used for genotyping RIP 2, 110 (38%) showed significantly ( $P \leq 0.01$ ) distorted segregation. The map of RIP 2 spanned 1,409.5 cM and contained 228 markers (158 AFLPs, 54 SSRs and 16 RFLPs) in 12 linkage groups. Anchor markers or AFLP markers common with RIP 1 indicated that linkage groups B and D were still split into two parts. Sixty three markers were unlinked or largely decreased the goodness-of-fit of the map. Markers which severely disturbed the mapping process were mostly RAPDs, dominantly scored AFLPs, or markers with highly distorted segregation. They were removed from the data set. The average

and maximal distances between two individual markers were 6.2 and 42.7 cM. Some clustering of AFLP markers was observed, and 16 positions in the genome were marked two-fold, one four-fold, and one six-fold. Large gaps (>25 cM) occurred on linkage groups A (1 gap), C (2), D (1), E (4), G (2) and I (1). In addition, there are two missing links in the linkage groups B and D. As with RIP 1, a high goodness-of-fit (mean  $\chi^2$  value <2.0 in the map construction of all linkage groups) indicated reliability of the linkage map for RIP 2. Loci with distorted segregation were found in all linkage groups, and mostly appeared in clusters. Linkage group G harbored the largest cluster of eight markers, spanning 52.4 cM, with significantly distorted segregation in favor of parent line N 13.


### Combined map of RIPs 1 and 2

The 67 common loci of the two individual maps were distributed over the linkage groups as follows: linkage group A: 10; B1: 0; B2: 8; C: 13; D: 4; E: 5; F: 6; G: 4; H: 4; I: 9; J: 4. The two individual maps generally showed conservation of loci order with four exceptions (on linkage groups A, C, E and I) where the position of tightly linked loci was exchanged. Heterogeneity of recombination frequencies was non-significant for the majority of common marker pairs. The numbers of marker pairs with significant ( $P \leq 0.01$ ) heterogeneity of recombination frequencies were distributed as follows over the individual linkage groups: A: 2; B2: 9; C: 2; D: 1; E: 0; F: 2; G: 0; H: 0; I: 0; J: 0. The combined map contained 339 markers (249 AFLPs, 63 SSRs, 24 RFLPs and 3 RAPDs) distributed over 11 linkage groups and spanning 1,423.9 cM (Fig. 1). Linkage group B was still split into two parts. The average and maximal distances between two individual markers were 4.2 and 44.9 cM. Marker clusters contained mainly AFLP loci. Twenty seven positions in the genome were marked two-fold, seven three-fold, two four-fold, and one position five-fold. Large gaps (>25 cM) occurred on linkage groups B (the missing link), C (one gap), E (3) and G (1). A high goodness-of-fit (mean  $\chi^2$  value <2.0 in the map construction of all linkage groups) indicated the reliability of the combined map.

## Discussion

### General features of the computed maps

The total lengths of the maps are within the range reported for sorghum in the literature (Table 1). The shorter map for RIP 1 (1,265 cM) compared to RIP 2 (1,410 cM) is probably due to lower genetic polymorphism between IS 9830 and E 36-1 (the parent lines of RIP 1) as compared to N 13 and E 36-1 (the parent lines of RIP 2). In the combined map, the average distance between markers (4.2 cM) is comparatively low. But there are still sev-



**Fig. 1** Combined map from two recombinant inbred populations of sorghum. Marker names are indicated to the right of each linkage group. Numbers to the left of the linkage groups indicate cumulative map distances in cM (Haldane). SSR markers are written in *normal font*; AFLP loci are marked with a *grey frame*; RFLP marker names are written in *italics*, and RAPD marker names are underlined. Linkage groups are named according to Bhattaramakki et al. (2000) via common SSR markers

eral gaps that need to be filled. Besides the missing link in linkage group B, the largest gaps (about 40 cM) appeared on linkage groups C, E and G. On the other hand, there are regions in the combined map with very high marker density. Obviously, the distribution of the AFLP markers is not uniform across the genome. The upper

part of linkage groups B and H, and the entire E linkage group are poorly covered with AFLPs. In the remaining linkage groups, the majority of AFLP markers tend to cluster in central regions. However, a few AFLPs are remarkable in that they mapped to unique, sometimes telomeric, positions. In the sorghum “AFLP RIL 379 genetic linkage map” of Boivin et al. (1999), AFLP markers also exhibited clustering and interspersions in the core RFLP map. Clusters of *Eco*RI AFLP markers have also been reported in other crops like maize (*Zea mays* L.; Castiglioni et al. 1999; Vuylsteke et al. 1999), ryegrass (*Lolium perenne* L.; Bert et al. 1999), rice (*Oryza sativa* L.; Maheswaran et al. 1997), soybean (*Glycine max* (L.) Merr.; Keim et al. 1997), sugar beet (*Beta vulgaris* L.; Schondelmaier et al. 1996) and barley (*Hordeum vulgare*




Fig. 1 (continued)

L.; Becker et al. 1995). Vuylsteke et al. (1999) showed that the *Eco*RI-AFLP clusters colocalized well with the putative centromeric regions of the maize chromosomes. In centromeric regions, crossing-over during meiosis is greatly reduced (Luckaszewski and Curtis 1993), causing the markers to cluster. The high frequency of *Eco*RI-AFLP markers in centromeres may possibly be explained by an AT-richness in the pericentromeric heterochromatin (as shown for *Arabidopsis* by Ross et al. 1996) and the AT-rich target sequences of the restriction enzymes *Eco*RI and *Mse*I (Vuylsteke et al. 1999). According to Boivin et al. (1999), heterochromatic regions around centromeres and at the chromosome tips could also be well-accessible to *Eco*RI-based AFLP markers because these markers are non-sensitive to CG methylation. Since, in our study, SSR and RFLP markers were primarily selected for particular map positions, their distribution in the maps is “artificial” and will not be further discussed. Their linear order was mostly as expected and is described in more detail below.

Alignment of the composite map with other sorghum genetic linkage maps

Peng et al. (1999) summarized the relationships among the linkage groups in their own and seven other published RFLP linkage maps of sorghum. Our combined map has markers in common with other recently published sorghum genetic linkage maps (Table 2). Our observations confirmed the relationships as summarized by Peng et al. (1999). The linkage group names in the maps of Bennetzen et al. (2001), Bhatramakki et al. (2000) and Kong et al. (2000), as well as those presented in this paper, correspond to those assigned initially by Peng et al. (1999).

Our combined map has 29 markers in common with the framework linkage map of sorghum proposed by Bennetzen et al. (2001). The order of common markers (where there are three or more in a linkage group) is fully consistent between the two maps. The largest difference between marker distances in the two maps appears on linkage group A: the distance between XTXP43 and XTXP 208 is 99.3 cM in our map but only about 55 cM in that of Bennetzen et al.

**Table 1** Overview of published sorghum genetic linkage maps

Reference	Size and type of mapping pop.	Parent lines	Mapped markers	Linkage groups	Genome length (cM) <sup>a</sup>
Hulbert et al. 1990	55 F <sub>2</sub> plants	Shanqui Red (koaliang from China) M91051 (zera zera from East Africa)	37 RFLP maize probes or cloned genes	8	283 R
Binelli et al. 1992	149 F <sub>2</sub> plants	IS 18729 (caudatum-bicolor from Texas) IS 24756 (durra-caudatum from Nigeria)	21 RFLP maize probes	5	440 U
Whitkus et al. 1992	81 F <sub>2</sub> plants	IS2482C ( <i>S. bicolor</i> ssp. <i>bicolor</i> ) IS18809 ( <i>S. bicolor</i> ssp. <i>arundinaceum</i> )	85 RFLP maize probes 7 Isoenzymes	13	949 H
Berhan et al. 1993	55 F <sub>2</sub> plants	Same parents as Hulbert et al. 1990	96 RFLP maize probes or cloned genes	15	709 R
Chittenden et al. 1994	56 F <sub>2</sub> plants	BTx 623 ( <i>S. bicolor</i> ) Unnamed accession of <i>S. propinquum</i>	256 RFLP sorghum probes 20 RFLP probes of maize, rice or oat	10	1,445 U
Pereira et al. 1994	78 F <sub>2</sub> plants	CK 60 ( <i>S. bicolor</i> ssp. <i>bicolor</i> ) PI229828 ( <i>S. bicolor</i> ssp. <i>drummondii</i> )	191 RFLP maize probes 10 RFLP sorghum probes	10	1,530 U
Xu et al. 1994	50 F <sub>2</sub> plants	BTx 623 (zera zera × kafir) IS 3620C (guinea line)	179 RFLP sorghum probes 11 RFLP maize probes	14	1,789 K
Ragab et al. 1994	93 F <sub>2,3</sub> families	BSC 35 BTx 631	38 RFLP sorghum probes 33 RFLP maize probes	15	633 H
Lin et al. 1995	370 F <sub>2</sub> plants	Same parents as Chittenden et al. 1994	78 RFLP sorghum probes 124 RFLP loci mapped by Chittenden et al. 1994	11	935 K
Tuinstra et al. 1996, 1997	98 RIL (F <sub>5;7-8</sub> )	TX 7078 B35 (durra sorghum from Ethiopia)	150 RAPD 20 RFLP probes from maize or sorghum	17	Ca 1,580 R
Dufour et al. 1997	110 RIL (F <sub>5</sub> )	IS 2807 (caudatum from Zimbabwe) 379 (guinea from South Africa)	126 RFLP maize probes 19 RFLP sugarcane probes 4 cloned genes, 2 morphological markers	13	977 H
	91 RIL (F <sub>5</sub> )	IS 2807 (caudatum from Zimbabwe) 249 (guinea from Burkina Faso)	115 RFLP maize probes 8 RFLP sugarcane probes 4 cloned genes, 1 morphological marker	12	878 H
	Composite map of the two populations		164 RFLP maize probes 19 RFLP sugarcane probes 3 cloned genes, 2 morphological markers	13	1,095 H
Taramino et al. 1997	68 F <sub>2</sub> plants	Same parents as Pereira et al. 1994	7 SSR added to map of Pereira et al. 1994	10	1,575 U
Ming et al. 1998	56 F <sub>2</sub> plants	Same parents as Chittenden et al. 1994	328 RFLP probes of sorghum and other cereals	10	1,750 K
Tao et al. 1998	120 RIL (F <sub>5</sub> )	QL39 QL41 (both elite lines from Australia)	68 RFLP sorghum probes 87 RFLP of maize, sugarcane, rice, oat, barley 8 SSR, 3 morphological traits	21	1,400 U
Boivin et al. 1999	Composite map with populations as Dufour et al. 1997		343 RFLP and morphological markers	11	1,352 H
	110 RIL (F <sub>5</sub> )	First population of Dufour et al. 1997	128 RFLP sorghum, rice, oat or barley probes 176 AFLP	11	1,899 H

**Table 1** (continued)

Reference	Size and type of mapping pop.	Parent lines	Mapped markers	Linkage groups	Genome length (cM) <sup>a</sup>
Boivin et al. 1999			151 RFLP loci mapped by Dufour et al. 1997		
Crasta et al. 1999	96 RIL (F <sub>6;7</sub> )	B35 (durra from Ethiopia) Tx 430	142 RFLP sorghum, maize, rice or wheat clones	14	1,602 K
Peng et al. 1999	137 RIL (F <sub>6-8</sub> )	Same parents as Xu et al. 1994	323 RFLP from sorghum, maize, rice, oat, barley	10	1,347 K
Bowers et al. 2000	65 F <sub>2</sub> plants	Same parents as Chittenden et al. 1994	2,399 Loci based on 1925 RFLP probes	10	1,200 U
Kong et al. 2000	137 RIL (F <sub>6-8</sub> )	Same parents as Xu et al. 1994	33 SSR	10	1,287 K
Tao et al. 2000	152 RIL (F <sub>5</sub> )	Same parents as Tao et al. 1998	111 RFLP probes from Peng et al. 1999 101 RFLP	14	1,871 U
Xu et al. 2000	98 RIL (F <sub>7</sub> )	B 35 (durra from Ethiopia) Tx 7000 (elite public line used in USA)	17 SSR 166 markers mapped by Tao et al. 1998 162 RFLP sorghum and maize probes, cloned genes or sequenced DNA probes	10	837 H
Bhatramakki et al. 2000	137 RIL (F <sub>6-8</sub> )	Same parents as Xu et al. 1994	116 SSR 354 RFLP or SSR markers mapped by Peng et al. 1999 or Kong et al. 2000	10	1,406 K
Bennetzen et al. 2001	Framework map derived from comparison of the maps of Kong et al. 2000, Peng et al. 1999, Pereira et al. 1994, and Berhan et al. 1993.		154 RFLP	10	1,450 U
Klein et al. 2001	125 RIL (F <sub>5</sub> )	RTx 430 (elite line from USA) Sureño	34 SSR, 10 morphological markers 44 SSR 85 AFLP, 1 morphological trait	10	970 K
Haussmann et al., present publication	225 RIL (F <sub>3;5</sub> )	IS 9830 (caudatum line from Sudan) E 36-1 (Ethiopian guinea-caudatum)	125 AFLP 45 SSR, 14 RFLP, 3 RAPD	10	1,265 H
	226 RIL (F <sub>3;5</sub> )	N 13 (Indian durra) E 36-1 (Ethiopian guinea-caudatum)	158 AFLP 54 SSR, 16 RFLP	12	1,410 H
	Composite map of the two populations		339 AFLP, SSR, RFLP and RAPD markers	11	1,424 H

<sup>a</sup> H, K = Map distances estimated using the mapping functions of Haldane (1919) and Kosambi (1944), respectively; U = mapping function not specified; R = Recombination frequency (%)

Sixty six markers are shared by our combined map and the map of Bhatramakki et al. (2000). The order of common markers is generally conserved, except for three marker pairs (XTXP97 and XTXP96 on linkage group B, XTXP289 and UMC12 on F, and XTXP217 and XTXP270 on G). These markers are closely linked in both maps but their order is reversed in our map. On our linkage group G, there is an additional reversal in the order of the markers XTXP141 and SB1-1 which are 17.4 distant in our map, and 29.3 cM apart in the map of Bhatramakki et al. (2000). The largest missing regions

in our combined map relative to the map of Bhatramakki et al. are: approximately 27 cM on linkage group A at the XTXP208 end, about 35 cM on linkage group B between XTXP50 and XTXP201, and about 46 cM on linkage group D at the XTXP12 end. Additional regions in our map (compared to the same map) were detected in linkage groups H (15 cM at the XTXP273 end) and I (11 cM at the XTXP6 end).

The number of common markers between the map of Klein et al. (2000) and our combined map is 20. In linkage group A, the order of the markers XTXP43 and

**Table 2** Linkage groups in our combined map *versus* corresponding linkage groups in selected, recently published sorghum maps. Numbers of markers shared with our map given in parentheses.

Reference	Linkage group in our combined map									
	A	B1+2	C	D	E	F	G	H	I	J
Bennetzen et al. 2001	A (4)	B (3)	C (3)	D (2)	E (2)	F (4)	G (2)	H (3)	I (4)	J (2)
Bhatramakki et al. 2000	A (9)	B(11)	C (7)	D (5)	E (6)	F (5)	G (6)	H (6)	I (5)	J (6)
Klein et al. 2000	A (3)	B (3)	C (1)	D (2)	E (2)	F (1)	G (0)	H (2)	I (3)	J (3)
Kong et al. 2000	A (2)	B (2)	C (1)	D (1)	E (1)	F (2)	G (1)	H (2)	I (1)	J (3)
Tao et al. 2000	C (1)	B (1)	–	F (3)	–	G (2)	I (2)	–	–	H (1)
Boivin et al. 1999	C (2)	F (3)	G (3)	D (2)	A (1)	E (3)	H (1)	I (2)	B (2)	J (2)
Peng et al. 1999	A (2)	B (2)	C (3)	D (0)	E (1)	F (2)	G (2)	H (0)	I (1)	J (0)

XTXP88, which are closely linked in both maps, is reversed in our map. In the other linkage groups with three common markers (B, I and J), marker order is completely conserved between the two maps. Distances between common markers are similar, except for linkage group J where our estimated distance between XTXP23 and XTXP303 is about 20-cM shorter.

Sixteen markers are common between our combined map and that of Kong et al. (2000). The distance between the common marker pair XTXP 37 and XTXP 43 on linkage group A is 12-cM shorter in our combined map than in that of Kong and his colleagues. On the other hand, the distances between common marker pairs on the linkage groups F (XTXP 10, UMC 12), H (XTXP 47, XTXP 18) and I (XTXP 65 and XTXP 15) are, respectively, 11-, 33-, and 26-cM longer in our map.

The map of Tao et al. (2000) shares ten markers with our combined map. A major difference is the order of common markers on our linkage group D (XTXP12–SB1–10–XTXP41) versus (SB1–10–XTXP12–XTXP41) in their linkage group F.

Twenty one markers are common with the composite map of Boivin et al. (1999). Distances between common marker pairs are in good agreement, with two exceptions. The distance between UMC 64 and BNL 5.47 is 49.0 cM in our map (linkage group F) but 89.6 cM in the linkage group E of Boivin et al.; and the distance between CSU 30 and PHP 20075 on linkage group J (in both maps) is 52.4 cM in our map but only 28.8 cM in the map of Boivin et al. (1999).

The map of Peng et al. (1999) contains 13 markers common to our combined map. The only major difference in distance between common markers occurred in linkage group A, between the markers BML 5.09 and UMC 166 (distance of 79.4 cM in our map versus about 100 cM in the map of Peng et al.).

In conclusion, the agreement of our combined map with other recently published sorghum maps is generally good, with a few inversions, deletions, and additional regions that we detected.

## Outlook

Efforts are needed to enhance the number of common markers in different sorghum maps, to allow for a more comprehensive comparison and integration. The search should continue for new, highly polymorphic SSRs that would map to regions in the sorghum genome which are currently poorly covered with SSRs. There is also an urgent need to relate the available genetic maps to a physical map. This would facilitate the map-based cloning of genes, comparative genome analysis, and provide sequence-ready clones for genome sequencing projects (Klein et al. 2000). Physical maps of sorghum are currently under construction at Texas A&M University in the group of P.E. Klein, R.R. Klein and J.E. Mullet (Klein et al. 2000), and at the University of Georgia/Texas A&M University in the group of A.H. Paterson (Bowlers et al. 2000).

**Acknowledgements** The authors sincerely thank Prof. Gary Hart (Texas A&M University) and his research group for providing SSR primer sequences and map positions even before the information was published. The generous financial support of the German Federal Ministry for Economic Cooperation and Development (BMZ) is gratefully acknowledged (Grant No. 94.7860.3-01.100).

## References

- Beavis WD, Grant D (1991) A linkage map based on information from four F<sub>2</sub> populations of maize (*Zea mays* L.). *Theor Appl Genet* 82:636–644
- Becker J, Vos P, Kuiper M, Salamini F, Heun M (1995) Combined mapping of AFLP and RFLP markers in barley. *Mol Gen Genet* 249:65–73
- Bennetzen JL, Subramanian V, Xu J, Salimath SS, Subramanian S, Bhatramakki D, Hart GE (2001) A framework genetic map of sorghum containing RFLP, SSR and morphological markers. In: Phillips RL, Vasil IK (eds) DNA-based markers in plants, 2nd edn, Chap 20. Kluwer Academic Publishers, Boston
- Berhan AM, Hulbert SH, Butler LG, Bennetzen JL (1993) Structure and evolution of the genomes of *Sorghum bicolor* and *Zea mays*. *Theor Appl Genet* 86:598–604
- Bert PF, Charmet G, Sourdille P, Hayward MD, Balfourier F (1999) A high-density molecular map for ryegrass (*Lolium perenne*) using AFLP markers. *Theor Appl Genet* 99:445–452
- Bhatramakki D, Dong J, Chhabra AK, Hart GE (2000) An integrated SSR and RFLP linkage map of *Sorghum bicolor* (L.) Moench. *Genome* 43:988–1002



- Binelli GL, Gianfranceschi L, Pe ME, Taramino G, Busso C, Stenhouse J, Ottaviano E (1992) Similarity of maize and sorghum genomes as revealed by maize RFLP probes. *Theor Appl Genet* 84:10–16
- Boivin K, Deu M, Rami JF, Trouche G, Hamon P (1999) Towards a saturated sorghum map using RFLP and AFLP markers. *Theor Appl Genet* 98:320–328
- Bowers JE, Schertz KF, Abbey C, Anderson S, Chang C, Chittenden LM, Draye X, Hoppe AH, Jessup R, Lennington J, Lin Y, Liu S, Ming R, Qiang D, Reischmann K, Skinner N, Wang Y, Paterson AH (2000) A high-density 2,399-locus genetic map of *Sorghum*. Final Abstracts Guide, Plant and Animal Genome VIII Conference, Jan 9–12, 2000, San Diego, California, <http://www.intl-pag.org/pag/8/abstracts/pag8712.html>
- Brown SM, Hopkins MS, Mitchell SE, Senior ML, Wang TY, Duncan RR, Gonzales-Candelas F, Kresovitch S (1996) Multiple methods for the identification of polymorphic simple sequence repeats (SSRs) in sorghum [*Sorghum bicolor* (L.) Moench]. *Theor Appl Genet* 93:190–198
- Burr B, Burr FA (1991) Recombinant inbreds for molecular mapping in maize: theoretical and practical considerations. *Trends Genet* 7:55–60
- Castiglioni P, Ajmone-Marsan P, van Wijk R, Motto M (1999) AFLP markers in a molecular linkage map of maize: codominant scoring and linkage group distribution. *Theor Appl Genet* 99:425–431
- Chittenden LM, Schertz KF, Lin YR, Wing RA, Paterson AH (1994) A detailed RFLP map of *Sorghum bicolor* × *S. propinquum*, suitable for high-density mapping, suggests ancestral duplication of *Sorghum* chromosomes or chromosomal segments. *Theor Appl Genet* 87:925–933
- Coe EH, Hoisington DA, Neuffer MG (1990) Linkage map of corn (maize) (*Zea mays* L.). In: O'Brien SJ (ed) Genetic maps. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, New York, pp 39–67
- Crasta OR, Xu WW, Rosenow DT, Mullet J, Nguyen HT (1999) Mapping of post-flowering drought resistance traits in grain sorghum: association between QTLs influencing premature senescence and maturity. *Mol Gen Genet* 262:579–588
- Dufour P, Deu M, Grivet L, D'Hont A, Paulet F, Bouet A, Lanaud C, Glaszmann JC, Hamon P (1997) Construction of a composite sorghum genome map and comparison with sugarcane, a related complex polyploid. *Theor Appl Genet* 94:409–418
- Gardiner JM, Coe EH, Melia-Hancock S, Hoisington DA, Chao S (1993) Development of a core RFLP map in maize using an immortalized F<sub>2</sub> population. *Genetics* 134:917–930
- Haldane JBS (1919) The combination of linkage values, and the calculation of distance between the loci of linked factors. *J Genet* 8:299–309
- Hulbert SH, Richter TE, Axtell JD, Bennetzen JL (1990) Genetic mapping and characterization of sorghum and related crops by means of maize DNA probes. *Proc Natl Acad Sci USA* 87:4251–4255
- Keim P, Schupp JM, Travis SE, Clayton K, Zhu T, Shi L, Ferreira A, Webb DM (1997) A high-density soybean genetic map based on AFLP markers. *Crop Sci* 37:537–543
- Klein PE, Klein RR, Morishige DT, Childs KL, Cartinhour SW, Mullet JE (2000) Construction of an integrated genetic and physical map of sorghum. Final Abstracts Guide, Plant and Animal Genome VIII Conference, Jan 9–12, 2000, San Diego, California, <http://www.intl-pag.org/pag/8/abstracts/pag8102.html>
- Klein RR, Rodriguez-Herrera R, Schlueter JA, Klein PE, Yu ZH, Rooney WL (2001) Identification of genomic regions that affect grain-mould incidence and other traits of agronomic importance in sorghum. *Theor Appl Genet* 102:307–319
- Kong L, Dong J, Hart GE (2000) Characteristics, linkage map positions, and allelic differentiation of *Sorghum bicolor* (L.) Moench DNA simple-sequence repeats (SSRs). *Theor Appl Genet* 101:438–448
- Kosambi DD (1944) The estimation of map distances from recombination values. *Ann Eugen* 12:172–175
- Luckaszewski P, Curtis CA (1993) Physical distribution of recombination in B-genome chromosomes of tetraploid wheat. *Theor Appl Genet* 85:121–127
- Maheswaran M, Subudhi PK, Nandi S, Xu JC, Parco A, Yang DC, Huang N (1997) Polymorphism, distribution, and segregation of AFLP markers in a double-haploid rice population. *Theor Appl Genet* 94:39–45
- Ming R, Liu S-C, Lin Y-R, da Silva J, Wilson W, Braga D, van Deynze A, Wenslaff TF, Wu KK, Moore PH, Burnquist W, Sorrells ME, Irvine JE, Paterson AH (1998) Detailed alignment of *Saccharum* and *Sorghum* chromosomes: comparative organization of closely related diploid and polyploid genomes. *Genetics* 150:1663–1682
- Peng Y, Schertz KF, Cartinhour S, Hart GE (1999) Comparative genome mapping of *Sorghum bicolor* (L.) Moench using an RFLP map constructed in a population of recombinant inbred lines. *Plant Breed* 118:225–235
- Pereira MG, Lee M, Bramel-Cox P, Woodman W, Doebley J, Whitkus R (1994) Construction of an RFLP map in sorghum and comparative mapping in maize. *Genome* 37:236–243
- Ragab RA, Dronavalli S, Maroof Saghai MA, Yu YG (1994) Construction of a sorghum RFLP linkage map using sorghum and maize DNA probes. *Genome* 37:590–594
- Ross K, Franz PF, Jones GH (1996) A light microscopic atlas of meiosis in *Arabidopsis thaliana*. *Chrom Res* 4:507–516
- Saghai Maroof MA, Soliman KM, Jorgenson RA, Allard RW (1984) Ribosomal DNA spacer length polymorphism in barley: Mendelian inheritance, chromosomal location and population dynamics. *Proc Natl Acad Sci USA* 81:8014–8018
- Schondelmaier J, Steinrucken G, Jung C (1996) Integration of AFLP markers into a linkage map of sugar beet (*Beta vulgaris* L.). *Plant Breed* 115:231–237
- Stam P, Van Ooijen JW (1995) JoinMap version 2.0: software for the calculation of genetic linkage maps. CPRO-DLO Wageningen, The Netherlands
- Tao YZ, Jordan DR, Henzell RG, McIntyre CL (1998) Construction of a genetic map in a sorghum RIL population using probes from different sources and its comparison with other sorghum maps. *Aust J Agric Res* 49:729–736
- Tao YZ, Henzell RG, Jordan DR, Butler DG, Kelly AM, McIntyre CL (2000) Identification of genomic regions associated with stay green in sorghum by testing RILs in multiple environments. *Theor Appl Genet* 100:1225–1232
- Taramino G, Tarchini R, Ferrario S, Lee M, Pe' ME (1997) Characterization and mapping of simple sequence repeats (SSRs) in *Sorghum bicolor*. *Theor Appl Genet* 95:66–72
- Tuinstra MR, Grote EM, Goldsbrough PB, Ejeta G (1996) Identification of quantitative trait loci associated with pre-flowering drought tolerance in sorghum. *Crop Sci* 36:1337–1344
- Tuinstra MR, Grote EM, Goldsbrough PB, Ejeta G (1997) Genetic analysis of post-flowering drought tolerance and components of grain development in *Sorghum bicolor* (L.) Moench. *Mol Breed* 3:439–448
- Vos P, Hogers R, Bleeker M, Reijans M, van de Lee T, Hornes M, Frijters A, Pot J, Peleman J, Kuiper M, Zabeau M (1995) AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Research* 23:4407–4414
- Vuytsteke M, Mank R, Antonise R, Bastiaans E, Senior ML, Stuber CW, Melchinger AE, Lübberstedt T, Xia XC, Stam P, Zabeau M, Kuiper M (1999) Two high-density AFLP linkage maps of *Zea mays* L.: analysis of distribution of AFLP markers. *Theor Appl Genet* 99:921–935
- Whitkus R, Doebley J, Lee M (1992) Comparative genome mapping of sorghum and maize. *Genetics* 132:1119–1130
- Xu GW, Magill CW, Schertz KF, Hart GE (1994) A RFLP linkage map of *Sorghum bicolor* (L.) Moench. *Theor Appl Genet* 89:139–145
- Xu W, Subudhi PK, Castra OR, Rosenow DT, Mullet JE, Nguyen HT (2000) Molecular mapping of QTLs conferring stay green on grain sorghum (*Sorghum bicolor* L. Moench). *Genome* 43:461–469