

T. Schnurbusch · S. Paillard · D. Fossati ·
M. Messmer · G. Schachermayr · M. Winzeler ·
B. Keller

Detection of QTLs for *Stagonospora glume blotch* resistance in Swiss winter wheat

Received: 3 March 2003 / Accepted: 11 June 2003 / Published online: 19 August 2003
© Springer-Verlag 2003

Abstract *Stagonospora nodorum* is the causal agent of the *Stagonospora glume blotch* disease in hexaploid wheat. The Swiss winter bread wheat cv. ‘Arina’ has a highly effective, durable and quantitative glume blotch resistance. We studied 240 single seed descent (SSD)-derived lines of an ‘Arina × Forno’ F_{5:7} population to identify and map quantitative trait loci (QTLs) for glume blotch resistance under natural infestation. Using composite interval mapping (CIM) and LOD>4.5, we detected two chromosomal regions on chromosome arms 3BS and 4BL which were specifically associated with glume blotch resistance. These identified QTLs were designated *QSng.sfr-3BS* and *QSng.sfr-4BL*, respectively. *QSng.sfr-3BS* peaked at the locus *Xgwm389* in the telomeric region of the short arm of chromosome 3B and explained 31.2% of the observed phenotypic variance for the resistance within the population. The responsible *QSng.sfr-3BS* allele originated from the resistant parent ‘Arina’. The QTL *QSng.sfr-4BL* (19.1%) mapped to chromosome arm 4BL (‘Forno’ allele) very close to two known genes, *TaMlo* and a catalase (*Cat*). Both QTL alleles combined could enhance the resistance level by about 50%. Additionally, they showed significant epistatic effects

(4.4%). We found PCR-based microsatellite markers closely linked to *QSng.sfr-3BS* (*gwm389*) and *QSng.sfr-4BL* (*gwm251*) which make marker-assisted selection (MAS) for *Stagonospora glume blotch* resistance feasible. We also found one resistance QTL, *QSng.sfr-5BL*, on the long arm of chromosome 5B which overlapped with QTLs for plant height as well as heading time.

Keywords *Triticum aestivum* · QTL · *Stagonospora nodorum* · Glume blotch · Resistance

Introduction

The leaf and glume blotch disease of bread wheat (*Triticum aestivum* L.) is caused by the fungus *Phaeosphaeria nodorum* (E. Müller) Hedjaroude, anamorph *Stagonospora nodorum* (Berk.) Castellani and Germano. The disease has recently been renamed according to Cunfer and Ueng (1999). *S. nodorum* survives on the field debris and is able to infest all above-ground plant organs of wheat. The major primary inoculum are the wind-dispersed ascospores (sexual form) in late autumn and early spring (Mittelstadt and Fehrmann 1987; Keller et al. 1997a; Bathgate and Loughman 2001). During the growing season, particularly under wet and warm weather conditions, the asexually generated and rain-splash-dispersed pycnidiospores successively infect different leaf levels of individual plants (Scharen 1966; Keller et al. 1997a). Wheat kernels are usually infected by the fungus and the infected seeds are possibly contributing to the spread of the fungus over longer distances (Shah and Bergstrom 1993).

S. nodorum is classified as one of the potentially most-devastating known plant pathogens in agriculture (McDonald and Linde 2002). The high risk potential is based on its mixed reproduction system and its high degree of gene and genotype diversity distributed on a small spatial scale, where almost every lesion on a leaf represents a genetically unique individual (McDonald and Linde 2002). The *Stagonospora blotch* disease regularly occurs

Communicated by H. C. Becker

T. Schnurbusch · B. Keller (✉)
Institute of Plant Biology,
University of Zürich,
Zollikerstrasse 107, 8008 Zürich, Switzerland
e-mail: bkeller@botinst.unizh.ch

S. Paillard · G. Schachermayr · M. Winzeler
Swiss Federal Research Station for Agroecology
and Agriculture (FAL),
Reckenholzstrasse 191, 8046 Zürich, Switzerland

D. Fossati
Swiss Federal Research Station for Plant Production,
Changins, Route de Duillier, P.O. Box 254, 1260 1 Nyon,
Switzerland

M. Messmer
Pharmaceutical Institute, University of Basel,
Benkenstrasse 254, 4108 Witterswil, Switzerland

in wheat-growing areas across Europe, North Africa, Near East, United States, Australia and Canada (Duczek et al. 1999; Halama 2002).

Genetic variability for *Stagonospora* blotch resistance is found within the wheat gene pool and its wild relatives (Polley and Thomas 1991; Ma and Hughes 1993; Cunfer and Johnson 1999; Loughman et al. 1999; Wicki et al. 1999). Breeding for *Stagonospora* blotch resistance is an effective strategy to reduce the yield losses caused by this disease. To-date, there is no evidence for immunity or gene-for-gene interaction in the wheat-*S. nodorum* pathosystem. Resistance to *Stagonospora* glume blotch is usually partial (Van der Plank 1963) and generally considered as a quantitative trait. A low correlation between leaf and spike resistance indicated various factors of resistance (Laubscher et al. 1966; Broennimann 1975; Nelson and Gates 1982; Fried and Meister 1987; Bostwick et al. 1993; Van Ginkel and Rajaram 1999; Wicki et al. 1999). As a consequence of the quantitative nature of the disease resistance, selecting resistant genotypes is time-consuming because of the complex interactions between host, pathogen and environment (Van Ginkel and Rajaram 1999). Additionally, plant height and heading time significantly influence the resistance level of *S. nodorum* on the leaf as well as on the spike (Broennimann et al. 1973; Scott et al. 1982; Van Ginkel and Rajaram 1999; Wicki et al. 1999).

Cytogenetic analyses for *Stagonospora* glume blotch resistance identified 16 chromosomes in the hexaploid wheat genome contributing to resistance in different lines and populations (Kleijer et al. 1977; Walag and Dziglo 1985, 1987; Auriou et al. 1988; Nicholson et al. 1993; Ma and Hughes 1995; Hu et al. 1996). So far, no QTLs for *Stagonospora* blotch resistance have been reported. In view of the difficulty to select resistant genotypes against glume blotch, marker-assisted selection (MAS) will be an ideal tool to efficiently introgress resistance QTLs into elite breeding material.

The objective of the present study was to analyze the *Stagonospora* glume blotch resistance within the Swiss winter wheat population 'Arina × Forno' under natural infestation, and to identify chromosomal regions which are specifically associated with glume blotch resistance.

Materials and methods

Plant material

Two hundred and forty single seed descent (SSD)-derived lines ($F_{5:7}$) from the intraspecific cross between the two adapted Swiss winter wheat (*T. aestivum* L.) cultivars 'Arina' and 'Forno' were evaluated in field trials for their *Stagonospora* glume blotch reaction. 'Arina' is a tall variety (1.25 m) with a loose spike. It was released in 1981 and has covered more than 40% of the wheat acreage of Switzerland since 1985. 'Arina' has maintained its excellent spike resistance against *Stagonospora* glume blotch in Switzerland and it can therefore be assumed that this resistance is durable. In contrast, 'Forno' is a shorter (1.05 m) Swiss winter bread-wheat variety highly susceptible to *Stagonospora* glume blotch under artificial inoculation conditions (Wicki et al. 1999).

Field experiments

Seven field trials were conducted in three locations in 2000 and 2001 representing the diverse wheat growing-areas in Switzerland. Three trials, ZH106 and ZH114 in 2000 as well as ZH109 in 2001, were conducted near Zürich, Switzerland, at the Federal Research Station for Agroecology and Agriculture (FAL-Reckenholz) at 443 m above sea level on loamy soils with an average precipitation of 1,000 mm and an average temperature of 7.9°C. There were two trials (Haag00 and Haag01) in Haag, Canton St. Gallen. Haag is located in the north-eastern part of Switzerland within the river Rhine valley at about 460 m above sea level. The field experiments were planted on heavy silty clayey soils with an average precipitation of 970 mm and an average temperature of 8.6°C. Additionally, in both years there was one trial in Vouvry, Canton Valais, which is located in the south-western part of Switzerland close to Lake Geneva, at the beginning of the Rhone valley. The two field trials (Vouvry00 and Vouvry01) were accomplished on loamy sandy soils at about 381 m above sea level with an average precipitation of 881 mm and an average temperature of 8.7°C.

The lines were grown together with the reciprocal F_1 , three standard cultivars and five replicated entries of the parental lines (267 entries) in a rectangular lattice design with three replications and nine genotypes per incomplete block. Each entry was sown as a one-row plot (1 m) with approximately 40 to 50 kernels.

Disease phenotyping and trait assessment

In 2000 and 2001, all field trials were evaluated for natural *Stagonospora* glume blotch infestation. All standard cultivars, the reciprocal F_1 , the parental lines and the lines of the 'Arina × Forno' population were phenotyped by estimating visually the average percentage of infected glume area per one-row plot according to the Broennimann scale (Broennimann 1968). The Broennimann scale displays the percentage of the infected glume area per spike starting with 0% or no visible lesions to 1, 5, 10, 25, 50, 75 and up to 100% infected glume area. In five environments, we scored twice with three replications in the growth stages of the BBCH-scale 71–77 (Meier 2001): ZH106 (26.06. and 30.06.2000), ZH114 (26.06. and 30.06.2000), Haag00 (28.06. and 03.07.2000), Vouvry01 (29.06. and 06.07.2001) and Haag01 (28.06. and 04.07.2001). In Vouvry00 and ZH109, only one late scoring for *Stagonospora* glume blotch resistance was made (04.07.2000 and 10.07.2001, respectively). Plant height (cm) in all seven environments as well as heading time (days after January 1st) in six environments were investigated with two replications for the 'Arina × Forno' population.

Genetic linkage map

Two hundred and forty lines of the 'Arina × Forno' cross were genotyped with molecular markers. A genetic linkage map based on microsatellite (SSR) and restriction fragment length polymorphism (RFLP) markers was established (Paillard et al. 2003). Three hundred and ninety four loci (186 RFLPs and 208 SSRs) were used on the 'Arina × Forno' genetic linkage map, and by linkage analysis 380 of these loci mapped to 27 linkage groups with an average marker density of around 8 cM and spanning a total of 3,086 cM. This represents a coverage of approximately 93% of the ITMI map (Paillard et al. 2003).

Statistical analysis

Phenotypic data

Lattice analysis of single environments and analysis of variance across environments were performed with the program PLAB-STAT, Version 2M (Utz 1995) on all evaluated traits. The obtained adjusted entry means from single environments were used to compute the analysis of variance (model I) across environments.

Components of variance were computed considering the effects of the environment and genotype as random. Estimates of variance components σ_G^2 (genetic variance), σ_E^2 (environment variance), $\sigma_{G \times E}^2$ (genotype \times environment interaction variance) and σ_{EIT}^2 (error variance) were calculated. Heritabilities were calculated on an entry mean basis according to Hallauer and Miranda Fo (1981). The distribution of the lines for *Stagonospora glume blotch* resistance was tested for normality using the SAS univariate procedure (SAS Institute 1991). Phenotypic correlation coefficients of *Stagonospora glume blotch* scores between the environments were calculated on an entry mean basis. The area under the disease progress curve (AUDPC) was calculated based on two *Stagonospora glume blotch* scorings per environment (Campbell and Madden 1990; Jeger and Viljanen-Rollinson 2001). Thus, the adjusted entry means for the AUDPC of *Stagonospora glume blotch* per environment were used to estimate the genotypic value across five environments. The estimation of the genotypic values for heading time as well as plant height are based on adjusted entry means from six and seven environments, respectively.

Genetic mapping and QTL analysis

For the construction of the genetic linkage map of the 'Arina \times Forno' cross, linkage analysis was performed using MAPMAKER 3.0b for MS-DOS (Lander et al. 1987) and assuming Haldane's mapping function (Haldane 1919). For a detailed description see Paillard et al. (2003).

QTL analysis was done for all single markers from the 'Arina \times Forno' map by a simple one-way ANOVA using the SAS glm procedure (SAS Institute 1991). Interval QTL analysis was carried out with the composite interval mapping (CIM) program PLABQTL, Version 1.1 (Utz and Melchinger 2000) which is based on multiple regression. Twenty five markers which were linked closer than 0.2 cM were excluded from the QTL analysis to prevent ill-conditioned equation systems and the generation of "synthetic" new markers by the program. In order to determine the significance of a QTL for simple interval mapping (SIM) and CIM, the critical LOD (logarithm of the odds) thresholds were determined executing a permutation test for each trait with 1,000 permutations. For SIM and CIM, the critical LOD thresholds were set to 2.5 and 4.5, respectively, because the individual critical LOD thresholds at a type I error rate of $\alpha=0.25$ (Beavis 1998) for AUDPC and plant height resulted in 2.27 and 2.38 for SIM, respectively, and 4.43 and 4.56 for CIM, respectively. After calculating SIM for each trait, a whole-genome scan with CIM was conducted using the automatic covariate selection statement ('cov select'). Selected covariates were checked individually for too tight linkage or accumulation close to a detected QTL. A five-fold cross-validation run was performed with 80% of the lines used for estimation of the QTLs (positions and effects) and with the remaining 20% used for validation. In all the interval mapping runs, we used the 'model AA' statement estimating only additive effects due to the small heterozygosity of our population. Detected epistatic effects (digenic QTL \times QTL interactions) were added to the additive effects in the model.

Results

Stagonospora glume blotch reaction under natural infestation

The AUDPC (Fig. 1A) of the lines across all tested environments was continuously distributed and differed significantly from normality ($p < 0.0001$). The average *Stagonospora glume blotch* infestation (in the percentage of the infected glume area) across environments was $6.2 \pm 1.7\%$ for the resistant parent 'Arina' and $19.4 \pm 5.0\%$ for 'Forno' (Table 1; $LSD_{5\%}=6.0$). No line exhibited a




Fig. 1A, B Different frequency distributions of *Stagonospora glume blotch* susceptibility for the area under the disease progress curve (AUDPC). **A** Distribution of the average *Stagonospora glume blotch* susceptibility across five environments in the 'Arina \times Forno' population. **B** Phenotypic distribution of lines carrying both resistance alleles (*QSnq.sfr-3BS* plus *QSnq.sfr-4BL*) and their phenotypic average ('R') compared to the phenotypic average of the susceptible allele combination ('S')

completely resistant phenotype and there was no line significantly more resistant than 'Arina' ($LSD_{5\%}=6.0$), whereas transgression for susceptibility occurred frequently (Fig. 1A). *Stagonospora glume blotch* susceptibility correlated to plant height and heading time with $r=-0.52$ ($p < 0.01$) and $r=-0.10$ (n.s.), respectively. The calculated heritabilities (h^2) for plant height, heading time and AUDPC were 0.97, 0.93 and 0.80, respectively. The phenotypic correlation coefficients (Spearman) for the AUDPC between all tested environments ranged from 0.59 to 0.84 ($p < 0.01$) reflecting the relatively high heritability of the trait. The performed ANOVA revealed highly significant ($p < 0.01$) differences of AUDPC among genotypes (G), environments (E) and the G \times E interaction as well as

Table 1 Average natural Stagonospora glume blotch infestation of the parental lines and the 240 phenotyped single seed descent (SSD)-derived lines from the 'Arina × Forno' cross

Environment		Average natural infestation (in % infected glume area)				
		Parental lines		'Arina × Forno' population		
		'Arina'	'Forno'	SSD lines	Min	Max
2000	ZH106	8.1±1.1	23.0±1.0	21.6±10.0	5.5	68.9
	ZH114	5.4±1.6	21.8±2.1	16.3±8.6	2.2	72.8
	Haag00	7.6±1.3	25.2±2.1	21.1±9.1	4.8	61.1
2001	Haag01	5.5±1.0	14.6±1.2	14.6±8.8	3.6	78.9
	Vouvry01	3.9±0.5	12.0±1.0	11.0±6.3	3.1	74.8
	Average	6.2±1.7	19.4±5.0	15.7±6.5	6.0	52.3

Table 2 Detected QTLs for Stagonospora glume blotch resistance. For each QTL the corresponding marker interval, chromosomal location, individual R^2 and LOD value is given

Marker interval	Chr.	2000+2001		2000						2001			
				Haag00		ZH106		ZH114		Haag01		Vouvry01	
		R^2	LOD	R^2	LOD	R^2	LOD	R^2	LOD	R^2	LOD	R^2	LOD
1 gwm389–cfd79c	3BS	31.2	18.0	23.0	12.6	–	–	12.8	6.6	24.9	13.7	18.5	9.8
2 gwm165–glk335	4BL	19.1	11.1	22.6	13.4	8.0	4.5	–	–	20.7	12.0	–	–
3 gwm371–gwm639a	5BL ^{PH b)} HT c)	9.0	4.9	14.4	8.0	9.1	5.0	–	–	–	–	–	–
4 glk317a–gwm639b	5AL ^{PH}	–	–	–	–	15.7	8.9	–	–	–	–	–	–
5 cfa2174b–gwm46	7BS ^{PH}	–	–	–	–	10.5	5.8	–	–	–	–	–	–
6 cfa2134b–gwm131b	3BL ^{PH}	–	–	9.4	5.2	–	–	–	–	–	–	–	–
7 cfd276a–cfa2086	2AL ^{PH}	–	–	–	–	9.1	4.9	–	–	–	–	–	–
Digenic effects,		1×2	4.4**^a	1×2	4.6**	2×7	1.9* ^a	–	–	1×2	11.5**	–	–
QTL × QTL interaction (%)													
Final simultaneous fit for the adjusted R^2 (%) and final LOD		38.9	28.8	40.5	30.2	19.4	15.8	6.2	3.8	42.0	29.5	12.4	7.3

^a * ** significant digenic QTL by QTL interactions are indicated by * $p < 0.05$ and ** $p < 0.01$

^b ^{PH} putative QTL intervals for resistance overlapping with QTL intervals for plant height; see also Table 3

^c ^{HT} putative QTL intervals for resistance overlapping with QTL intervals for heading time; see also Table 3

a significant difference between the averages of the 2 years ($p < 0.1$). We observed a reduced level of disease severity in 2001 compared to 2000 (Table 1). Vouvry01 yielded the lowest disease severity among all year-locations (Table 1). This was most likely due to the usually early dry weather conditions at this particular location which does not favor fungal spread (Scharen 1966).

QTLs for Stagonospora glume blotch resistance

In total, we detected seven QTLs for Stagonospora glume blotch resistance using CIM with a LOD threshold of 4.5 (Table 2). In two cases (on chromosome arms 3BS and 4BL), the QTL intervals were not associated with a QTL for a morphological trait such as plant height or heading time. Therefore, these QTLs were considered as resistance loci and designated according to McIntosh et al. (1998) as *Q_{Sng.sfr-3BS}* and *Q_{Sng.sfr-4BL}*, respectively (Sng = *S. nodorum* glume blotch).

The resistance QTLs *Q_{Sng.sfr-3BS}* and *Q_{Sng.sfr-4BL}* accounted together for 38.9% (adjusted R^2) of the observed phenotypic variance within the population across all tested environments (Table 2). Comparing the single environments, the individual adjusted R^2 values ranged from 6.2 to 42.0%. In two environments (ZH114 and Vouvry01), we found one QTL (always *Q_{Sng.sfr-3BS}*,

Table 2). By contrast, in ZH106 we detected five QTLs for Stagonospora glume blotch resistance (Table 2). From these five QTLs, only *Q_{Sng.sfr-4BL}* was not overlapping with QTL intervals for plant height and/or heading time (Table 2 and Table 3). In Haag00 and Haag01, *Q_{Sng.sfr-3BS}* and *Q_{Sng.sfr-4BL}* were simultaneously detectable with a LOD larger than 4.5, which resulted in the highest explainable phenotypic variance for resistance compared to the other environments (Table 2; 40.5% and 42.0%, respectively). In these two environments we also found high epistatic effects between the two QTLs which explained an additional 4.6% and 11.5%, respectively, of the observed phenotypic variation (Table 2).

The average resistance of the lines carrying the *Q_{Sng.sfr-3BS}* allele plus the *Q_{Sng.sfr-4BL}* allele ('R' = average resistant alleles), compared with those lines which did not have any of these alleles ('S' = average susceptible alleles), was significantly increased by 50%. For the average resistant allele combination ('R'), we found on average 58.3±18.8 for the AUDPC, in comparison to 117.0±43.8 for the lines with the susceptible alleles ('S') (LSD_{5%}=45.8; Fig. 1B).

The most consistent QTL explaining the largest phenotypic variance was *Q_{Sng.sfr-3BS}* and it was localized on the telomeric region of the short arm of chromosome 3B, peaking at *Xgwm389* being 0.4 cM distal from the RFLP

Table 3 Detected QTLs for *Stagonospora glume blotch* resistance. LOD value, is listed. QTLs for plant height (Ht) and heading time (Eet) are also given
 For single environments, each QTL with its corresponding marker interval, peak position, support interval, individual R^2 as well as

Chr.	Detected QTL (parental allele)	Marker interval	QTL peak (cM) ^a	Supp.-interval (cM) ^b	Environment	R^2 (%)	LOD
2AL	<i>QSng.sfr-2AL (Arina)</i> <i>QHt.sfr-2AL (Arina)</i>	cfd276a–cfa2086	188	186–196	ZH106	9.1	4.9
			186	174–190	Haag01	8.6	4.5
			186	180–190	Vouvry01	26.8	15.7
			186	182–188	ZH109	38.7	24.7
			186	180–190	Haag00	21.9	12.7
			188	180–192	Vouvry00	12.8	6.9
			186	182–190	ZH106	21.9	12.7
			186	180–190	ZH114	34.8	21.8
2BS*	<i>QSng.sfr-2BS* (Forno)</i>	*OA102–psr933b	*0	*0–6	Haag00*	*3.6	*7.0
2BL*	<i>QSng.sfr-2BL* (Forno)</i> <i>QHt.sfr-2BL (Forno)</i>	*glk600–psr644b gwm526b–psr644b	*158	*154–162	ZH114*	*7.8	*4.1
			176	170–176	Haag01	15.9	4.8
			176	168–176	Vouvry01	22.5	7.2
			176	168–176	Haag00	23.8	7.9
			172	164–176	Vouvry00	23.3	7.4
			176	166–176	ZH106	14.9	4.7
			*176	*166–176	ZH114*	*13.8	*4.3
3AS*	<i>QSng.sfr-3AS* (Arina)</i>	*cfd79a–gwm369	*0	*0–4	ZH106*	*7.1	*3.5
			*0	*0–4	ZH114*	*8.3	*4.1
3BS	<i>QSng.sfr-3BS (Arina)</i>	gwm389–cfd79c	0	0–2	Haag01	24.9	13.7
			0	0–2	Vouvry01	18.5	9.4
			0	0–2	Haag00	23.0	12.6
			0	0–2	ZH114	12.8	6.6
3BL	<i>QSng.sfr-3BL (Arina)</i> <i>QHt.sfr-3BL (Arina)</i>	cfa2134b–gwm131b cfa2134b–gwm131b	86	78–96	Haag00	9.4	5.2
			78	70–86	ZH109	8.5	4.6
4BL	<i>QSng.sfr-4BL (Forno)</i>	gwm165–glk335	14	12–16	Haag01	20.7	12.0
			16	14–20	Haag00	22.6	13.4
			16	14–20	ZH106	8.0	4.4
			*16	*14–24	Vouvry01*	*7.5	*4.1
			*16	*14–20	ZH114*	*7.0	*3.8
5AL	<i>QSng.sfr-5AL (Arina)</i> <i>QHt.sfr-5AL (Arina)</i>	glk317a–gwm639b glk317a–gwm639b	68	62–72	ZH106	15.7	8.9
			70	64–74	ZH109	27.7	16.5
			68	62–72	Haag00	15.7	8.9
			68	64–72	Vouvry00	15.4	8.6
			68	64–72	ZH106	12.2	6.7
			66	64–70	ZH114	24.3	14.2
5BL	<i>QSng.sfr-5BL (Forno)</i>	gwm371–gwm639a	64	62–64	Haag00	14.4	8.0
			58	50–64	ZH106	9.1	5.0
			62	*60–64	Vouvry01*	*7.5	*4.1
	<i>QHt.sfr-5BL (Forno)</i>	gwm371–gwm639a	62	60–64	Vouvry01	13.0	7.1
			62	60–64	Haag00	9.3	5.1
			62	60–64	ZH114	17.0	9.6
	<i>QEet.sfr-5BL (Forno)</i>	gwm371–gwm639a	64	60–64	Vouvry01	19.7	11.3
			62	60–64	ZH109	9.5	5.2
			64	60–64	Haag00	12.9	7.1
			62	60–64	Vouvry00	18.9	10.6
		64	60–64	ZH106	19.4	10.7	
5DL*	<i>QSng.sfr-5DL* (Arina)</i>	*cfd81–cfd266	*40	*26–52	Haag00*	*7.3	*4.0
6BL*	<i>QSng.sfr-6BL* (Arina)</i> <i>QHt.sfr-6BL* (Arina)</i>	*fba81–psr924 *fba81–psr924	*46	*40–48	ZH114*	*6.9	*3.6
			*44	*36–48	Vouvry01*	*7.9	*4.0
6Dc	<i>QSng.sfr-6Dc (Arina)</i> <i>QHt.sfr-6Dc (Arina)</i>	cfd19–gwm55b gwm55b–cfd47	108	102–110	Haag00	9.2	5.0
			120	118–120	ZH109	22.0	11.8
			118	114–120	Haag00	11.0	5.6
			120	118–120	ZH106	19.2	10.3
			120	118–120	ZH114	15.9	8.3
	<i>QEet.sfr-6Dc (Arina)</i>	gwm55b–cfd47	114	112–116	Vouvry01	29.8	18.4
			116	114–120	ZH109	12.0	6.6
			116	114–118	Vouvry00	22.1	12.5
			116	114–120	ZH106	19.5	10.8
7BS	<i>QSng.sfr-7BS (Forno)</i> <i>QHt.sfr-7BS (Forno)</i>	cfa2174b–gwm46 cfa2174b–gwm46	26	20–30	ZH106	10.5	5.8
			26	20–30	ZH109	9.7	5.2
			22	18–28	ZH114	9.1	4.8

^a Peak of the QTL in cM referring to the chromosome in the A×F map (Paillard et al. 2003)

^b 1-LOD-support-interval providing the approximate QTL position on the chromosome in cM according to the A×F map (Paillard et al. 2003)

* Putative QTL intervals having a LOD between 3.50 and 4.50 are marked with an asterisk

marker *bcd907b* (Fig. 2A). The resistant *QSng.sfr-3BS* allele originated from the resistant parent ‘Arina’ and was significant in four environments ranging in R^2 values from 12.8% to 24.9% (Table 2). Across environments, *QSng.sfr-3BS* was found with a LOD of 18.0 and accounted for 31.2% of the phenotypic variance for Stagonospora glume blotch resistance (Table 2).

The most probable map position for *QSng.sfr-4BL* peaked 1.1 cM distal from the SSR marker *gwm251* and 0.8 cM distal to the RFLP marker *psr914b* (Fig. 2B). *QSng.sfr-4BL* was detected in all environments with a $\text{LOD} > 3.5$ with R^2 values ranging from 7.0 to 22.6%, respectively (Table 3). With a $\text{LOD} > 4.5$, it was still found in three environments (Haag00, ZH106 and Haag01; Table 2). Across environments, this QTL was detected with a LOD of 11.1 and explained 19.1% of the phenotypic variance for Stagonospora glume blotch resistance (Table 2). The resistant *QSng.sfr-4BL* allele was derived from the susceptible parent ‘Forno’.

The weakest detected QTL explaining the lowest phenotypic variance across all tested environments was *QSng.sfr-5BL* (R^2 9.0% and LOD 4.9; Table 2). It was localized on the long arm of chromosome 5B peaking between *Xcfd7b* and *Xgwm639a* (Fig. 2C). The *QSng.sfr-5BL* allele was derived from the susceptible parent ‘Forno’ and was found in three environments with a $\text{LOD} > 3.5$ ranging from 7.5% to 14.4% for the R^2 value (Supplementary table). With a $\text{LOD} > 4.5$, this QTL was detected only in two environments (Haag00 and ZH106; Table 2). Across all environments, *QSng.sfr-5BL* was the only detected Stagonospora glume blotch resistance QTL associated with plant height as well as heading time (Fig. 2C). The explained phenotypic variance for *QHt.sfr-5BL* and *QEet.sfr-5BL* was 11.2% and 11.5%, respectively.

Cross validation for the AUDPC

In the five-fold cross validation run (CIM, $\text{LOD} > 4.5$) *QSng.sfr-3BS* and *QSng.sfr-4BL* were always detected, whereas *QSng.sfr-5BL* only occurred once in split number 2 (Table 4). A similar situation was found for the 100-fold cross validation (CIM, $\text{LOD} > 4.5$). *QSng.sfr-3BS* and *QSng.sfr-4BL* occurred in all 100 splits but with *QSng.sfr-5BL* only in 35 cases (data not shown). The mean phenotypic values (R^2 in %) for the five-fold and 100-fold cross validation did not differ greatly for the calibration (192 lines) with 33.8 and 35.6, respectively, and for the validation (48 lines) with 37.1 and 31.5, respectively (Table 4).

Discussion

Stagonospora glume blotch resistance is quantitatively inherited

It was known from field trials with artificial inoculations of *S. nodorum* that the two Swiss winter bread wheat




Fig. 2A–C Composite interval analysis of QTLs for the AUDPC on various linkage groups. Linkage groups corresponding to chromosome arms 3BS (A), 4BL (B) and 5B (C). LOD curves were calculated with CIM, $\text{LOD} > 4.5$. Each dash on the cM scale represents one cM. The estimated centromere position is indicated by a black arrow

varieties ‘Arina’ and ‘Forno’ differ significantly in their level of resistance (Wicki et al. 1999). This difference was also observed in field trials with our segregating population. The relatively low disease score for ‘Forno’ and the low average score for the lines reflect the

Table 4 Five-fold cross validation (CV) of QTLs for the AUDPC across all environments

Cross validation split (CV)	Calibration for AUDPC (192 lines)			Validation for AUDPC (48 lines)
	Detected QTLs	LOD	Adjusted R^2 (%)	adjusted phenotypic variance explained (%)
1.	3BS 4BL	16.2	30.3	45.2
2.	3BS 4BL 5BL	21.0	36.5	35.6
3.	3BS 4BL	21.1	38.1	25.0
4.	3BS 4BL	16.2	30.4	44.8
5.	3BS 4BL	18.1	33.5	34.8
5-fold CV: mean phenotypic R^2 :			33.8	37.1
100-fold CV: mean phenotypic R^2 :			35.6	31.5

moderate disease pressure under natural conditions. Nevertheless, we found high correlations between environments and high heritabilities (see Results). This clearly reflects that the evaluation of phenotype and experimental design were appropriate.

In our population we observed a continuous phenotypic distribution for *Stagonospora glume blotch* reaction, confirming its quantitative nature involved in an oligo- or polygenic inheritance pattern. None of the lines showed a completely resistant phenotype and alleles from the susceptible parent also contributed to the resistance. These findings confirm several other genetic studies for resistance against this spike disease (Fried and Meister 1987; Bostwick et al. 1993; Wicki et al. 1999).

Chromosomal regions associated with glume blotch resistance

We found two QTLs, *QSng.sfr-3BS* and *QSng.sfr-4BL*, for *Stagonospora glume blotch* resistance which were not associated with plant height or heading time.

The most consistent QTL explaining the largest phenotypic variance ($R^2=31.2\%$) was *QSng.sfr-3BS* which was found in four out of five environments. For this reason, we consider *QSng.sfr-3BS* to be a major resistance QTL for *Stagonospora glume blotch*. Chromosome arm 3BS has not yet been reported to contribute to glume blotch resistance, although the 3BL arm was shown to affect flag leaf and spike resistance to *S. nodorum* (Hu et al. 1996). Moreover, chromosome 3B conferred seedling resistance against *Stagonospora leaf blotch* (Nicholson et al. 1993). *QSng.sfr-3BS* was localized in a highly interesting chromosomal region peaking at *Xgwm389* on chromosome arm 3BS. Recently, a major *Fusarium head blight* (FHB) QTL (*QFhs.ndsu-3B*) from wheat was mapped in different genetic backgrounds at about 8 cM proximal to *Xgwm389* within the marker interval *gwm533–barc147* (Anderson et al. 2001; Zhou et al. 2002). After CIM analyses in our population, *QSng.sfr-3BS* peaked at *Xgwm389* supporting the hypothesis that it is probably distal to *QFhs.ndsu-3B*.

QSng.sfr-4BL shows that the susceptible variety 'Forno' also contributed to glume blotch resistance. Combined with *QSng.sfr-3BS*, these two QTLs reduced glume blotch susceptibility in our population by about

50%. This is in agreement with the observation that the combination of a few genes may be sufficient to confer partial resistance to the *Stagonospora blotch* disease in wheat (Van Ginkel and Rajaram 1999).

QSng.sfr-4BL was localized on the long arm of chromosome 4B. So far, there were no indications for chromosome 4BL to be involved in *Stagonospora blotch* resistance in wheat at all, neither for the leaf nor for the spike. Only 5 cM distal from *Xpsr914* a catalase gene, designated as *Xpsr484(Cat)-4B*, mapped in the consensus map of hexaploid wheat (Gale et al. 1995). Catalases are activated during abiotic or biotic stress responses in plants. They belong to one of the major reactive oxygen species (ROS)-scavenging mechanisms in plant cells which are localized in the peroxisomes and their primary ROS is hydrogen peroxide (H_2O_2) (Willekens et al. 1997; Mittler 2002). Another candidate gene for *QSng.sfr-4BL* adjacent to *Xpsr484(Cat)-4B* is *TaMlo*, the wheat ortholog of the *Mlo* gene in barley. The presence of the *MLO* protein provides cell death protection in responses to biotic and abiotic stresses (Piffanelli et al. 2002). *Mlo* maps to the long arm of chromosome 4H 2.9 cM distal to the catalase, *Cat2*, gene (Skadsen et al. 1995). In wheat, *TaMlo* was recently mapped 1.2 cM distal to *Xcdo1312-4BL* (Elliott et al. 2002). In the ITMI population, it mapped approximately 3 cM proximal to *Xgwm251-4BL*, and *QSng.sfr-4BL* peaked 1.9 cM distal to *Xgwm251-4BL* in our cross. Further high-resolution mapping is required to test whether *QSng.sfr-4BL* maps to *Xpsr484(Cat)-4B* or *TaMlo*.

QTLs for plant height often coincide with QTLs for glume blotch resistance

One out of three detected QTLs for *Stagonospora glume blotch* resistance, *QSng.sfr-5BL*, across all environments simultaneously, overlapped with a QTL for plant height (*QHT.sfr-5BL*) as well as a QTL for heading time (*QEet.sfr-5BL*).

Resistance to *Stagonospora leaf and glume blotch* is genetically tightly linked or pleiotropic to plant height and heading time (Scott et al. 1982; Wicki et al. 1999). In other host-pathogen interactions such as FHB in barley, it was found that QTLs for morphological characters like plant height or heading time influenced resistance and

resulted in overlapping QTL intervals for FHB resistance (Zhu et al. 1999; Ma et al. 2000). We found a high negative phenotypic correlation between plant height and glume blotch susceptibility under natural infection. This correlation is in agreement with the natural epidemiology of *S. nodorum*, as the natural history of disease initially starts from field debris on the ground. Then, during the growing season the fungus successively infests all above-ground plant organs, including glumes as a last resort. Under such conditions, taller plants would have an advantage because their glumes are higher and not easy for the fungus to reach. In this respect, tallness functions as a sort of escape resistance to the fungus. For *QSng.sfr-5BL*, we can conclude that heading time is most likely not affecting resistance due to the low phenotypic correlation. Thus, the resistance effect of *QSng.sfr-5BL* is probably closely linked or pleiotropic to plant height because both LOD-curves peak at the same map location.

Environmental influence on Stagonospora glume blotch resistance

From population genetic studies, *S. nodorum* exhibited a high genetic similarity among natural populations from different geographical origins, suggesting widespread gene flow and the absence of pathogen adaptation to specific wheat varieties (Keller et al. 1997a, b). Thus, the observed environmental differences for Stagonospora glume blotch resistance in our population should not result from pathogen races. In fact, the differences might have occurred from uncontrolled infection conditions under natural infestation. Some environments seemed to favor the association of morphological traits such as plant height and resistance (ZH106) and, therefore, these locations are less valuable in breeding programs. In contrast, the results from both years in Haag confirmed the observation that this particular location is very favorable to select for Stagonospora blotch resistance (Broennimann 1968). Despite these problems, the obtained genotypic values across all environments allowed the detection of three QTLs for Stagonospora glume blotch resistance. These findings were confirmed by the five-fold and 100-fold cross-validation runs which analyzed the influence of genotypic and environmental sampling, and also determined the bias and sampling error of the explained variance. Based on these results, we conclude that at least 30% of the observed phenotypic variance within the population could be explained by the detected resistance QTLs.

Outlook

Considering the labor of selecting for glume blotch resistance in adult plants, we think that MAS will be an excellent tool for breeders to introduce identified major resistance QTLs into elite short-straw breeding material to enhance the level of partial resistance against *S. nodorum*.

As we could identify closely linked molecular markers for two major resistance QTLs, MAS for Stagonospora glume blotch resistance seems to be feasible. Moreover, mapping of additional markers will result in a better genetic resolution and more tightly linked markers for *QSng.sfr-3BS* and *QSng.sfr-4BL*. Successful MAS and cloning of the major resistance QTL *QSng.sfr-3BS* will crucially depend on the generation of new flanking markers on chromosome 3BS.

Acknowledgements We thank B. Senger and P. Streckeisen for their excellent technical assistance, and R. Enderlin for conducting the field trials in Haag, Canton St. Gallen, Switzerland. We are grateful to Prof. Dr. H.F. Utz for providing the computer packages PLABSTAT and PLABQTL. This work was supported by the University of Zürich. The authors declare that the experiments conducted for this publication comply with the current laws of Switzerland.

References

- Anderson JA, Stack RW, Liu S, Waldron BL, Fjeld AD, Coyne C, Moreno-Sevilla B, Fetch JM, Song QJ, Cregan PB, Frohberg RC (2001) DNA markers for Fusarium head blight resistance QTLs in two wheat populations. *Theor Appl Genet* 102:1164–1168
- Auriau P, Rapilly F, Cauderon Y (1988) Monosomic analysis of resistance to *Septoria nodorum* Berk. in soft red wheat [*Triticum aestivum* L. em Thell (Vill) M.K.]. *Agronomie* 8:71–77
- Bathgate JA, Loughman R (2001) Ascospores are a source of inoculum of *Phaeosphaeria nodorum*, *P. avenaria* f. sp. *avenaria* and *Mycosphaerella graminicola* in Western Australia. *Aust Plant Pathol* 30:317–322
- Beavis WD (1998) QTL analyses: power, precision, and accuracy. In: Paterson AH (ed) *Molecular dissection of complex traits*. CRC Press, Boca Raton, New York, pp 145–162
- Bostwick DE, Ohm HW, Shaner G (1993) Inheritance of Septoria glume blotch resistance in wheat. *Crop Sci* 33:439–443
- Broennimann A (1968) Untersuchungen über *Septoria nodorum* Berk. des Weizens. *Mitteil Schweiz Landw* 16:65–76
- Broennimann A (1975) Contributions to genetics of tolerance towards *Septoria nodorum* Berk. in wheat (*Triticum aestivum* L.). *J Plant Breed* 75:138–160
- Broennimann A, Fossati A, Hani F (1973) Spreading of *Septoria nodorum* Berk. and damage to artificially induced mutants of winter wheat cultivar Zenith (*Triticum aestivum* L.). *J Plant Breed* 70:230–245
- Campbell CL, Madden LV (1990) *Introduction to plant disease epidemiology*. John Wiley and Sons, New York, USA
- Cunfer BM, Johnson JW (1999) Soft red winter wheat with resistance to *Stagonospora nodorum* and other foliar pathogens. In: van Ginkel M, McNab A, Krupinsky JM (eds) *Septoria and Stagonospora diseases of cereals: a compilation of global research*. CIMMYT, Mexico, D.F., pp 159
- Cunfer BM, Ueng PP (1999) Taxonomy and identification of *Septoria* and *Stagonospora* species on small-grain cereals. *Annu Rev Phytopathol* 37:267–284
- Duczek LJ, Sutherland KA, Reed SL, Bailey KL, Lanford GP (1999) Survival of leaf spot pathogens on crop residues of wheat and barley in Saskatchewan. *Can J Plant Pathol* 21:165–173
- Elliott C, Zhou FS, Spielmeier W, Panstruga R, Schulze-Lefert P (2002) Functional conservation of wheat and rice *Mlo* orthologs in defense modulation to the powdery mildew fungus. *Mol Plant-Microbe Interact* 15:1069–1077

- Fried PM, Meister E (1987) Inheritance of leaf and head resistance of winter wheat to *Septoria nodorum* in a diallel cross. *Phytopathology* 77:1371–1375
- Gale MD, Atkinson MD, Chinoy CN, Harcourt RL, Jia J, Li QY, Devos KM (1995) Genetic maps of hexaploid wheat. Proc 8th Int Wheat Genetics Symposium, China Agricultural Sciencetech Press, Beijing, China, pp 29–40
- Halama P (2002) Mating relationships between isolates of *Phaeosphaeria nodorum*, (anamorph *Stagonospora nodorum*) from geographical locations. *Eur J Plant Pathol* 108:593–596
- Haldane JBS (1919) The combination of linkage values, and the calculation of distances between loci of linked factors. *J Genet* 8:299–309
- Hallauer AR, Miranda Fo JB (1981) Quantitative genetics in maize breeding. The Iowa University Press, Ames, Iowa, USA
- Hu XY, Bostwick D, Sharma H, Ohm H, Shaner G (1996) Chromosome and chromosomal arm locations of genes for resistance to *Septoria glume blotch* in wheat cultivar Cotipora. *Euphytica* 91:251–257
- Jeger MJ, Viljanen-Rollinson SLH (2001) The use of the area under the disease progress curve (AUDPC) to assess quantitative disease resistance in crop cultivars. *Theor Appl Genet* 102:32–40
- Keller SM, McDermott JM, Pettway RE, Wolfe MS, McDonald BA (1997a) Gene flow and sexual reproduction in the wheat glume blotch pathogen *Phaeosphaeria nodorum* (Anamorph *Stagonospora nodorum*). *Phytopathology* 87:353–358
- Keller SM, Wolfe MS, McDermott JM, McDonald BA (1997b) High genetic similarity among populations of *Phaeosphaeria nodorum* across wheat cultivars and regions in Switzerland. *Phytopathology* 87:1134–1139
- Kleijer G, Bronnimann A, Fossati A (1977) Chromosomal location of a dominant gene for resistance at seedling stage to *Septoria nodorum* Berk. in wheat variety Atlas66. *J Plant Breed* 78:170–173
- Lander ES, Green P, Abrahamson J, Barlow A, Daley MJ, Lincoln SE, Newburg L (1987) MAPMAKER: an interactive computer package for constructing primary genetic linkage maps of experimental and natural populations. *Genomics* 1:174–181
- Laubscher FX, von Wechmar B, von Schalkwyk D (1966) Heritable resistance of wheat varieties to glume blotch (*Septoria nodorum* Berk.). *Phytopathology Z* 56:260–264
- Loughman R, Wilson RE, Goss IM, Foster DT, Murphy NEA (1999) Varieties and advanced lines resistant to *Septoria* diseases of wheat in Western Australia. In: van Ginkel M, McNab A, Krupinsky JM (eds) *Septoria and Stagonospora diseases of cereals: a compilation of global research*. CIMMYT, Mexico, D.F., pp 145–148
- Ma H, Hughes GR (1993) Resistance to *Septoria nodorum* blotch in several *Triticum* species. *Euphytica* 70:151–157
- Ma H, Hughes GR (1995) Genetic control and chromosomal location of *Triticum timopheevii*-derived resistance to *Septoria nodorum* blotch in durum wheat. *Genome* 38:332–338
- Ma ZP, Steffenson BJ, Prom LK, Lapitan NLV (2000) Mapping of quantitative trait loci for *Fusarium* head blight resistance in barley. *Phytopathology* 90:1079–1088
- McDonald BA, Linde C (2002) Pathogen population genetics, evolutionary potential, and durable resistance. *Annu Rev Phytopathol* 40:349–379
- McIntosh RA, Hart GE, Devos KM, Gale MD, Rogers WJ (1998) Catalogue of gene symbols for wheat. Proc 9th Int Wheat Genetics Symposium, Saskatoon, Saskatchewan, University Extension Press, University of Saskatchewan, vol 5
- Meier U (2001) Growth stages of mono- and dicotyledonous plants. 2nd, Federal Biological Research Centre for Agriculture and Forestry (BBA), Braunschweig, Germany
- Mittelstadt A, Fehrmann H (1987) The occurrence of the perfect stage of *Septoria nodorum* in the Federal Republic of Germany. *J Plant Dis Protect* 94:380–385
- Mittler R (2002) Oxidative stress, antioxidants and stress tolerance. *Trends Plant Sci* 7:405–410
- Nelson LR, Gates CE (1982) Genetics of host plant resistance of wheat to *Septoria nodorum*. *Crop Sci* 22:771–773
- Nicholson P, Rezanoor HN, Worland AJ (1993) Chromosomal location of resistance to *Septoria nodorum* in a synthetic hexaploid wheat determined by the study of chromosomal substitution lines in Chinese Spring wheat. *Plant Breed* 110:177–184
- Paillard S, Schnurbusch T, Abderhalden O, Messmer M, Sourdille P, Winzeler M, Keller B, Schachermayr G (2003) An integrative genetic linkage map of winter wheat (*Triticum aestivum* L.). *Theor Appl Genet* (in press)
- Piffanelli P, Zhou FS, Casais C, Orme J, Jarosch B, Schaffrath U, Collins NC, Panstruga R, Schulze-Lefert P (2002) The barley *MLO* modulator of defense and cell death is responsive to biotic and abiotic stress stimuli. *Plant Physiol* 129:1076–1085
- Polley RW, Thomas MR (1991) Surveys of diseases of winter wheat in England and Wales, 1976 to 1988. *Ann Appl Biol* 119:1–20
- SAS Institute (1991) SAS language and procedures: usage 2, version 6, first edition. SAS Institute Inc, Cary, N.C., USA
- Scharen AL (1966) Cyclic production of pycnidia and spores in dead wheat tissue by *Septoria nodorum*. *Phytopathology* 56:580
- Scott PR, Benedikz PW, Cox CJ (1982) A genetic study of the relationship between height, time of ear emergence and resistance to *Septoria nodorum* in wheat. *Plant Pathol* 31:45–60
- Shah D, Bergstrom GC (1993) Assessment of seed-borne *Stagonospora nodorum* in New York soft white winter wheat. *Plant Dis* 77:468–471
- Skadsen RW, Schulze-Lefert P, Herbst JM (1995) Molecular cloning, characterization and expression analysis of two catalase isozyme genes in barley. *Plant Mol Biol* 29:1005–1014
- Utz HF (1995) PLABSTAT—a computer program for statistical analysis of plant breeding experiments. Version 2.0M. Institute of Plant Breeding, Seed Science and Population Genetics, University of Hohenheim, Stuttgart, Germany
- Utz HF, Melchinger AE (2000) PLABQTL: a computer program to map QTL. Version 1.1. Institute of Plant Breeding, Seed Science, and Population Genetics, University of Hohenheim, Stuttgart, Germany
- Van der Plank JE (1963) *Plant diseases: epidemics and control*. Academic Press, New York, USA
- Van Ginkel M, Rajaram S (1999) Breeding for resistance to the *Septoria/Stagonospora* blights of wheat. In: van Ginkel M, McNab A, Krupinsky JM (eds) *Septoria and Stagonospora diseases of cereals: a compilation of global research*. CIMMYT, Mexico, D.F., pp 117–126
- Walag H, Dzieglo A (1985) Monosomic location of genes for resistance to *Septoria nodorum* (Berk.) in the spring wheat (*Triticum aestivum* L.) variety Sappo. *Hodowla Roslin Aklimatyzacja I Nasiennictwo* 29:11–19
- Walag H, Dzieglo A (1987) Location of resistance genes to *Septoria nodorum* (Berk.) of summer wheat of the Weibull 15440 variety (*Triticum aestivum* L.). *Hodowla Roslin Aklimatyzacja I Nasiennictwo* 31:1–6
- Wicki W, Winzeler M, Schmid JE, Stamp P, Messmer M (1999) Inheritance of resistance to leaf and glume blotch caused by *Septoria nodorum* Berk. in winter wheat. *Theor Appl Genet* 99:1265–1272
- Willekens H, Chamnongpol S, Davey M, Schraudner M, Langebartels C, VanMontagu M, Inze D, VanCamp W (1997) Catalase is a sink for H₂O₂ and is indispensable for stress defence in C-3 plants. *Embo J* 16:4806–4816
- Zhou WC, Kolb FL, Bai GH, Shaner G, Domier LL (2002) Genetic analysis of scab resistance QTLs in wheat with microsatellite and AFLP markers. *Genome* 45:719–727
- Zhu H, Gilchrist L, Hayes P, Kleinhofs A, Kudrna D, Liu Z, Prom L, Steffenson B, Toojinda T, Vivar H (1999) Does function follow form? Principal QTLs for *Fusarium* head blight (FHB) resistance are coincident with QTLs for inflorescence traits and plant height in a doubled-haploid population of barley. *Theor Appl Genet* 99:1221–1232