

Sabine Lezar · A. A. Myburg · D. K. Berger ·
M. J. Wingfield · B. D. Wingfield

Development and assessment of microarray-based DNA fingerprinting in *Eucalyptus grandis*

Received: 28 August 2003 / Accepted: 9 June 2004 / Published online: 29 July 2004
© Springer-Verlag 2004

Abstract Development of improved *Eucalyptus* genotypes involves the routine identification of breeding stock and superior clones. Currently, microsatellites and random amplified polymorphic DNA markers are the most widely used DNA-based techniques for fingerprinting of these trees. While these techniques have provided rapid and powerful fingerprinting assays, they are constrained by their reliance on gel or capillary electrophoresis, and therefore, relatively low throughput of fragment analysis. In contrast, recently developed microarray technology holds the promise of parallel analysis of thousands of markers in plant genomes. The aim of this study was to develop a DNA fingerprinting chip for *Eucalyptus grandis* and to investigate its usefulness for fingerprinting of eucalypt trees. A prototype chip was prepared using a partial genomic library from total genomic DNA of 23 *E. grandis* trees, of which 22 were full siblings. A total of 384 cloned genomic fragments were individually amplified and arrayed onto glass slides. DNA fingerprints were obtained for 17 individuals by hybridizing labeled genome representations of the individual trees to the 384-element chip. Polymorphic DNA fragments were identified by evaluating the binary distribution of their background-corrected signal intensities across full-sib individuals. Among 384 DNA fragments on the chip, 104 (27%) were found to be polymorphic. Hybridization of these polymorphic fragments was highly repeatable ($R^2 > 0.91$) within the *E. grandis* individuals, and they allowed us to identify

all 17 full-sib individuals. Our results suggest that DNA microarrays can be used to effectively fingerprint large numbers of closely related *Eucalyptus* trees.

Introduction

Eucalyptus spp. are widely planted as exotics in many tropical and subtropical regions of the world (Eldridge et al. 1993). Because many of these plantations are commonly developed using vegetative propagation, the routine identification of clones and selection of elite genotypes have become increasingly important. Until recently, tree breeders have had to rely on detailed pedigree information and careful labeling to identify individual trees in breeding programs. However, incorrect identification is common and poses a major problem in forestry operations (Keil and Griffin 1994). DNA-based molecular markers have provided a solution to this problem. Several studies have thus shown that individual genotypes can be discriminated using molecular markers (Eppelen et al. 1991; Nybom 1991; Weising et al. 1991).

A variety of molecular-marker techniques can be used for DNA fingerprinting. These techniques include restriction fragment length polymorphisms [(RFLPs) Botstein et al. 1980], simple sequence repeats [(SSRs) Weber and May 1989], random amplified polymorphic DNAs [(RAPDs) Williams et al. 1990], and amplified fragment length polymorphisms [(AFLPs) Vos et al. 1995]. Despite the high throughput afforded by some of these methods, they are all constrained by their dependence on gel electrophoresis. This hampers the processing of a large number of samples or markers in parallel (Smith and Beavis 1996). Furthermore, several of these methods require processing with many independent restriction enzymes or probes to achieve low error rates.

Originally designed for analysis of gene expression, DNA microarrays permit the parallel processing of large numbers of DNA fragments immobilized on a solid-state surface (Schna et al. 1995). To adopt microarray technology for fingerprinting and diversity studies, Jacoud et al. (2001) recently reported the development of Diversity Array Technology (DArT) in rice, while

Communicated by O. Savolainen

S. Lezar · A. A. Myburg · B. D. Wingfield (✉)
Department of Genetics, University of Pretoria,
Pretoria, 0020, South Africa
e-mail: brenda.wingfield@fabi.up.ac.za
Tel.: +27-12-4203946
Fax: +27-12-4203947

D. K. Berger
Department of Botany, University of Pretoria,
Pretoria, 0020, South Africa

M. J. Wingfield
Forestry and Agricultural Biotechnology Institute (FABI),
University of Pretoria,
Pretoria, 0020, South Africa

Borevitz et al. (2003) reported the use of oligonucleotide arrays to detect and genotype single-feature polymorphisms in *Arabidopsis*. No oligonucleotide arrays are available for *Eucalyptus* and therefore, the DARt technique is the only microarray-based genotyping method that would be applicable for these trees. DARt is a solid-state fingerprinting method similar to AFLP and enables analysis of large numbers of marker loci without any DNA sequence information. Microarray-based genotyping as implemented in the DARt technique is a two-dye approach and relies on the detection of DNA fragments in a complex mixture of selectively amplified restriction fragments. Reduction of complexity by selective amplification allows comparison of polymorphic fragments among genotypes. This is achieved by hybridizing DNA to an array containing a large number of DNA fragments, derived from total genomic DNA of an organism. However, plant genomes contain large amounts of highly repetitive DNA sequences, and it is not clear how this feature might affect the rigor of hybridization-based fingerprinting.

The aim of this study was to develop a prototype microarray chip to evaluate the potential of DNA microarrays for fingerprinting closely related *Eucalyptus* clones. In this study, we evaluate the reproducibility of microarray hybridization profiles in *Eucalyptus grandis* and provide recommendations for using this technology in plantation forestry.

Materials and methods

Plant material and DNA extraction

A total of 15 full-sib progeny of *E. grandis* clone ZG14 (Mondi Forests, South Africa) were fingerprinted in this study. Clone ZG14 was used in a controlled cross with *E. grandis* clone TAG-S (Mondi Forests), from which 22 cloned progeny (clones 44D, 32A, 67D, 36E, 31C, 62D, 74C, 53B, 12C, 17C, 10D, 28D, 18C, 60D, 30B, 4D, 13C, 44C, 17D, 74C, 16C, and 56E) were selected for the generation of a genomic representation of the whole full-sib family (described below). Genomic DNA was extracted from tree ZG14 and one ramet of each tree, as described by Murray and Thompson (1980). The second parent tree (TAG-S) was lost during the early stages of this study and plant material was not available for it. A DNA sample was, therefore, obtained from tree TAG-5, a putative sibling relative of TAG-S.

Generation of genome representations

The method used for preparation of genome representations (Fig. 1) was essentially the same as that described by Jaccoud et al. (2001). DNA samples were pooled from 23 trees (144 ng DNA in total from 22 full-sib progeny and parental tree ZG14). The DNA in the pool was digested with 20 U *Pst*I, using buffer H (Roche Diagnostics, Mannheim, Germany) in a reaction volume of 50 μ l. The reactions were incubated at 37°C for 3 h, and the restriction enzyme removed using an equal volume of phenol:chloroform. The DNA fragments were then precipitated with 100% EtOH and 100 mM NaCl. The precipitated DNA was washed with 70% EtOH and resuspended in 20 μ l deionized water to a final concentration of 30 ng/ μ l.

Purified DNA was ligated to *Pst*I-specific adapters (Jaccoud et al. 2001) in a total volume of 30 μ l at 10°C overnight. The ligation mixture consisted of 1 \times ligation buffer, 2 U T₄ DNA ligase (Roche Diagnostics), 10 ng/ μ l BSA (Amersham Biosciences, Piscataway, USA), 1.0 mM ATP (Amersham Biosciences), and 10 μ M *Pst*I adapters. After ligation, 0.2 mM EDTA was added, and the samples




Fig. 1 Schematic representation of the microarray-based genotyping method used in this study. Note that a pooled DNA sample was used to prepare the genome representation that was printed on the array. Also, the length of the selectively amplified restriction fragments determine the number of incorporated dye molecules per fragment, and therefore the average intensity of the corresponding spot on the array

were heated at 70°C for 5 min to inactivate the ligase. The mixture was then diluted to 100 μ l with water, and 2 μ l was used as a template in a subsequent selective PCR reaction.

The PCR was performed in 50 μ l containing 0.8 μ M PCR primer (adapter +T), 0.25 mM of each dNTP, 1 U *Taq* polymerase, and 1 \times reaction buffer (Roche Diagnostics). The PCR amplification consisted of 30 cycles of 94°C for 30 s, 53°C for 45 s, and 72°C for 1 min, with an initial denaturation step of 94°C for 5 min and a final extension step of 72°C for 8 min.

Cloning, PCR amplification, and sequencing of genomic fragments from representations

The amplified products were inserted into a PCR 2.1-TOPO vector, using a T/A cloning kit (Invitrogen, Carlsbad, Calif., USA). After transforming *Escherichia coli* TOP 10F' host cells with ligation products, single colonies were grown overnight at 37°C in LB medium containing 50 μ g/ml ampicillin. Recombinant *E. coli* clones were diluted in 1 vol of 50% glycerol and stored at -80°C. From each culture, 10 μ l was transferred to 10 μ l water and boiled for 10 min to disrupt the cells and release plasmid DNA into the growth medium. A 1- μ l aliquot of this solution was used in a 100- μ l PCR reaction with M13 forward (-20) and M13 reverse primers (Invitrogen). The reaction mix contained 1 \times PCR buffer, 1 U *Taq* polymerase (Roche Diagnostics), 0.25 mM of each dNTP, and 0.4 μ M of each primer. The PCR amplification consisted of 30 cycles of 94°C for 30 s, 53°C for 30 s, and 72°C for 1 min, with an initial denaturation step of 95°C for 5 min and a final extension step of 72°C for 7 min. Aliquots of the PCR products were separated on a 1.4% agarose gel for quality control. The remainder of each sample was then precipitated in 90% ethanol and 0.9 mM NaAc (pH 5.2) to exclude low-molecular-weight fragments. The precipitate was collected by centrifugation at 3,600 g for 30 min. Pellets

were washed in 70% ethanol, dried, and then resuspended in deionized water at ~250 ng/ μ l.

Of the 384 amplified clones, 40 were sequenced. The insert sequences were subjected to similarity searches in GenBank using BLASTN and BLASTX. BLAST alignments were used to estimate the number of repetitive clones in the library that could result in cross-hybridization or uninformative spots on the array.

Array printing and processing

Equal volumes (10 μ l each) of purified PCR product and 100% DMSO were transferred into a 384-well plate (Amersham Pharmacia Biotech). Eight replicates per fragment were arrayed on each slide at 250- μ m spacing onto Vapour Phase Coated Glass Slides (Amersham Pharmacia Biotech), using a Molecular Dynamics Gen III spotter at the African Centre for Gene Technologies (ACGT) Microarray Facility, University of Pretoria, Pretoria, South Africa (<http://fabinet.up.ac.za/microarray>). Following printing, the slides were allowed to dry at 45–50% relative humidity overnight. Spotted DNA was then bound to the slides by UV cross-linking at 250 mJ and baking at 80°C for 2 h.

Preparation of labeled probes

For microarray hybridizations, genome representations from parent tree ZG14 and 15 full-sib progeny were used. We also included tree TAG-5, the putative relative of parent TAG-S. Probe DNA from individual plants was prepared by restriction-enzyme digestion of genomic DNA (144 ng per tree), ligation of restriction fragments to adapters, and subsequent amplification following the protocol described above. Amplicons were precipitated in 1 vol isopropanol to remove excess dNTPs. Labeling of the amplified fragments was carried out using the Klenow fragment of DNA Polymerase I (Roche Diagnostics). Each labeling reaction contained 5 μ g amplified DNA, 1.8 mM dNTP mix (0.3 mM of dATP, dGTP, dCTP each, 0.8 mM of dTTP), 0.1 mM Cy3-dUTP (Amersham Biosciences, Buckinghamshire, UK), 1x hexanucleotide mix (Roche Diagnostics), and 8 U Klenow enzyme (Roche Diagnostics). The reaction was incubated at 37°C overnight. After labeling, the DNA was column-purified (QIAquick PCR Purification Kit, Qiagen, Germany).

Hybridization and washing

Microarray slides were pre-hybridized for 20 min at 60°C in a solution containing 3.5 \times SSC, 0.2% SDS, and 1% BSA (Roche Diagnostics). Slides were rinsed three times in deionized water and dried with N₂ gas. The Cy3-labeled probe was then dissolved in hybridization solution containing 50% formamide (SIGMA), 25% 2 \times hybridization buffer (Amersham Pharmacia Biotech), and 25% deionized water. The mixture was denatured at 92°C for 5 min and quickly cooled on ice. The denatured probe (approximately 35 μ l) was pipetted directly onto the microarray surface and covered with a glass coverslip (24 \times 60 mm, No.1, Marienfeld, Germany). Slides were placed in a custom-made hybridization chamber (N.B. Engineering Works, Pretoria, South Africa) and incubated for 16–18 h in a 42°C water bath.

After hybridization, slides were washed once in 1 \times SSC, 0.2% SDS at 37°C for 4 min, twice in 0.1 \times SSC, 0.2% SDS at 37°C for 4 min, twice in 0.1 \times SSC at room temperature for 1 min, and then rinsed in deionized water for 2 s. Slides were dried using N₂ gas.

Scanning, image processing, and data analysis

Slides were scanned using a GenePix 4000B Scanner (Molecular Dynamics, USA). The mean pixel intensity within each spot and the local background the spot were determined using Array Vision, version 6.0, software (Imaging Research, Molecular Dynamics, USA). All signal intensities were background corrected. Abnormal

spots (e.g., high background, dust, irregularities, etc.) were manually flagged for removal. Anomalous spots not detected through manual inspection were removed if the signal intensity of a spot varied more than 10% from the mean of the eight replicates on each slide. The mean background-corrected spot intensity of the remaining replicates of each DNA fragment was used in subsequent data analyses. The single-dye (Cy3) data were normalized across slides by regression on the spot-intensity data for tree ZG14, which was used as a reference for normalization of all progeny data. The normalized data were then converted into log₂ intensity values.

Identification of polymorphic fragments

Polymorphic DNA fragments were identified in Microsoft Excel, based on the bimodal distribution of their normalized intensity values across slides, consistent with their segregation as dominant PCR-based testcross (*Aa:aa*=1:1) or intercross (*A:aa*=3:1) markers. Relative intensity values were obtained by scaling the signal intensities to that of the DNA fragment with the highest intensity value across slides (set to 1.0). The ranked spot intensities were plotted for each DNA fragment, and identification of DNA fragments with bimodal distribution was based on the presence of two clearly defined intensity classes with mean relative intensity values differing by at least 0.5. A binary scoring table of polymorphic spots was developed for all the *Eucalyptus* trees analyzed. The data for all the polymorphic spots were used to calculate the relative “distances” between the hybridization profiles of individual *Eucalyptus* trees, using Spearman correlation and hierarchical clustering (CLUSTER, available at <http://rana.lbl.gov/>). The clustering results were visualized with TreeView (Eisen et al. 1998).


Verification of DNA polymorphisms

Two of the DNA polymorphisms detected in the array experiment were analyzed by Southern hybridization. *Pst*I-digested total genomic DNA of nine individual trees was resolved on agarose gels and transferred to nylon membranes. Probes representing two of the polymorphic DNA fragments were labeled and hybridized to the *Pst*I-digested DNA on the nylon membranes, using the DIG High Prime DNA Labeling and Detection Starter Kit I (Roche Diagnostics).

Reproducibility of DNA microarray fingerprints

We tested the reproducibility of hybridization profiles, starting from independently prepared genome representations and that of stripping and re-hybridization of the same slides. Repeated stripping and re-hybridization of slides allows for multiple rounds of hybridization on the same slides. To test the reproducibility of the hybridization fingerprints obtained from stripped slides, slides were treated using the protocol of Dolan et al. (2001), with minor modifications. Used slides were immersed four times in stripping buffer (2.5 mM Na₂HPO₄, 0.1% SDS) at 95°C for 25 s. Slides were then washed in deionized water at room temperature for 2 s and dried using N₂ gas. Stripped slides were scanned to verify that all signal had been removed. The stripped slides were then used for a repeat of the same hybridization as before, but with independently labeled DNA. Data analysis was performed as described above. Independent replicates were also prepared from fresh leaf samples of the genome representations of tree ZG14. These genome representations were labeled and hybridized to new slides. Signal-intensity values of the replicate hybridizations were plotted against each other in Microsoft Excel.

Fig. 2a–d Microarray hybridization patterns of two different *Eucalyptus* individuals on the same section of the slide. Each column represents four replicates of the same spot. **a** Hybridization fingerprint of *Eucalyptus* individual 67D and **b** parent ZG14. **c, d** Hybridization of the same individuals on replicate, stripped arrays



Results

DNA microarray analysis

To consider the potential use of microarrays for fingerprinting *Eucalyptus* clones, a prototype DNA microarray chip was constructed with selectively amplified restriction fragments of pooled genomic DNA of an *E. grandis* full-sib family. The technique used to generate a genome representation of the full-sib family and of each *Eucalyptus* tree employs the principle of AFLP (Vos et al. 1995). The complexity of each genomic DNA sample was reduced 16-fold by using +1/+1 selective nucleotides for PCR amplification of genomic restriction fragments. PCR amplicons prepared in this way ranged from 0.2 kb to 1.5 kb, with an average insert size of 700 bp. Sequencing of 40 of the cloned PCR products revealed that there was a low proportion (17%) of “repeat” clones (i.e., clones with microsatellite or other simple repeat sequences, or multiple copies of the same genomic DNA fragment) in the *Eucalyptus* library generated (data not shown).

Proportion of polymorphic fragments useful for fingerprinting

To determine the proportion of polymorphic DNA fragments on the fingerprinting chip, tree ZG14 and 15 full-sib progeny were used in single-dye experiments (Fig. 2). While many of the array features were common (monomorphic) to all individuals (58%), or showed no hybridization signal (15%), many (27%) were clearly polymorphic among individuals. However, only 55 of these spots (15%) were selected for further analyses. The analysis was limited to these 55 spots, because clear threshold values (difference of 0.5 in relative intensity between two intensity classes) could be assigned for them (Fig. 3a), and they were easily convertible into a binary scoring table (results not shown). In contrast, non-polymorphic spots, including both clearly monomorphic loci and loci that were not possible to score as either monomorphic or polymorphic (Fig. 3b), exhibited a greater proportion of high relative-intensity values. This can be attributed to the fact that monomorphic loci share the same signal intensities. Polymorphic spots for which no clear threshold values could be assigned are responsible for the lower relative-intensity values.

The CLUSTER software program allowed us to visualize the relationships of the hybridization profiles using




Fig. 3a, b Examples of signal-intensity distributions of log-transformed hybridization data among 17 *Eucalyptus* individuals. **a** Distribution of relative (normalized) log intensities of four random polymorphic fragments that show a clear bimodal distribution across slides. **b** Non-polymorphic spots show a unimodal distribution

TreeView (Eisen et al. 1998; Fig. 4). The branching orders of duplicate experiments were all identical, and duplicate experiments clustered as nearest neighbors. However, depending on which similarity metric setting was used, the overall branching order varied substantially. Because the Spearman correlation analysis provides a more conservative and reliable estimation of the relationship between hybridization profiles (Murray et al. 2001), this correlation was used for data analysis. The dendrogram generated merely provides a means to visualize the relationship of fingerprints and should not be seen as representative of genetic relationships between the full-sib progeny.




Fig. 4 TreeView (Eisen et al. 1998) representation of relationships of hybridization profiles among 17 *Eucalyptus* individuals based on microarray analysis with the 384-probe array. *Columns* represent hybridization profiles of individuals (or replicates) and *rows* represent the mean log intensities for labeled DNA–DNA hybridizations across individuals. *Red bars* and *green bars* indicate high and low mean log intensity values, *black bars* indicate intermediate values, and *grey bars* show missing data. Nine of the hybridizations were performed in replicate (*rep*). The replicate for ZG14 is a biological replicate, i.e., starting from independently obtained leaf samples of the same tree

All of the hybridization profiles were unique and allowed unambiguous discrimination of the full-sib individuals. The probability of obtaining a particular 55-locus fingerprint is 2.7×10^{-17} , assuming no linkage among polymorphic spots. This provides an upper estimate of the discriminating power of our data. We also randomly selected small subsets of polymorphic DNA fragments and

determined that as few as seven polymorphisms were sufficient to discriminate among full-sib progeny.

Reproducibility

To assess the reproducibility of the experimental procedure, replicate experiments were performed for nine individuals (Figs. 2, 4). Signal intensities of the experimental replicates exhibited regression coefficients (R^2) ranging from 0.90 to 0.93 (Table 1). These are considered to reflect acceptable levels of reproducibility for microarray analysis (Hertzberg et al. 2001). These values were compared to the repeatability of binary scores obtained from the same hybridizations. Repeatability of binary scores of replicate experiments were on average 1.5% higher than regression coefficients of signal intensities.

The regression of the hybridization (normalized signal intensity) data obtained from two different sources of DNA (Fig. 5) for the parent ZG14 revealed a linear R^2 of 0.91. This was not significantly different from the regression coefficient obtained for the experimental replicates of the same tree ($R^2 > 0.93$), suggesting that independent DNA sampling did not introduce much additional experimental variance.

Validation of DNA polymorphisms

Two polymorphic DNA fragments (nos. 227 and 229) were analyzed by Southern hybridization. When probe




Fig. 5 Log plot of the microarray hybridization signals of *Eucalyptus* individual (ZG14). The signal intensity obtained with ZG14 (*x-axis*) was plotted against its biological replicate

Table 1 Repeatability of hybridization profiles and binary scores. Regression coefficient (R^2) values are based on two separate labeling reactions and hybridizations, starting from a single genome representation of each individual

<i>Eucalyptus</i> individual no.	Hybridization profile (R^2) ^a	Repeatability of binary scores ^b
ZG14 (parent tree)	93.53	98.18
ZG14 (parent tree—biological replicate) ^c	91.47	94.55
TAG-5 (relative)	91.72	96.37
18C	92.34	96.37
28C	91.79	94.55
53B	93.95	98.18
36E	92.52	96.37
30B	93.09	96.37
67D	91.86	94.55
74C	93.97	98.18

^aBased on the spot intensities in two replicate experiments of all 384 features on the array

^bProportion of polymorphic probes (55 total) with same binary score across experimental replicates $[1-(\text{number of misscores}/55)] \times 100\%$

^cFor tree ZG14, in addition to a direct experimental replicate, an independently obtained DNA sample and genome representation were used as biological replicate

Table 2 Hybridization patterns of individual restriction fragment length polymorphisms (RFLP) alleles and microarray features. Hybridization patterns were only determined for replicated individuals (see Table 1)

	ZG14	74C	18C	28C	53B	36E	30B	67D	TAG-5
Probe 227									
RFLP allele (300 bp)	–	–	–	–	+	+	–	+	+
Microarray hybridizations	–	–	–	–	+	+	–	+	+
Probe 229									
RFLP allele (350 bp)	–	–	+	+	+	+	+	+	+
Microarray hybridizations	–	–	+	+	+	+	+	+	+

227 of the genomic library was hybridized to a blot of the representations, trees 67D, TAG-S, 36E and 53B produced a band 300 bp in size, while a band of 430 bp was detected for the other genotypes. The genomic Southern blot of probe 229 resulted in a band of 500 bp in the case of trees ZG14, 74C, and a band of 350 bp in size for the other genotypes. These RFLP banding patterns were converted to absence/presence of a band. These RFLPs were consistent with the bimodal-hybridization pattern observed for these two probes in the microarray experiment (Table 2).

Stripping and re-use of slides

Coefficients of determination, which are a measure of the correlation between two variables (experiments), were observed to be higher than 0.90 in replicate hybridization experiments on stripped slides (data included in Table 1). This confirmed that re-used slides resulted in reproducible data. Although the signal intensities decreased on average by 10% after each successive hybridization (Fig. 2), spot-signal intensities remained detectable and were quantifiable and proportionally accurate.

Discussion

In this study, we have shown that microarray technology can be used for genome-wide fingerprinting of closely related *Eucalyptus* trees. Several features of the DNA

microarray technology make it attractive for this purpose. The DNA for hybridization is prepared by selective PCR amplification of short restriction fragments. This means that <250 ng of total genomic DNA provides essentially unlimited starting material for future genotyping of the same trees. This technique—like AFLP analysis—also allows genomic fingerprinting of organisms such as *Eucalyptus* tree species with no prior DNA sequence information (Jaccoud et al. 2001). Most importantly, analysis of the polymorphic fragments is not restricted by the need for gel electrophoresis, and thousands of polymorphic loci in each tree genome can potentially be analyzed in a single assay. Gel electrophoresis, in contrast, is limited in throughput and suffers from difficulties in precisely matching allelic variants of the same size on different gels (Ticknor et al. 2001).

Despite the recent progress that has been made towards the application of microarray technology for DNA fingerprinting and high-throughput genotyping in plants (Jaccoud et al. 2001; Borevitz et al. 2003), cross-hybridization remains a problem. The highly repetitive DNA content of plant genomes undoubtedly results in cross-hybridization of DNA fragments to printed-probe DNA. This increases the overall spot intensity of many probes, and it masks potential polymorphisms. It has been demonstrated that small regions of similarity can lead to cross-hybridization on oligonucleotide microarrays. Kane et al. (2000) found that in 50mer oligonucleotide arrays, cross-hybridization occurred between fragments of relatively low sequence similarity. This has also been observed on microarrays with PCR-based probes (Wren et al. 2002). In general,

cross-hybridization of many different genomic fragments will result in the conversion of polymorphic probes into monomorphic probes. However, a much more serious problem is presented by background segregation of a small number of strongly cross-hybridizing fragments, which will result in mixed hybridization patterns and incorrect marker phenotypes. This problem can be detected at the locus level in segregating progeny, but not in population or fingerprinting studies.

The prototype microarray chip developed in this study for fingerprinting *Eucalyptus* clones allowed for the discrimination among full-sib progeny and thus, very closely related individuals. The hybridization profiles obtained for *E. grandis* individuals were highly repeatable ($R^2 > 0.9$) and allowed us to identify distinct intensity classes (bimodal-intensity distributions) for 55 (14.3%) of the 384 printed probes (Fig. 3). An additional 49 of the probes showed bimodal-intensity distributions, but the overlap between the two intensity classes for these probes was such that it was easy to assign them to presence or absence classes. The total proportion of bimodal probes (27%) and polymorphisms (14.3%) that could be scored was somewhat lower than the rate of polymorphisms often reported for gel-based AFLP markers in outcrossed *Eucalyptus* pedigrees (up to 50%, Myburg et al. 2003). The lower rate of scorable polymorphisms is most probably the result of cross-hybridization obscuring polymorphic features. This is in addition to the "normal" inaccuracies introduced during labeling and hybridization.

In an outcrossed pedigree, the majority of restriction fragment polymorphisms would be expected to segregate in testcross configuration ($Aa:aa=1:1$), while a smaller proportion are expected to segregate in intercross configuration ($AA:Aa:aa=1:2:1$ or $3:1$). The majority of fragments will segregate as testcross fragments because a higher heterozygosity is expected in an outcrossing pedigree. Our pedigree set (15 full sibs) was not sufficiently large to reliably distinguish between intercross and testcross segregation patterns or to determine whether these fragments can be scored in a dosage dependent (co-dominant) fashion on microarrays. Therefore, the bimodal-intensity distribution shown in Fig. 3a probably contains a mixture of testcross and intercross fragments, which may explain the width and height of the plot.

Signal-intensity differences among genotypes can be compared across arrays using either single-dye or two-dye color detection. The DArT technique as described by Jaccoud et al. (2001) represents a two-dye approach. Differences among genotypes (presence or absence of fragments) are detected by comparing the Cy3 signal of each array element to the Cy5 signal of a reference (another genome representation, or a labeled vector fragment). Polymorphic spots show a bimodal distribution of log ratios relative to the reference. The use of a vector-based reference therefore provides an internal standard for each spot and a way to control for differences in the amount of DNA spotted on each array. However, if the same amount of DNA is spotted in each position across arrays, as can be expected for spots printed with the same pin, the value of the reference channel has to be balanced against the additional cost of labeling. We typically do not

observe significant variation in printing across arrays and therefore used a single-dye approach and normalized signal intensities rather than signal ratios. The normalized intensities were used to identify polymorphic spots, based on their bimodal-frequency distribution across individuals.

Reproducibility is essential in genotyping and fingerprinting. We tested for reproducibility of fingerprinting profiles at the experimental and biological level and found that the R^2 of normalized mean signal intensities was always higher than 0.90 in duplicate experiments, even when different sources of genomic DNA were used. The observed variability in signal intensities of 6–9% between replicates of the same individual (in different labeling and hybridization reactions) can be ascribed to variability in the experimental process. Spot variability probably resulted from inaccuracies introduced in labeling, array hybridizations, signal detection and quantification, or low hybridization signal. We also observed a higher frequency of errors at lower signal intensities, due to signals being close to the background noise (Hertzberg et al. 2001). In comparison to the mean signal intensities, the (dominant) binary scores obtained from the same hybridization data were more repeatable (>95%). This was because correct scores could still be obtained when signal intensities varied within signal-intensity classes and due to the low occurrence of spots that varied sufficiently to be erroneously classified. In addition, the repeatability of the hybridization profiles based on the 55 scored polymorphic probes was on average approximately 1.5% higher than that based on the full set of 384 probes (data not shown).

The power of microarray-based fingerprinting lies in its ability to compare different genomes at large numbers of loci in a single assay. In this context, direct comparison of signal-intensity profiles may allow accurate identification of individuals, if proper normalization procedures are followed. Our results suggest that binary scores based on underlying hybridization patterns are only marginally more repeatable than the hybridization data. However, binary (or ideally co-dominant) scores are required to determine allelic frequencies in target populations and in order to calculate probabilities of misidentification for forensic purposes. Binary scores are also required for linkage analysis in mapping pedigrees. Although the use of this technology for linkage mapping still remains to be tested, our results suggest that the technology is useful for rapid genome-wide comparison of closely related germplasm. We found that the branching orders of replicate hybridization fingerprints were all identical and replicate fingerprints all clustered as nearest neighbors. This allowed for the unambiguous identification of *E. grandis* individuals and the identification of two unknown samples (included as blind test samples).

Microarray-based fingerprints may allow the identification of genomic regions shared between related individuals, or identification of genomic regions inherited from specific parents in outcrossed pedigrees. Borevitz et al. (2003) recently demonstrated the use of oligonucleotide probes to demarcate recombination events along chromosomes of recombinant inbred lines of *Arabidopsis*. In our case, map information is not available, but in the future the

internal sequences of probes will be useful to link polymorphisms to a genome sequence when that becomes available for *Eucalyptus*. The clustering of probes into columns according to levels of similarity based on their hybridization (or segregation) patterns across individuals suggests the presence of major linkage groups. This approach may allow ordering of polymorphic markers if the population size is increased adequately.

No dedicated software products are currently available to define hybridization-based DNA fingerprints or to extract binary scores from hybridization data. The majority of available microarray software is designed for two-color expression profiling studies. For single-color fingerprinting applications, such as the one used in this study, the presence or absence of fragments (dominant scoring) or signal intensity (for co-dominant scoring) has to be determined to construct a fingerprint, and quality values need to be assigned to each data point to evaluate the reliability of the combined fingerprint. Kingsley et al. (2001) used the automated peak extraction algorithm to measure spot intensities, and to determine whether a spot is "on" or "off." This algorithm has advantages over the software used in the present study, and should be considered for future work.

The long-term objective of the research presented in this study is to develop a larger array, or set of arrays, with informative probes that can be used for genome-wide fingerprinting of most commercially planted *Eucalyptus* tree species. This will require multiple rounds of selection of polymorphic probes within *E. grandis* and selection of polymorphic probes in other species or interspecific mapping pedigrees. Such an array of polymorphic probes will be useful to saturate existing genetic linkage maps, and may also allow comparative mapping of many eucalypt genomes. Future fingerprinting arrays may be based on oligonucleotides residing in genes (Borevitz et al. 2003) or on genomic restriction fragments such as those cloned in this study.

References

- Borevitz JO, Liang D, Plouffe D, Chang H-S, Zhu T, Weigel D, Berry CC, Winzeler E, Chory J (2003) Large-scale identification of single-feature polymorphisms in complex genomes. *Genome Res* 13:513–523
- Botstein D, White R, Skolnick M, Davis R (1980) Construction of a genetic linkage map in man using restriction fragment length polymorphisms. *Am J Hum Genet* 32:314–331
- Dolan PL, Wu Y, Ista LK, Metznerberg RL, Nelson MA, Lopez GP (2001) Robust and efficient synthetic method for forming DNA microarrays. *Nucleic Acids Res* 29(21):e107
- Eisen MB, Spellmann PT, Brown, PO, Botstein D (1998) Cluster analysis and display of genome-wide expression patterns. *Proc Natl Acad Sci USA* 95:14863–14868
- Eldridge KJ, Davidson J, Harwood C, van Wyk G (1993) *Eucalypt* domestication and breeding. Oxford University Press, Oxford
- Epplen JT, Ammer H, Epplen C, Kammerbauer C, Mitreiter R, Roewer L, Schwaiger W, Steinle V, Zischler H, Albert E, Andreas A, Beuermann B, Meyer W, Buitkamp J, Nanda I, Schmid M, Nuernberg P, Pena SDJ, Poeche H, Sprecher W, Scharl M, Weising K, Yassouridis A (1991) Oligonucleotide fingerprinting using single repeat motifs: a convenient, ubiquitously applicable method to detect hypervariability for multiple purposes. In: Burke T, Dolf G, Jeffreys AJ, Wolff R (eds) *DNA fingerprinting approaches and applications*. Birkhaeuser, Basel, pp 50–69
- Hertzberg M, Sievertzon M, Aspeborg H, Nilsson P, Sandberg G, Lundeberg J (2001) cDNA microarray analysis of small plant tissue samples using a cDNA tag target amplification protocol. *Plant J* 25(4):1–9
- Jaccoud D, Peng K, Feinstein D, Kilian A (2001) Diversity arrays: a solid state technology for sequence independent genotyping. *Nucleic Acids Res* 29(4):e25
- Kane MD et al (2000) Assessment of the sensitivity and specificity of oligonucleotide (50mer) microarrays. *Nucleic Acids Res* 28:4552–4557
- Keil M, Griffin AR (1994) Use of random amplified polymorphic DNA (RAPD) markers in the discrimination and verification of genotypes in *Eucalyptus*. *Theor Appl Genet* 89:442–450
- Kingsley MT, Straub TM, Call DR, Daly DS, Wunschel SC, Chandler DP (2001) Fingerprinting closely related *Xanthomonas* pathovars with random nonamer oligonucleotide microarrays. *Appl Environ Microbiol* 68(12):6361–6370
- Murray MG, Thompson WF (1980) Rapid isolation of high molecular weight plant DNA. *Nucleic Acids Res* 8:4321–4325
- Murray AE, Lies D, Li G, Nealon K, Zhou J, Tiedje JM (2001) DNA/DNA hybridization to microarrays reveals gene-specific differences between closely related microbial genomes. *Proc Natl Acad Sci USA* 98(17):9853–9858
- Myburg AA, Griffin AR, Sederoff RR, Whetten RW (2003) Comparative genetic linkage maps of *E. grandis*, *E. globulus* and their F₁ hybrid based on a double pseudo-backcross mapping approach. *Theor Appl Genet* 107:1028–1042
- Nybom H (1991) Applications of DNA fingerprinting in plant breeding. In: Burke T, Dolf G, Jeffreys AJ, Wolff R (eds) *DNA fingerprinting approaches and applications*. Birkhaeuser, Basel, pp 294–331
- Schena M, Shalon D, Davis RW, Brown PO (1995) Quantitative monitoring of gene expression patterns with a complementary DNA microarray. *Science* 270:467–470
- Smith S, Beavis W (1996) Molecular marker assisted breeding in a company environment. In: Sobral BWS (ed) *The impact of plant molecular genetics*. Birkhaeuser, Boston
- Ticknor LO, Kolsto A-B, Hill KK, Keim P, Laker MT, Tonks M, Jackson PJ (2001) Fluorescent amplified fragment length polymorphism analysis of Norwegian *Bacillus cereus* and *Bacillus thuringiensis* soil isolates. *Appl Environ Microbiol* 67:4863–4873
- Vos P, Hogers R, Bleeker M, Reijans M, van de Lee T, Hornes M, Fritjers A, Pot J, Peleman J, Kuiper M, Zabeau M (1995) AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Res* 23:4407–4414
- Weber J, May P (1989) Abundant class of human DNA polymorphisms which can be typed using the polymerase chain reaction. *Am J Hum Genet* 44:388–396
- Weising K, Ramser J, Kaemmer D, Kahl G, Epplen JT (1991) Oligonucleotide fingerprinting in plant and fungi. In: Burke T, Dolf G, Jeffreys AJ, Wolff R (eds) *DNA fingerprinting approaches and applications*. Birkhaeuser, Basel, pp 312–329
- Williams J, Kubelik A, Livak K, Rafalski J, Tingey S (1990) DNA polymorphisms amplified by arbitrary primers are useful as genetic markers. *Nucleic Acids Res* 18:6531–6535
- Wren JD, Kulkarni A, Joslin J, Butow RA, Garner HR (2002) Cross-hybridization on PCR-spotted microarrays. *IEEE Eng Med Biol* 21:71–75