

Linkage disequilibrium in two European F_2 flint maize populations under modified recurrent full-sib selection

K. C. Falke · H. P. Maurer · A. E. Melchinger ·
H.-P. Piepho · C. Flachenecker · M. Frisch

Published online: 26 May 2007
© Springer-Verlag 2007

Erratum to: Theor Appl Genet
DOI 10.1007/s00122-007-0544-y

Due to a typesetting error, one sentence in the chapter “Assessment of LD in selection cycles” (Materials and methods) is incorrect. The first line of the second paragraph should read: LD between the favorable alleles was assessed by (1) the linkage disequilibrium coefficient D (Eq. 1) and (2) the composite linkage disequilibrium measure Δ (Weir 1996).

The online version of the original article can be found under
doi:[10.1007/s100122-007-0544-y](https://doi.org/10.1007/s100122-007-0544-y).

K. C. Falke · H. P. Maurer · A. E. Melchinger (✉) ·
C. Flachenecker · M. Frisch
Institute of Plant Breeding, Seed Science,
and Population, Genetics,
University of Hohenheim,
70593 Stuttgart, Germany
e-mail: melchinger@uni-hohenheim.de

H.-P. Piepho
Institute for Crop Production and Grassland Research,
University of Hohenheim, 70593 Stuttgart, Germany