

THE DOUBLY PERIODIC SOLUTIONS OF POISSON'S EQUATION
IN TWO INDEPENDENT VARIABLES*

BY

MAX MASON

The only doubly periodic solution † of LAPLACE'S equation

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0$$

is $u = c$, where c is a constant. For if u be such a solution, and v the conjugate potential to u , then $u + iv$ would be a complex analytic function which has a value under a fixed finite limit for all values of x, y . But, as is well known, such a function is necessarily a constant.

It is the object of this paper to investigate, by the use of methods analogous to those of the potential theory, the doubly periodic solutions of POISSON'S equation,

$$(1) \quad \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = f(x, y),$$

where $f(x, y)$ is continuous and periodic in x and in y with the periods a and b respectively. ‡ A "doubly periodic GREEN'S function," G , will be formed from known functions, and the desired solution of (1) found by quadrature from G and f .

* Presented to the Society December 29, 1904. Received for publication November 26, 1904.

† A function u will be called a solution of the differential equation within a region Ω , provided that u satisfies the differential equation at every point within Ω . This definition requires the existence of the second derivatives of u at every point in Ω , and therefore the continuity of the first derivatives. By a doubly periodic solution we shall mean a doubly periodic function which is a solution of the equation in the period rectangle, and therefore in the entire plane. Such a function, in particular, has a value less than a fixed finite number for all values of x, y .

‡ In a recent article (*Journal de Mathématiques*, ser. 5, vol. 10 (1904), p. 445) I have considered by a different method the existence of periodic solutions of the equation

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \lambda A(x, y)u = f(x, y),$$

where λ is a parameter.

§ 1. *A doubly periodic Green's function and its law of reciprocity.*

Let \Re denote the real part of the term before which it is written, and consider the function

$$\Re \log \frac{\sigma(z - \zeta)}{\sigma(z - \gamma)}; \quad z = x + iy, \quad \zeta = \xi + i\eta, \quad \gamma = \alpha + i\beta,$$

where $\sigma(z)$ is the Sigma function of WEIERSTRASS, formed with the periods a, ib ; and ξ, η are two points in the interior of the period rectangle Ω bounded by the lines $x = 0, y = a, x = a, y = b$. This function is a solution of Laplace's equation within Ω , except at the points (ξ, η) and (α, β) , and has the form

$$\log \frac{1}{\sqrt{(x - \xi)^2 + (y - \eta)^2}} - \log \frac{1}{\sqrt{(x - \alpha)^2 + (y - \beta)^2}} + g,$$

where g is a solution of LAPLACE'S equation throughout Ω .

Since for any integers m, n , the function σ obeys the law

$$\sigma(z + ma + inb) = (-1)^{mn+m+n} e^{(m\eta_1+n\eta_3)(2z+ma+inb)} \sigma(z),$$

where η_1, η_3 are certain complex constants, * we have

$$(2) \quad \Re \log \frac{\sigma(z + ma + inb - \zeta)}{\sigma(z + ma + inb - \gamma)} = \Re \log \frac{\sigma(z - \zeta)}{\sigma(z - \gamma)} = m\Re 2\eta_1(\zeta - \gamma) - n\Re 2\eta_3(\zeta - \gamma).$$

Define a real function V by the equation

$$V(x, y, \xi, \eta, \alpha, \beta) = \Re \log \frac{\sigma(z - \zeta)}{\sigma(z - \gamma)} + \frac{x}{a} \Re 2\eta_1(\zeta - \gamma) + \frac{y}{b} \Re 2\eta_3(\zeta - \gamma).$$

Since the last two terms are linear in x and y , V has the form

$$V(x, y, \xi, \eta, \alpha, \beta) = \log \frac{1}{\sqrt{(x - \xi)^2 + (y - \eta)^2}} - \log \frac{1}{\sqrt{(x - \alpha)^2 + (y - \beta)^2}} + S(x, y, \xi, \eta, \alpha, \beta),$$

where S is a known solution of

$$\frac{\partial^2 S}{\partial x^2} + \frac{\partial^2 S}{\partial y^2} = 0$$

within Ω . Furthermore V is doubly periodic in x, y with the periods a, b , since from (2) the equation results:

*See e. g., BURKHARDT, *Elliptische Functionen*, p. 53.

$$\begin{aligned}
 V(x + ma, y + nb, \xi, \eta, a, \beta) &= \Re \log \frac{\sigma(z + ma + inb - \xi)}{\sigma(z + ma + inb - \gamma)} \\
 &\quad + \frac{x + ma}{a} \Re 2\eta_1(\xi - \gamma) + \frac{y + nb}{b} \Re 2\eta_3(\xi - \gamma) \\
 &= \Re \log \frac{\sigma(z - \xi)}{\sigma(z - \gamma)} + \frac{x}{a} \Re 2\eta_1(\xi - \gamma) + \frac{y}{b} \Re 2\eta_3(\xi - \gamma) \\
 &= V(x, y, \xi, \eta, a, \beta)^*.
 \end{aligned}$$

We shall call the function

$$G(x, y, \xi, \eta, a, \beta) = V(x, y, \xi, \eta, a, \beta) - S(a, \beta, \xi, \eta, a, \beta)$$

the doubly periodic Green's function for the periods a, b . This function has the following characteristics:

1°. Except at (ξ, η) and (a, β) , G is, within Ω , a solution of the equation

$$\frac{\partial^2 G}{\partial x^2} + \frac{\partial^2 G}{\partial y^2} = 0.$$

2°. G has the form

$$G = \log \frac{1}{\sqrt{(x - \xi)^2 + (y - \eta)^2}} - \log \frac{1}{\sqrt{(x - a)^2 + (y - \beta)^2}} + R(x, y, \xi, \eta, a, \beta),$$

where R is a known function, which, with respect to the variables x, y , is a solution of LAPLACE'S equation within Ω , and which satisfies for all values of ξ, η in Ω , the equation

$$R(a, \beta, \xi, \eta, a, \beta) = 0.$$

3°. G is doubly periodic in x, y with the periods a, b .

The functions G and R obey the following laws of reciprocity:

$$\begin{aligned}
 G(x, y, \xi, \eta, \alpha, \beta) + \log \frac{1}{\sqrt{(x - \alpha)^2 + (y - \beta)^2}} &= G(\xi, \eta, x, y, \alpha, \beta) \\
 &\quad + \log \frac{1}{\sqrt{(\xi - \alpha)^2 + (\eta - \beta)^2}},
 \end{aligned}$$

$$R(x, y, \xi, \eta, \alpha, \beta) = R(\xi, \eta, x, y, \alpha, \beta).$$

* In the same manner may be formed a doubly periodic function

$$V(x, y, \xi_1, \eta_1, \dots, \xi_n, \eta_n) = \sum_{i=1}^n c_i \log \sqrt{(x - \xi_i)^2 + (y - \eta_i)^2} + S(x, y, \xi_1, \eta_1, \dots, \xi_n, \eta_n)$$

with any number of logarithmic singularities, where S is a solution of LAPLACE'S equation in Ω with respect to the variables x, y , provided that $c_1 + c_2 + \dots + c_n = 0$.

To prove these laws apply GREEN's theorem,

$$\iint (v\Delta u - u\Delta v) dx dy = \int \left(v \frac{\partial u}{\partial n} - u \frac{\partial v}{\partial n} \right) ds,$$

where n is the outward drawn normal, to the region Ω' formed by excluding from Ω the circles $c(\xi, \eta)$, $c(\xi', \eta')$, $c(\alpha, \beta)$ of radius r about the points (ξ, η) , (ξ', η') , (α, β) respectively, and choose

$$u = G(x, y, \xi, \eta, \alpha, \beta), \quad v = G(x, y, \xi', \eta', \alpha, \beta).$$

Since u and v are, within Ω' , solutions of LAPLACE's equation the double integral over Ω' is zero. Furthermore, since u and v are doubly periodic in x, y , each assumes equal values at opposite points of the bounding lines of the rectangle Ω , while at these points the normal derivative of each assumes values numerically equal but opposite in sign. Therefore the line integral over the sides of the rectangle Ω is zero, and we have, replacing ds by $rd\theta$,

$$\begin{aligned} & \int_{c(\xi, \eta)} \frac{1}{r} G(x, y, \xi', \eta', \alpha, \beta) rd\theta - \int_{c(\xi', \eta')} \frac{1}{r} G(x, y, \xi, \eta, \alpha, \beta) rd\theta \\ & + \int_{c(\alpha, \beta)} \frac{1}{r} \{ G(x, y, \xi, \eta, \alpha, \beta) - G(x, y, \xi', \eta', \alpha, \beta) \} rd\theta + h = 0, \end{aligned}$$

where

$$\lim_{r=0} h = 0.$$

But

$$G(x, y, \xi, \eta, \alpha, \beta) - G(x, y, \xi', \eta', \alpha, \beta) =$$

$$\begin{aligned} & \log \frac{1}{\sqrt{(x - \xi)^2 + (y - \eta)^2}} - \log \frac{1}{\sqrt{(x - \xi')^2 + (y - \eta')^2}} \\ & + R(x, y, \xi, \eta, \alpha, \beta) - R(x, y, \xi', \eta', \alpha, \beta), \end{aligned}$$

and

$$R(\alpha, \beta, \xi, \eta, \alpha, \beta) = 0, \quad R(\alpha, \beta, \xi', \eta', \alpha, \beta) = 0.$$

We obtain therefore in the limit $r = 0$, by well known methods,

$$G(\xi, \eta, \xi', \eta', \alpha, \beta) - G(\xi', \eta', \xi, \eta, \alpha, \beta)$$

$$+ \log \frac{1}{\sqrt{(\alpha - \xi)^2 + (\beta - \eta)^2}} - \log \frac{1}{\sqrt{(\alpha - \xi')^2 + (\beta - \eta')^2}} = 0,$$

or writing x, y for ξ', η' ,

$$\begin{aligned} & G(x, y, \xi, \eta, \alpha, \beta) + \log \frac{1}{\sqrt{(x - \alpha)^2 + (y - \beta)^2}} \\ & = G(\xi, \eta, x, y, \alpha, \beta) + \log \frac{1}{\sqrt{(\xi - \alpha)^2 + (\eta - \beta)^2}}. \end{aligned}$$

which is the law of reciprocity for G . If we replace G in this equation by its expression from 2° we have immediately,

$$R(x, y, \xi, \eta, \alpha, \beta) = R(\xi, \eta, x, y, \alpha, \beta).$$

Thus R is symmetrical with respect to x, y and ξ, η and is therefore a solution of

$$\frac{\partial^2 R}{\partial \xi^2} + \frac{\partial^2 R}{\partial \eta^2} = 0$$

within Ω .

§ 2. The doubly periodic solutions of

$$(1) \quad \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = f(x, y).$$

Suppose a doubly periodic solution u of equation (1) exists, for the periods a, b , where f is continuous and doubly periodic with the same periods. If a second solution of the same nature existed, the difference of the two would be a doubly periodic solution of LAPLACE'S equation, and therefore a constant. It follows that a doubly periodic solution of (1) for the periods a, b is uniquely determined if its value at a fixed point is given.

Apply GREEN'S theorem to the period rectangle Ω , choosing for u a doubly periodic solution of (1) and taking $v = 1$. Since the integral over the boundary vanishes, the following equation results:

$$\int_0^b \int_0^a \Delta u \, dx \, dy = \int_0^b \int_0^a f(x, y) \, dx \, dy = 0.$$

This equation is a necessary condition for the existence of a doubly periodic solution of (1). We shall now show that it is also sufficient. Consider the function

$$\begin{aligned} u(\xi, \eta) &= -\frac{1}{2\pi} \int_0^b \int_0^a G(x, y, \xi, \eta, \alpha, \beta) f(x, y) \, dx \, dy \\ &= -\frac{1}{2\pi} \int_0^b \int_0^a \log \frac{1}{\sqrt{(x-\xi)^2 + (y-\eta)^2}} f(x, y) \, dx \, dy \\ &\quad + \frac{1}{2\pi} \int_0^b \int_0^a \log \frac{1}{\sqrt{(x-\alpha)^2 + (y-\beta)^2}} f(x, y) \, dx \, dy \\ &\quad - \frac{1}{2\pi} \int_0^b \int_0^a R(x, y, \xi, \eta, \alpha, \beta) \, dx \, dy. \end{aligned}$$

Since

$$\frac{\partial^2 R}{\partial \xi^2} + \frac{\partial^2 R}{\partial \eta^2} = 0,$$

we see at once, from the potential theory, that $u(\xi, \eta)$ is a solution of

$$\frac{\partial^2 u}{\partial \xi^2} + \frac{\partial^2 u}{\partial \eta^2} = f(\xi, \eta).$$

Furthermore, putting $\xi = \alpha$, $\eta = \beta$ we have

$$u(\alpha, \beta) = 0,$$

since

$$R(x, y, \alpha, \beta, \alpha, \beta) = R(\alpha, \beta, x, y, \alpha, \beta) = 0.$$

From the law of reciprocity for G we have

$$\begin{aligned} & G(x, y, \xi + ma, \eta + nb) - G(x, y, \xi, \eta, \alpha, \beta) \\ &= \log \frac{1}{\sqrt{(\xi + ma - \alpha)^2 + (\eta + nb - \beta)^2}} - \log \frac{1}{\sqrt{(\xi - \alpha)^2 + (\eta - \beta)^2}}, \end{aligned}$$

and therefore

$$\begin{aligned} & u(\xi + ma, \eta + nb) - u(\xi, \eta) \\ &= -\frac{1}{2\pi} \log \sqrt{\frac{(\xi - \alpha)^2 + (\eta - \beta)^2}{(\xi + ma - \alpha)^2 + (\eta + nb - \beta)^2}} \int_0^b \int_0^a f(x, y) dx dy. \end{aligned}$$

Therefore, if

$$\int_0^b \int_0^a f(x, y) dx dy = 0,$$

the function u possesses the periods a , b . We have therefore proved the theorems:

The necessary and sufficient condition for the existence of a doubly periodic solution (periods a , b) of the equation

$$(1) \quad \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = f(x, y),$$

where f is a continuous doubly periodic function with the periods a , b , is that f satisfy the equation

$$\int_0^b \int_0^a f(x, y) dx dy = 0.$$

If this condition is satisfied, then the doubly periodic solution of (1), with periods a , b , which assumes the value C at $x = \alpha$, $y = \beta$ is uniquely determined, and is given by the formula:

$$u(\xi, \eta) = -\frac{1}{2\pi} \int_0^b \int_0^a G(x, y, \xi, \eta, \alpha, \beta) f(x, y) dx dy + C,$$

where G is a known function, expressible in terms of Sigma functions.