

TATE COHOMOLOGY OF PERIODIC K -THEORY WITH REALITY IS TRIVIAL

LISBETH FAJSTRUP

ABSTRACT. We calculate the $RO(\mathbb{Z}/2)$ -graded spectrum for Atiyah's periodic K -theory with reality and the Tate cohomology associated to it. The latter is shown to be trivial.

1. INTRODUCTION

Let KR be the periodic $\mathbb{Z}/2$ -spectrum representing Atiyah's Real K -theory or K -theory with reality [A].

In this note we prove that the generalized Tate spectrum associated to it (in the sense of Greenlees and May [GM]) is trivial. This is quite surprising, since the Tate spectrum for ordinary equivariant K -theory is not contractible for any group G .

The proof is quite simple however; it just relies on the fact that the generator η of $KR(\mathbb{R}P^1)$ is nilpotent. We give a more general statement to emphasize this in 3.1.

It is a pleasure to thank Marcel Bökstedt and John Greenlees, who shortcut my own lengthy calculation proving Theorem 5.2 by pointing out that the essential ingredient was the fact that $\eta^3 = 0$; and the referee who gave many valuable suggestions to make the exposition more readable.

2. G -SPECTRA AND G -COHOMOLOGY THEORIES

In this paper we will be working in the equivariant stable category of [LMS], and we begin by recalling some definitions. We let G be a finite group, since we do not need the Lie group setup of [LMS].

A complete universe U is an infinite dimensional real inner product space with G acting through isometries such that U contains a countably infinite direct sum of regular representations of G as a subspace.

A G -spectrum k_G indexed on a given U associates a based G -space $k_G(V)$ to each finite dimensional G -subspace $V \subset U$ such that for any two G -subspaces V and W of U with $V \subset W$ the usual transitive system of structure maps $k_G(V) \rightarrow \Sigma^{V-W} k_G(W)$ are G -homeomorphisms.

Received by the editors October 12, 1993 and, in revised form, March 24, 1994; originally communicated to the *Proceedings of the AMS* by Thomas Goodwillie.

1991 *Mathematics Subject Classification*. Primary 55N15, 55N20, 55R91; Secondary 55P91, 55P42, 55R25.

Key words and phrases. K -theory with reality, Tate cohomology, G -spectrum.

©1995 American Mathematical Society
0002-9947/95 \$1.00 + \$.25 per page

Let $U = \bigoplus (V_i)^\infty$ for a set of distinct irreducible representations V_i . Then $RO(G, U)$ is the free abelian group generated by the V_i .

Given a G -spectrum k_G indexed on U , we define the associated $RO(G, U)$ -graded homology and cohomology theory:

For any virtual representation $a = V - W$ with V and W in $RO(G, U)$ there are sphere G -spectra $S^a = \sum^{-W} S^V$, and we let

$$k_G^a(X) = [X \wedge S^{-a}, k_G]^G \quad \text{and} \quad k_a^G(X) = [S^a, X \wedge k_G]^G$$

for any G -spectrum X . For a G -space Y , let $k_G^a(Y) = k_G^a(\sum^\infty Y)$ and similarly for homology; here $\sum^\infty Y$ is the $RO(G, U)$ -graded suspension spectrum of Y as in [LMS, p. 14]. These theories have suspension isomorphisms $k_G^V(X) \cong k_G^{V \oplus W}(S^W \wedge X)$ and similarly for homology, induced by the structure maps in the spectrum.

On the other hand, given an $RO(G, U)$ -graded cohomology theory on G -spectra indexed over U (for a definition of this, see [May, Chapter 19], or [LMS, p. 34]), there is a spectrum which classifies this theory as above. Furthermore if the cohomology theory is only defined on G -spaces, it has an extension to G -spectra, which is unique up to nonunique isomorphism, and hence it gives rise to a classifying G -spectrum.

3. THE TATE SPECTRUM FOR SOME $\mathbb{Z}/2$ -SPECTRA

In [GM] a generalized Tate cohomology theory for a G -spectrum k_G is introduced for any compact Lie group G . We repeat this construction for the convenience of the reader. Let X_+ be the disjoint union of the G -space X with a fixed base point and let EG be a contractible free G -space. Let \widetilde{EG} be the unreduced suspension of EG . Then there is a cofibering

$$(1) \quad EG_+ \rightarrow S^0 \rightarrow \widetilde{EG}.$$

Let $F(EG_+, k_G)$ be the function G -spectrum of maps from EG_+ to k_G . The projection $EG_+ \rightarrow S^0$ induces a map of G -spectra

$$\varepsilon: k_G = F(S^0, k_G) \rightarrow F(EG_+, k_G).$$

Smashing ε with the cofibering (1) yields the following map of cofiberings of G -spectra:

$$(2) \quad \begin{array}{ccccc} k_G \wedge EG_+ & \longrightarrow & k_G & \longrightarrow & k_G \wedge \widetilde{EG} \\ \varepsilon \wedge 1 \downarrow & & \varepsilon \downarrow & & \varepsilon \wedge 1 \downarrow \\ F(EG_+, k_G) \wedge EG_+ & \longrightarrow & F(EG_+, k_G) & \longrightarrow & F(EG_+, k_G) \wedge \widetilde{EG} \end{array}$$

The Tate G -spectrum associated to k_G is then the spectrum in the lower right hand corner of this diagram: $t(k_G) = F(EG_+, k_G) \wedge \widetilde{EG}$. The associated homology and cohomology theories are the Tate homology and cohomology of k_G .

When $G = \mathbb{Z}/2$ and k_G is a ring spectrum, the following special case of [GM, §16] gives a method for calculating the upper right-hand corner of (2).

Let m denote the trivial real m -dimensional representation of $\mathbb{Z}/2$ and let $k\xi$ be \mathbb{R}^k with involution given by multiplication by -1 . We allow m and k to be infinite and let \mathbb{R}^∞ have the direct limit topology as usual.

When $G = \mathbb{Z}/2$, our universe will always be $U = \infty\xi \oplus \infty$.

A filtration of $\widetilde{E\mathbb{Z}/2}$ can be obtained from the fact that $B(\infty\xi)/S(\infty\xi)$ is a model for $\widetilde{E\mathbb{Z}/2}$, which implies that the one point compactifications $S^{p\xi}$ give a filtration. We get canonical isomorphisms in homology for any X and for any subspace V of our universe; in cohomology this works for finite X :

$$(3) \quad (k_{\mathbb{Z}/2} \wedge \widetilde{E\mathbb{Z}/2})^V(X) \cong \operatorname{colim}(k_{\mathbb{Z}/2} \wedge S^{p\xi})^V(X) \cong \operatorname{colim}(k_{\mathbb{Z}/2}^{V \oplus p\xi}(X)),$$

$$(4) \quad (k_{\mathbb{Z}/2} \wedge \widetilde{E\mathbb{Z}/2})_V(X) \cong \operatorname{colim}(k_{\mathbb{Z}/2} \wedge S^{p\xi})_V(X) \cong \operatorname{colim}(k_{V-p\xi}^{\mathbb{Z}/2}(X)).$$

Here the maps are induced by the inclusion $S^{p\xi} \rightarrow S^{(p+1)\xi}$. Studying these maps a little closer, we find that the maps giving the latter colimits may also be described as multiplication by a certain element of $k_{\mathbb{Z}/2}^\xi(S^0)$, namely the image of the identity element of $k_{\mathbb{Z}/2}^0(S^0)$ under the isomorphism $k_{\mathbb{Z}/2}^0(S^0) \rightarrow k_{\mathbb{Z}/2}^\xi(S^\xi)$ composed with the map induced by the inclusion of S^0 in S^ξ

$$i^*: k_{\mathbb{Z}/2}^\xi(S^\xi) \rightarrow k_{\mathbb{Z}/2}^\xi(S^0).$$

This element is called the Euler class of ξ and is denoted χ_ξ . This leads to the following lemma:

Lemma 3.1. *Let $k_{\mathbb{Z}/2}^*(-)$ be an $RO(\mathbb{Z}/2, U)$ -graded cohomology theory such that χ_ξ is nilpotent and the classifying spectrum $k_{\mathbb{Z}/2}$ is a ring spectrum. Then the projection p of $E\mathbb{Z}/2_+$ to S^0 induces a weak $\mathbb{Z}/2$ homotopy equivalence $p^*: k_{\mathbb{Z}/2} \rightarrow F(E\mathbb{Z}/2_+, k_{\mathbb{Z}/2})$, and the Tate spectrum $t(k_{\mathbb{Z}/2})$ is trivial.*

Proof. The long exact sequence induced by the cofibration (1) and the fact that $k_{\mathbb{Z}/2}^0(E\mathbb{Z}/2_+ \wedge X) = [X, F(E\mathbb{Z}/2_+, k_{\mathbb{Z}/2})]^{\mathbb{Z}/2}$ implies that for a proof of the first statement it suffices to see that $k_{\mathbb{Z}/2}^*(\widetilde{E\mathbb{Z}/2} \wedge X) = 0$ for X compact. There is a Milnor \lim^1 -exact sequence [M]

$$0 \rightarrow \lim^1 k_{\mathbb{Z}/2}^{*-1}(S^{p\xi} \wedge X) \rightarrow k_{\mathbb{Z}/2}^*(\widetilde{E\mathbb{Z}/2} \wedge X) \rightarrow \lim k_{\mathbb{Z}/2}^*(S^{p\xi} \wedge X) \rightarrow 0.$$

The maps in the inverse system are the same as in (3), namely

$$i^*: k_G^*(S^{k\xi} \wedge X) \rightarrow k_G^*(S^{(k-1)\xi} \wedge X)$$

or equivalently $i^*: k_G^{*-k\xi}(X) \rightarrow k_G^{*-(k-1)\xi}(X)$, and they are multiplication by χ_ξ . Since χ_ξ is nilpotent by assumption, we conclude that the inverse limit and the \lim^1 are both trivial.

For the second part of the lemma, observe that we have just seen that the middle arrow in (2) is a $\mathbb{Z}/2$ -homotopy equivalence; the leftmost arrow is always an equivalence and hence we see that $t(k_{\mathbb{Z}/2})$ is $\mathbb{Z}/2$ -equivalent to $k_{\mathbb{Z}/2} \wedge \widetilde{E\mathbb{Z}/2}$.

From (4) it now follows that the Tate homology groups $t(k_{\mathbb{Z}/2}^{\mathbb{Z}/2})_V(X)$ are trivial for any X , since this is a direct limit over multiplication with the nilpotent element χ_ξ , and hence the spectrum classifying this theory, $t(k_{\mathbb{Z}/2})$, is trivial. \square

4. THE REAL K-THEORY SPECTRUM

In the following we state some facts about the $RO(\mathbb{Z}/2, U)$ -graded spectrum representing Atiyah's Real K -theory.

Definition 4.1 (Atiyah [A]). A Real vector bundle is a complex vector bundle $E \downarrow X$, where (E, τ) and (X, t) are $\mathbb{Z}/2$ spaces such that the involutions commute with the projection map and with τ antilinear, i.e. $\tau(ze) = \bar{z}\tau(e)$, $z \in \mathbb{C}$.

The Real K -theory of (X, t) , $KR(X)$, is then the Grothendieck group of the isometry classes of Real vector bundles over X .

The kernel of the complex dimension map gives reduced KR -theory, $\widetilde{KR}(X)$, and this is the theory which we study here. The dimension map is split, so $KR(X)$ is isomorphic to $\widetilde{KR}(X) \oplus \mathbb{Z}$.

The classifying space for reduced Real K -theory is BU with $\mathbb{Z}/2$ -action given by complex conjugation [tD]. A model for this is the infinite Grassmannian $GR(\mathbb{C}^\infty) = \bigcup GR_n(\mathbb{C}^\infty)$ of complex subspaces of \mathbb{C}^∞ with action induced by complex conjugation on \mathbb{C}^∞ .

An n -dimensional Real bundle is a $(\mathbb{Z}/2 \times_\alpha U(n))$ -bundle in the sense of tom Dieck [tD], where $\alpha: \mathbb{Z}/2 \rightarrow \text{Aut}(U(n))$ is complex conjugation, whereas equivariant K or KO would have trivial α in this description. Another way of stating this is that a Real bundle is a $\mathbb{Z}/2$ -bundle with total group $\Gamma = \mathbb{Z}/2 \times_\alpha U(n)$, structural group $U(n)$ and fiber \mathbb{C} in the sense of [LMS, p. 175].

In [A], Atiyah introduces a $\mathbb{Z} \oplus \mathbb{Z}$ -graded cohomology theory on $\mathbb{Z}/2$ -spaces as follows:

$$KR^{p,q}(X, Y) = KR(X \times B^{p,q}, X \times S^{p,q} \cup Y \times B^{p,q}) = KR(S^{p\xi+q} \wedge X/Y)$$

where $B^{p,q}$ and $S^{p,q}$ are the ball, respectively the sphere of the representation $p\xi + q$, and $KR(X, A) = KR(X/A)$.

This theory has two kinds of periodicity, namely 1-1-periodicity: $KR^{p,q}(X) \cong KR^{p+1,q+1}(X)$, [A, Theorem 2.3], and 8-periodicity:

$$KR^{p,q}(X) \cong KR^{p+8,q}(X)$$

[A, Theorem 3.10]. Thus it can be extended to negative p and q .

As in §3, we work in the complete universe $U = \infty\xi \oplus \infty$. For KR^* to be an $RO(\mathbb{Z}/2, U)$ -graded cohomology theory, we want suspension isomorphisms $KR^V(S^V \wedge X_+) \cong KR(X)$ for any subspace V in U . For this to be true we define

$$KR^V(X_+) = KR^{-p,-q}(X)$$

for $V = p\xi \oplus q$. The reader should beware of the change of signs, since this is quite confusing.

We let KR denote the $\mathbb{Z}/2$ -spectrum representing this theory, and we state some of Atiyah's results in this language.

Let H be the canonical line bundle over \mathbb{CP}^1 and let $b = [H] - 1 \in KR(\mathbb{CP}^1) \cong KR^{-(\xi \oplus 1)}(*)$. The 1-1 periodicity theorem states

Theorem 4.2 (Atiyah). *Multiplication by b induces an equivariant homotopy equivalence between $KR \wedge S^V$ and $KR \wedge S^{V-\xi \oplus 1}$ for $V \subset U$ and $\xi \oplus 1 \subset V$.*

Proof. By the 1-1 periodicity theorem [A, Theorem 2.3], multiplication with b gives an isomorphism between $KR^V(X)$ and $KR^{V-\xi \oplus 1}(X)$. Since this is true for any $\mathbb{Z}/2$ -space, we get the desired $\mathbb{Z}/2$ homotopy equivalence. \square

Now let $\alpha(\lambda)$ be the generator of $KR^{8\xi}(*)$. The 8-periodicity theorem is

Theorem 4.3 (Atiyah). *Multiplication by $\alpha(\lambda)$ induces an equivariant homotopy equivalence between KR and $KR \wedge S^{8\xi}$.*

Proof. The Real periodicity theorem [A, Theorem 3.10] says, that multiplication by $\alpha(\lambda)$ induces an isomorphism between $KR(X)$ and $KR^{8\xi}(X)$ for any X . \square

We state some consequences of the periodicity theorems: For $p \geq q$

$$KR \wedge S^{p\xi \oplus q} \simeq KR \wedge S^{(p-q)\xi}$$

and for any X

$$KR^{p\xi \oplus q}(X) \cong KR^{(p-q)\xi}(X).$$

Now $\sum^{-V}(S^V \wedge KR) \simeq KR$ and $KR \wedge S^V \simeq \sum^{-(\xi \oplus 1)}(KR \wedge S^V)$, so

$$KR \wedge S^{p\xi \oplus q} \simeq \sum^{-(p\xi \oplus p)}(S^{p\xi \oplus q} \wedge KR) \simeq \sum^{-(p-q)}(KR).$$

For $p \leq q$, by the same arguments

$$KR \wedge S^{p\xi \oplus q} \simeq \sum^{-(q-p)\xi}(KR) \simeq KR \wedge S^{q-p}$$

and

$$KR^{p\xi \oplus q}(X) \cong KR^{q-p}(X).$$

The fixed points of KR yield periodic real K -theory, KO , since $KR(Y) = KO(Y)$ for any $\mathbb{Z}/2$ -fixed space Y and nonequivariantly KR is homotopy equivalent to periodic K -theory. Thus Real K -theory provides an example of a non-split G -spectrum, since a splitting [GM, Definition 0.2] is a homotopy equivalence $KU \rightarrow KO \rightarrow KU$ mapping the naive G -spectrum KR considered nonequivariantly (KU) through the fixed point spectrum (KO) and via the inclusion to KU . This can never be a homotopy equivalence, one reason being that $\pi_6(KO)$ is trivial and $\pi_6(KU)$ is not.

5. CALCULATION OF $t(KR)$

We will see that KR fulfills the conditions of 3.1.

Let η be the generator of $KR^\xi(S^0) \cong KR(\mathbb{R}P^1) \cong KO(\mathbb{R}P^1)$, which represents the reduced Hopf bundle over $\mathbb{R}P^1$.

Lemma 5.1. *Let $i: S^0 \rightarrow S^{p\xi}$ be the inclusion. Then the induced map $i^*: KR^*(S^{p\xi}) \rightarrow KR^*(S^0)$ is multiplication by $(-\eta)^p$.*

Proof. In [A, 3.2], Atiyah studied a map induced by the inclusion of the ball $(B(p\xi)_+, +)$ in $(B(p\xi)_+, S(p\xi)_+)$. The induced map in KR -theory is equivalent to the map induced by $i: S^0 \rightarrow S^{p\xi}$, via equivariant homotopy equivalences of the spaces, and Atiyah proved that it is multiplication by $(-\eta)^p$. \square

This is all we need to prove our main theorem.

Theorem 5.2. *The Tate spectrum $t(KR)$ is trivial, and the projection of $E\mathbb{Z}/2_+$ to S^0 induces a $\mathbb{Z}/2$ -homotopy equivalence between KR and $F(E\mathbb{Z}/2_+, KR)$.*

Proof. This will follow from 3.1 once we show that χ_ξ is nilpotent, since KR is a ring spectrum via tensor product of real bundles. Now Lemma 5.1 implies that $\chi_\xi = -\eta$ and $\eta^3 = 0$ by [A]. \square

By the discussion in [GM, §5], this implies that the homotopy orbit spectrum for KR is equivalent to the homotopy fixed point spectrum, which by (5.2) is the fixed point spectrum KO , but we will not go into that here.

REFERENCES

- [A] M. F. Atiyah, *K-theory and reality*, Quart. J. Math. Oxford Ser. (2) **17** (1966), 367–386.
- [tD] T. tom Dieck, *Faserbündel mit Gruppenoperation*, Arch. Math. (Basel) **20** (1969), 136–143.
- [GM] J. P. C. Greenlees and J. P. May, *Generalized Tate, Borel and coBorel cohomology*, Preprint, Univ. of Chicago, 1993.
- [LMS] L. G. Lewis, J. P. May and M. Steinberger, *Equivariant stable homotopy theory*, Lecture Notes in Math., vol. 1213, Springer, 1986.
- [May] J. P. May, *Lecture notes from the NSF/CBMS regional research conference on equivariant homotopy and cohomology theory*, in Fairbanks Alaska, August, 1993.
- [M] J. W. Milnor, *Axiomatic homology theory*, Pacific J. Math. **12** (1962), 337–341.

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, AALBORG UNIVERSITY, FR. BAJERS-
VEJ 7E, 9220 AALBORG ØST, DENMARK
E-mail address: fajstrup@iesd.auc.dk