

HYPONORMALITY AND SPECTRA OF TOEPLITZ OPERATORS

DOUGLAS R. FARENICK AND WOO YOUNG LEE

ABSTRACT. This paper concerns algebraic and spectral properties of Toeplitz operators T_φ , on the Hardy space $H^2(\mathbb{T})$, under certain assumptions concerning the symbols $\varphi \in L^\infty(\mathbb{T})$. Among our algebraic results is a characterisation of normal Toeplitz operators with polynomial symbols, and a characterisation of hyponormal Toeplitz operators with polynomial symbols of a prescribed form. The results on the spectrum are as follows. It is shown that by restricting the spectrum, a set-valued function, to the set of all Toeplitz operators, the spectrum is continuous at T_φ , for each quasicontinuous φ . Secondly, we examine under what conditions a classic theorem of H. Weyl, which has extensions to hyponormal and Toeplitz operators, holds for all analytic functions of a single Toeplitz operator with continuous symbol.

INTRODUCTION

An elegant and useful theorem of C. Cowen [7] characterises the hyponormality of a Toeplitz operator T_φ on the Hardy space $H^2(\mathbb{T})$ of the unit circle $\mathbb{T} \subset \mathbb{C}$ by properties of the symbol $\varphi \in L^\infty(\mathbb{T})$. This result makes it possible to answer an algebraic question coming from operator theory – namely, is T_φ hyponormal? – by studying the function φ itself. In a recent paper [18] of T. Nakazi and K. Takahashi, Cowen's method is carried out to obtain substantial new information about hyponormal Toeplitz operators and their symbols. In the present paper we study the hyponormality of T_φ in the cases where φ is a trigonometric polynomial $\varphi(e^{i\theta}) = \sum_{-m}^N a_n e^{in\theta}$; the goal here is to find conditions on the coefficients a_n that are necessary and sufficient for T_φ to be hyponormal. This problem is still rather complicated in general; however, in §1 we are able to offer necessary and sufficient conditions for the normality and hyponormality of T_φ in the cases where the Fourier coefficients of φ satisfy certain extremal and symmetry properties.

In 1909 H. Weyl examined the spectra of all compact perturbations $A + K$ of a single hermitian operator A and discovered that $\lambda \in \sigma(A + K)$ for every compact operator K if and only if λ is not an isolated eigenvalue of finite multiplicity in $\sigma(A)$. Today this result is known as Weyl's theorem, and it has been extended

Received by the editors May 23, 1995.

1991 *Mathematics Subject Classification*. Primary 47B20, 47B35; Secondary 47A10.

Key words and phrases. Toeplitz operators, hyponormality, spectrum.

The work of the first author is supported in part by a grant from The Natural Sciences and Engineering Research Council of Canada.

The work of the second author is supported in part by The Basic Science Research Institute Program, Ministry of Education, 1995, Project No. BSRI-95-1420, KOSEF 94-0701-02-3 and the GARC-KOSEF, 1995.

from hermitian operators A to hyponormal operators and to Toeplitz operators by L. Coburn [4], and to seminormal operators by S. Berberian [1]. In §3 of this paper we determine properties of continuous functions φ that imply that Weyl's theorem holds for all analytic functions of the Toeplitz operator T_φ . This analysis entails an interesting new fact, which seems to be absent from the literature, concerning the continuity of the spectrum: when restricted to the linear manifold of all Toeplitz operators, the spectrum is a continuous (set-valued) function at every Toeplitz operator T_φ with quasicontinuous symbol φ . In fact, somewhat more general results are true, and these form the basis of our work in §2 of this paper.

Let $L(H)$ and $K(H)$ denote the algebra of bounded linear operators and the ideal of compact operators on a complex Hilbert space H , and let π denote the canonical map $L(H) \rightarrow L(H)/K(H)$. If $T \in L(H)$ is a Fredholm operator, that is if $\pi(T)$ is invertible in $L(H)/K(H)$, then $\ker T$ and $\ker T^*$ are finite-dimensional and the index of T is the integer

$$\text{ind } T = \dim \ker T - \dim \ker T^* .$$

The subset of $\sigma(T)$ that is stable under compact perturbations is denoted by $w(T)$ and is called the Weyl spectrum of T . Those Fredholm operators that have index zero are called Weyl operators. The essential spectrum $\sigma_e(T)$ and the Weyl spectrum $w(T)$ are described succinctly as follows [14], [15]:

$$\begin{aligned} \sigma_e(T) &= \{\lambda \in \mathbb{C} : T - \lambda 1 \text{ is not a Fredholm operator}\}, \\ w(T) &= \{\lambda \in \mathbb{C} : T - \lambda 1 \text{ is not a Weyl operator}\} . \end{aligned}$$

Evidently $\sigma_e(T) \subseteq w(T) \subseteq \sigma(T)$, although unlike σ_e and σ , the Weyl spectrum of T need not satisfy the spectral mapping theorem. The most general result in this direction [12] states that if f is an analytic function on an open set containing $\sigma(T)$, then

$$(0.1) \quad w(f(T)) \subseteq f(w(T)) ;$$

but if T is hyponormal, then [17]

$$(0.2) \quad w(f(T)) = f(w(T)) .$$

It is of interest to know which classes of operators T satisfy (0.2), and for some time it was thought that the Toeplitz operators with continuous symbols may be one of these classes. In §3 we will show that this conjecture is false; conditions on $\varphi \in C(\mathbb{T})$ will then be sought so that $w(f(T_\varphi)) = f(w(T_\varphi))$ for every function f analytic on an open neighbourhood of $\sigma(T_\varphi)$. We arrive at our results in §3 by comparing the spectra of the operators $T_{f \circ \varphi}$ and $f(T_\varphi)$.

We review here a few essential facts concerning Toeplitz operators with continuous symbols that we will need to begin with, using [8]. The Hilbert space $L^2(\mathbb{T})$ has a canonical orthonormal basis given by the trigonometric functions $e_n(z) = z^n$, for all $n \in \mathbb{Z}$, and the Hardy space $H^2(\mathbb{T})$ is the closed linear span of $\{e_n : n = 0, 1, \dots\}$. An element $f \in L^2$ is referred to as analytic if $f \in H^2$ and coanalytic if $f \in L^2 \ominus H^2$. If P denotes the projection operator $L^2 \rightarrow H^2$, then for every $\varphi \in L^\infty(\mathbb{T})$, the operator T_φ on H^2 defined by

$$(0.3) \quad T_\varphi g = P(\varphi g) \quad \text{for all } g \in H^2$$

is called the Toeplitz operator with symbol φ . Every Toeplitz operator has connected spectrum and essential spectrum, and by [4],

$$(0.4) \quad \sigma(T_\varphi) = w(T_\varphi) .$$

The sets $C(\mathbb{T})$ of all continuous complex-valued functions on the unit circle \mathbb{T} and $H^\infty(\mathbb{T}) = L^\infty \cap H^2$ are Banach algebras, and it is well-known that every Toeplitz operator with symbol $\varphi \in H^\infty$ is subnormal. The C^* -algebra \mathfrak{A} generated by all Toeplitz operators T_φ with $\varphi \in C(\mathbb{T})$ has an important property which is very useful for spectral theory: the commutator ideal of \mathfrak{A} is the ideal $K(H^2)$ of compact operators on H^2 . As $C(\mathbb{T})$ and $\mathfrak{A}/K(H^2)$ are $*$ -isomorphic C^* -algebras, then for every $\varphi \in C(\mathbb{T})$,

$$(0.5) \quad T_\varphi \text{ is a Fredholm operator if and only if } \varphi \text{ is invertible,}$$

$$(0.6) \quad \text{ind } T_\varphi = -wn(\varphi) ,$$

$$(0.7) \quad \sigma_e(T_\varphi) = \varphi(\mathbb{T}) ,$$

where $wn(\varphi)$ denotes the winding number of φ with respect to the origin. Finally, we make note that if $\varphi \in C(\mathbb{T})$ and if f is an analytic function defined on an open set containing $\sigma(T_\varphi)$, then $f \circ \varphi \in C(\mathbb{T})$ and $f(T_\varphi)$ is well-defined by the analytic functional calculus.

1. HYONORMALITY OF TOEPLITZ OPERATORS WITH TRIGONOMETRIC POLYNOMIAL SYMBOLS

An operator T is said to be hyponormal if its selfcommutator $[T^*, T] = T^*T - TT^*$ is positive (semidefinite). Normal Toeplitz operators were characterised by a property of their symbol in the early 1960's by A. Brown and P.R. Halmos [3], and so it is somewhat of a surprise that 25 years passed before the exact nature of the relationship between the symbol $\varphi \in L^\infty$ and the positivity of the selfcommutator $[T_\varphi^*, T_\varphi]$ was understood (via Cowen's theorem [7]). As Cowen notes in his survey paper [6], the intensive study of subnormal Toeplitz operators in the 1970's and early 80's is one explanation for the relatively late appearance of the sequel to the Brown-Halmos work. The characterisation of hyponormality in [7] requires one to solve a certain functional equation in the unit ball of H^∞ (see below); in this section we solve this functional equation for trigonometric polynomials φ under certain assumptions about the coefficients of φ . Very recently K. Zhu has studied the same problem from a different point of view [27]; in his work he reformulates Cowen's theorem, in the case of a polynomial φ , so that the hyponormality of T_φ can be decided by applying Schur's algorithm to the Schur function Φ_N . Our results here are more readily applicable, but they apply to only a subclass of all possible trigonometric polynomials φ inducing hyponormal Toeplitz operators. The case of arbitrary trigonometric polynomials φ , though solved in principle by Cowen's theorem or Schur's algorithm, is in practice very complicated. Indeed it may not even be possible to find tractable necessary and sufficient conditions for the hyponormality of T_φ in terms of the Fourier coefficients of a trigonometric polynomial φ unless certain assumptions are made about φ .

For each $\varphi \in L^\infty$ let $\mathcal{E}(\varphi) = \{k \in H^\infty : \|k\|_\infty \leq 1 \text{ and } \varphi - k\bar{\varphi} \in H^\infty\}$. Cowen's theorem can be stated as follows (see [18, Lemma 1]): a Toeplitz operator T_φ is hyponormal if and only if the subset $\mathcal{E}(\varphi)$ of H^∞ is nonempty. Suppose that φ is

the trigonometric polynomial $\varphi(e^{i\theta}) = \sum_{n=-N}^N a_n e^{in\theta}$, where $a_N \neq 0$. If a function $k \in H^\infty$ satisfies $\varphi - k\overline{\varphi} \in H^\infty$, then k necessarily satisfies

$$(1.0) \quad k \sum_{n=1}^N \overline{a_n} e^{-in\theta} - \sum_{n=1}^N a_{-n} e^{-in\theta} \in H^\infty .$$

From (1.0) one computes the Fourier coefficients $\hat{k}(0), \dots, \hat{k}(N-1)$ of k to be $\hat{k}(n) = c_n$, for $n = 0, 1, \dots, N-1$, where c_0, c_1, \dots, c_{N-1} are determined uniquely from the coefficients of φ by the recurrence relation

$$(1.1) \quad \begin{aligned} c_0 &= \frac{a_{-N}}{a_N}, \\ c_n &= (\overline{a_N})^{-1} \left(a_{-N+n} - \sum_{j=0}^{n-1} c_j \overline{a_{N-n+j}} \right), \text{ for } n = 1, \dots, N-1 . \end{aligned}$$

Therefore if $k_1, k_2 \in \mathcal{E}(\varphi)$, then $c_n = \hat{k}_1(n) = \hat{k}_2(n)$ for all $n = 0, 1, \dots, N-1$, and $k_p(z) = \sum_{j=0}^{N-1} c_j z^j$ is the unique (analytic) polynomial of degree less than N satisfying $\varphi - k\overline{\varphi} \in H^\infty$. Conversely, if k_p is the polynomial $k_p(z) = \sum_{j=0}^{N-1} c_j z^j$, where c_0, c_1, \dots, c_{N-1} are determined from the recurrence relation (1.1), then for every integer $n > 0$, the Fourier coefficients $\widehat{\varphi - k\overline{\varphi}}(-n)$ of $\varphi - k\overline{\varphi}$ satisfy

$$\begin{aligned} \widehat{\varphi - k\overline{\varphi}}(-n) &= a_{-n} - \sum_{j=0}^{N-n} c_j \overline{a_{n+j}} \\ &= \left(a_{-n} - \sum_{j=0}^{N-n-1} c_j \overline{a_{n+j}} \right) - c_{N-n} \overline{a_N} \\ &= 0 , \end{aligned}$$

which implies that $\varphi - k_p\overline{\varphi} \in H^2$. But since $\varphi - k_p\overline{\varphi}$ is a polynomial, it follows that $\varphi - k_p\overline{\varphi} \in H^\infty$. However despite the fact that the recurrence relation (1.1) can always be solved uniquely to produce an analytic polynomial k_p satisfying $\varphi - k_p\overline{\varphi} \in H^\infty$, the polynomial k_p need not be contained in the set $\mathcal{E}(\varphi)$, even if $\mathcal{E}(\varphi)$ is known to be nonempty; the problem here is that it is possible for the norm $\|k_p\|_\infty > 1$. Consider, for example, the trigonometric polynomial $\varphi(e^{i\theta}) = e^{-i2\theta} + 2e^{-i\theta} + e^{i\theta} + 2e^{i2\theta}$. Solving the recurrence relation (1.1) produces the polynomial $k_p(z) = \frac{1}{2} + \frac{3}{4}z$, which has norm $\|k_p\|_\infty = \frac{5}{4} > 1$, making k_p ineligible for membership in $\mathcal{E}(\varphi)$. On the other hand, a straightforward calculation shows that the linear fractional transformation

$$b(z) = \frac{z + \frac{1}{2}}{1 + \frac{1}{2}z}$$

satisfies $\varphi - b\overline{\varphi} \in H^\infty$; as b maps the unit circle onto itself, b has norm $\|b\|_\infty = 1$. Thus $b \in \mathcal{E}(\varphi)$ and so T_φ is hyponormal. We note here that the Fourier series of b , namely

$$\begin{aligned} b(e^{i\theta}) &\sim \frac{1}{2} + \frac{3}{4}e^{i\theta} - \frac{3}{2} \sum_{j=2}^\infty \left(\frac{-1}{2}\right)^j e^{ij\theta} \\ &= k_p(e^{i\theta}) + h(e^{i\theta}) , \end{aligned}$$

converges uniformly on \mathbb{T} to b , and that b is a finite Blaschke product. (The existence of a such a Blaschke product in $\mathcal{E}(\varphi)$ is predicted by Theorem 10 of [18].)

The discussion above views the solution c_0, \dots, c_{N-1} to the recurrence relation (1.1) as the Fourier coefficients of every possible candidate k for membership in $\mathcal{E}(\varphi)$. In [27], Zhu applies the Schur functions Φ_N to the N complex numbers c_0, \dots, c_{N-1} to obtain his formulation of Cowen’s theorem (for trigonometric polynomials).

Before continuing further, we record here a condition that φ must necessarily satisfy in order for T_φ to be a hyponormal operator.

A Condition Necessary for Hyponormality. *Suppose that φ is a trigonometric polynomial of the form $\varphi(e^{i\theta}) = \sum_{n=-m}^N a_n e^{in\theta}$, where a_{-m} and a_N are nonzero. If T_φ is hyponormal, then $m \leq N$ and $|a_{-m}| \leq |a_N|$.*

Proof. Proofs that $m \leq N$ can be found in [18, Cor. 5] and [27, Cor. 2]. Let c_0, \dots, c_{N-1} be the solution to the recurrence relation (1.1); because $|a_N| \neq 0$, we have $|c_{N-m}| = |a_{-m}|/|a_N|$. There is a function $k \in \mathcal{E}(\varphi)$ such that $\hat{k}(N - m) = c_{N-m}$; thus $1 \geq \|k\|_\infty \geq |c_{N-m}| = |a_{-m}|/|a_N|$, which implies that $|a_{-m}| \leq |a_N|$. \square

The necessary condition above shows that the cases where $|a_{-m}| = |a_N|$ are, in some sense, extremal among all possibilities for hyponormality. Theorem 1.4 treats such cases, and the result will show that one further feature, namely a symmetry property, is also present.

Proposition 1.1 shows that under strong enough conditions, the polynomial k_p will be an element of $\mathcal{E}(\varphi)$.

Proposition 1.1. *If $\varphi(e^{i\theta}) = \sum_{n=-N}^N a_n e^{in\theta}$, where $a_N \neq 0$, and if $c_0, c_1, \dots, c_{N-1} \in \mathbb{C}$ are obtained from the coefficients of φ by solving the recurrence relation (1.1), then the Toeplitz operator T_φ is hyponormal whenever*

$$(1.1.1) \quad \sum_{j=0}^{N-1} |c_j| \leq 1 .$$

Proof. As we know, the polynomial $k_p(z) = \sum_{j=0}^{N-1} c_j z^j$ satisfies $\varphi - k_p \bar{\varphi} \in H^\infty$. From $\|k_p\|_\infty \leq \sum_{j=0}^{N-1} |c_j| \leq 1$ we have that $k_p \in \mathcal{E}(\varphi)$, and so T_φ is hyponormal. \square

Remark 1.2. If $\varphi(e^{i\theta}) = \sum_{n=-N}^N a_n e^{in\theta}$, where $|a_j| \leq |a_N|$, for all $j = 2, \dots, N - 1$, then from the recurrence relation (1.1) we have that

$$\sum_{j=0}^{N-1} |c_j| \leq |c_0| + |a_N|^{-2} \sum_{n=1}^{N-1} 2^{n-1} |D_n| ,$$

where $D_n = \det \begin{pmatrix} a_{-n} & a_{-N} \\ \bar{a}_n & \bar{a}_N \end{pmatrix}$. Therefore if

$$(1.2.1) \quad \sum_{n=1}^{N-1} 2^{n-1} |D_n| + |a_{-N} a_N| \leq |a_N|^2 ,$$

then by Proposition 1.1, T_φ is hyponormal. Because the left-hand side of (1.2.1) depends on \bar{a}_{-N} and a_N and the right-hand side depends on $|a_N|^2$, it follows that

T_φ is hyponormal whenever $|a_N|$ is sufficiently large. In particular, the Toeplitz operator with symbol $\varphi + \lambda e^{iN\theta}$ is hyponormal whenever $\lambda \in \mathbb{C}$ is such that

$$|\lambda| \geq \sum_{n=1}^{N-1} 2^{n-1}(|a_{-n}| + |a_n|) + |a_{-N}| + |a_N|.$$

Remark 1.3. If $a_{-N} = \dots = a_{-2} = 0$, then the solution to the recurrence relation (1.1) is $c_0 = \dots = c_{N-2} = 0$ and $c_{N-1} = a_{-1}/\overline{a_N}$; thus the analytic polynomial $k_p \in H^\infty$ is $k_p(z) = (a_{-1}/\overline{a_N})z^{N-1}$. Therefore the norm of every $k \in H^\infty$ that satisfies $\varphi - k\overline{\varphi} \in H^\infty$ is such that

$$\|k\|_\infty \geq \left| \frac{a_{-1}}{\overline{a_N}} \right| = \|k_p\|_\infty.$$

Therefore, T_φ is hyponormal if and only if $|a_{-1}| \leq |a_N|$ (which was shown earlier in [11]).

The following theorem and its corollary concern the extremal cases: $|a_{-m}| = |a_N| \neq 0$. Equations (1.4.1) and (1.5.1) below emphasize the symmetry underlying the hyponormality and normality of these operators.

Theorem 1.4. *Suppose that $\varphi(e^{i\theta}) = \sum_{n=-m}^N a_n e^{in\theta}$, where $m \leq N$ and $|a_{-m}| = |a_N| \neq 0$, and let $\mathcal{E}(\varphi) \subset H^\infty$ be the subset of all $k \in H^\infty$ for which $\|k\|_\infty \leq 1$ and $\varphi - k\overline{\varphi} \in H^\infty$. The following statements are equivalent.*

1. *The Toeplitz operator T_φ is hyponormal.*
2. *For all $k = 1, \dots, N - 1$, $\det \begin{pmatrix} a_{-(m-k)} & a_{-m} \\ \overline{a_{(N-k)}} & \overline{a_N} \end{pmatrix} = 0$.*
3. *The following equation in \mathbb{C}^m holds:*

$$(1.4.1) \quad \overline{a_N} \begin{pmatrix} a_{-1} \\ a_{-2} \\ \vdots \\ \vdots \\ a_{-m} \end{pmatrix} = a_{-m} \begin{pmatrix} \overline{a_{N-m+1}} \\ \overline{a_{N-m+2}} \\ \vdots \\ \vdots \\ \overline{a_N} \end{pmatrix}.$$

4. $\mathcal{E}(\varphi) = \{a_{-m}(\overline{a_N})^{-1}z^{N-m}\}$.

Moreover, if T_φ is hyponormal, then the rank of $[T_\varphi^*, T_\varphi]$ is $N - m$.

Proof. Let c_0, \dots, c_{N-1} be the solution to the recurrence relation (1.1); because $|a_{-m}| = |a_N| \neq 0$, we have $|c_{N-m}| = 1$. Note that if $m < N$, then $c_0 = \dots = c_{N-m-1} = 0$.

If a function $k \in H^\infty$ satisfies $\varphi - k\overline{\varphi} \in H^\infty$, then the Fourier series expansion of k is

$$k = \sum_{j=0}^{N-1} c_j e^{ij\theta} + \sum_{n=N}^{\infty} b_n e^{in\theta} \quad \text{for some set of } b_n \in \mathbb{C}.$$

From the fact that $\|k\|_\infty \geq \|k\|_2$ we have that $\|k\|_\infty \geq |c_{N-m}| = 1$; if for some $j > (N - m)$ or $n \geq N$ there is a nonzero Fourier coefficient c_j or b_n of k , then

$$\|k\|_\infty \geq \max \left\{ \sqrt{|c_{N-m}|^2 + |c_j|^2}, \sqrt{|c_{N-m}|^2 + |b_n|^2} \right\} > 1.$$

Thus $\|k\|_\infty = 1$ if and only if c_{N-m} is the only nonzero Fourier coefficient of k . Therefore $\mathcal{E}(\varphi)$ can have at most one element: namely $c_{N-m}z^{N-m}$. Hence, statements (1) and (4) are equivalent. We now proceed to prove the equivalence of statements (1) and (2); obviously (2) and (3) are the exact same statement.

Suppose that T_φ is hyponormal. Then there exists $k \in \mathcal{E}(\varphi)$ and by the discussion above, $k(z) = c_{N-m}z^{N-m}$. Hence, for every $k = 1, \dots, m - 1$,

$$\begin{aligned} 0 = |c_{N-m+k}| &= \left| \frac{1}{a_N} (a_{-(m-k)} - c_{N-m} \overline{a_{N-k}}) \right| \\ &= \left| \frac{1}{a_N} \right|^2 \left| \det \begin{pmatrix} a_{-(m-k)} & a_{-m} \\ a_{(N-k)} & a_N \end{pmatrix} \right|. \end{aligned}$$

Conversely, if $\det \begin{pmatrix} a_{-(m-k)} & a_{-m} \\ a_{(N-k)} & a_N \end{pmatrix} = 0$ for all $k = 1, \dots, N - 1$, then

$$|c_{N-m+1}| = \left| \frac{1}{a_N} (a_{-(m-1)} - c_{N-m} \overline{a_{N-1}}) \right| = \left| \frac{1}{a_N} \right|^2 \left| \det \begin{pmatrix} a_{-(m-1)} & a_{-m} \\ a_{(N-1)} & a_N \end{pmatrix} \right| = 0$$

and hence

$$\begin{aligned} |c_{N-m+2}| &= \left| \frac{1}{a_N} (a_{-(m-2)} - c_{N-m} \overline{a_{N-2}} - c_{N-m+1} \overline{a_{N-1}}) \right| \\ &= \left| \frac{1}{a_N} \right|^2 \left| \det \begin{pmatrix} a_{-(m-2)} & a_{-m} \\ a_{(N-2)} & a_N \end{pmatrix} \right| = 0. \end{aligned}$$

Inductively, we obtain $c_k = 0$ for all $k = 1, \dots, N - m - 1$. As $c_0 = \dots = c_{N-m-1} = 0$ if $m < N$, and $|c_{N-m}| = 1$, we have that the analytic polynomial $k_p(z) = \sum_{j=0}^{N-1} c_j z^j$ is of the form $k_p(z) = c_{N-m}z^{N-m}$ and therefore $k_p \in \mathcal{E}(\varphi)$. This completes the proof that statements (1) and (2) are equivalent.

Lastly, if T_φ is hyponormal, then $\mathcal{E}(\varphi) = \{ \frac{a_{-m}}{a_N} z^{N-m} \}$. Because the selfcommutator $[T_\varphi^*, T_\varphi]$ has finite rank, Theorem 10 of [18] states that there exists a finite Blaschke product $b \in \mathcal{E}(\varphi)$ of degree equal to the rank of $[T_\varphi^*, T_\varphi]$. In our case there is only one element in $\mathcal{E}(\varphi)$: $b(z) = \frac{a_{-m}}{a_N} z^{N-m}$, which is a finite Blaschke product of degree $N - m$. □

The following necessary and sufficient condition for normality is to be expected, given that every normal Toeplitz operator is a translation and rotation of a hermitian Toeplitz operator [3].

Corollary 1.5. *If $\varphi(e^{i\theta}) = \sum_{n=-m}^N a_n e^{in\theta}$, then T_φ is normal if and only if $m = N$, $|a_{-N}| = |a_N|$, and*

$$(1.5.1) \quad \overline{a_N} \begin{pmatrix} a_{-1} \\ a_{-2} \\ \vdots \\ \vdots \\ a_{-N} \end{pmatrix} = a_{-N} \begin{pmatrix} \overline{a_1} \\ \overline{a_2} \\ \vdots \\ \vdots \\ \overline{a_N} \end{pmatrix}.$$

Proof. If $m = N$, $|a_{-m}| = |a_N|$, and $\det \begin{pmatrix} a_{-(m-k)} & a_{-m} \\ a_{(N-k)} & a_N \end{pmatrix} = 0$ for all $k = 1, \dots, N-1$, then by Theorem 1.4, T_φ is hyponormal and $\text{rank} [T_\varphi^*, T_\varphi] = N - m = 0$; that is, T_φ is normal. Conversely, if T_φ is normal, then by the Brown-Halmos theorem [3], there are scalars $\alpha, \beta \in \mathbb{C}$ and a real-valued $\psi \in L^\infty$ such that $T_\varphi = \alpha T_\psi + \beta 1$. As T_ψ is a hermitian Toeplitz operator, the Fourier coefficients of ψ satisfy $\hat{\psi}(n) = \overline{\hat{\psi}(-n)}$ for all n ; in particular $|\alpha| |a_N| = |\hat{\psi}(N)| = |\hat{\psi}(-N)| = |\alpha| |a_{-N}|$, showing that $|a_{-N}| = |a_N|$. Thus, $N = m$ and (1.5.1) holds. \square

Remark 1.6. For trigonometric polynomials φ satisfying the assumptions of Theorem 1.4, the question of whether or not the Toeplitz operator T_φ is hyponormal is completely independent of the values the coefficients a_0, \dots, a_{N-m} of φ . This interesting fact does not appear to be a coincidence, for it is noted as well by Zhu [27] under weaker assumptions.

Example 1.7. Consider the following two trigonometric polynomials:

$$\begin{aligned} \varphi_1(e^{i\theta}) &= e^{-i2\theta} + e^{i3\theta} + e^{i4\theta}, \\ \varphi_2(e^{i\theta}) &= e^{-i2\theta} + e^{-i\theta} + e^{i3\theta} + e^{i4\theta}. \end{aligned}$$

Intuition suggests that φ_2 is less likely than φ_1 to induce a hyponormal Toeplitz operator, as φ_2 is “less analytic” in that the (coanalytic) term $e^{-i\theta}$ is present in φ_2 but not in φ_1 . However the opposite is true: Theorem 1.4 shows that T_{φ_2} is hyponormal (with rank-2 selfcommutator) whereas T_{φ_1} is not.

With the following result, we relax the condition that $|a_{-N}| = |a_N|$, however we retain some symmetry. In the case where $N = 2$ below, the result reduces to Theorem 1 of [10].

Theorem 1.8. Suppose that $\varphi(e^{i\theta}) = \sum_{n=-N}^N a_n e^{in\theta}$, where $N \geq 2$, $|a_N| \neq 0$, and the coefficients of φ satisfy

$$(1.8.1) \quad \overline{a_N} \begin{pmatrix} a_{-2} \\ a_{-3} \\ \vdots \\ \vdots \\ a_{-N} \end{pmatrix} = a_{-N} \begin{pmatrix} \overline{a_2} \\ \overline{a_3} \\ \vdots \\ \vdots \\ \overline{a_N} \end{pmatrix}.$$

Then we have that

$$\begin{aligned} (1.8.2) \quad & |a_N|^2 - |a_{-N}|^2 \geq \left| \det \begin{pmatrix} a_{-1} & a_{-N} \\ \overline{a_1} & \overline{a_N} \end{pmatrix} \right| \\ & \Rightarrow T_\varphi \text{ is hyponormal} \\ & \Rightarrow |a_N|^2 - |a_{-N}|^2 \geq \sqrt{\left| \det \begin{pmatrix} a_{-1} & a_{-N} \\ \overline{a_1} & \overline{a_N} \end{pmatrix} \right|^2 + d^2} - d, \end{aligned}$$

where $d = \frac{1}{2}(1 - |a_{-N}|^2 |a_N|^{-2}) \sum_{n=2}^{N-1} |a_n|^2$ and d is taken to be 0 if $N = 2$.

Proof. To begin with, assume that the first inequality (1.8.2) holds; we are to prove that T_φ is hyponormal. Solving the recurrence relation (1.1) under the condition

(1.8.1) produces the analytic polynomial $k_p(z) = c_0 + c_{N-1}z^{N-1}$, where

$$c_0 = a_{-N}/\overline{a_N} \quad \text{and} \quad c_{N-1} = (\overline{a_N})^{-2} \det \begin{pmatrix} a_{-1} & a_{-N} \\ a_1 & a_N \end{pmatrix}.$$

The first inequality (1.8.2) implies that

$$(1.8.3) \quad 1 - |c_0|^2 \geq |c_{N-1}|.$$

The right-hand side of (1.8.3) is nonnegative, and so $|c_0| \leq 1$. Now if $|c_0| = 1$, then $|c_{N-1}| = 0$ and T_φ is normal; assume, therefore, that $|c_0| < 1$. Let $k \in H^2$ be the function with Fourier series expansion

$$k = k_p(e^{i\theta}) + c_{N-1} \sum_{n=1}^{\infty} (-1)^n \left(\frac{c_{N-1}\overline{c_0}}{|c_{N-1}|} \right)^n e^{i(N-1)(n+1)\theta}.$$

As $\hat{k}(n) = c_n$ for $n = 0, \dots, N - 1$, it remains only to prove that k is in the unit ball of H^∞ . Let $\alpha = \frac{c_{N-1}\overline{c_0}}{|c_{N-1}|}$, which is a complex number of modulus $|\alpha| = |c_0| < 1$. Then

$$\begin{aligned} k(z) &= c_0 + c_{N-1}z^{N-1} + c_{N-1} \sum_{n=1}^{\infty} (-1)^n \left(\frac{c_{N-1}\overline{c_0}}{|c_{N-1}|} \right)^n (z^{N-1})^{n+1} \\ &= \frac{c_{N-1}}{1 - |c_0|^2} \left(\frac{c_0\overline{c_{N-1}}}{|c_{N-1}|} + (1 - |c_0|^2)z^{N-1} \right) + \left(1 - \frac{|c_{N-1}|}{1 - |c_0|^2} \right) c_0 \\ &\quad + \frac{c_{N-1}}{1 - |c_0|^2} \sum_{n=1}^{\infty} \left(-\frac{\overline{c_{N-1}c_0}}{|c_{N-1}|} \right)^n (1 - |c_0|^2)(z^{N-1})^{n+1} \\ &= \frac{c_{N-1}}{1 - |\alpha|^2} \left(-\alpha + \sum_{n=0}^{\infty} \overline{\alpha}^n (1 - |\alpha|^2)(z^{N-1})^{n+1} \right) + \left(1 - \frac{|c_{N-1}|}{1 - |\alpha|^2} \right) c_0 \\ &= \frac{c_{N-1}}{1 - |\alpha|^2} \left(\frac{z^{N-1} - \alpha}{1 - \overline{\alpha}z^{N-1}} \right) + \left(1 - \frac{|c_{N-1}|}{1 - |\alpha|^2} \right) c_0. \end{aligned}$$

Because the function $w \mapsto (w - \alpha)(1 - \overline{\alpha}w)^{-1}$ is a linear fractional transformation, mapping \mathbb{T} onto itself, we obtain the estimate

$$\begin{aligned} \|k\|_\infty &\leq \frac{|c_{N-1}|}{1 - |\alpha|^2} \left| \frac{z^{N-1} - \alpha}{1 - \overline{\alpha}z^{N-1}} \right| + \left(1 - \frac{|c_{N-1}|}{1 - |\alpha|^2} \right) |c_0| \\ &\leq \frac{|c_{N-1}|}{1 - |\alpha|^2} + \left(1 - \frac{|c_{N-1}|}{1 - |\alpha|^2} \right) = 1, \end{aligned}$$

which proves that $k \in \mathcal{E}(\varphi)$.

Suppose now that T_φ is hyponormal. With respect to the orthonormal basis $\{z^n : n = 0, 1, \dots\}$ of H^2 , the selfcommutator of T_φ is a matrix with (μ, ν) -entry given by

$$\alpha_{\mu\nu} = \sum_{j=0}^{\infty} (\overline{a_{j-\mu}}a_{j-\nu} - a_{\mu-j}\overline{a_{\nu-j}}), \quad \text{where } \mu, \nu = 0, 1, 2, \dots$$

Thus, in particular,

$$\begin{aligned} \alpha_{00} &= \sum_{n=1}^N (|a_n|^2 - |a_{-n}|^2), \\ \alpha_{N-1\ N-1} &= |a_N|^2 - |a_{-N}|^2, \\ \alpha_{0\ N-1} &= \overline{\alpha_{N-1\ 0}} = \overline{a_N}a_1 - a_{-N}\overline{a_{-1}}. \end{aligned}$$

The operator $[T_\varphi^*, T_\varphi]$ is positive and, therefore, so is its 2×2 principal submatrix

$$\begin{pmatrix} \alpha_{00} & \alpha_{0\ N-1} \\ \alpha_{N-1\ 0} & \alpha_{N-1\ N-1} \end{pmatrix}.$$

Hence α_{00} and $\alpha_{N-1\ N-1}$ are nonnegative and

$$\begin{aligned} 0 &\leq \det \begin{pmatrix} \alpha_{00} & \alpha_{0\ N-1} \\ \alpha_{N-1\ 0} & \alpha_{N-1\ N-1} \end{pmatrix} = \alpha_{00}\alpha_{N-1\ N-1} - |\alpha_{0\ N-1}|^2 \\ &= \left(\sum_{n=1}^N (|a_n|^2 - |a_{-n}|^2) \right) (|a_N|^2 - |a_{-N}|^2) - |\overline{a_N}a_1 - a_{-N}\overline{a_{-1}}|^2. \end{aligned}$$

The symmetry condition (1.8.1) yields $|a_{-n}| = |a_{-N}/a_N| |a_n|$ for $n = 2, \dots, N - 1$. Direct computation reveals that

$$(|a_1|^2 - |a_{-1}|^2)(|a_N|^2 - |a_{-N}|^2) + |\overline{a_N}a_{-1} - a_{-N}\overline{a_1}|^2 = |\overline{a_N}a_1 - a_{-N}\overline{a_{-1}}|^2,$$

and so

$$\begin{aligned} 0 &\leq \det \begin{pmatrix} \alpha_{00} & \alpha_{0\ N-1} \\ \alpha_{N-1\ 0} & \alpha_{N-1\ N-1} \end{pmatrix} \\ &\leq (|a_N|^2 - |a_{-N}|^2)^2 + (|a_1|^2 - |a_{-1}|^2)(|a_N|^2 - |a_{-N}|^2) \\ &\quad - |\overline{a_N}a_1 - a_{-N}\overline{a_{-1}}|^2 + (|a_N|^2 - |a_{-N}|^2) \sum_{n=2}^{N-1} (|a_n|^2 - |a_{-n}|^2) \\ &= (|a_N|^2 - |a_{-N}|^2)^2 - |\overline{a_N}a_{-1} - a_{-N}\overline{a_1}|^2 \\ &\quad + (|a_N|^2 - |a_{-N}|^2) \left(1 - \left| \frac{a_{-N}}{a_N} \right|^2 \right) \sum_{n=2}^{N-1} |a_n|^2. \end{aligned}$$

Therefore,

$$|a_N|^2 - |a_{-N}|^2 \geq \sqrt{\left| \det \begin{pmatrix} a_{-1} & a_{-N} \\ \overline{a_1} & \overline{a_N} \end{pmatrix} \right|^2 + d^2} - d,$$

where $d = \frac{1}{2}(1 - |a_{-N}|^2|a_N|^{-2}) \sum_{n=2}^{N-1} |a_n|^2$. □

Theorem 1.8 can be applied to show that the Toeplitz operator with symbol

$$\varphi(e^{i\theta}) = e^{-i5\theta} - e^{-i4\theta} + e^{-i2\theta} + e^{-i\theta} + 2e^{i2\theta} - 2e^{i4\theta} + 2e^{i5\theta},$$

whose coefficients satisfy the symmetric relation (1.8.1), but for which there is no symmetry involving a_{-1} and a_1 , is hyponormal. However with full symmetry, meaning that a_{-1} and a_1 are related as well, we obtain the following interesting necessary and sufficient condition for hyponormality, which is, in some sense, dual to Theorem 1.4 but comparable to Corollary 1.5. The result is a generalisation of the fact that $U^* + \lambda U$, where U is the unilateral shift operator, is hyponormal if and only if $|\lambda| \geq 1$.

Corollary 1.9. *If $\varphi(e^{i\theta}) = \sum_{n=-N}^N a_n e^{in\theta}$ is such that*

$$(1.9.1) \quad \overline{a_N} \begin{pmatrix} a_{-1} \\ a_{-2} \\ \vdots \\ \vdots \\ a_{-N} \end{pmatrix} = a_{-N} \begin{pmatrix} \overline{a_1} \\ \overline{a_2} \\ \vdots \\ \vdots \\ \overline{a_N} \end{pmatrix},$$

then T_φ is hyponormal if and only if $|a_{-N}| \leq |a_N|$.

2. SPECTRAL VARIATION WITHIN THE MANIFOLD \mathfrak{T}
OF TOEPLITZ OPERATORS

Let \mathcal{K} denote the set, equipped with the Hausdorff metric, of all compact subsets of \mathbb{C} . The spectrum can be viewed as a function $\sigma : L(H) \rightarrow \mathcal{K}$, mapping each operator T to its spectrum $\sigma(T)$. It is well-known that the function σ is upper-semicontinuous and that σ does have points of discontinuity. Of interest, therefore, is the identification of points of spectral continuity, as in [5], and of classes \mathfrak{C} of operators for which σ becomes continuous when restricted to \mathfrak{C} . Perhaps the most accessible result in the latter direction is the one of J. Newburgh [19]: when restricted to the set of normal operators, σ is a continuous function. As noted in Solution 104 of [13], Newburgh’s argument uses the fact that normal operators have normal resolvents and that normal operators are *normaloid* (i.e., the spectral radius is the same as the norm). Although Toeplitz operators are normaloid, their inverses need not, in general, be normaloid. Of course, if φ is analytic or coanalytic, and if T_φ is invertible, then its inverse T_φ^{-1} is also a Toeplitz operator $T_{\frac{1}{\varphi}}$ [25, Theorem II] and, hence, normaloid. In this case, the arguments of Newburgh apply to show that σ is continuous when restricted to the manifolds of analytic Toeplitz operators and co-analytic Toeplitz operators.

Let \mathfrak{T} denote the subset of $L(H^2)$ consisting of all Toeplitz operators. In this section we study the continuity properties of σ as a function $\sigma : \mathfrak{T} \rightarrow \mathcal{K}$; that is, we restrict σ to the set of Toeplitz operators. Although it is open in regards to whether or not the function $\sigma : \mathfrak{T} \rightarrow \mathcal{K}$ is continuous, we are able to establish points of spectral continuity at Toeplitz operators with quasicontinuous symbols. In fact we shall demonstrate that under fairly general assumptions on $\varphi \in L^\infty$, the operator T_φ is a point of continuity for the spectral function $\sigma : \mathfrak{T} \rightarrow \mathcal{K}$.

We require the use of certain closed subspaces and subalgebras of $L^\infty(\mathbb{T})$, which are described in further detail in [9] and Appendix 4 of [20]. Recall that the subspace $H^\infty(\mathbb{T}) + C(\mathbb{T})$ is a closed subalgebra of L^∞ . The elements of the closed selfadjoint subalgebra QC , which is defined to be

$$QC = (H^\infty(\mathbb{T}) + C(\mathbb{T})) \cap \overline{(H^\infty(\mathbb{T}) + C(\mathbb{T}))},$$

are called quasicontinuous functions. The subspace PC is the closure in $L^\infty(\mathbb{T})$ of the set of all piecewise continuous functions on \mathbb{T} . Thus $\varphi \in PC$ if and only if it is right continuous and has both a left- and right-hand limit at every point. There are certain algebraic relations among Toeplitz operators whose symbols come from these classes, including

$$(2.0.1) \quad T_\psi T_\varphi - T_{\psi\varphi} \in K(H^2) \text{ for every } \varphi \in H^\infty(\mathbb{T}) + C(\mathbb{T}) \text{ and } \psi \in L^\infty(\mathbb{T}),$$

and

$$(2.0.2) \quad \text{the commutator } [T_\varphi, T_\psi] \text{ is compact for every } \varphi, \psi \in PC.$$

We add to these relations the following one.

Lemma 2.1. *If T_φ is a Toeplitz operator with quasicontinuous symbol φ , and if f is an analytic function on an open set containing $\sigma(T_\varphi)$, then $T_{f \circ \varphi} - f(T_\varphi)$ is a compact operator.*

Proof. Assume that $\varphi \in QC$. Recall from [8, p. 188] that if $\psi \in H^\infty + C(\mathbb{T})$, then T_ψ is Fredholm if and only if ψ is invertible in $H^\infty + C(\mathbb{T})$. Therefore for every $\lambda \notin \sigma(T_\varphi)$, both $\varphi - \lambda$ and $\overline{\varphi - \lambda}$ are invertible in $H^\infty + C(\mathbb{T})$; hence, $(\varphi - \lambda)^{-1} \in QC$. Using this fact together with (2.0.1) we have that, for $\psi \in L^\infty$ and $\lambda, \mu \in \mathbb{C}$,

$$T_{\varphi - \mu} T_\psi T_{\varphi - \lambda}^{-1} - T_{(\varphi - \mu)\psi(\varphi - \lambda)^{-1}} \in K(H^2) \quad \text{whenever } \lambda \notin \sigma(T_\varphi).$$

The arguments above extend to rational functions to yield: if r is any rational function with all of its poles outside of $\sigma(T_\varphi)$, then $r(T_\varphi) - T_{r \circ \varphi} \in K(H^2)$. Suppose that f is an analytic function on an open set containing $\sigma(T_\varphi)$. By Runge’s theorem there exists a sequence of rational functions r_n such that the poles of each r_n lie outside of $\sigma(T_\varphi)$ and $r_n \rightarrow f$ uniformly on $\sigma(T_\varphi)$. Thus $r_n(T_\varphi) \rightarrow f(T_\varphi)$ in the norm-topology of $L(H^2)$. Furthermore, because $r_n \circ \varphi \rightarrow f \circ \varphi$ uniformly, we have $T_{r_n \circ \varphi} \rightarrow T_{f \circ \varphi}$ in the norm-topology. Hence, $T_{f \circ \varphi} - f(T_\varphi) = \lim(T_{r_n \circ \varphi} - r_n(T_\varphi))$, which is compact. \square

Lemma 2.1 does not extend to piecewise continuous symbols $\varphi \in PC$, as we cannot guarantee that $T_\varphi^n - T_{\varphi^n} \in K(H^2)$ for each $n \in \mathbb{Z}^+$. For example, if $\varphi(e^{i\theta}) = \chi_{\mathbb{T}_+} - \chi_{\mathbb{T}_-}$, where $\chi_{\mathbb{T}_+}$ and $\chi_{\mathbb{T}_-}$ are characteristic functions of, respectively, the upper semicircle and the lower semicircle, then $T_\varphi^2 - I$ is not compact.

Corollary 2.2. *If T_φ is a Toeplitz operator with quasicontinuous symbol φ , then for every analytic function f on an open set containing $\sigma(T_\varphi)$,*

- (1) $w(f(T_\varphi)) = \sigma(T_{f \circ \varphi})$, and
- (2) $f(T_\varphi)$ is essentially normal and is unitarily equivalent to a compact perturbation of $f(T_\varphi) \oplus M_{f \circ \varphi}$, where $M_{f \circ \varphi}$ is the operator of multiplication by $f \circ \varphi$ on $L^2(\mathbb{T})$.

Proof. Because the Weyl spectrum is stable under the compact perturbations, it follows from Lemma 2.1 that $w(f(T_\varphi)) = w(T_{f \circ \varphi}) = \sigma(T_{f \circ \varphi})$, which proves statement (1). To prove (2), observe that because QC is a closed algebra, the composition of the analytic function f with $\varphi \in QC$ produces a quasicontinuous function $f \circ \varphi \in QC$. Moreover, by (2.0.1), every Toeplitz operator with quasicontinuous symbol is essentially normal. The (normal) Laurent operator $M_{f \circ \varphi}$ on $L^2(\mathbb{T})$ has its spectrum contained within the spectrum of the (essentially normal) Toeplitz operator $T_{f \circ \varphi}$. Thus there is the following relationship involving the essentially normal operators $f(T_\varphi)$ and $M_{f \circ \varphi} \oplus f(T_\varphi)$:

$$\sigma_e(f(T_\varphi) \oplus M_{f \circ \varphi}) = \sigma_e(f(T_\varphi)) \quad \text{and} \quad \mathcal{SP}(f(T_\varphi)) = \mathcal{SP}(f(T_\varphi) \oplus M_{f \circ \varphi}),$$

where $\mathcal{SP}(T)$ denotes the spectral picture of an operator T . (The spectral picture $\mathcal{SP}(T)$ is the structure consisting of the set $\sigma_e(T)$, the collection of holes and pseudoholes in $\sigma_e(T)$, and the Fredholm indices associated with these holes and pseudoholes.) Thus it follows from the Brown-Douglas-Fillmore theorem [23] that there exists a unitary operator W and a compact operator K such that

$$W(f(T_\varphi) \oplus M_{f \circ \varphi})W^* + K = f(T_\varphi). \quad \square$$

Corollary 2.2 (1) can be viewed as saying that $\sigma(f(T_\varphi)) \setminus \sigma(T_{f \circ \varphi})$ consists of holes with winding number zero.

Continuity modulo the compact operators will be a key to our study of spectral variation. The first result is an easy application of a theorem of Newburgh.

Lemma 2.3. ([19, Theorem 4]) *If $\{T_n\}_n$ is a sequence of operators converging to an operator T and such that $[T_n, T]$ is compact for each n , then $\lim \sigma_e(T_n) = \sigma_e(T)$.*

Proof. Newburgh’s theorem is stated as follows: if in a Banach algebra A , $\{a_i\}_i$ is a sequence of elements commuting with $a \in A$ and such that $a_i \rightarrow a$, then $\lim \sigma(a_i) = \sigma(a)$. If π denotes the canonical homomorphism of $L(H)$ onto the Calkin algebra $L(H)/K(H)$, then the assumptions give that $\pi(T_n) \rightarrow \pi(T)$ and $[\pi(T_n), \pi(T)] = 0$ for each n . Hence, $\lim \sigma(\pi(T_n)) = \sigma(\pi(T))$; that is, $\lim \sigma_e(T_n) = \sigma_e(T)$. \square

Theorem 2.4. *Suppose that $T, T_n \in L(H)$, for $n \in \mathbb{Z}^+$, are such that T_n converges to T . Suppose that f is any analytic function whose domain is an open set V containing $\sigma(T)$. If $[T_n, T] \in K(H)$ for each n , then*

$$(2.4.1) \quad \lim w(f(T_n)) = w(f(T)).$$

Remark. Because $T_n \rightarrow T$, by the upper-semicontinuity of the spectrum, there is a positive integer N such that $\sigma(T_n) \subseteq V$ whenever $n > N$. Thus, in the left-hand side of (2.4.1) it is to be understood that $n > N$.

Proof of Theorem 2.4. If T_n and T commute modulo the compact operators then, whenever T_n^{-1} and T^{-1} exist, T_n, T, T_n^{-1} and T^{-1} all commute modulo the compact operators. Therefore $r(T_n)$ and $r(T)$ also commute modulo $K(H)$ whenever r is a rational function with no poles in $\sigma(T)$ and n is sufficiently large. Using Runge’s theorem we can approximate f on compact subsets of V by rational functions r whose poles lie off of V . So there exists a sequence of rational functions r_i whose poles lie outside of V and $r_i \rightarrow f$ uniformly on compact subsets of V . If $n > N$, then by the functional calculus,

$$f(T_n)f(T) - f(T)f(T_n) = \lim_i (r_i(T_n)r_i(T) - r_i(T)r_i(T_n)),$$

which is compact for each n . Furthermore,

$$\begin{aligned} \|f(T_n) - f(T)\| &= \left\| \frac{1}{2\pi i} \int_\Gamma f(\lambda) ((\lambda - T_n)^{-1} - (\lambda - T)^{-1}) d\lambda \right\| \\ &\leq \frac{1}{2\pi i} \ell(\Gamma) \max_{\lambda \in \Gamma} |f(\lambda)| \cdot \max_{\lambda \in \Gamma} \|(\lambda - T_n)^{-1} - (\lambda - T)^{-1}\|, \end{aligned}$$

where Γ is the boundary of V and $\ell(\Gamma)$ denotes the arc length of Γ . The right-hand side of the above inequality converges to 0, and so $f(T_n) \rightarrow f(T)$. By Lemma 2.3, $\lim \sigma_e(f(T_n)) = \sigma_e(f(T))$. The arguments used by J.B. Conway and B.B. Morrel in Proposition 3.11 of [5] can now be used here to obtain the conclusion $\lim w(f(T_n)) = w(f(T))$. \square

We now are ready to prove the main result in this section.

Theorem 2.5. *The restriction of σ to the manifold \mathfrak{T} of all Toeplitz operators is continuous at every Toeplitz operator with quasicontinuous symbol. Moreover, if $\varphi \in QC$, $\varphi_n \in L^\infty$, and $\|T_{\varphi_n} - T_\varphi\| \rightarrow 0$, then $\lim w(f(T_{\varphi_n})) = \sigma(T_{f \circ \varphi})$.*

Proof. Suppose $\varphi \in QC$, $\varphi_n \in L^\infty$, and $\|T_{\varphi_n} - T_\varphi\| \rightarrow 0$. Then by (2.0.1), $[T_{\varphi_n}, T_\varphi] \in K(H^2)$. Therefore by Theorem 2.4, $\lim w(T_{\varphi_n}) = w(T_\varphi)$, and hence $\lim \sigma(T_{\varphi_n}) = \sigma(T_\varphi)$. Also, because $f \circ \varphi \in QC$ and $f \circ \varphi_n \rightarrow f \circ \varphi$, it follows from Lemma 2.1 that $\lim w(f(T_{\varphi_n})) = \lim \sigma(T_{f \circ \varphi_n}) = \sigma(T_{f \circ \varphi})$. \square

The argument of Theorem 2.5 is limited to quasicontinuous symbols, as we need to ensure that $[T_{\varphi_n}, T_\varphi]$ is compact for every n . If one imposes more requirements on the functions $\varphi_n \in L^\infty$, then Theorem 2.5 can be made more general. This occurs, in particular, if each φ_n is an element of PC .

Corollary 2.6. *The restriction of σ to \mathfrak{T}_{PC} is continuous, where \mathfrak{T}_{PC} is the set of all Toeplitz operators having symbols that are uniform limits of piecewise continuous functions.*

Proof. This follows from (2.0.2) and Theorem 2.4. \square

With a piecewise continuous function φ , one can obtain a continuous curve $\varphi^\#$ by joining $\varphi(e^{i\theta-0})$ and $\varphi(e^{i\theta})$ ($0 \leq \theta < 2\pi$) by the line segment $[\varphi(e^{i\theta-0}), \varphi(e^{i\theta})]$. Widom [25] showed that for every $\varphi \in PC$, $\sigma_e(T_\varphi) = \varphi^\#(\mathbb{T})$ and $\sigma(T_\varphi)$ consists of $\varphi^\#(\mathbb{T})$ together with some of its holes. This work is described in [8] and [20] as well. In a footnote on page 23 of his monograph [9], R.G. Douglas observes that the results he had been developing for Toeplitz operators with piecewise continuous symbols in fact hold, more generally, for symbols $\varphi \in L^\infty(\mathbb{T})$ having the property that

$$(2.7.1) \quad V_{\lambda_0}(\varphi) = \bigcap_{\epsilon > 0} \text{cl}[\varphi(\lambda_0 - \epsilon, \lambda_0 + \epsilon)]$$

is contained in some line segment L_{λ_0} for each $\lambda_0 \in \mathbb{T}$. In this case,

$$(2.7.2) \quad \sigma_e(T_\varphi) = \bigcup_{\lambda_0 \in \mathbb{T}} \text{conv } V_{\lambda_0}(\varphi).$$

We shall call functions φ satisfying (2.7.1) *Douglas functions*; let $D(\mathbb{T})$ denote the set of all Douglas functions in $L^\infty(\mathbb{T})$. Our aim is to extend Corollary 2.6 so that \mathfrak{T}_{PC} is replaced by a more general class in which every Toeplitz operator with symbol $\varphi \in D(\mathbb{T})$ is a point of spectral continuity (see Theorem 2.11). This general class of operators will be the Toeplitz operators whose symbols are pseudo-piecewise continuous functions.

Definition 2.8. Let $\hat{\cdot} : L^\infty \rightarrow C(\tilde{\partial}H^\infty)$ denote the Gelfand transform, where $\tilde{\partial}H^\infty$ is the Šilov boundary of $H^\infty(\mathbb{T})$ (i.e. $\tilde{\partial}H^\infty$ is the maximal ideal space of L^∞). If $\varphi \in L^\infty$, then by the Gelfand theory, $\hat{\varphi}(\tilde{\partial}H^\infty)$ is the spectrum of φ , as an element of L^∞ ; namely, $\hat{\varphi}(\tilde{\partial}H^\infty)$ is the essential range $\text{ess-ran } \varphi$ of φ . Now given $\varphi \in L^\infty(\mathbb{T})$, let $V_{\lambda_0}(\varphi)$ be as in (2.7.1). If φ has the property that $\partial \text{conv } V_{\lambda_0}(\varphi) \subseteq \hat{\varphi}(\tilde{\partial}H^\infty)$, or that $\partial \text{conv } V_{\lambda_0}(\varphi)$ is contained in some line segment L_{λ_0} , for each $\lambda_0 \in \mathbb{T}$, then φ will be called *pseudo-piecewise continuous*. Write PPC for the set of all pseudo-piecewise continuous functions in L^∞ .

For every $\lambda_0 \in \mathbb{T}$ and $\varphi \in D(\mathbb{T})$, $\text{conv } V_{\lambda_0}(\varphi) = \partial \text{conv } V_{\lambda_0}(\varphi)$, and so $D(\mathbb{T}) \subseteq PPC$. If $\varphi \in PPC$, then (2.7.2) (together with the fact that T_φ is not a Fredholm operator whenever φ cannot be inverted in $L^\infty(\mathbb{T})$) gives

$$(2.8.1) \quad \bigcup_{\lambda_0 \in \mathbb{T}} \partial \text{conv } V_{\lambda_0}(\varphi) \subseteq \sigma_e(T_\varphi).$$




FIGURE 1

The following example shows that the inclusion $D(\mathbb{T}) \subseteq PPC$ is proper.

Example 2.9. *There exists $\varphi \in L^\infty(\mathbb{T})$ such that $\varphi \in PPC \setminus D(\mathbb{T})$.*

Proof. Set

$$\varphi(e^{i\theta}) = \begin{cases} e^{i\pi(1+\frac{1}{2}\sin\frac{\theta}{3})} & (0 < \theta < \frac{2}{3}\pi), \\ (\frac{1}{\pi} + i) + \frac{1}{\pi}e^{i\frac{3\pi^2}{6\pi^2-s}(2\pi-\frac{2}{\pi}-\theta)} & (\frac{2}{3}\pi \leq \theta \leq 2\pi - \frac{2}{\pi}), \\ 2\pi - \theta + i \sin \frac{1}{2\pi-\theta} & (2\pi - \frac{2}{\pi} < \theta < 2\pi). \end{cases}$$

At $\lambda_0 = 0$, the graphs of $\varphi(\mathbb{T})$ and $V_0(\varphi)$ are in Figure 1. Therefore $\text{conv } V_0(\varphi)$ is contained in no line segment and hence $\varphi \notin D(\mathbb{T})$. But evidently, $\partial \text{conv } V_{\lambda_0}(\varphi) = V_{\lambda_0}(\varphi)$ for each $\lambda_0 \in \mathbb{T}$. In fact,

$$\bigcup_{\lambda_0 \in \mathbb{T}} V_{\lambda_0}(\varphi) = \{\hat{\varphi}(\gamma) : \gamma \in \tilde{\partial}H^\infty\}.$$

Therefore $\varphi \in PPC$. □

Theorem 2.10. *If $\varphi_n \in PPC, \varphi \in D(\mathbb{T})$ and $\|T_{\varphi_n} - T_\varphi\| \rightarrow 0$, then*

$$\partial\sigma(T_\varphi) \subseteq \partial(\liminf \sigma(T_{\varphi_n})).$$

Proof. Since $\liminf \sigma(T_{\varphi_n}) \subseteq \sigma(T_\varphi)$ and hence $\text{int}(\liminf \sigma(T_{\varphi_n})) \subseteq \text{int} \sigma(T_\varphi)$, it suffices to show that $\partial\sigma(T_\varphi) \subseteq \liminf \sigma(T_{\varphi_n})$. Assume $\lambda \notin \liminf \sigma(T_{\varphi_n})$. Then

there exists a neighborhood $\mathcal{N}_1(\lambda)$ of λ that does not intersect in finitely many $\sigma(T_{\varphi_n})$. Thus we choose a subsequence $\{\varphi_{n_i}\}$ of $\{\varphi_n\}$ such that $T_{\varphi_{n_i}} - \mu$ is invertible for each $\mu \in \mathcal{N}_1(\lambda)$, which says that $\varphi_{n_i}(\mathbb{T}) \cap \mathcal{N}_1(\lambda) = \emptyset$ for each n_i . Since $\|\varphi_n - \varphi\|_\infty = \|T_{\varphi_n} - T_\varphi\| \rightarrow 0$, there exists a neighborhood $\mathcal{N}_2(\lambda)$ of λ such that $\varphi(\mathbb{T}) \cap \mathcal{N}_2(\lambda) = \emptyset$ and $\mathcal{N}_2(\lambda) \subseteq \mathcal{N}_1(\lambda)$. We now claim that

$$\lambda \notin \bigcup_{\lambda_0 \in \mathbb{T}} \text{conv } V_{\lambda_0}(\varphi).$$

On the contrary, we assume that $\lambda \in \text{conv } V_{\lambda_0}(\varphi)$ for some $\lambda_0 \in \mathbb{T}$. Since $\varphi(\mathbb{T}) \cap \mathcal{N}_2(\lambda) = \emptyset$, and $\varphi \in D(\mathbb{T})$, λ must lie in some line segment $L_{\lambda_0}(\varphi)$ such that $L_{\lambda_0}(\varphi) \cap \varphi(\mathbb{T}) \neq \emptyset$. Since $\|\varphi_{n_i} - \varphi\| \rightarrow 0$, we have $V_{\lambda_0}(\varphi_{n_i}) \rightarrow V_{\lambda_0}(\varphi)$ and hence $\partial \text{conv } V_{\lambda_0}(\varphi_{n_i}) \rightarrow \partial \text{conv } V_{\lambda_0}(\varphi)$. But since $\partial \text{conv } V_{\lambda_0}(\varphi)$ is contained in a line segment and, by (2.8.1), $\partial \text{conv } V_{\lambda_0}(\varphi_{n_i}) \subseteq \sigma_e(T_{\varphi_{n_i}})$, it follows that for each neighborhood $\mathcal{N}(\lambda)$, there exists a $\mu \in \mathcal{N}(\lambda)$ such that $T_{\varphi_{n_i}} - \mu$ is not Fredholm, which gives a contradiction. Therefore $\lambda \notin \bigcup_{\lambda_0 \in \mathbb{T}} \text{conv } V_{\lambda_0}(\varphi)$. Thus by (2.7.2), $T_\varphi - \lambda$ is Fredholm. Now because for every $T \in L(H)$, $\partial\sigma(T) \setminus \sigma_e(T)$ consists of isolated points of $\sigma(T)$, we can conclude $\lambda \notin \partial\sigma(T_\varphi)$ because $\sigma(T_\varphi)$ is connected. This completes the proof. \square

We now have our extension of Corollary 2.6 with the following result.

Theorem 2.11. *The restriction of σ to the set of all Toeplitz operators with pseudo-piecewise continuous symbols is continuous at each Toeplitz operator with Douglas symbol.*

Proof. Suppose $\varphi_n \in \text{PPC}$, $\varphi \in D(\mathbb{T})$ and $\|T_{\varphi_n} - T_\varphi\| \rightarrow 0$. By Theorem 2.10

$$\sigma(T_\varphi)^\wedge = (\liminf \sigma(T_{\varphi_n}))^\wedge,$$

where K^\wedge denotes the polynomial-convex hull of K . Consequently, the passage from $\liminf \sigma(T_{\varphi_n})$ to $\sigma(T_\varphi)$ is the filling of some holes of $\liminf \sigma(T_{\varphi_n})$. Thus if $\liminf \sigma(T_{\varphi_n})$ has no holes, then evidently $\sigma(T_\varphi) = \liminf \sigma(T_{\varphi_n})$. Suppose now $\liminf \sigma(T_{\varphi_n})$ has a hole Ω . Since $\bigcup_{\lambda_0 \in \mathbb{T}} \partial \text{conv } V_{\lambda_0}(\varphi_n) \rightarrow \bigcup_{\lambda_0 \in \mathbb{T}} \partial \text{conv } V_{\lambda_0}(\varphi)$, we see that for sufficiently large n , φ_n behaves like a Douglas function locally on \mathbb{T} . Thus we can conclude that $\partial\Omega$ can be regarded as a ‘‘local closed curve’’ (see [9]) determined by $\partial \text{conv } V_\lambda(\varphi)$. Since $\partial \text{conv } V_{\lambda(\varphi)} = \text{conv } V_\lambda(\varphi)$, we have that

$$\partial\Omega = \bigcup_{\lambda_0 \in \mathbb{S}} \text{conv } V_{\lambda_0}(\varphi) \quad \text{for some subset } \mathbb{S} \text{ of } \mathbb{T}.$$

Thus the index theory for continuous symbols can be applied for this local closed curve ([9]). But $\|\varphi_n - \varphi\|_\infty \rightarrow 0$ and so, for sufficiently large n ,

$$-\text{ind}(T_\varphi - \lambda) = \text{wn}(\varphi - \lambda) = \text{wn}(\varphi_n - \lambda) = -\text{ind}(T_{\varphi_n} - \lambda) \quad \text{for each } \lambda \in \Omega.$$

Hence $\sigma(T_\varphi) \setminus \liminf \sigma(T_{\varphi_n})$ has no hole with non-zero winding number, and consequently $\sigma(T_\varphi) = \liminf \sigma(T_{\varphi_n})$. \square

We were unable to decide whether or not, in Theorem 2.11, $D(\mathbb{T})$ can be replaced by PPC . (If we could have equality in (2.8.1), then the answer would be yes.) More interesting still is the the following open problem.

Problem A. Is the restriction of σ to the set of all Toeplitz operators continuous?

3. WEYL'S THEOREM FOR ANALYTIC FUNCTIONS OF TOEPLITZ OPERATORS

We follow [4] in saying that *Weyl's theorem holds for T* if

$$w(T) = \sigma(T) \setminus \pi_{00}(T),$$

where $\pi_{00}(T)$ is the set of isolated points of $\sigma(T)$ that are eigenvalues of finite multiplicity. The set of operators for which Weyl's theorem holds includes all seminormal operators and all Toeplitz operators [1], [4], [21]. The following old question of K. Oberai [22] led to the work in this section: *if T_φ is a Toeplitz operator, then does Weyl's theorem hold for T_φ^2 ?*

To answer the Oberai's question, we begin with a spectral property of Toeplitz operators with continuous symbols.

Lemma 3.1. *Suppose that φ is continuous and that f is an analytic function defined on some open set containing $\sigma(T_\varphi)$. Then*

$$(3.1.1) \quad \sigma(T_{f \circ \varphi}) \subseteq f(\sigma(T_\varphi)),$$

and equality occurs if and only if *Weyl's theorem holds for $f(T_\varphi)$.*

Proof. By Corollary 2.2, $\sigma(T_{f \circ \varphi}) = w(f(T_\varphi)) \subseteq \sigma(f(T_\varphi)) = f(\sigma(T_\varphi))$. Because $\sigma(T_\varphi)$ is connected, so is $f(\sigma(T_\varphi)) = \sigma(f(T_\varphi))$; therefore the set $\pi_{00}(f(T_\varphi))$ is empty. Again by Corollary 2.2, $w(f(T_\varphi)) = \sigma(T_{f \circ \varphi})$ and so

$$w(f(T_\varphi)) = \sigma(f(T_\varphi)) \setminus \pi_{00}(f(T_\varphi)) \quad \text{if and only if } \sigma(T_{f \circ \varphi}) = f(\sigma(T_\varphi)).$$

□

Remark 3.2. If φ is not continuous, it is possible for Weyl's theorem to hold for some $f(T_\varphi)$ without $\sigma(T_{f \circ \varphi})$ being equal to $f(\sigma(T_\varphi))$. One example is as follows. Let $\varphi(e^{i\theta}) = e^{\frac{i\theta}{3}}$ ($0 \leq \theta < 2\pi$), a piecewise continuous function. The operator T_φ is invertible but T_{φ^2} is not; hence $0 \in \sigma(T_{\varphi^2}) \setminus \{\sigma(T_\varphi)\}^2$. However $w(T_\varphi^2) = \sigma(T_\varphi^2)$, and $\pi_{00}(T_\varphi^2)$ is empty (see Figure 2); therefore Weyl's theorem holds for T_φ^2 .

We can now answer Oberai's question: the answer is no.

Example 3.3. *There exists a continuous function $\varphi \in C(\mathbb{T})$ such that $\sigma(T_{\varphi^2}) \neq \{\sigma(T_\varphi)\}^2$.*




FIGURE 2




FIGURE 3

Proof. Let φ be defined by

$$\varphi(e^{i\theta}) = \begin{cases} -e^{2i\theta} + 1 & (0 \leq \theta \leq \pi), \\ e^{-2i\theta} - 1 & (\pi \leq \theta \leq 2\pi). \end{cases}$$

The orientation of the graph of φ is shown in Figure 3. Evidently, φ is continuous and, in Figure 3, φ has winding number $+1$ with respect to the hole of C_1 ; the hole of C_2 has winding number -1 . Thus we have $\sigma_e(T_\varphi) = \varphi(\mathbb{T})$ and $\sigma(T_\varphi) = \text{conv } \varphi(\mathbb{T})$. On the other hand, a straightforward calculation shows that $\varphi^2(\mathbb{T})$ is the cardioid $r = 2(1 + \cos \theta)$. In particular, $\varphi^2(\mathbb{T})$ traverses the cardioid once in a counterclockwise direction and then once in a clockwise direction. Thus $wn(\varphi^2 - \lambda) = 0$ for each λ in the hole of $\varphi^2(\mathbb{T})$. Hence $T_{\varphi^2 - \lambda}$ is a Weyl operator and is, therefore, invertible for each λ in the hole of $\varphi^2(\mathbb{T})$. This implies that $\sigma(T_{\varphi^2})$ is the cardioid $r = 2(1 + \cos \theta)$. But because $\{\sigma(T_\varphi)\}^2 = \{\text{conv } \varphi(\mathbb{T})\}^2 = \{(r, \theta) : r \leq 2(1 + \cos \theta)\}$, it follows that $\sigma(T_{\varphi^2}) \neq \{\sigma(T_\varphi)\}^2$. \square

Remark 3.4. It is instructive to observe that Lemma 3.1 gives a necessary condition for T_φ to be hyponormal. We recall [17] that if $T \in L(H)$ is hyponormal, then Weyl’s theorem holds for every $f(T)$. In conjunction with Lemma 3.1, this is to say that if T_φ is hyponormal, then $\sigma(T_{f \circ \varphi}) = f(\sigma(T_\varphi))$. But this necessary condition is not sufficient, for a slight extension of Theorem 1 in [17] shows that Weyl’s theorem holds for $f(T_\varphi)$, where T_φ is the cohyponormal Toeplitz operator with symbol $\varphi(e^{i\theta}) = e^{-i\theta}$; hence $\sigma(T_{f \circ \varphi}) = f(\sigma(T_\varphi))$.

We conclude our work by studying continuous symbols φ that have the property that Weyl’s theorem holds for $f(T_\varphi)$, for every analytic function f on a neighbourhood of $\sigma(T_\varphi)$.

Theorem 3.5. *If $\varphi \in C(\mathbb{T})$ is such that $\sigma(T_\varphi)$ has planar Lebesgue measure zero, then $\sigma(T_{f \circ \varphi}) = f(\sigma(T_\varphi))$ for every analytic function f defined on an open set containing $\sigma(T_\varphi)$.*

Proof. As φ is continuous, so is $f \circ \varphi$, and thus $\sigma_e(T_\varphi) = \varphi(\mathbb{T})$ and $\sigma_e(f(T_\varphi)) = \sigma_e(T_{f \circ \varphi}) = f \circ \varphi(\mathbb{T})$. The planar measure of $\sigma(T_\varphi)$ is zero; because $\sigma(T_\varphi)$ is a compact connected set consisting of $\varphi(\mathbb{T})$ and some of its holes, it follows that $\partial\sigma(T_\varphi) = \sigma_e(T_\varphi) = \varphi(\mathbb{T})$, which is just a continuous curve. Furthermore, as analytic functions map open connected sets onto open connected sets, we have that $\partial\sigma(f(T_\varphi)) = \sigma_e(f(T_\varphi)) = \sigma(f(T_\varphi))$. Thus $\sigma(f(T_\varphi)) \subseteq \sigma(T_{f \circ \varphi})$, which together with (3.1.1) gives the result. \square

Remark 3.6. We note here that Toeplitz operators whose symbol satisfies the hypothesis of Theorem 3.5 are essentially normal of the type “normal + compact.” To

see this, let D be a diagonal operator whose spectrum is $\varphi(\mathbb{T})$. Because T_φ and D are both essentially normal and $\mathcal{SP}(T_\varphi) = \mathcal{SP}(D)$, it follows from the Brown-Douglas-Fillmore theorem that T_φ and D are compalant; that is, $T_\varphi = N + K$ for some normal operator N on H^2 and some $K \in K(H^2)$. This observation is of interest because if $\sigma(T_\varphi)$ has planar Lebesgue measure zero and, further, if T_φ is hyponormal, then by Putnam's inequality T_φ is normal and $\varphi(\mathbb{T})$ must be a line segment.

Theorem 3.7. *If the winding number of $\varphi \in C(\mathbb{T})$ with respect to each hole of $\varphi(\mathbb{T})$ is nonnegative (or is nonpositive), then $\sigma(T_{f \circ \varphi}) = f(\sigma(T_\varphi))$ for every analytic function f defined on an open set containing $\sigma(T_\varphi)$.*

Proof. Suppose that the holes of $\varphi(\mathbb{T})$ have only nonnegative winding numbers. Since φ is continuous, it follows that $\sigma_e(T_\varphi) = \varphi(\mathbb{T})$ and

$$(3.7.1) \quad \sigma_e(T_{f \circ \varphi}) = \sigma_e(f(T_\varphi)) = f(\sigma_e(T_\varphi)).$$

If $\varphi(\mathbb{T})$ has no holes or has holes of winding number zero only, then $\sigma(T_\varphi) = \sigma_e(T_\varphi)$; thus

$$f(\sigma(T_\varphi)) = f(\sigma_e(T_\varphi)) = \sigma_e(f(T_\varphi)) = \sigma_e(T_{f \circ \varphi}) \subseteq \sigma(T_{f \circ \varphi}),$$

which together with (3.1.1) gives $\sigma(T_{f \circ \varphi}) = f(\sigma(T_\varphi))$. Now assume that there exists at least a hole Ω of $\varphi(\mathbb{T})$ such that $wn(\varphi - \lambda) \neq 0$ for all $\lambda \in \Omega$. Namely, $wn(\varphi - \lambda) = w > 0$ for all $\lambda \in \Omega$. In view of (3.7.1), it suffices to show that $f(\sigma(T_\varphi)) \setminus f(\sigma_e(T_\varphi)) \subseteq \sigma(T_{f \circ \varphi}) \setminus \sigma_e(T_{f \circ \varphi})$. Thus the proof is completed by showing that if $\lambda \in \Omega$, then $f(\lambda) \in \sigma(T_{f \circ \varphi})$. Suppose that $\lambda \in \Omega$; thus $T_\varphi - \lambda$ is Fredholm with $\text{ind}(T_\varphi - \lambda) = -wn(\varphi - \lambda) = -w < 0$. Write

$$f(z) - f(\lambda) = (z - \lambda)(z - \mu_1)^{\alpha_1} \cdots (z - \mu_n)^{\alpha_n} F(z),$$

where $\alpha_i \in \mathbb{Z}^+$, $\mu_i \in \sigma(T_\varphi)$ ($1 \leq i \leq n$) and $F(z)$ is analytic and has no zeros in $\sigma(T_\varphi)$. We have

$$f \circ \varphi - f(\lambda) = (\varphi - \lambda)(\varphi - \mu_1)^{\alpha_1} \cdots (\varphi - \mu_n)^{\alpha_n} F \circ \varphi.$$

From (3.7.1), $T_{f \circ \varphi - f(\lambda)}$ is Fredholm and hence $f \circ \varphi - f(\lambda)$ is invertible on \mathbb{T} . So each $\varphi - \mu_i$ ($1 \leq i \leq n$) and $F \circ \varphi$ vanish nowhere on \mathbb{T} . Therefore $T_{\varphi - \mu_i}$ and $T_{F \circ \varphi}$ are all Fredholm. By assumption, $wn(\varphi - \mu_i) \geq 0$, and because $F \circ \varphi$ has no zeros in $\sigma(T_\varphi)$, $wn(F \circ \varphi) = 0$. Thus

$$\begin{aligned} \text{ind}(T_{f \circ \varphi - f(\lambda)}) &= -wn\{(\varphi - \lambda)(\varphi - \mu_1)^{\alpha_1} \cdots (\varphi - \mu_n)^{\alpha_n} F(\varphi)\} \\ &= -wn(\varphi - \lambda) - \sum_{i=1}^n \alpha_i wn(\varphi - \mu_i) < 0, \end{aligned}$$

which shows that $T_{f \circ \varphi - f(\lambda)}$ is not a Weyl operator and hence is not invertible. We conclude that $f(\lambda) \in \sigma(T_{f \circ \varphi})$. The proof in the case of nonpositive winding numbers is similar. \square

Example 3.8. *If φ is of the form $p(\frac{a}{z} + bz)$, where $a, b \in \mathbb{R}$ and p is any polynomial, then $\sigma(T_{f \circ \varphi}) = f(\sigma(T_\varphi))$.*

Proof. If $a = b$, then T_φ is hermitian and the desired conclusion is evident. If $a \neq b$, set $\psi = \frac{a}{z} + bz$. Then

$$\psi(\mathbb{T}) = \{(u, v) \in \mathbb{C} : (\frac{u}{b+a})^2 + (\frac{v}{b-a})^2 = 1\},$$




FIGURE 4

which is a circle or an ellipse. Thus $\varphi(\mathbb{T}) = (p \circ \psi)(\mathbb{T}) = p(\psi)(\mathbb{T})$, which has no holes or has exactly one hole (because polynomials map continuous curves onto continuous curves and open sets onto open sets). The conclusion now follows from Theorem 3.7. \square

Remark 3.9. Lemma 3.1 and Theorems 3.5, 3.7 hold for a quasicontinuous symbol φ . In this case, if T_φ is Fredholm, then the index of T_φ is the negative of the winding number with respect to the origin of the curve $\hat{\varphi}(re^{i\theta})$ for $1 - \delta < r < 1$, and

$$\sigma_e(T_\varphi) = \bigcap_{0 < \delta < 1} \text{cl} \{ \hat{\varphi}(re^{i\theta}) : 1 - \delta < r < 1 \},$$

where $\hat{\varphi}$ is the harmonic extension of φ to the open unit disk \mathbb{D} (cf. [8]).

Remark 3.10. The index of a hyponormal operator is always nonpositive and therefore, in general, the holes of the essential spectrum of a hyponormal operator cannot have negative winding numbers. This fact may lead one to believe that if $\varphi(\mathbb{T})$ has no hole with negative winding number (in particular, if φ is a trigonometric polynomial), then T_φ is hyponormal. But such is not the case. For example, if

$$\varphi_1(e^{i\theta}) = e^{-2i\theta} + e^{i\theta} + e^{2i\theta} \quad \text{and} \quad \varphi_2(e^{i\theta}) = e^{-2i\theta} - e^{-i\theta} + e^{i\theta} + e^{2i\theta},$$

then $\varphi_1(\mathbb{T})$ has just one essential hole whose winding number is $+1$, and $\varphi_2(\mathbb{T})$ has no hole, as shown in Figure 4. But by Theorem 1.4, T_{φ_1} and T_{φ_2} both fail to be hyponormal.

Remark 3.11. Recall [23] that an operator $T \in L(H)$ is quasitriangular if there exists an increasing sequence $\{P_n\}$ of projections of finite rank in $L(H)$ that converges strongly to the identity and satisfies $\|P_n T P_n - T P_n\| \rightarrow 0$. By the work of Apostol, Foias, and Voiculescu, it is known that T is quasitriangular if and only if $\mathcal{SP}(T)$ contains no hole or pseudohole with negative winding number. Rewrite Theorem 3.7 as follows: if T_φ or (T_φ^*) is a quasitriangular Toeplitz operator with continuous symbol φ , then $\sigma(T_{f \circ \varphi}) = f(\sigma(T_\varphi))$. In Remark 3.10 we showed that even if T_φ^* is a quasitriangular Toeplitz operator (with trigonometric polynomial symbol φ), T_φ may fail to be hyponormal. In spite of this, it would be interesting to have a method by which one could determine the winding numbers of curves given by trigonometric polynomials with respect to the various holes these polynomials produce. We expect the solution will make extensive use of Theorem 3.7. The following open problem is of particular interest in operator theory.

Problem B. If φ is a trigonometric polynomial, find necessary and sufficient conditions, in terms of the coefficients of φ , for the Toeplitz operator T_φ^* to be quasi-triangular.

Acknowledgements. The work of the second author was undertaken during a sabbatical leave at The University of Regina. He would like to take this opportunity to thank the Department of Mathematics and Statistics, University of Regina, for the hospitality he received during his stay.

REFERENCES

1. S.K. Berberian, *An extension of Weyl's theorem to a class of not necessarily normal operators*, Michigan Math. J. **16** (1969), 273–279. MR **40**:3335
2. S.K. Berberian, *The Weyl spectrum of an operator*, Indiana Univ. Math. J. **20** (1970), 529–544. MR **43**:5344
3. A. Brown and P.R. Halmos, *Algebraic properties of Toeplitz operators*, J. Reine Angew. Math. **213** (1963/1964), 89–102. MR **28**:3350
4. L.A. Coburn, *Weyl's theorem for nonnormal operators*, Michigan Math. J. **13** (1966), 285–288. MR **34**:1846
5. J.B. Conway and B.B. Morrel, *Operators that are points of spectral continuity*, Integral Equations Operator Theory **2** (1979), 174–198. MR **80h**:47004
6. C. Cowen, *Hyponormal and subnormal Toeplitz operators*, Surveys of Some Recent Results in Operator Theory, I (J.B. Conway and B.B. Morrel, eds.), Pitman Research Notes in Mathematics, Volume 171, Longman, 1988, pp. (155–167). MR **90j**:47022
7. C. Cowen, *Hyponormality of Toeplitz operators*, Proc. Amer. Math. Soc. **103** (1988), 809–812. MR **89f**:47038
8. R.G. Douglas, *Banach algebra techniques in operator theory*, Academic Press, New York (1972). MR **50**:14335
9. R.G. Douglas, *Banach algebra techniques in the theory of Toeplitz operators*, CBMS 15, Providence: AMS, 1973. MR **50**:14336
10. P. Fan, *Remarks on hyponormal trigonometric Toeplitz operators*, Rocky Mountain J. Math. **13** (1983), 489–493. MR **84j**:47044
11. D.R. Farenick and R. McEachin, *Toeplitz operators hyponormal with the unilateral shift*, Integral Equations Operator Theory **22** (1995), 273–280. CMP 95:14
12. B. Gramsch and D. Lay, *Spectral mapping theorems for essential spectra*, Math. Ann. **192** (1972), 17–32. MR **45**:936
13. P.R. Halmos, *A Hilbert space problem book*, Springer, New York, 1982. MR **84e**:47001
14. R.E. Harte, *Fredholm, Weyl and Browder theory*, Proc. Roy. Irish Acad. **85A** (1985), 151–176. MR **87h**:47029
15. R.E. Harte, *Invertibility and singularity for bounded linear operators*, Dekker, New York, 1988. MR **89d**:47001
16. T. Ito and T.K. Wong, *Subnormality and quasinormality of Toeplitz operators*, Proc. Amer. Math. Soc. **34** (1972), 157–164. MR **46**:2472
17. W.Y. Lee and S.H. Lee, *A spectral mapping theorem for the Weyl spectrum*, Glasgow Math. J. **38** (1996), 61–64. CMP 96:8
18. T. Nakazi and K. Takahashi, *Hyponormal Toeplitz operators and extremal problems of Hardy spaces*, Trans. Amer. Math. Soc. **338** (1993), 753–769. MR **93j**:47040
19. J.D. Newburgh, *The variation of spectra*, Duke Math. J. **18** (1951), 165–176. MR **14**:481b
20. N.K. Nikolskii, *Treatise on the shift operator*, Springer, New York, 1986. MR **87i**:47042
21. K.K. Oberai, *On the Weyl spectrum*, Illinois J. Math. **18** (1974), 298–212. MR **48**:12086
22. K.K. Oberai, *On the Weyl spectrum (II)*, Illinois J. Math. **21** (1977), 84–90. MR **55**:1102
23. C.M. Pearcy, *Some recent developments in operator theory*, CBMS 36, Providence:AMS, 1978. MR **58**:7120
24. J.G. Stampfli, *Hyponormal operators and spectral density*, Trans. Amer. Math. Soc. **117** (1965), 469–476; erratum, *ibid.* **115** (1965), 550. MR **30**:3375, **33**:4686
25. H. Widom, *Inversion of Toeplitz matrices (II)*, Illinois J. Math. **4** (1960), 88–99. MR **24**:A432

26. H. Widom, *On the spectrum of a Toeplitz operator*, Pacific J. Math. **14** (1964), 365–375. MR **29**:476
27. K. Zhu, *Hyponormal Toeplitz operators with polynomial symbols*, Integral Equations Operator Theory **21** (1995), 376–381. MR **95m**:47044

DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF REGINA, REGINA, SASKATCHEWAN S4S 0A2, CANADA

E-mail address: `farenick@abel.math.uregina.ca`

DEPARTMENT OF MATHEMATICS, SUNG KYUN KWAN UNIVERSITY, SUWON 440-746, KOREA

E-mail address: `wylee@yurim.skku.ac.kr`