

A Coherent Approach to Spacetime Foam

R. Garattini

Facoltà di Ingegneria, Università di Bergamo, Viale Marconi 5, 24044 Dalmine (Bergamo), Italy.

Reprint requests to Dr. R. G.; E-mail: Garattini@mi.infn.it

Z. Naturforsch. **56 a**, 200–201 (2001); received February 11, 2001

Presented at the 3rd Workshop on Mysteries, Puzzles and Paradoxes in Quantum Mechanics, Gargnano, Italy, September 17 - 23, 2000.

A coherent superposition of N Schwarzschild wormholes is proposed as a model for spacetime foam. Following the subtraction procedure for manifolds with boundaries, we calculate by variational methods the Casimir energy. A proposal for an alternative foamy model formed by N Schwarzschild-Anti-de Sitter wormholes is here considered. Finally, a conjecture about the foam evolution is proposed.

Key words: Spacetime Foam; Black Holes; Quantum Gravity.

The term *Spacetime Foam* was used for the first time by J. A. Wheeler to indicate that spacetime may be subjected to quantum fluctuations in topology and metric at the Planck scale [1]. In a series of papers we have proposed a model for such a *foamy space* made by N coherent Schwarzschild wormholes [2 - 4]. The relevance of this proposal is based on a Casimir energy computation showing that quantum fluctuations of the gravitational metric shift the minimum of the effective energy from flat space (the classical minimum of the energy) to a multi-wormhole configuration. Indeed the final energy contribution to one loop is

$$\Delta_{N_w} E(M) \sim -N_w \frac{V}{64 \pi^2} \frac{\Lambda^4}{e}, \quad (1)$$

where V is the volume of the system, Λ the U. V. cut-off and N_w is the wormholes number [2]. This expression shows that a non-trivial vacuum of the multi-wormhole type is favoured with respect to flat space. It is important to remark that the N -coherent superposition of wormholes is privileged with respect to flat space, and not the single wormhole, because the single wormhole energy contribution has an imaginary contribution in its spectrum: a clear sign of an instability. Nevertheless the presence of an unstable mode is necessary to have a transition from one vacuum (the false one) to the other one (the true vacuum) [5]. Three consequences of this multiply connected spacetime are:

1. The event horizon area of a black hole is quantized and by means of the Bekenstein-Hawking relation [6, 7], also the entropy of a black hole is quantized. In particular for a Schwarzschild black hole

$$M = \frac{\sqrt{N_w}}{2 l_p} \sqrt{\frac{\ln 2}{\pi}}, \quad (2)$$

the black hole mass is quantized. Here l_p is the Planck length in natural units.

2. A cosmological constant is induced by vacuum fluctuations as shown by (1), whose value is

$$\Lambda_c = \frac{\Lambda^4 l_p^2}{N_w 8 e \pi}. \quad (3)$$

When the area-entropy relation is applied to the de Sitter geometry, we obtain

$$\frac{3 \pi}{\ln 2 l_p^2 N_w} = \Lambda_c. \quad (4)$$

3. Combining (3) and (4), one gets

$$\Lambda_c = \frac{\Lambda^4 l_p^2}{N_w 8 e \pi} = \frac{3 \pi}{\ln 2 l_p^2 N_w}. \quad (5)$$

0932-0784 / 01 / 0100-0200 \$ 06.00 © Verlag der Zeitschrift für Naturforschung, Tübingen · www.znaturforsch.com



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

This means that we have found a constraint on the U. V. cut-off

$$\Lambda^4 = \frac{24 e \pi^2}{\ln 2 l_p^4}. \quad (6)$$

It is interesting to note that (1) can be obtained at least for $N_w = 1$ even for Schwarzschild-Anti-de Sitter wormholes (S-AdS) [8], whose line element is

$$ds^2 = -f(r) dt^2 + f(r)^{-1} dr^2 + r^2 d\Omega^2, \quad (7)$$

where

$$f(r) = 1 - \frac{2MG}{r} + \frac{r^2}{b^2}. \quad (8)$$

Λ_{AdS} is the negative cosmological constant and $b = \sqrt{-3/\Lambda_{\text{AdS}}}$. To consider a large N_w approach to spacetime foam even with S-AdS wormholes, one has to consider the rescaling

$$R_{\pm} \rightarrow R_{\pm}/N_w, \quad l_p^2 \rightarrow N_w l_p^2, \quad \Lambda_{\text{AdS}} \rightarrow \Lambda_{\text{AdS}}/N_w^2, \quad (9)$$

where R_{\pm} are the boundaries related to the single wormhole. This rescaling is a consequence of the boundary reduction related to the coherent superposition of wormholes wave functionals leading to the stabilization of the system. This means that a selection rule has to emerge to compare the quantity

$$\Gamma_{\text{N-S-AdS holes}} = \frac{P_{\text{N-S-AdS holes}}}{P_{\text{AdS}}} \simeq \frac{P_{\text{foam}}}{P_{\text{AdS}}} \quad (10)$$

with

$$\Gamma_{\text{N-S holes}} = \frac{P_{\text{N-S holes}}}{P_{\text{flat}}} \simeq \frac{P_{\text{foam}}}{P_{\text{flat}}}. \quad (11)$$

In both cases we find a non-vanishing probability that a non-trivial vacuum has to be considered. Moreover

the rescaling in (9) leaves the potential (8) invariant, and when N_w is very large $\Lambda_{\text{AdS}} \rightarrow 0$. This seems to suggest that not only (10) and (11) have to be compared, but that a hierarchical mechanism of the type

$$\begin{array}{c} \frac{P_{\text{N-S-AdS holes}}}{P_{\text{AdS}}} \simeq \frac{P_{\text{foam}}}{P_{\text{AdS}}} \\ \downarrow \Lambda_{\text{AdS}} \rightarrow 0 \\ \text{Flat Space} \\ \downarrow \\ \frac{P_{\text{N-S holes}}}{P_{\text{flat}}} \simeq \frac{P_{\text{foam}}}{P_{\text{flat}}} \end{array} \quad (12)$$

is likely when $N_w \rightarrow \infty$. Although this picture has to be examined in detail, and (12) is only at the conjecture level, an important question comes into play: why has a spacetime formed by wormholes to be preferred with respect to flat spacetime, when the last one is the one we observe. The answer that at this stage can only be conjectured, is that, if we consider the expectation value on the foam state to be

$$\frac{\langle \Psi_F | \hat{g}_{ij} | \Psi_F \rangle}{\langle \Psi_F | \Psi_F \rangle} \quad (13)$$

when the number of wormholes is large enough, i. e. the scale is sufficiently large, we should have

$$\frac{\langle \Psi_F | \hat{g}_{ij} | \Psi_F \rangle}{\langle \Psi_F | \Psi_F \rangle} \rightarrow \eta_{ij}, \quad (14)$$

where η_{ij} is the flat space metric. This is a test that this foamy model has to pass if phenomenological aspects are to be considered.

Acknowledgements

I would like to thank Prof. R. Bonifacio who has given me the opportunity to participate in the Conference.

- [1] J. A. Wheeler, Ann. Phys. **2**, 604 (1957); J. A. Wheeler, Geometrodynamics, Academic Press, New York 1962.
- [2] R. Garattini, A Spacetime Foam approach to the cosmological constant and entropy, to appear in Int. J. Mod. Phys. **D**; gr-qc/0003090.
- [3] R. Garattini, Phys. Lett. **B 446**, 135 (1999); hep-th/9811187.

- [4] R. Garattini, Phys. Lett. **B 459**, 461 (1999); hep-th/9906074.
- [5] S. Coleman, Nucl. Phys. **B 298**, 178 (1988).
- [6] J. Bekenstein, Phys. Rev. **D 7**, 2333 (1973).
- [7] S. Hawking, Phys. Lett. **B 134**, 403 (1984).
- [8] R. Garattini, Class. Quant. Grav. **17**, 3335 (2000); gr-qc/0006076.