

## Production of Be<sup>7</sup> from O<sup>16</sup> with Bremsstrahlung up to 57 MeV

H. ARTUS

Institut für Technische Kernphysik  
der Technischen Hochschule, Darmstadt  
(Z. Naturforschg. **20 a**, 320 [1965]; eingegangen am 29. Januar 1965)

In the course of earlier experiments<sup>1</sup> on the photo-production of Be<sup>7</sup> from beryllium, boron and carbon, the observation of the reaction O<sup>16</sup>( $\gamma$ , x)Be<sup>7</sup> was also reported. In the meantime this reaction has been studied in more detail. The activity of Be<sup>7</sup> was measured relative to that of C<sup>11</sup> from the C<sup>12</sup>( $\gamma$ , n)C<sup>11</sup> process with a 3''  $\times$  3''  $\phi$  NaJ(Tl)-scintillation spectrometer. The samples consisted of bidistilled water and reactor graphite. They were irradiated in various geometries at the Darmstadt 60 MeV electron linear accelerator.

The measured ratios of the yields are shown in Fig. 1 as a function of the maximum bremsstrahlung energy  $E_m$ . The yield is defined by

$$Y(E_m) = \int \sigma(E) S(E) dE$$

where  $\sigma(E)$  is the cross section and  $S(E)$  stands for the number of  $\gamma$ -quanta per energy interval. Those points in Fig. 1 with large uncertainties in the energy determination were obtained from irradiations with bremsstrahlung from the unanalyzed linac beam, the others from beams well defined in energy. The production cross section  $\sigma_y$ , given by

$$\sigma_y = E_m Y(E_m) / \int S(E) E dE,$$

is 0.012 mbarn at  $E_m = 57$  MeV. Drawing a smooth curve through the measured points and calculating the cross section for the O<sup>16</sup>( $\gamma$ , x)Be<sup>7</sup> reaction as a function of photon energy yields a maximum at about

46 MeV and an integrated cross section of about 0.7 MeV mbarn to 57 MeV. The data given are based on the cross section for the C<sup>12</sup>( $\gamma$ , n)C<sup>11</sup> process as published by BARBER et al.<sup>2</sup>




Fig. 1. Measured ratios of the yields  $Y(\text{Be}^7)/Y(\text{C}^{11})$  as a function of electron kinetic energy  $E_m$ .

Recently, the O<sup>16</sup>( $\gamma$ , x)Be<sup>7</sup> reaction has also been observed by NEFKENS et al.<sup>3</sup> for bremsstrahlung energies between 45 MeV and 275 MeV. These authors state a production cross section of 0.03 mbarn.

The O<sup>16</sup>( $\gamma$ , x)Be<sup>7</sup> process does not contribute significantly to the dipole sum for photon absorption. However, because the lowest threshold energy equals 31.8 MeV, it gives information on the nuclear photo-effect above the giant dipole resonance in the theoretically interesting nucleus O<sup>16</sup>.

I wish to thank Professor P. BRIX for his advice and help. This work was sponsored by the Bundesministerium für Wissenschaftliche Forschung.

<sup>1</sup> H. ARTUS, G. FRICKE, and D. E. VON STEIN, Z. Phys. **178**, 109 [1964].

<sup>2</sup> W. C. BARBER, W. D. GEORGE, and D. D. REAGAN, Phys. Rev. **98**, 73 [1955].

<sup>3</sup> B. M. K. NEFKENS, G. MOSCATI, and J. TODOROFF, Abstract 4d/C421 of Contributed Papers to the „Congrès International de Physique Nucléaire“, Paris, July 1964.

## BERICHTIGUNG

Zu H. W. DRAWIN, Besetzungsdichten angeregter He I-Atome in einem nicht-thermischen Plasma, Band **19 a**, 1451 [1964].

Die Gl. (7) auf S. 1455 für  $Q_i$  lautet richtig

$$Q_i = \frac{\tilde{\omega}_i}{2 \sum_+} \cdot \frac{h^3}{(2 \pi m_e k T)^{3/2}} e^{+u_i} S_i,$$

$\tilde{\omega}_i$  = statistisches Gewicht.

Die numerischen Rechnungen wurden mit der richtigen Formel durchgeführt.

Nachdruck — auch auszugsweise — nur mit schriftlicher Genehmigung des Verlages gestattet  
Verantwortlich für den Inhalt: A. KLEMM  
Satz und Druck: Konrad Triltsch, Würzburg



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.