

## “Frozen” Properties and Cut-off Criteria of High Temperature Gases

M. CAPITELLI and E. FICOCELLI V.

Centro di studio per la Chimica dei plasmi del C. N. R.  
Istituto di Chimica Generale ed Inorganica dell'Università  
Bari (Italy)

(Z. Naturforsch. **25** a, 977–979 [1970]; received 14 March 1970)

The importance of the contribution coming from electronic excitation for the evaluation of the “frozen” properties of a plasma and the strong dependence of the electronic contribution on the adopted cut-off criterion are shown.

In the evaluation of electronic partition functions  $Q_{ej}$  of a species  $j$

$$Q_{ej} = \sum_n^{n_{\max}} g_{nj} \exp\{-E_{nj}/kT\}$$

or

$$Q_{ej} = \sum_0^{E_{nj\max}} g_{nj} \exp\{-E_{nj}/kT\}$$

one has to resort to some cut-off criterion<sup>1</sup> of the  $Q_{ej}$  in order to avoid the divergence of the summation. The following criteria have most extensively been used in recent times:

### The Criterion of MARGENAU and LEWIS<sup>2,3</sup>

The partition function is cut at a value of the quantum number  $n$  given by

$$n_{\max} = 36.11 \cdot 10^3 \cdot \left( \frac{Z_j^2 \text{eff.} \cdot T}{\sum n_j z_j^2} \right)^{1/4} \quad (2)$$

where  $Z_j \text{eff.}$  is the effective charge of the  $j$ -th species and  $n_j$  is the number density ( $\text{cm}^{-3}$ ) of the species with charge  $z_j$ .

### The criterion of GRIEM<sup>4,5</sup>

The partition function is cut at a maximum value of  $E_n$  given by

$$E_{nj\max} = E_{0j} - \Delta E_{0j} \quad (3)$$

with

$$\Delta E_0 = 2(z_j + 1) e^3 (\pi/kT)^{1/2} (\sum n_j z_j^2)^{1/2}$$

where  $E_{0j}$  and  $\Delta E_{0j}$  are the ionization potential and lowering of ionization potential respectively.

## The ground state criterion

$$Q_{ej} = g_{0j}$$

Once the cut-off criterion has been selected the procedure for the calculation of equilibrium compositions and thermodynamic properties is straightforward as already discussed in previous works<sup>5,6</sup>. The purpose of the present work is to emphasize the importance of the contribution coming from electronic excitation in evaluating the “frozen” properties of a plasma and the strong dependence of the electronic contribution on the cut-off criterion adopted.

Frozen properties are important quantities indeed for the analysis of heat transfer in plasmas where their values are needed for the evaluation of the adimensional PRANDTL and LEWIS numbers.

The observation<sup>5,7-9</sup> that thermodynamic properties and total specific heats of high temperature gases are practically insensitive to the cut-off criteria adopted has often led to the neglect<sup>10,11</sup> of the contribution coming from electronic excitation in the evaluation of the equilibrium properties of plasmas.

The results obtained according to this procedure are however accurate only as far as total thermodynamic quantities are concerned (i. e. properties which include ionization effects). The contribution of electronic excitation to these quantities is indeed small when compared with the contribution coming from ionization.

However, when “frozen” properties are calculated, which do not include the contribution from ionization, the neglect of the electronic contribution can lead to very inaccurate results, as will be shown below. This point has been overlooked in recent calculations<sup>11</sup>.

Helium plasma ( $\text{He}$ ,  $\text{He}^+$ ,  $e$ ) in the temperature range 10,000–35,000 °K and in the pressure interval 0.1–10 atmospheres will be taken as an example.

Results obtained for  $\text{N}_2$  plasma ( $\text{N}_2$ ,  $\text{N}_2^+$ ,  $\text{N}$ ,  $\text{N}^+$ ,  $\text{N}^{++}$ ,  $\text{N}^{+++}$ ,  $e$ ) will also be presented. The conclusions reached from analysis of these systems should be considered of general validity for high temperature gases.

### Frozen properties and cut-off criteria

The frozen enthalpy ( $H_f$ ) of a mixture of monatomic gases composed of atoms, ions and electrons, is made up of a translational contribution ( $H_{tr}$ ) and a term coming from electronic excitation ( $H_{int}$ ). In the par-

Reprints request to Dr. M. CAPITELLI, Istituto di Chimica Generale ed Inorganica dell'Università di Bari, Via Amendola 173, I-70 125 Bari (Italy).

<sup>1</sup> H. W. DRAWIN and P. FELENBOK, Data for Plasmas in Local Thermodynamics Equilibrium, Gauthier-Villars, Paris 1965.

<sup>2</sup> H. MARGENAU and M. LEWIS, Rev. Mod. Phys. **31**, 594 [1959].

<sup>3</sup> K. S. DRELLISHAK, C. F. KNOPP, and A. B. CAMBEL, Phys. Fluids **6**, 1280 [1963].

<sup>4</sup> H. R. GRIEM, Phys. Rev. **128**, 997 [1962].

<sup>5</sup> M. CAPITELLI and E. MOLINARI, J. Plasma Phys., in press.

<sup>6</sup> M. CAPITELLI, E. FICOCELLI, and E. MOLINARI, Equi-

librium Compositions and Thermodynamic Properties of Mixed Plasmas: He– $\text{N}_2$ , Ar– $\text{N}_2$  and Xe– $\text{N}_2$  Plasmas at one Atmosphere, between 5000 °K and 35 000 °K, 1969 (Report available from the authors on request).

<sup>7</sup> P. FAUCHAIS and N. MANSON, XX. Meeting of IUPAC, Moscow 1965. — P. FAUCHAIS, Rev. Int. Hautes Temp. Refractaires **6**, 77 [1969].

<sup>8</sup> J. C. HESTER and K. G. SEWELL, J. Appl. Phys. **35**, 729 [1964].

<sup>9</sup> J. ARTMANN, Z. Physik **183**, 65 [1965].

<sup>10</sup> B. ALBHORN and M. SALVAT, Z. Naturforsch. **22** a, 260 [1967].

<sup>11</sup> C. H. BACK, Phys. Fluids **10**, 807 [1967].



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.




Fig. 1. Equilibrium composition of Helium plasma ( $p=1$  atm) as a function of temperature.




Fig. 2. Electronic partition function and its derivatives as a function of temperature for the He species ( $p=1$  atm) (for the  $\text{He}^+$  species:  $Q_{ei}=2$ ;  $\dot{Q}_{ei}=\dot{Q}_{ei}=0$ ).




Fig. 3. "Frozen" enthalpy as a function of temperature for a Helium plasma.




Fig. 4. "Frozen" specific heats as a function of temperature for a Helium plasma.

ticular case of the Helium plasma ( $\text{He } \text{He}^+ \text{e}$ ), for the temperature and pressure ranges considered in the present work, the contribution coming from electronic excitation is simply due to the neutral species, because excitation energies of the  $\text{He}^+$  are too high.

One can write for this case:

$$H_f = H_{tr} + H_{int} \quad (4)$$

$$= (5/2) k T (N_n + N_i + N_e) + k T \left( \frac{d \ln Q_{en}}{d \ln T} \right) N_n.$$

Deriving Eq. (4) with respect to temperature, at constant composition, one obtains the frozen specific heat:

$$C_{pt} = \left( \frac{\partial H_f}{\partial T} \right)_{p, N_j} = C_{p \text{ tr}} + C_{p \text{ int}} \quad (5)$$

$$= (5/2) k (N_n + N_i + N_e) + k \left[ \frac{d \ln Q_{en}}{d \ln T} + \frac{d^2 \ln Q_{en}}{d^2 \ln T} \right] N_n$$

where  $N_j$  is the number of particles of  $j$ -th species per gram of plasma. From Eqs. (4) and (5) we can see that the quantities  $H_f$  and  $C_{pt}$  can be influenced by the different cut-off criteria either through  $H_{tr}$  and  $C_{p \text{ tr}}$  or through  $H_{int}$  and  $C_{p \text{ int}}$ .

Differences in  $H_{tr}$  and  $C_{p \text{ tr}}$  can arise from variations in the  $N_j$ 's. The equilibrium composition is in fact determined by the Saha equation<sup>12</sup>

$$\frac{n_i n_e}{n_n} = \frac{2 (2 \pi m_e k T)^{3/2}}{h^2} \frac{Q_{ei}}{Q_{en}} \exp - \{E_{0n} - \Delta E_{0n} / k T\}. \quad (6)$$

Fig. 1 shows that the criterion adopted is felt by the minority species only. The meaning of these results is that the partition function of He becomes appreciably different from its ground state value,  $g_{0\text{He}}$ , only when the concentration of this species is markedly diminishing and its contribution becomes therefore small.

Differences in  $H_{tr}$  and  $C_{p \text{ tr}}$  calculated according the Margenau-Lewis criterion and the ground state method do not differ by more than 3%. The quantities  $H_{int}$  and  $C_{p \text{ int}}$  depend upon

$$\dot{Q}_{en} \equiv d \ln Q_{en} / d \ln T \quad \text{and} \quad \ddot{Q}_{en} \equiv d^2 \ln Q_{en} / d^2 \ln T.$$

Values of  $\dot{Q}_{en}$  and  $\ddot{Q}_{en}$  calculated at atmospheric pressure according to the three cut-off criteria have been plotted in Fig. 2 as a function of temperature, along with  $Q_{en}$ . One sees that  $\dot{Q}_{en}$  increases in a temperature range where the He species is sharply decreasing. The contribution of the terms  $k T N_n Q_{en}$  in Eq. (4) is therefore rather small and values of  $H_f$  do not differ by more than 10% (Fig. 3). The plots of  $\dot{Q}_{en}$  do on the contrary show the presence of a maximum in a temperature region where He is not a minority species. A marked contribution to  $C_{p \text{ int}}$  is therefore to be expected (Fig. 4). When use is made of the Griem's method

the maxima of the quantities  $\dot{Q}_{en}$  and  $\ddot{Q}_{en}$  are shifted toward higher temperatures (Fig. 2). This is immediately reflected by the frozen quantities, as can be seen from Fig. 4.

Results for the  $\text{N}_2$  plasma given in Fig. 5 are in general agreement with those obtained for the Helium plasma.

$C_{pt}$  (cal/g °K)





Fig. 5. "Frozen" specific heats as a function of temperature for a Nitrogen Plasma.

The behaviour of the frozen quantities as illustrated by the results for the He and the  $\text{N}_2$  plasmas should be of a rather general validity. It is in fact a rather general situation with equilibria of the type



that the  $z$ -charged species is excited and ionized in a temperature range where the energy levels of the  $(z+1)$  ion are too high to be considerably populated.

The authors wish to thank Prof. E. MOLINARI for his many helpful suggestions and discussions during the course of this work.

<sup>12</sup> The following expressions have been used for the calculations of the  $\Delta E_{0j}$ 's:

- a) MARGENAU-LEWIS:  $\Delta E_{0j} = E_{0j} / n_{\text{max}}^2$ ;
- b) Griem: Eq. (3); c) Ground state:  $\Delta E_{0j} = 0$ .